



HAL
open science

Mental health symptoms are not correlated with peripheral inflammatory biomarkers concentrations in COVID-19-ARDS survivors

Dara Chean, Samuel Benarroch, Frédéric Pochard, Nancy Kentish-Barnes, Élie Azoulay, Matthieu Resche-Rigon, Bruno Mégarbane, Danielle Reuter, Vincent Labbé, Alain Cariou, et al.

► **To cite this version:**

Dara Chean, Samuel Benarroch, Frédéric Pochard, Nancy Kentish-Barnes, Élie Azoulay, et al.. Mental health symptoms are not correlated with peripheral inflammatory biomarkers concentrations in COVID-19-ARDS survivors. *Intensive Care Medicine*, 2022, 10.1007/s00134-022-06936-2. hal-03888549

HAL Id: hal-03888549

<https://hal.science/hal-03888549>

Submitted on 9 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mental health symptoms are not correlated with peripheral inflammatory biomarkers concentrations in COVID-19 ARDS survivors

Dara Cheadan^{1,2}, Samuel Benarroch^{1,2}, Frederic Pochard^{1,2}, Nancy Kentish-Barnes^{1,2} and Elie Azoulay^{1,2*} on behalf of the Famirea study group

Collaborators

- **Famirea study group:**

[Matthieu Resche-Rigon](#), [Bruno Megarbane](#), [Danielle Reuter](#), [Vincent Labbé](#), [Alain Cariou](#), [Guillaume Géri](#), [Guillaume Meersch](#), [Achille Kouatchet](#), [Olivier Guisset](#), [Fabrice Bruneel](#), [Jean Reignier](#), [Virginie Souppart](#), [François Barbier](#), [Laurent Argaud](#), [Jean-Pierre Quenot](#), [Laurent Papazian](#), [Bertrand Guidet](#), [Guillaume Thiéry](#), [Kada Klouche](#), [Olivier Lesieur](#), [Alexandre Demoule](#), [Christophe Guitton](#), [Gilles Capellier](#), [Bruno Mourvillier](#), [Lucie Biard](#)


*Correspondence: elie.azoulay@aphp.fr

1 Médecine Intensive et Réanimation, APHP, Hôpital Saint Louis, Paris Cité University, 1 Avenue Claude Vellefaux, 75010 Paris, France

Dear Editor,

Clinically significant and long-lasting symptoms of anxiety, depression, and post-traumatic stress disorder (PTSD) are common in acute respiratory distress syndrome (ARDS) survivors [1]. Post-ARDS psychiatric morbidity is associated with younger age, female sex, prior psychological distress, unemployment, greater alcohol and opioid use, and the number of traumatic memories.

In patients survived after ARDS due to coronavirus disease 2019 (COVID-19), high levels of acute stress disorders have been reported [2]. Previous studies assessing mental health symptoms in ARDS survivors did not report any significant difference in the prevalence and the severity of PTSD-related symptoms between COVID-19- ARDS and other ARDS etiologies [3]. In general, studies of PTSD have consistently found increased concentrations of inflammatory cytokines and imbalances in immune cell proportions [4, 5]. Along this line, plasma concentration of C-reactive protein (CRP) has been associated with the development of PTSD symptoms in war zone-deployed Marines [4], with each tenfold increment in CRP concentration being associated with an odds ratio of PTSD symptoms of 1.51 (95% CI 1.15–1.97). However, in quarantined COVID-19 patients in whom very high rates of symptoms of anxiety and depression were self-reported [5], CRP concentrations and other inflammatory markers did not show statistical difference between patients with and without psychological symptoms.



Identifying patients at higher risk of mental health symptoms would offer the opportunity of preventive measures to avoid or alleviate the prevalence and severity of psychological burden. We sought to correlate peripheral inflammatory biomarkers to the development of symptoms of anxiety, depression and PTSD in critically ill patients with severe COVID-19 ARDS admitted to 23 intensive care units (ICUs) in France during the first COVID-19 wave (March and April 2020). These patients were reported elsewhere [3]. CRP, procalcitonin (PCT) and ferritin were sampled at ICU admission. During the day-90 telephone interviews, the patients completed the Impact of Event Scale Revised (IES-R) to assess PTSD-related symptoms and the Hospital Anxiety and Depression Scale (HADS) to assess the prevalence of symptoms of anxiety and depression. The Pearson correlation coefficient (R) was used to measure the strength and the direction of the linear relationship between the concentrations of inflammatory biomarker and the IES-R, or the anxiety and the depression subscales of the HADS.

Among the 178 ICU patients with COVID-19 ARDS, assessment of mental health symptoms was available in 156 (88%, median age 59 (IQR 50–68), 118 (76%) men and 38 (24%) women). Mechanical ventilation was required in 114 (73%) patients and 42 (27%) required high flow nasal oxygen only. Symptoms of anxiety (HADS-anxiety subscale > 7), depression (HADSdepression subscale > 7), and PTSD (IES-R > 22) were found in 25%, 22% and 22% of the patients, respectively. Symptoms of anxiety and depression were found in 55% and 42% of patients with PTSD, respectively, as compared to 9% and 8% of patients without PTSD, respectively. CRP concentration at admission was 157 mg/l (IQR 86–246), procalcitonin was 0.37 ng/ml (0.16–1.17), and ferritin was 1280 µg/l (830–2605). Pearson correlation coefficient for the association between CRP and the anxiety subscale of the HADS was 0.03, showing a poor association between the two scores (Fig. 1). Similar figures were found for the association between CRP and the depression subscale of the HADS or the IES-R (R = 0.009 and R = 0.06, respectively). Similarly, PCT concentrations at admission were poorly correlated with the anxiety or depression subscales of the HADS or with the IES-R (R = 0.02, R = 0.1, and R = -0.04, respectively). Last, ferritin concentrations at admission were also poorly correlated with mental health symptoms (R = 0.05 for the anxiety subscale of the HADS, R = 0.03 for the depression subscale of the HADS, and R = 0.11 for the IES-R).

In conclusion, peripheral inflammatory biomarkers are poorly correlated with further occurrence of symptoms of anxiety, depression or PTSD in survivors of COVID-19 severe ARDS. Until PTSD-specific biomarkers can be identified validated in this setting, early screening of mental health symptoms in ARDS survivors should be offered regardless the concentrations of initial peripheral inflammatory biomarkers.

Author details

1 Médecine Intensive et Réanimation, APHP, Hôpital Saint Louis, Paris Cité University, 1 Avenue Claude Vellefaux, 75010 Paris, France. 2 Psychiatry Department, Lariboisière Fernand-Widal University Hospital, Paris, France.

Acknowledgements

The Famirea study group: Matthieu Resche-Rigon: Clinical Research Unit, APHP, Saint Louis University Hospital, Paris, France; Bruno Megarbane: Medical Intensive Care Unit, APHP, Lariboisière University Hospital, Paris, France; Danielle Reuter: Medical-Surgical Intensive Care Unit, CH Sud Francilien, Corbeil, France; Vincent Labbé: Medical-Surgical Intensive Care Unit, APHP, Tenon University Hospital, Paris, France; Alain Cariou: Medical Intensive Care Unit, Cochin University Hospital, APHP, Centre-Université de Paris, Paris, France; Guillaume Géri: Medical-Surgical Intensive Care Unit, APHP, Ambroise Paré University Hospital, Boulogne, France; Guillaume Van der Meersch: Medical-Surgical Intensive Care Unit, APHP, Avicenne University Hospital, Bobigny, France; Achille Kouatchet: Medical Intensive Care Unit, Angers Teaching Hospital, Angers, France; Olivier Guisset: Medical Intensive Care Unit, Saint-André Hospital, Bordeaux, France; Fabrice Bruneel: Intensive Care Unit, André Mignot Hospital, Le Chesnay, France; Jean Reignier: Medical Intensive Care Unit, University Hospital Centre, Nantes, France.; Virginie Souppart: Famirea Study Group, Medical Intensive Care Unit, APHP, Saint Louis University Hospital, Paris, France; François Barbier: Medical Intensive Care Unit, La Source Hospital, CHR Orléans, Orléans, France; Laurent Argaud: Medical Intensive Care Department, Edouard Herriot Hospital, Hospices Civils de Lyon, Lyon, France; Jean Pierre Quenot: Medical Intensive Care Department, University Hospital, Dijon, France; Laurent Papazian: Respiratory and Infectious Diseases Intensive Care Unit, APHM Hôpital Nord, Marseille, France; Bertrand Guidet: Medical Intensive Care Unit, APHP, Saint-Antoine University Hospital, Paris, France; Guillaume Thiéry: Medical Intensive Care Unit, Saint-Etienne, University Hospital, Paris, France; Kada Klouche: Department of Intensive Care Medicine, Lapeyronie Hospital, Montpellier, France; Olivier Lesieur: Medical-Surgical Intensive Care Unit, La Rochelle Hospital, La Rochelle, France; Alexandre Demoule: AP-HP, Groupe Hospitalier Universitaire APHP-Sorbonne Université, Pitié-Salpêtrière site, Service de Pneumologie, Médecine Intensive et Réanimation (Département R3S) and Sorbonne Université, INSERM, UMRS1158 Neurophysiologie Respiratoire Expérimentale et Clinique, F-75005 Paris, France; Christophe Guitton: Medical Intensive Care Unit, Le Mans Hospital, Le Mans, France; Gilles Capellier: Medical Intensive Care Unit, Besançon, University Hospital, Besançon, France; Bruno Mourvillier: Medical Intensive Care Unit, Reims University Hospital, Reims, France; Lucie Biard: Clinical Research Unit, APHP, Saint Louis University Hospital, Paris, France.

Funding

All financial support for the research was provided by the French Ministry of Health (Grant COVID2020).

Declarations

Conflicts of interest

Authors declare no conflict of interest in relation with this manuscript.

References

1. Bienvenu OJ et al (2018) Psychiatric symptoms after acute respiratory distress syndrome: a 5-year longitudinal study. *Intensive Care Med* 44:38–47
2. Mongodi S et al (2021) High prevalence of acute stress disorder and persisting symptoms in ICU survivors after COVID-19. *Intensive Care Med* 47:616–618
3. Azoulay E et al (2022) Association of COVID-19 acute respiratory distress syndrome with symptoms of posttraumatic stress disorder in family members after ICU discharge. *JAMA* 327:1042–1050
4. Eraly SA et al (2014) Assessment of plasma C-reactive protein as a biomarker of posttraumatic stress disorder risk. *JAMA Psychiat* 71:423–431

5. Guo Q et al (2020) Immediate psychological distress in quarantined patients with COVID-19 and its association with peripheral inflammation: a mixed-method study. *Brain Behav Immun* 88:17–27