



HAL
open science

Approximate control of parabolic equations with on-off shape controls by Fenchel duality

Camille Pouchol, Emmanuel Trélat, Christophe Zhang

► **To cite this version:**

Camille Pouchol, Emmanuel Trélat, Christophe Zhang. Approximate control of parabolic equations with on-off shape controls by Fenchel duality. 2022. hal-03889865v1

HAL Id: hal-03889865

<https://hal.science/hal-03889865v1>

Preprint submitted on 8 Dec 2022 (v1), last revised 11 Dec 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Approximate control of parabolic equations with on-off shape controls by Fenchel duality

Camille Pouchol^a, Emmanuel Trélat^{b,d}, and Christophe Zhang^c

^aLaboratoire MAP5 UMR 8145, Université de Paris UFR Mathématiques & Informatique, Campus Saint-Germain-des-Prés, 75006 PARIS. Email address: emmanuel.trelat@sorbonne-universite.fr

^bSorbonne Université, Université de Paris, CNRS, Laboratoire Jacques-Louis Lions, 75005 Paris, France. email address: camille.pouchol@u-paris.fr

^cSPHINX, INRIA, Faculté des Sciences et Technologies Campus, Boulevard des Aiguillettes 54506 Vandœuvre-lès-Nancy, France. Email address: christophe.zhang@polytechnique.org

^dCAGE, INRIA, Paris, France.

Abstract

We consider the internal control of linear parabolic equations through *on-off shape* controls, *i.e.*, controls of the form $M(t)\chi_{\omega(t)}$ with $M(t) \geq 0$ and $\omega(t)$ with a prescribed maximal measure.

We establish small-time approximate controllability towards all possible final states allowed by the comparison principle with nonnegative controls. We manage to build controls with constant amplitude $M(t) \equiv \bar{M}$. In contrast, if the moving control set $\omega(t)$ is confined to evolve in some region of the whole domain, we prove that approximate controllability fails to hold for small times.

The method of proof is constructive. Using Fenchel-Rockafellar duality and the bathtub principle, the on-off shape control is obtained as the bang-bang solution of an optimal control problem, which we design by relaxing the constraints.

Our optimal control approach is outlined in a rather general form for linear constrained control problems, paving the way for generalisations and applications to other PDEs and constraints.

Contents

1	Introduction	2
1.1	Constrained internal control	2
1.2	Main results	3
1.3	General results	5
1.4	Proof strategy and related works	7
1.5	Extensions and perspectives	8
2	Building the optimal control problem	10
2.1	Convex analytic framework	10
2.2	Approximate controllability by Fenchel duality ([24])	10
2.3	Convex analytic interpretation of the bathtub principle	12
2.4	From the static bathtub principle to the dual problem and its corresponding cost	13

3	Approximate controllability results	14
3.1	Strong duality	15
3.2	Coercivity of $J_{T,\varepsilon}$, nonnegative approximate controllability	16
3.3	Characterisation of the minimisers	16
3.4	Uniqueness	18
4	Obstructions to controllability	19
5	Further comments	20
5.1	Properties of the value function in the general case	20
5.2	Obstructions	22
5.2.1	Obstruction to reachability and small-time controllability	22
5.2.2	Characterisation of minimal time controls	23
A	Convex analysis	25
A.1	Core properties of Fenchel conjugation	25
A.2	Some properties of indicator and support functions	26
A.3	Technical lemmas	27
A.4	Fenchel-Rockafellar duality	27
A.5	Parametric convex optimisation	29
B	The classical bathtub principle	30

1 Introduction

1.1 Constrained internal control

This article is devoted to the internal approximate controllability problem at time $T > 0$ for linear parabolic equations on a domain Ω by means of *on-off shape controls*, *i.e.*, internal controls taking the form

$$\forall t \in (0, T), \forall x \in \Omega, \quad u(t, x) = M(t)\chi_{\omega(t)}(x),$$

where, at a given time $t \in (0, T)$,

- $M(t) > 0$ is the **nonnegative** amplitude of the control
- $\chi_{\omega(t)}$ is the characteristic function of the set $\omega(t) \subset \Omega$, *i.e.*, $\chi_{\omega(t)}(x) := \begin{cases} 1 & \text{if } x \in \omega(t), \\ 0 & \text{otherwise} \end{cases}$.

Both the amplitude and location may be subject to constraints. This problem is a paradigmatic simplification of many practical situations where one can act on a complex system with on-off devices that can be moved in time, while their shape can also be modified.

Along the introduction, we expose our results for general operators A , while first illustrating them in the case of the controlled linear heat equation with Dirichlet boundary conditions

$$\begin{cases} y_t - \Delta y = u \text{ in } \Omega, \\ y = 0 \text{ on } \partial\Omega, \\ y(0) = y_0 \text{ in } \Omega. \end{cases} \quad (1)$$

In this setting, Ω is an open connected bounded subset of \mathbb{R}^d , with C^2 boundary, and $y_0 \in L^2(\Omega)$.

Control without constraints. When constraints are removed, generic parabolic equations are well-known to be approximately controllable [2, 42], and even null-controllable [13, 22] in arbitrarily small time by means of internal controls, acting only on an arbitrary fixed measurable subset $\omega \subset \Omega$ of positive measure.

This more precisely means that for any time $T > 0$, any measurable set $\omega \subset \Omega$ of positive measure, any $\varepsilon > 0$, any $y_0 \in L^2(\Omega)$ and target $y_f \in L^2(\Omega)$, there holds

$$\exists u \in L^2((0, T) \times \Omega), \text{ such that } \forall t \in (0, T), \text{ supp}(u(t, \cdot)) \subset \omega \text{ and } \|y(T) - y_f\|_{L^2(\Omega)} \leq \varepsilon,$$

where $\text{supp}(u)$ refers to the essential support of a function $u \in L^2(\Omega)$.

Constrained control. In view of applications where unilateral or bilateral or L^∞ constraints naturally appear, constrained controllability has been an active area of research [1, 8, 34], whether in finite or infinite dimension.

In various contexts, control constraints have been shown to lead to controllability obstructions, even for unilateral constraints. Some states are out of reach, regardless of how large $T > 0$ may be [37, 41]. On the other hand, some states are reachable but only for T large enough: constraints may lead to the appearance of a minimal time of controllability [25, 26, 27, 36].

In the case of unilateral constraints for linear control problems in finite dimension, these obstructions can be categorised thanks to Brunovsky's normal form as done in [26], leading to the existence of a positive minimal time. In infinite dimension, however, we are only aware of obstructions based on the comparison principle (see [25] and [37]). The present work uncovers another type of obstruction, already hinted at in [36].

1.2 Main results

As our results require different sets of hypotheses and in order to give a quick glance at the main ideas, we first present them in the simplified context of the heat equation (1).

Given a constraint set $\mathcal{U}_+ \subset L^2(\Omega)$, we will be considering control constraints of the form

$$\forall t \in (0, T), \quad u(t) \in \mathcal{U}_+.$$

Here, the notation \mathcal{U}_+ emphasises that we will always deal with constraints that include the nonnegativity constraint, *i.e.*, sets $\mathcal{U}_+ \subset \{u \in L^2(\Omega), u \geq 0\}$.

Now, when the control u satisfies $u \geq 0$, it follows from *the parabolic comparison principle* satisfied by the Dirichlet Laplacian [16] that

$$\forall t \geq 0, \quad y(t) \geq e^{t\Delta} y_0, \tag{2}$$

where $(e^{t\Delta})_{t \geq 0}$ denotes the heat semigroup with Dirichlet boundary conditions. Hence, targets y_f which do not satisfy $y_f \geq e^{T\Delta} y_0$ cannot be reached with nonnegative controls, let alone on-off shape controls.

Taking into account the obstruction to controllability given by the inequality (2), we adapt the usual definition of approximate controllability to the context of nonnegative controls.

More precisely, we say that system (1) is *nonnegatively approximately controllable with controls in \mathcal{U}_+ in time $T > 0$* , if for all $\varepsilon > 0$, and all $y_0, y_f \in L^2(\Omega)$ such that $y_f \geq e^{T\Delta} y_0$, there exists a control $u \in L^2((0, T) \times \Omega)$ with values in \mathcal{U}_+ such that the corresponding solution to (1) satisfies $\|y(T) - y_f\|_{L^2(\Omega)} \leq \varepsilon$.

On-off shape control. For our first main result, we focus on nonnegative approximate controllability with on-off shape controls: for a fixed $L \in (0, 1)$, we consider the constraint set

$$\mathcal{U}_L^{\text{shape}} := \{M\chi_\omega, \quad \omega \subset \Omega, \quad |\omega| \leq L|\Omega|, \quad M > 0\} \subset L^2(\Omega).$$

Within the above class of on-off shape controls, we establish nonnegative approximate controllability in arbitrary time (see Theorem 3.1 for the precise and general statement), whatever the value of $L \in (0, 1)$.

Theorem A. For any $L \in (0, 1)$, $T > 0$, system (1) is nonnegatively approximately controllable with controls in $\mathcal{U}_{\text{shape}}^L$ in time T .

To establish this result, we draw from Lions's strategy in [24], which develops a constructive approach in studying the approximate controllability of a linear wave equation. The idea is to consider the requirement $\|y(T) - y_f\|_{L^2(\Omega)} \leq \varepsilon$ as a constraint. With $L_T u := \int_0^T e^{(T-t)\Delta} u(t) dt$ and since $y(T) = L_T u + S_T y_0$, Lions considers the constrained optimal control problem

$$\pi := \inf \left\{ \frac{1}{2} \|u\|_{L^2((0,T) \times \Omega)}^2, \|e^{T\Delta} y_0 + L_T u - y_f\|_{L^2(\Omega)} \leq \varepsilon \right\}.$$

The infimum satisfies $\pi < +\infty$ if and only if there exists $u \in L^2((0, T) \times \Omega)$ steering y_0 to a closed ε -ball around y_f . To find minimisers, *i.e.*, to build controls, note that

$$\pi = \inf_{u \in L^2((0,T) \times \Omega)} \frac{1}{2} \|u\|_{L^2((0,T) \times \Omega)}^2 + G_{T,\varepsilon}(L_T u) = \inf_{u \in L^2((0,T) \times \Omega)} F_T(u) + G_{T,\varepsilon}(L_T u),$$

with $F_T(u) = \frac{1}{2} \|u\|_{L^2((0,T) \times \Omega)}^2$ and

$$G_{T,\varepsilon}(y) = \begin{cases} 0 & \text{if } \|e^{T\Delta} y_0 + y - y_f\|_{L^2(\Omega)} \leq \varepsilon, \\ +\infty & \text{otherwise.} \end{cases}$$

From this optimisation problem, one computes its Fenchel dual optimisation problem, which reads

$$d := - \inf_{p_f \in L^2(\Omega)} F_T^*(L_T^* p_f) + G_{T,\varepsilon}^*(-p_f) = - \inf_{p_f \in L^2(\Omega)} \frac{1}{2} \|L_T^* p_f\|_{L^2((0,T) \times \Omega)}^2 + G_{T,\varepsilon}^*(-p_f),$$

where $F_T^*(= F_T)$ and $G_{T,\varepsilon}^*$ are the Fenchel conjugates of F_T and $G_{T,\varepsilon}$, respectively, and L_T^* is the adjoint of the linear bounded operator $L_T : L^2((0, T) \times \Omega) \rightarrow L^2(\Omega)$. Recall that for a given $p_f \in L^2(\Omega)$, $p = L_T^* p_f$ is the solution to the adjoint equation ending at p_f , *i.e.*, it solves

$$\begin{cases} p_t + \Delta p = 0, \\ p = 0 \text{ on } \partial\Omega, \\ p(T) = p_f \text{ on } \Omega. \end{cases} \quad (3)$$

Under suitable hypotheses, the Fenchel-Rockafellar theorem [40] ensures that $\pi = d$. As a result, one can then study the dual functional to establish that $\pi = d < +\infty$, and that its minimum is attained. Furthermore, the cost function F_T is differentiable in this case and the first order optimality condition for the (unique) variable p_f^* minimising the dual functional then reads $L_T L_T^* p_f^* = y_f - \varepsilon \frac{p_f^*}{\|p_f^*\|_{L^2(\Omega)}}$. The optimal control $u^* := L_T^* p_f^*$ is thus constructed from the minimiser of the dual problem p_f^* .

Accordingly, in this paper we reframe constrained approximate controllability as an optimal control problem, replacing $\frac{1}{2} \|u\|_{L^2((0,T) \times \Omega)}^2$ of [24] with a suitable cost functional F_T . This constitutes a novel generalisation of Lions's method.

As the detailed statements in Theorem 3.1 and Proposition 3.7 show:

- Instead of using the L^2 norm as in the optimal control problem studied in [24], we will consider the cost functional:

$$F_T(u) := \frac{1}{2} \sup_{t \in [0, T]} \max \left(\|u(t, \cdot)\|_{L^\infty(\Omega)}, \frac{\|u(t, \cdot)\|_{L^1(\Omega)}}{L|\Omega|} \right)^2 + \delta_{\{u \geq 0\}}(u), \quad (4)$$

where $\delta_{\{u \geq 0\}}(u) = 0$ if $u \geq 0$ and $+\infty$ otherwise. The rather unusual form of the minimisation criterion (4) is finely designed so as to handle nonnegativity and the other (bound, volume) constraints we are dealing with.

- The optimal controls have constant amplitude in time, *i.e.*, $M(t) \equiv \overline{M}$.
- The proof is constructive: the optimal control u^* can be computed from a unique dual optimal variable p_f^* solving the corresponding Fenchel dual problem. This computation generalises what is done in [24] to the broader case of costs that are not differentiable but still convex. More precisely, u^* is given by

$$u^*(t, \cdot) = \overline{M} \chi_{\{p^*(t, \cdot) > h(p^*(t, \cdot))\}}, \quad \overline{M} = \int_0^T \int_{\{p^*(t, \cdot) > h(p^*(t, \cdot))\}} p^*(t, x) dx dt,$$

where $h : L^2(\Omega) \rightarrow \mathbb{R}$ is a function that will be defined in Section 2.3, and p^* solves the adjoint equation (3) with $p^*(T) = p_f^*$.

Obstructions to nonnegative controllability. In the spirit of the unconstrained case, one may wonder whether nonnegative approximate controllability can be achieved with controls acting only in some prescribed time-independent subdomain ω . We emphasise that our first result does not *a priori* prevent the control from visiting the whole domain Ω .

Our second result proves that visiting the whole Ω is necessary in the following sense: if the sets $\omega(t)$, $t \in (0, T)$ do not intersect some fixed open subset of Ω , nonnegative approximate controllability is lost for small times.

Theorem B. *Assume that the constraint set \mathcal{U}_+ satisfies the following property: there exists a ball $B(x, r) \subset \Omega$ with $x \in \Omega$ and $r > 0$ such that*

$$\forall u \in \mathcal{U}_+, \quad \text{supp}(u) \cap B(x, r) = \emptyset.$$

Then, there exists $T^ > 0$ such that the control system (1) is not nonnegatively approximately controllable with controls in \mathcal{U}_+ in time $T \leq T^*$.*

We refer to Theorem 4.1 for the complete statement. Let us mention that obstructions of this type have been reported for similar problems in [36].

Amplitude and time optimal control. In Section 5, we gather several further results regarding the dependence of the amplitude $\overline{M} = \overline{M}(T, y_0, y_f, \varepsilon)$ with respect to its arguments. Using duality once more, we study its dependence on the final time T .

Focusing on the case $y_0 = 0$, we then establish an equivalence between the optimal control problem and the related minimal time problem

$$\inf\{T > 0, \quad \exists u \in L^2((0, T) \times \Omega), \quad \|L_T u - y_f\|_{L^2(\Omega)} \leq \varepsilon, \quad F_T(u) \leq \lambda\}, \quad \lambda > 0.$$

1.3 General results

Theorems A and B above have been stated for the heat equation with Dirichlet boundary conditions, in order to provide the reader with a quick overview of our main results. In fact, they all hold for more general semigroups under suitable hypotheses presented hereafter.

The underlying general setting is that of linear control problems of the form

$$\begin{cases} y_t - Ay = u, \\ y(0) = y_0 \text{ in } \Omega \end{cases} \quad (5)$$

where Ω is an open subset of \mathbb{R}^d , and $A : D(A) \rightarrow L^2(\Omega)$ is an operator generating a C_0 semigroup $(S_t)_{t \geq 0}$ on $L^2(\Omega)$ [14, 35].

In this more general context, we define nonnegative approximate controllability as follows.

Definition 1.1. Given a constraint set of **nonnegative** controls $\mathcal{U}_+ \subset L^2(\Omega)$, we say that system (5) is *nonnegatively approximately controllable with controls in \mathcal{U}_+ in time T* if for all $\varepsilon > 0$, and all $y_0, y_f \in L^2(\Omega)$ such that $y_f \geq S_T y_0$, there exists a control $u \in L^2((0, T) \times \Omega)$ with values in \mathcal{U}_+ such that the corresponding solution to (5) satisfies $\|y(T) - y_f\|_{L^2(\Omega)} \leq \varepsilon$.

General hypotheses for Theorem A. We have previously presented Theorem A for the heat equation as a paradigmatic example. Nevertheless, the underlying hypotheses on which some of our proofs rely are much more general in nature; we review them below.

- First, we consider the (unusual) unique-continuation like property

$$\forall y \in L^2(\Omega), \quad \exists t \in (0, T), \quad S_t y \text{ is constant over } \Omega \quad \implies \quad y = 0. \quad (\text{GUC})$$

This property is satisfied as soon as the three assumptions below hold:

- for $y \in L^2(\Omega)$, $S_t y \in D(A)$ for all $t > 0$ (for instance, this is true if $(S_t)_{t \geq 0}$ is analytic [35]),
- the only constant function in $D(A)$ is the zero function,¹
- S_t is injective for all $t > 0$.²
- Second, we will be interested in *analytic-hypoellipticity*: $\partial_t - A$ is said to be analytic-hypoelliptic if any distributional solution y to $\partial_t y - Ay = f$ on $\Omega \times (0, T)$ with f analytic on Ω is analytic on Ω , where analyticity refers to real-analyticity.
- Third, we will say that $(S_t)_{t \geq 0}$ satisfies the *comparison principle* if

$$\forall y \in L^2(\Omega), \quad y \geq 0 \implies \forall t > 0, \quad S_t y \geq 0. \quad (6)$$

The first two properties are sufficient for the generalisation of Theorem A, see Theorem 3.1. The third will play an important role when it comes to minimal controllability times, and is in line with our definition of nonnegative approximate controllability.

Elliptic operators. As a generalisation of the Dirichlet Laplacian, let us discuss a large class of uniformly elliptic operators that do satisfy these properties and to which our obstruction result Theorem B generalises (see Theorem 4.1).

Let us assume that Ω is a bounded, open, connected subset of \mathbb{R}^d , with C^2 boundary. Defining $D(A) := H_0^1(\Omega) \cap H^2(\Omega)$, we introduce operators of the form

$$\forall y \in D(A), \quad Ay := \sum_{1 \leq i, j \leq d} \partial_{x_j} (a_{ij}(x) \partial_{x_i} y) - \sum_{i=1}^d b_i(x) \partial_{x_i} y + c(x)y. \quad (7)$$

When referring to operators of the form (7), we will always assume that the functions $a_{ij} = a_{ji}$, b_i are in $W^{1, \infty}(\Omega)$, c is in $L^\infty(\Omega)$, and that the operator is uniformly elliptic, *i.e.*, there exists $\theta > 0$ such that

$$\forall x \in \Omega, \quad \forall \xi \in \mathbb{R}^d, \quad \sum_{1 \leq i, j \leq d} a_{ij}(x) \xi_i \xi_j \geq \theta |\xi|^2.$$

The adjoint of A is given by

$$\forall p \in D(A^*), \quad A^* p = \sum_{1 \leq i, j \leq d} \partial_{x_i} (a_{ij}(x) \partial_{x_j} p) + \sum_{i=1}^d b_i(x) \partial_{x_i} p + \left(c(x) - \sum_{i=1}^d \partial_{x_i} (b_i(x)) \right) p,$$

¹This is the case for the Dirichlet Laplacian with domain $D(A) = H^2(\Omega) \cap H_0^1(\Omega)$ if Ω has a C^2 boundary.

²This is the case for groups, such as the wave equation, and for parabolic equations thanks to the parabolic maximum principle. This is also true for analytic semigroups: if $S_t y = 0$ for some $t > 0$, then $S_s y = 0$ for all $s \geq t$ and by analyticity $S_s y = 0$ for all $s \geq 0$, which for $s = 0$ yields $y = 0$.

and we have $D(A^*) = D(A)$.

Both A and A^* satisfy the parabolic comparison principle [16], hence they satisfy the comparison principle (6). They also satisfy the three conditions sufficient for the **(GUC)** property to hold.³ Furthermore, both $\partial_t - A$ and $\partial_t - A^*$ are analytic-hypoelliptic as soon as all functions a_{ij} , b_i and c are analytic [33].

1.4 Proof strategy and related works

In the unconstrained case, approximate controllability of the heat equation is a consequence of the unique continuation property, thanks to a general property of linear control problems (see for example [12, Section 2.3]). In the case of heat equations, the latter property can be obtained by the Holmgren Uniqueness Theorem [2]. In contrast to these existence results, the variational approach developed in [24] (see Section 1.2), handles approximate controllability in a *constructive manner*.

Our strategy consists in extending this approach to the constrained case: the main idea is to find a suitable cost function F_T such that optimal controls must satisfy the constraint $u \in \mathcal{U}_L^{\text{shape}}$. A remarkable feature of our strategy lies in how we design the cost function: we do so by building an adequate Fenchel dual function, instead of trying to find the cost function directly.

Constrained controllability. Constrained control problems in infinite dimension have been studied in papers such as [3, 4, 5, 15, 19]. In [15], sufficient conditions (in the form of unique continuation properties) for controllability results are derived when the control and states are constrained to some prescribed subspaces, but at the expense of controlling only a finite-dimensional subpart of the final state. In [19], the authors deal with a form of approximate controllability of the heat equation akin to ours, focusing on minimal time problems. They derive bang-bang type necessary optimality conditions for minimal time controls, and then build such controls using an auxiliary optimisation problem.

The papers [3, 4, 5] address constrained exact controllability through modified observability inequalities, thus giving abstract necessary and/or sufficient conditions. One key difference with our work is that constraint sets are assumed to be convex. In fact, all examples handled by [3, 4, 5] feature isotropic constraints.

It is noteworthy that all the above references introduce so-called dual functionals, drawing from the variational formulation of the Hilbert Uniqueness Method. However, the formalism of Fenchel-Rockafellar duality in itself, as developed in [24], has increasingly been abandoned in the literature. Some notable exceptions are [44] in the context of stabilisation, and, to some extent, [4], which uses Fenchel duality to study null-controllability under some hypotheses.

We fully exploit the ideas hinted at in the latter paper by choosing a different type of functional, which allows us to handle anisotropic, non-convex constraints. In contrast with the aforementioned trend in the literature, we work with Fenchel duality, but in a rather unusual way, in that we will focus mainly on the dual problem. The nature of the actual primal problem (optimal control problem) being solved follows effortlessly. To perform the necessary computations, we will make extensive use of convex analysis. Doing so bypasses many technical difficulties thanks to properties of subdifferentials and Fenchel conjugates, among others, and allows for the use of costs which are not differentiable but still smooth in the convex analytic sense.

Bathtub principle for appropriate costs. The second main idea is what underlies our choice of cost function F_T , forcing optimal controls to satisfy the required the on-off shape constraint. As the set of on-off shape controls is a non-convex cone, we are led to relaxation, *i.e.*, to consider the closure of its convex hull. In order to build relevant costs, we then rely on the so-called bathtub principle (actually, a relaxed version of it) [23].

³The analyticity of the semigroup is well known for this class of elliptic operators on open domains with C^2 boundary. There are clearly no nonzero constant functions in $H^2 \cap H_0^1$. Finally, injectivity follows from the comparison principle (see above footnote).

For a given function $v \in L^2(\Omega)$, the latter principle solves

$$\sup_{u \in \bar{\mathcal{U}}_L} \int_{\Omega} u(x)v(x) dx, \quad \bar{\mathcal{U}}_L := \left\{ u \in L^2(\Omega), 0 \leq u \leq 1 \text{ and } \int_{\Omega} u \leq L|\Omega| \right\}.$$

This optimisation problem comes up naturally in some control problems similar to ours [21, 28], or in shape optimisation problems [38].

Interpreting the bathtub principle as a Fenchel conjugate leads us to design the unusual cost functional (4). This allows us to design dual problems such that optimal controls exist, and are characterised as maximisers of some bathtub principle. Then, using analyticity properties for solutions of the dual problem, we prove their uniqueness and hence their extremality, thereby uncovering that they are on-off shape controls.

Bang-bang property of optimal controls. Bang-bang controls (*i.e.*, controls that saturate their constraints) are a common feature in time optimal control problems. A growing literature on the heat equation alone [29, 31, 43, 45, 47] shows that this property extends well to some infinite-dimensional systems. In our case, we will see that the on-off shape controls we have constructed can be understood as time-optimal controls. As these controls are bang-bang, this yields another occurrence of the bang-bang property in the time-optimal control of the heat equation.

Note, however, that in the references cited above, the controls are constrained to lie in balls of specific function spaces, whereas we consider non-negative constraints on the controls, which is an anisotropic constraint. Moreover, the bang-bang property is usually established separately using optimality conditions, having established controllability at the onset. In our case, the Fenchel-Rockafellar duality approach allows to do all those things simultaneously.

1.5 Extensions and perspectives

Operator, boundary conditions. The **(GUC)** property and analytic-hypoellipticity are two key sufficient properties for approximate controllability by on-off shape controls. We have highlighted second-order elliptic operators with analytic coefficients Dirichlet boundary conditions as an example. Our results apply to such operators with Robin boundary conditions of the form $a(x)y + b(x)\partial_{\nu}y = 0$ over $\partial\Omega$ (with a, b analytic) as soon as the function a does not vanish on the whole of $\partial\Omega$ (more generally, as soon as a is nontrivial on any connected component of $\partial\Omega$). This excludes the important case of Neumann boundary conditions, which remains open.

Our approach also accommodates subelliptic operators. This includes a large class of Hörmander operators, *i.e.*, operators of the form $A = \sum_{i=1}^m X_i^2 + X_0 + V\text{Id}$ with vector fields X_1, \dots, X_m generating a Lie algebra that equals \mathbb{R}^d on the whole of Ω . Under general regularity assumptions and boundary conditions, such an operator and its adjoint generate a strongly continuous semigroup on $L^2(\Omega)$, satisfy the comparison principle [6], all three conditions sufficient for the **(GUC)** property, and are analytic-hypoelliptic for instance if the characteristic manifold is an analytic symplectic manifold (see [32]).

Control operator. Our results have been stated with the identity control operator. They extend to the nonnegative control of

$$\begin{cases} y_t - Ay = \varphi u, \\ y(0) = y_0 \text{ in } \Omega \end{cases}$$

where $\varphi \in L^{\infty}(\Omega)$ is positive, analytic.

An interesting perspective is to follow our proof strategy with boundary control operators, where on-off shape controls now refer to characteristic functions over the boundary $\partial\Omega$.

Exact nonnegative controllability. In the case of unconstrained controllability with a control acting in some fixed subset ω , any function that can be reached exactly is (at least) analytic in $\Omega \setminus \omega$, preventing exact controllability to hold true.

On the one hand, this argument for (non)-exact controllability by on-off shape controls fails since the control may act everywhere. On the other hand, our approach heavily relies on targeting a ball $\bar{B}(y_f, \varepsilon)$ with $\varepsilon > 0$. As a result, exact nonnegative controllability by on-off shape controls is an open and seemingly difficult question.

A related matter is that of the cost of approximate controllability as a function of $\varepsilon \rightarrow 0$.

Abstract constrained control. The strategy of proof developed in this article hints at generalisations, where the method is applied to abstract linear control problems with abstract constraint sets \mathcal{U} .

In particular, we expect it to lead to necessary and sufficient conditions for controllability when \mathcal{U} is convex. When \mathcal{U} is not convex as is the case for on-off shape controls, this requires to study the convex hull of \mathcal{U} , following the relaxation approach. This abstract setting should allow us to discern how one can design a cost function F_T , analogous to (4), tailored to a given \mathcal{U} .

Further sufficient conditions should be derived to ensure that optimal controls in the convex hull of \mathcal{U} actually are in the original constraint set \mathcal{U} . In the present work, analytic-hypoellipticity and the **(GUC)** property play that role in the case of on-off shape controls.

This will be the subject of an ulterior article.

Regularity of the sets $\omega(t)$. Another problem is to analyse the complexity of the sets $\omega(t)$ occupied by optimal controls over time. For instance, how smooth (*BV* regularity, number of connected components, etc) are the sets $\omega(t)$ achieving approximate controllability?

In view of applications, these are important issues for the controls to be implementable in practice. For example, if the sets $\omega(t)$ are constrained to depend on a few parameters, or if they are restricted to rigid movements, controllability is a totally open question.

Numerical approximation of optimal controls. Optimal controls are given explicitly in terms of optimisers of the dual problem: the constructive nature of our approach means that optimal controls may be numerically computed, at least on paper.

Providing reliable and efficient methods to compute optimal controls is a difficult issue which has been studied in the case of Lions's cost functional with $\varepsilon = 0$ (*i.e.*, exact controllability) [7, 20]. Similar results in a generalised setting with our Fenchel-Rockafellar-based approach would be valuable.

Contrary to Lions's cost functional, we note that ad hoc algorithms are required in order to cope with functions that are not necessarily differentiable, as is the case in the present paper. Recent primal-dual algorithms designed for optimisation problems with objective functions of the form $F(u) + G(L_T u)$ are likely to be good candidates [10].

Outline of the paper. First, Section 2 lays out the convex analytic framework, that of Fenchel-Rockafellar duality, and how it may be applied to constrained approximate controllability. We then introduce the bathtub principle and interpret it in terms of Fenchel conjugation in order to design a relevant optimal control problem for our purposes. Section 3 is dedicated to the proof of our nonnegative approximate controllability result given by Theorem 3.1, and Section 4 to that of the obstruction result, Theorem 4.1. Finally, Section 5 gathers our results about further obstructions when the control amplitude is bounded, along with our analysis of the corresponding minimal time control problem.

2 Building the optimal control problem

2.1 Convex analytic framework

Let H be a Hilbert space. We let $\Gamma_0(H)$ be the set of functions from H to $]-\infty, +\infty]$ that are convex, lower semicontinuous (abbreviated lsc) and proper (*i.e.*, not identically $+\infty$). For $f \in \Gamma_0(H)$, we let

$$\text{dom}(f) = \{x \in H, f(x) < +\infty\}$$

be its *domain*.

Fenchel conjugate. For a proper function $f : H \rightarrow]-\infty, +\infty]$, we denote $f^* : H \rightarrow]-\infty, +\infty]$ its convex conjugate, given by the convex lsc function

$$f^*(y) := \sup_{x \in H} (\langle y, x \rangle - f(x)), \quad \forall y \in H.$$

Support and indicator functions. Given a subset $C \subset H$, the *indicator function* of C is the function defined by

$$\delta_C(x) := \begin{cases} 0 & \text{if } x \in C \\ +\infty & \text{if } x \notin C \end{cases}, \quad \forall x \in H,$$

and the *support function* of C is defined by

$$\sigma_C(p) := \sup_{x \in C} \langle p, x \rangle = \delta_C^*(p), \quad \forall p \in H,$$

i.e., the Fenchel conjugate function of the indicator function of C .

Subdifferentials. For $f \in \Gamma_0(H)$, we let

$$\partial f(x) := \{p \in H, \forall y \in H, f(y) \geq f(x) + \langle p, y - x \rangle\},$$

be its subdifferential at a point $x \in H$.

Various common properties of Fenchel conjugates, support functions and subdifferentials are used throughout the article. These are all recalled in Appendix A, where a few additional lemmas are proved.

2.2 Approximate controllability by Fenchel duality ([24])

Let us explain how the approximate controllability problem is reformulated in the context of Fenchel-Rockafellar duality [40] (see A.4 for a general presentation), following the strategy introduced by Lions in [24]. We work with the control problem (5), with the control space $E := L^2((0, T) \times \Omega)$ and the state space $L^2(\Omega)$.

By Duhamel's formula $y(T) = S_T y_0 + L_T u$, the inclusion $y(T) \in \overline{B}(y_f, \varepsilon)$ (where the closed ball of center y_f and radius ε is with respect to the $L^2(\Omega)$ -norm) can equivalently be written as $L_T u \in \overline{B}(y_f - S_T y_0, \varepsilon)$.

Given some cost functional $F_T : E \rightarrow [0, +\infty] \in \Gamma_0(E)$, consider the optimal control problem (which we will refer to as the *primal* problem)

$$\pi := \inf_{u \in E} F_T(u) + G_{T, \varepsilon}(L_T u).$$

where

$$G_{T, \varepsilon} := \delta_{\overline{B}(y_f - S_T y_0, \varepsilon)} \in \Gamma_0(L^2(\Omega)).$$

Now consider the Fenchel *dual* to the above problem, which writes

$$d = - \inf_{p_f \in L^2(\Omega)} J_{T,\varepsilon}(p_f), \quad J_{T,\varepsilon}(p_f) := F_T^*(L_T^* p_f) + G_{T,\varepsilon}^*(-p_f). \quad (8)$$

Thanks to the formulae for conjugates, we find

$$G_{T,\varepsilon}^*(z) = \langle y_f - S_T y_0, z \rangle_{L^2} + \varepsilon \|z\|_{L^2},$$

leading to

$$J_{T,\varepsilon}(p_f) = F_T^*(L_T^* p_f) - \langle y_f - S_T y_0, p_f \rangle_{L^2} + \varepsilon \|p_f\|_{L^2}.$$

We recall that $p = L_T^* p_f$ solves the adjoint equation

$$\begin{cases} p_t + A^* p = 0, \\ p(T) = p_f \text{ in } \Omega. \end{cases} \quad (9)$$

Strong duality. Weak duality $\pi \geq d$ always holds. According to the Fenchel-Rockafellar duality theorem recalled in Appendix A.4, the existence of $p_f \in \text{dom}(G_{T,\varepsilon}^*)$ such that F_T^* is continuous at $L_T^* p_f$ is a sufficient condition for strong duality $\pi = d$ to hold. Since $\text{dom}(G_{T,\varepsilon}^*) = L^2(\Omega)$, this condition reduces to the existence of a point of continuity of the form $L_T^* p_f$ for F_T^* . In the cases covered here, we shall check that the chosen F_T^* is continuous at 0. When strong duality obtains, it is therefore equivalent to work with the dual problem, which is easier to handle especially when it has full domain, *i.e.*, its objective function is finite everywhere.

Non-trivial strong duality. Furthermore, the primal value π is attained if finite, *i.e.*, if this equality is not the trivial $+\infty = +\infty$ (the uncontrollable case). Thus, if d is finite, π is finite as well and attained: we may speak of optimal controls.

This requirement that d be finite is by far the subtlest one. It may be tackled by proving that the functional $J_{T,\varepsilon}$ underlying the dual problem (written in infimum form $\inf_{p_f \in L^2(\Omega)} J_{T,\varepsilon}(p_f)$) has a minimum. In practice, we will always find this to be the case, as the dual problem is usually unconstrained (depending on the choice of F_T), unlike the primal problem. Hence, both π and d will be attained and, from Proposition A.8, any optimal dual variable p_f^* is such that any optimal control u^* satisfies

$$u^* \in \partial F_T^*(L_T^* p_f^*). \quad (10)$$

Proposition 2.1. *Assume that, for any $y_0, y_f \in L^2(\Omega)$ such that $y_f \geq S_T y_0$ and any $\varepsilon > 0$,*

- *there exists $p_f \in L^2(\Omega)$ such that F_T^* is continuous at $L_T^* p_f$,*
- *$d \neq +\infty$.*

If for any dual optimal variable p_f^ , the controls characterised by (10) are in \mathcal{U}_+ , then the control system (5) is nonnegatively approximately controllable with controls in \mathcal{U}_+ in time T .*

This shows how the choice of the cost F_T impacts the existence and properties of optimal controls. More precisely, it must be pointed out that all the hypotheses of Proposition 2.1 are formulated with respect to the dual problem. Accordingly, from the next section onwards, our strategy will be to determine an adequate optimal control problem by designing its dual problem.

Finally, we emphasise that (10) is only a necessary condition for the optimality of u^* . It becomes sufficient only when $\partial F_T^*(L_T^* p_f^*)$ is reduced to a singleton, which will occur in our case.

2.3 Convex analytic interpretation of the bathtub principle

Starting from the set of on-off shape controls of amplitude 1,

$$\mathcal{U}_L := \{\chi_\omega, \quad \omega \subset \Omega, \quad |\omega| \leq L|\Omega|\}, \quad (11)$$

where $|\cdot|$ denotes the Lebesgue measure, we define the closure of its convex hull (which is also its weak-* closure for the $L^\infty(\Omega)$ -topology)

$$\bar{\mathcal{U}}_L := \left\{ u \in L^2(\Omega), \quad 0 \leq u \leq 1 \text{ and } \int_\Omega u \leq L|\Omega| \right\}. \quad (12)$$

Given a fixed $v \in L^2(\Omega)$, we consider the (static) maximisation problem

$$\sup_{u \in \bar{\mathcal{U}}_L} \int_\Omega u(x)v(x) dx. \quad (13)$$

This is a relaxed version of the so-called bathtub principle, which gives the maximum value as well as a characterisation of maximisers. For the sake of readability, we introduce the necessary results for what follows, but refer to Appendix B for a more detailed statement. For a given $v \in L^2(\Omega)$, we let

$$\Phi_v(r) := |\{v > r\}|. \quad (14)$$

and its pseudo-inverse function

$$\Phi_v^{-1}(s) := \inf_{r \in \mathbb{R}} \{\Phi_v(r) \leq s\} = \inf_{r \in \mathbb{R}} \{|\{v > r\}| \leq s\}. \quad (15)$$

Finally, we set

$$h(v) := \max(0, \Phi_v^{-1}(L|\Omega|)). \quad (16)$$

Remark 2.2. *The function Φ_v^{-1} is the Schwarz radial rearrangement of v , see [18].*

Lemma 2.3 (relaxed bathtub principle). *Let $v \in L^2(\Omega)$. The maximum in (13) equals*

$$\int_0^{\min(\Phi_v(0), L|\Omega|)} \Phi_v^{-1}.$$

Furthermore, if all the level sets of the function v have measure zero, the maximum equals $\int_{\{v > h(v)\}} v$ and is uniquely attained by

$$u^* := \chi_{\{v > h(v)\}},$$

We refer to Lemma B.2 for the comprehensive statement of the relaxed bathtub principle. We may interpret the above results as a formula for the support function of $\bar{\mathcal{U}}_L$ in $L^2(\Omega)$:

$$\sigma_{\bar{\mathcal{U}}_L}(v) = \sup_{u \in \bar{\mathcal{U}}_L} (\langle u, v \rangle_{L^2} - \delta_{\bar{\mathcal{U}}_L}(u)) = \sup_{u \in \bar{\mathcal{U}}_L} \int_\Omega u(x)v(x) dx = \int_0^{\min(\Phi_v(0), L|\Omega|)} \Phi_v^{-1}. \quad (17)$$

First, using the characterisation of the subdifferential given in Appendix A, we arrive at the following characterisation for the solutions to the maximisation problem given in Lemma 2.3:

Proposition 2.4. *Let $v \in L^2(\Omega)$. The maximisers of the relaxed bathtub problem are given by the elements of $\partial\sigma_{\bar{\mathcal{U}}_L}(v)$.*

Proof. We have, for $v \in L^2(\Omega)$,

$$\arg \max_{u \in \bar{\mathcal{U}}_L} \langle u, v \rangle_{L^2} = \arg \max_{u \in L^2} \langle u, v \rangle_{L^2} - \delta_{\bar{\mathcal{U}}_L}(u) = \{u \in L^2, \quad \langle u, v \rangle_{L^2} - \delta_{\bar{\mathcal{U}}_L}(u) = \sigma_{\bar{\mathcal{U}}_L}(v)\} = \partial\sigma_{\bar{\mathcal{U}}_L}(v),$$

where we have used $(\delta_{\bar{\mathcal{U}}_L})^* = \sigma_{\bar{\mathcal{U}}_L}$ along with (46) given in Appendix A.1. \square

Remark 2.5. Proposition 2.4 implies that for any maximiser u of the relaxed bathtub problem,

$$v \in \partial \delta_{\overline{U}_L}(u).$$

Proposition A.4 in Appendix A.2 shows that this implies $u \in \partial \overline{U}_L$. Propositions 2.3 and 2.4 characterise exactly which elements of the boundary $\partial \overline{U}_L$ are involved.

2.4 From the static bathtub principle to the dual problem and its corresponding cost

Following Section 2.2 and recalling Proposition 2.1 and (10), we are looking for a cost function F_T such that the corresponding optimal controls are on-off shape controls, and we have established that it suffices to find a conjugate functional F_T^* satisfying two key properties. First, if there exists $p_f \in L^2(\Omega)$ such that F_T^* is continuous at $L_T^* p_f$, and if we can provide the existence of a minimiser p_f^* of $J_{T,\varepsilon}$, then π is attained and there exists at least one optimal control. Second, any optimal control u^* should satisfy (10) so F_T^* should be chosen so that the subdifferential $\partial F_T^*(L_T^* p_f^*)$ contains only characteristic functions. Given Proposition A.4 and Section 2.3, elements of

$$\partial \sigma_{\overline{U}_L}(v), \quad v \in L^2(\Omega)$$

are bang-bang, in the sense that they are characteristic functions, under some mild conditions that v must satisfy.

To go from the static optimisation problem to the adequate dual problem, we add a time dependency. Moreover, to ensure coercivity of the dual problem, we add a quadratic exponent. All in all, we choose the following conjugate:

$$F_T^*(p) := \frac{1}{2} \left(\int_0^T \sigma_{\overline{U}_L}(p(t)) dt \right)^2 = \frac{1}{2} \left(\int_0^T \int_0^{\min(\Phi_{p(t)}(0), L|\Omega|)} \Phi_{p(t)}^{-1}(s) ds dt \right)^2, \quad \forall p \in E. \quad (18)$$

Since the controllability problem corresponds to $G_{T,\varepsilon}^* := \sigma_{\overline{B}(y_f - S_T y_0, \varepsilon)}$, this defines a dual problem of the form (8). As pointed out in Section 2.2, we are now dealing with an unconstrained optimisation problem (*i.e.*, the domain of the functions involved is the whole space $L^2(\Omega)$).

We can now derive the corresponding constrained optimisation problem, by compute the actual cost F_T associated to the choice (18) for F_T^* . We find, as announced by (4) in the introduction:

Lemma 2.6. *The function F_T^* defined by (18) satisfies $F_T^* \in \Gamma_0(E)$. Defining*

$$\mathcal{M}(u) := \max \left(\|u\|_{L^\infty}, \frac{\|u\|_{L^1}}{L|\Omega|} \right), \quad \forall u \in L^2(\Omega),$$

its Fenchel conjugate $(F_T^)^* = F_T$ is given for $u \in E$ by*

$$F_T(u) = \frac{1}{2} \left(\sup_{t \in [0, T]} \mathcal{M}^2(u(t, \cdot)) \right) + \delta_{\{u \geq 0\}}(u) = \frac{1}{2} \left(\sup_{t \in [0, T]} \max \left(\|u(t, \cdot)\|_{L^\infty}, \frac{\|u(t, \cdot)\|_{L^1}}{L|\Omega|} \right)^2 \right) + \delta_{\{u \geq 0\}}(u).$$

Proof. Lemma A.5 in Appendix A.3 shows that $F_T^* \in \Gamma_0(E)$. We proceed by computing $(F_T^*)^*$. We have $F_T^* = \frac{1}{2} H^2$, with $H(p) := \int_0^T \sigma_{\overline{U}_L}(p(t, \cdot)) dt$. Since $\sigma_{\overline{U}_L} \in \Gamma_0(L^2(\Omega))$, the definition of the support function together with Lemma A.5 in Appendix show that $H \in \Gamma_0(E)$ with

$$H^*(u) = \int_0^T \sigma_{\overline{U}_L}^*(u(t, \cdot)) dt = \int_0^T \delta_{\overline{U}_L}(u(t, \cdot)) dt.$$

Furthermore, we find the conjugate of $\frac{1}{2} H^2$ by using (45) in Appendix A.1, which leads to

$$\left(\frac{1}{2} H^2 \right)^*(u) = \min_{\alpha > 0} \left(\frac{1}{2} \alpha^2 + \alpha H^* \left(\frac{u}{\alpha} \right) \right),$$

where we used that $\text{dom}(H) = E$. Clearly,

$$H^* \left(\frac{u}{\alpha} \right) = 0 \quad \text{if } u \geq 0 \quad \text{and} \quad \sup_{t \in [0, T]} \mathcal{M}(u(t, \cdot)) \leq \alpha,$$

and is $+\infty$ otherwise.

We end up with

$$\left(\frac{1}{2} H^2 \right)^* (u) = \min_{\alpha > 0} \left(\frac{1}{2} \alpha^2 + \delta_{\{\sup_{t \in [0, T]} \mathcal{M}(u(t, \cdot)) \leq \alpha\}} (u) \right) + \delta_{\{u \geq 0\}} (u) = F_T(u).$$

The lemma is proved. \square

Note that F_T^* is (positively)-homogeneous of degree 2. Indeed, $v \mapsto \sigma_{\bar{u}_L}$ is positively-homogeneous of degree 1, i.e., $\sigma_{\bar{u}_L}(\lambda v) = \lambda \sigma_{\bar{u}_L}(v)$ for all $\lambda > 0$, $v \in L^2(\Omega)$.

We end this subsection by establishing a crucial property satisfied by F_T^* . It will play a role similar to that of the unique continuation property in proving that the dual functional is coercive.

Lemma 2.7. *For all $p_f \in L^2(\Omega)$, if $F_T^*(L_T^* p_f) = 0$, then $p_f \leq 0$.*

Proof. By definition of F_T^* and using (17), the equality $F_T^*(L_T^* p_f) = 0$ entails

$$\sup_{u \in \bar{u}_L} \langle u, p(t, \cdot) \rangle_{L^2} = \int_0^{\min(\Phi_{p(t, \cdot)}(0), L|\Omega|)} \Phi_{p(t, \cdot)}^{-1} = 0, \quad \text{for a.e. } t \in (0, T),$$

where p is the solution to the adjoint equation (9) such that $p(T) = p_f$.

For a fixed $t \in (0, T)$, it is easily seen that the supremum on the left-hand side is positive as soon as $p(t, \cdot) > 0$ on a set of positive measure, by appropriately choosing u supported in this set. We have proved that $p(t, \cdot) \leq 0$ for a.e. $t \in (0, T)$ and in particular that $p_f \leq 0$ since $p \in C([0, T]; L^2(\Omega))$. \square

3 Approximate controllability results

In this section, we state and prove our main result on approximate controllability. The full statement for our Theorem A is given with more details below, for general linear operators, satisfying the properties given in Section 1.3.

We are considering the following optimal control problem:

$$\pi = \inf_{u \in E} F_T(u) + G_{T, \varepsilon}(L_T u) = \inf_{u \in E} \left\{ \frac{1}{2} \sup_{t \in [0, T]} \max \left(\|u(t)\|_{L^\infty}, \frac{\|u(t)\|_{L^1}}{L|\Omega|} \right)^2 + \delta_{\bar{B}(y_f - S_T y_0, \varepsilon)}(L_T u) \right\}, \quad (19)$$

whose dual problem is

$$d = - \inf_{p_f \in L^2(\Omega)} J_{T, \varepsilon}(p_f) = - \inf_{p_f \in L^2(\Omega)} \left\{ \frac{1}{2} \left(\int_0^T \sigma_{\bar{u}_L}(L_T^* p_f(t)) dt \right)^2 - \langle y_f - S_T y_0, p_f \rangle_{L^2} + \varepsilon \|p_f\|_{L^2} \right\}. \quad (20)$$

Theorem 3.1. *Assume that A^* satisfies the (GUC) property and that $\partial_t - A^*$ is analytic-hypoelliptic. Then for the cost function F_T defined by (4),*

- *strong duality $\pi = d$ holds,*
- *the dual problem (20) is attained at a unique minimiser $p_f^* \in L^2(\Omega)$,*
- *there exists a unique optimal control $u^* \in E$ for the primal problem (19).*

Furthermore, if the target is not reached by the trivial control $u \equiv 0$, i.e., if

$$y_f \notin \overline{B}(S_T y_0, \varepsilon), \quad (21)$$

then the optimal control is given by

$$u^*(t, \cdot) = \overline{M} \chi_{\{p^*(t, \cdot) > h(p^*(t, \cdot))\}}, \quad \overline{M} = \int_0^T \int_{\{p^*(t, \cdot) > h(p^*(t, \cdot))\}} p^*(t, x) dt dx, \quad (22)$$

where h is defined by (16), and where $p^* = L_T^* p_f^*$ is the solution of the adjoint equation (9) satisfying $p^*(T) = p_f^*$.

Remark 3.2. As mentioned in the introduction, Theorem 3.1 holds for uniformly elliptic operators of the form (7) with analytic coefficients, and in particular the classical heat equation with Dirichlet boundary conditions, on a bounded, open, connected domain with C^2 boundary.

Throughout this section, we assume the hypotheses sufficient for Theorem 3.1, i.e., that A^* satisfies the (GUC) property and that $\partial_t - A^*$ is analytic-hypoelliptic. The proof is then scattered into the section as follows:

- First, we establish that strong duality holds.
- Second, we prove that the corresponding dual functional is coercive: hence, the dual functional attains its minimum (the dual problem attains its maximum).
- Third we prove (22).
- Finally, we investigate the uniqueness of optimal variables.

Remark 3.3. As the proofs show, the first two steps and the uniqueness of dual optimal variables are valid for any operator A . In particular, they do not require that A^* satisfy the (GUC) property and that $\partial_t - A^*$ be analytic-hypoelliptic. Hence, strong duality and existence of optimal controls does not require any specific assumption the semigroup must satisfy. This remark will be of importance in the next subsection where we manipulate optimal controls without making these two hypotheses.

3.1 Strong duality

Lemma 3.4. F_T^* is continuous at $0 = L_T^* 0$.

Proof. By the Cauchy-Schwarz inequality,

$$\forall u \in \overline{U}_L, \quad \langle u, v \rangle_{L^2} \leq \|u\|_{L^2} \|v\|_{L^2} \leq |\Omega|^{1/2} \|v\|_{L^2},$$

which leads to

$$\sigma_{\overline{U}_L}(v) = \int_0^{\min(\Phi_v(0), L|\Omega|)} \Phi_v^{-1} \leq |\Omega|^{1/2} \|v\|_{L^2}.$$

As a result, we may bound with the Cauchy-Schwarz inequality again

$$0 \leq F_T^*(p) \leq \frac{1}{2} |\Omega| \left(\int_0^T \|p(t, \cdot)\|_{L^2} dt \right)^2 \leq \frac{1}{2} T |\Omega| \|p\|_E^2,$$

hence the continuity of F_T^* at $0 = L_T^* 0$. □

The above lemma shows that the first condition of Proposition 2.1 is satisfied, i.e., strong duality holds.

3.2 Coercivity of $J_{T,\varepsilon}$, nonnegative approximate controllability

Proposition 3.5. *The functional $J_{T,\varepsilon}$ defined by*

$$J_{T,\varepsilon}(p_f) = \left(\frac{1}{2} \int_0^T \sigma_{\mathcal{U}_L}(L_T^* p_f) dt \right)^2 - \langle y_f - S_T y_0, p_f \rangle_{L^2} + \varepsilon \|p_f\|_{L^2}. \quad (23)$$

is coercive on $L^2(\Omega)$, i.e.,

$$J_{T,\varepsilon}(p_f) \xrightarrow{\|p_f\|_{L^2} \rightarrow \infty} \infty,$$

and thus attains its minimum.

Proof. Since we know that $J_{T,\varepsilon}$ is convex, proper, strongly lsc, if $J_{T,\varepsilon}$ is coercive then $\inf_{p_f \in L^2(\Omega)} J_{T,\varepsilon}(p_f) \neq -\infty$, and that it is actually attained.

We will actually prove a stronger condition than coercivity, namely

$$\liminf_{\|p_f\|_{L^2} \rightarrow \infty} \frac{J_{T,\varepsilon}(p_f)}{\|p_f\|_{L^2}} > 0.$$

Our method of proof follows that of [19, 30]. Take a sequence $\|p_f^n\|_{L^2} \rightarrow \infty$. We denote $q_f^n := \frac{p_f^n}{\|p_f^n\|_{L^2}}$, and $q^n \in E$ the corresponding solution of the adjoint equation (9), i.e., such that $q^n(T) = q_f^n$, which by linearity is $\frac{p^n}{\|p_f^n\|_{L^2}}$, where $p^n = L_T^* p_f^n$ is the solution of (9) such that $p^n(T) = p_f^n$. By positive homogeneity of F_T^* (of degree 2), we have

$$\frac{J_{T,\varepsilon}(p_f^n)}{\|p_f^n\|_{L^2}} = \|p_f^n\|_{L^2} F_T^*(L_T^* q_f^n) - \langle y_f - S_T y_0, q_f^n \rangle_{L^2} + \varepsilon$$

and hence if $\liminf_{n \rightarrow \infty} F_T^*(L_T^* q_f^n) > 0$, then

$$\liminf_{n \rightarrow \infty} \frac{J_{T,\varepsilon}(p_f^n)}{\|p_f^n\|_{L^2}} = +\infty.$$

Let us now treat the remaining case where $\liminf_{n \rightarrow \infty} F_T^*(L_T^* q_f^n) = 0$. Since $\|q_f^n\|_{L^2} = 1$, upon extraction of a subsequence, we have $q_f^n \rightharpoonup q_f$ weakly in $L^2(\Omega)$ for some $q_f \in L^2(\Omega)$. Since $L_T^* \in L(L^2(\Omega), E)$, we have $L_T^* q_f^n \rightharpoonup L_T^* q_f$ weakly in E .

Now, since F_T^* is convex and strongly lsc on E , it is (sequentially) weakly lsc and taking the limit we obtain $F_T^*(L_T^* q_f) = 0$. By Lemma 2.7, we infer that $q_f \leq 0$.

Then, recalling that the target satisfies $y_f - S_T y_0 \geq 0 \Leftrightarrow y_f \geq S_T y_0$, we end up with

$$\liminf_{n \rightarrow \infty} \frac{J_{T,\varepsilon}(p_f^n)}{\|p_f^n\|_{L^2}} \geq -\langle y_f - S_T y_0, q_f \rangle_{L^2} + \varepsilon \geq \varepsilon > 0,$$

which concludes the proof. □

3.3 Characterisation of the minimisers

In this section, we assume that the target is not reached with the trivial control $u = 0$, i.e.,

$$y_f \notin \overline{B}(S_T y_0, \varepsilon)$$

Note that, if (21) is not satisfied, the control $u = 0$ steers y_0 to the target, and is indeed a control in $\mathcal{U}_{\text{shape}}^L$.

We first remark the following fact:

Lemma 3.6. *Under Assumption (21), any minimiser p_f^* of (20) satisfies $p_f^* \neq 0$.*

Proof. Suppose $p_f^* = 0$. Then, $d = 0$ and by strong duality, $\pi = 0$. By Proposition 3.5, this value is attained: there exists some optimal control u^* such that

$$F_T(u^*) + G_{T,\varepsilon}(L_T u^*) = 0.$$

This implies that $F_T(u^*) = G_{T,\varepsilon}(L_T u^*) = 0$. On the one hand, this leads to $u^*(t, \cdot) = 0$ for a.e. $t \in (0, T)$, i.e., $u = 0$, and on the other hand $G_{T,\varepsilon}(L_T u^*) = G_{T,\varepsilon}(0) = 0$, which is equivalent to $0 \in \overline{B}(y_f - S_T y_0, \varepsilon)$. This contradicts Assumption (21). \square

Proposition 3.7. *Under Assumption (21), any optimal control for (19) is of the form (22), with p^* the solution of the adjoint equation (9) such that $p^*(T) = p_f^*$, where p_f^* is any dual optimal variable.*

Proof. Thanks to Proposition 3.5, we know that $J_{T,\varepsilon}$ defined by (20) attains its minimum. Let p_f^* be a minimiser for $J_{T,\varepsilon}$, i.e., an optimal dual variable. From Lemma 3.6, we have $p_f^* \neq 0$. We denote p^* the solution of the adjoint equation (9) such that $p^*(T) = p_f^*$.

Let u^* be an optimal control. Thanks to Lemma 3.4, we can apply the first identity of (50) in Proposition A.8 (see Appendix A.4) to obtain $u^* \in \partial F_T^*(L_T^* p_f^*) = \partial F_T^*(p^*)$. Using again the notation $H(p) := \int_0^T \sigma_{\overline{U}_L}(p(t)) dt$, so that $F_T^* = \frac{1}{2} H^2$, we have $H(p^*) \geq 0$ and $\text{dom}(H) = L^2(\Omega)$.

Then, applying the generalised chain rule (see [11, Theorem 2.3.9, point (ii)]) with the functions $x \mapsto \frac{1}{2} x^2$ and H , we compute the subdifferential of the convex functional F_T^* : $u^* \in H(p^*) \partial H(p^*)$. Applying Lemma A.5 to H , we find $u^*(t, \cdot) \in \overline{M} \partial \sigma_{\overline{U}_L}(p^*(t, \cdot))$ for a.e. $t \in (0, T)$, with $\overline{M} := H(p^*)$.

Now, let $t \in (0, T)$ be fixed and let us justify that all level sets of $p^*(t, \cdot)$ are of measure zero, i.e.,

$$|\{p^*(t, \cdot) = \lambda\}| = 0, \quad \forall \lambda \in \mathbb{R},$$

Indeed, since the operator $\partial_t - A^*$ is analytic-hypoelliptic, we know that $p^*(t, \cdot)$ is analytic on Ω . Hence, its level sets are of measure zero unless $p^*(t, \cdot) = S_{T-t}^* p_f^*$ is constant. Using the **(GUC)** property, this leads to $p_f^* = 0$, contradicting (21).

Applying Propositions 2.3 and 2.4, and recalling that $\partial \sigma_{\overline{U}_L}(p^*(t, \cdot)) = \{\chi_{\{p^*(t, \cdot) > h(p^*(t, \cdot))\}}\}$, we obtain the result. \square

Remark 3.8. *As evidenced by the proof, a weaker (but less workable) property than analytic-hypoellipticity is sufficient to infer that optimal controls are on-off shape controls. Indeed, it suffices to require either one of the following conditions (in decreasing order of strength):*

- (i) *All solutions $t \mapsto p(t)$ of the adjoint equation such that $p(T) \neq 0$ have zero-measure level sets.*
- (ii) *For all solutions $t \mapsto p(t)$ of the adjoint equation such that $p(T) \neq 0$, the level sets $\{p(t, \cdot) = h(p(t, \cdot))\}$ (see (16) for the definition of $h(p)$) have measure 0.*
- (iii) *For all solutions $t \mapsto p(t)$ of the adjoint equation such that $p(T) \neq 0$,*

$$\begin{cases} |\{p(t, \cdot) = h(p(t, \cdot))\}| = L|\Omega| - |\{p(t, \cdot) > h(p(t, \cdot))\}|, & \text{if } h(p(t, \cdot)) \neq 0 \\ |\{p(t, \cdot) = h(p(t, \cdot))\}| = 0, & \text{if } h(p(t, \cdot)) = 0 \end{cases} \quad \text{for a.e. } t \in [0, T].$$

Note that requirement (iii) is minimal (see Lemma B.2 and Remark B.3).

Finally, an even weaker requirement would be to restrict any of the above (i), (ii) or (iii) to a single solution $t \mapsto p^(t)$ of the adjoint equation, namely that with $p^*(T) = p_f^*$ where p_f^* is the unique dual optimal variable (see below for the uniqueness of optimal variables).*

3.4 Uniqueness

Our first uniqueness statement below (*i.e.*, that of the dual optimal variable) is a consequence of Fenchel-Rockafellar duality, and the fact that we work with a Hilbert space, rather than specific properties of the evolution equation under consideration.

Remark 3.9. *Still applying Proposition A.8, we get*

$$L_T u^* \in \partial G_{T,\varepsilon}^*(-p_f^*) = \partial \sigma_{\overline{B}(y_f - S_T y_0, \varepsilon)}(-p_f^*).$$

Flipping the subdifferentials, we get $-p_f^ \in \partial \delta_{\overline{B}(y_f - S_T y_0, \varepsilon)}(L_T u^*)$. Thanks to Proposition A.4, this means that $L_T u^*$ lies at the boundary of the closed ball $\overline{B}(y_f - S_T y_0, \varepsilon)$.*

Proposition 3.10. *Under the assumptions of Theorem 3.1, the primal-dual optimal pairs (u^*, p_f^*) are unique.*

Proof. Uniqueness of the dual optimal variable. First note that if Assumption (21) does not hold, then 0 is the unique optimal control, *i.e.*,

$$\{L_T u^*, u^* \text{ is optimal}\} = \{0\}. \quad (24)$$

On the other hand, if Assumption (21) holds, according to Remark 3.9, and since the set of minimisers of a convex function is convex, the set $\{L_T u^*, u^* \text{ is optimal}\}$ is a convex subset of the sphere $S(y_f - S_T y_0, \varepsilon)$. The closed ball being strictly convex since we are working in the Hilbert space $L^2(\Omega)$, there exists some $y^* \in \overline{B}(y_f - S_T y_0, \varepsilon)$ with $\|y^* - (y_f - S_T y_0)\|_{L^2} = \varepsilon$ such that

$$\{L_T u^*, u^* \text{ is optimal}\} = \{y^*\}. \quad (25)$$

Thus, in any case, the set of targets reached by optimal controls is always reduced to a single point.

Now, let p_f^* be a dual optimal variable, and u^* an optimal control. Then, as strong duality holds, Proposition A.7 implies that the pair (u^*, p_f^*) satisfies the two optimality conditions from (25). We then have

$$p_f^* \in -\partial G_{T,\varepsilon}(L_T u^*) = -\partial \delta_{\overline{B}(y_f - S_T y_0, \varepsilon)}(L_T u^*). \quad (26)$$

If Assumption (21) does not hold, then (26) and (24) imply $p_f^* \in -\partial \delta_{\overline{B}(y_f - S_T y_0, \varepsilon)}(0)$. If $\|y_f - S_T y_0\|_{L^2} < \varepsilon$, then $0 \in B(y_f - S_T y_0, \varepsilon)$ and

$$p_f^* \in -\partial \delta_{\overline{B}(y_f - S_T y_0, \varepsilon)}(0) = \{0\}. \quad (27)$$

Otherwise, $0 \in \partial B(y_f - S_T y_0, \varepsilon)$ and (47) yield

$$p_f^* \in \left\{ \lambda \frac{y_f - S_T y_0}{\varepsilon}, \lambda \geq 0 \right\} = \{\lambda(y_f - S_T y_0), \lambda \geq 0\}.$$

Restricting the function $J_{T,\varepsilon}$ defining the primal problem (19) to the above half-line, using the homogeneities of each of its terms, and the fact that $\|y_f - S_T y_0\|_{L^2} = \varepsilon$, we get

$$\gamma_0(\lambda) := J_{T,\varepsilon}(\lambda(y_f - S_T y_0)) = a_0 \lambda^2, \quad \lambda \geq 0. \quad (28)$$

It is clear that 0 is the unique minimiser of γ_0 . From (27) and (28), 0 is the unique dual optimal variable if (21) does not hold.

If Assumption (21) holds, then (26) and (25) imply

$$p_f^* \in -\partial \delta_{\overline{B}(y_f - S_T y_0, \varepsilon)}(y^*) = -\partial \delta_{\overline{B}(0,1)}\left(\frac{y^* - (y_f - S_T y_0)}{\varepsilon}\right).$$

Since y^* lies at the boundary of $\overline{B}(y_f - S_T y_0, \varepsilon)$, formula (47) yields

$$p_f^* \in \left\{ \lambda \left(\frac{y_f - S_T y_0 - y^*}{\varepsilon} \right), \lambda \geq 0 \right\} = \{ \lambda (y_f - S_T y_0 - y^*), \lambda \geq 0 \}.$$

Restricting $J_{T,\varepsilon}$ to the above half-line as previously, we find

$$\gamma(\lambda) := J_{T,\varepsilon}(\lambda(y_f - S_T y_0 - y^*)) = a\lambda^2 + b\lambda, \quad \lambda \geq 0,$$

where, using $\|y_f - S_T y_0 - y^*\|_{L^2} = \varepsilon$ and the homogeneities involved $a = F_T^*(L_T^*(y_f - S_T y_0 - y^*))$ and $b = -\langle y_f - S_T y_0, y_f - S_T y_0 - y^* \rangle_{L^2} + \varepsilon^2$. By coercivity, $a > 0$, and given Lemma 3.6, we have $b < 0$.

Thus, γ has a unique minimiser $\lambda^* := -b/2a > 0$. Hence, $p_f^* = \lambda^*(y_f - S_T y_0 - y^*)$, and the dual optimal variable is unique.

Uniqueness of the optimal control. If Assumption (21) does not hold, then 0 is the unique optimal control.

Now, suppose that Assumption (21) holds. We know from the proof of Proposition 3.7 that a given dual optimal variable uniquely determines one optimal control. Moreover, as we have proved that strong duality holds, we can apply Proposition A.7: for any pair of primal and dual optimal variables, the relations (48) are satisfied. That is, any optimal control u^* is uniquely determined by the unique dual optimal variable p_f^* through the identity $u^* \in \partial F_T^*(L_T^* p_f^*)$. \square

4 Obstructions to controllability

We here prove Theorem B, through the more general result below in the case of second-order uniformly elliptic operators of the form (7). We use the notation $A \subset\subset B$ to mean that there exists a compact set K such that $A \subset K \subset B$.

Theorem 4.1. *Let $\mathcal{U}_+ \subset L^2(\Omega)$ be a constraint set of nonnegative controls. Assume that there exists a ball $B(x, r) \subset \Omega$ such that*

$$\forall u \in \mathcal{U}_+, \quad \text{supp}(u) \cap B(x, r) = \emptyset.$$

Let A be a second-order uniformly elliptic operator of the form (7). Let $y_0 = 0$ and $y_f \in L^2(\Omega)$ be any target such that $y_f \geq S_T y_0 = 0$, $y_f \neq 0$ and $\text{supp}(y_f) \subset B(x, r)$. Then there exist $T^ > 0$ and $\varepsilon > 0$ such that for any time $T \leq T^*$, no control with values in \mathcal{U}_+ can steer 0 to $\overline{B}(y_f, \varepsilon)$.*

The proof relies on the following lemma, inspired by [36].

Lemma 4.2. *Let $B(x, r) \subset\subset \Omega$. Under the assumptions of Theorem 4.1, for any $K \subset B(x, r)$ compact, there exists $p_f \in L^2(\Omega)$ such that*

- (i) $p_f < 0$ on K ,
- (ii) $\exists T^* > 0$ such that for all $t \in (0, T^*)$, $p(t, \cdot) \geq 0$ on $\Omega \setminus B(x, r)$, where p solves the adjoint equation (9) with $p(T) = p_f$.

Proof. Let us build p_f such that for all $1 < r < +\infty$, $p_f \in W^{2,r}(\Omega) \cap W_0^{1,r}(\Omega)$, with $p_f < 0$ on K , $p_f > 0$ on $\Omega \setminus B(x, r)$, $p_f = 0$ on $\partial\Omega$, and $\partial_\nu p_f < 0$ on $\partial\Omega$.

To that end, we denote φ_1 the first eigenfunction of the Dirichlet Laplacian on Ω , which satisfies $\varphi_1 > 0$ on Ω and $\partial_\nu \varphi_1 < 0$ on $\partial\Omega$ and since Ω is of class C^2 , $\varphi_1 \in W^{2,r}(\Omega) \cap W_0^{1,r}(\Omega)$ for all $1 < r < +\infty$ [9][Theorem 9.32]. We then set $p_f = \xi \varphi_1$ where $\xi \in C^\infty(\Omega)$ is chosen to satisfy $\xi = 1$ on $\Omega \setminus B(x, r)$ and $\xi = -1$ on K . The function p_f satisfies all the required properties (note that $p_f = \varphi_1$ locally around $\partial\Omega$ since $B(x, r) \subset\subset \Omega$, hence $\partial_\nu p_f = \partial_\nu \varphi_1 < 0$ on $\partial\Omega$).

We now set $q(t) = p(T - t)$, so that q solves the (forward) adjoint equation (9) with $q(0) = p_f$. Then, by parabolic regularity, we both have $q \in C([0, T] \times \overline{\Omega})$ and $\partial_\nu q \in C([0, T] \times \partial\Omega)$ [36][Theorem 8.1]. As

a result, by continuity there exists T^* such that $\partial_\nu q < 0$ over $[0, T^*] \times \partial\Omega$, there exists some compact set K_1 containing $B(x, r)$ such that $q \geq 0$ on $[0, T^*] \times (\Omega \setminus K_1)$.

Then, upon reducing T^* if necessary and by continuity again, we have $q \geq 0$ over $[0, T^*] \times (K_1 \setminus B(x, r))$, which concludes the proof. \square

Proof of Theorem 4.1. Upon reducing r , we may without loss of generality assume that $B(x, r) \subset\subset \Omega$. Letting $K := \text{supp}(y_f)$, we consider p_f as given by Lemma 4.2, and the corresponding T^* .

Let $T \leq T^*$ be fixed. For any control $u \in E$, any $y_0, y_f \in L^2(\Omega)$, any solution to the adjoint equation (9) such that $p(T) = p_f$, we have $\frac{d}{dt} \langle y(t), p(t) \rangle_{L^2} = \langle p(t), u(t) \rangle_{L^2}$. As a result and owing to $y_0 = 0$,

$$\langle y(T), p_f \rangle_{L^2} = \int_0^T \langle p(t), u(t) \rangle_{L^2} dt. \quad (29)$$

We now assume by contradiction that, for any $\varepsilon > 0$ there exists a nonnegative control $u_\varepsilon \in E$ satisfying $\forall t \in (0, T)$, $\text{supp}(u_\varepsilon(t)) \cap B(x, r) = \emptyset$ and steering $y_0 = 0$ to the ball $\bar{B}(y_f, \varepsilon)$ in time T . We inspect the sign of the equality (29) along the controls u_ε , $\varepsilon > 0$.

On the one hand, because of the condition (ii) in Lemma 4.2 satisfied by p , and owing to $u_\varepsilon \geq 0$, the right-hand side of (29) is nonnegative, *i.e.*,

$$\langle y(T), p_f \rangle_{L^2} \geq 0. \quad (30)$$

On the other hand, the left-hand side of (29) satisfies

$$\langle y(T), p_f \rangle_{L^2} = \langle y_f, p_f \rangle_{L^2} + \langle y(T) - y_f, p_f \rangle_{L^2} \leq \langle y_f, p_f \rangle_{L^2} + \varepsilon \|p_f\|_{L^2}$$

Now, $\langle y_f, p_f \rangle_{L^2} < 0$, because of (i) in Lemma 4.2. As a result, there exists $\alpha > 0$ such that $p_f \leq -\alpha$ on K , so that

$$\langle y_f, p_f \rangle_{L^2} \leq -\alpha \int_K y_f < 0,$$

because y_f is nonnegative and nontrivial on K by assumption.

Hence, for $\varepsilon > 0$ small enough, $\langle y(T), p_f \rangle_{L^2} < 0$, which contradicts (30). \square

Remark 4.3. *As the proof shows, the obstruction to nonnegative approximate controllability in \mathcal{U}_+ does not rely on the comparison principle, but is of dual nature as evidenced by the core idea behind it, *i.e.*, to construct p_f and y_f violating equality (29). The proof of Theorem 4.1 follows directly from the existence of p_f satisfying the assumptions of Lemma 4.2. Hence, this obstruction to nonnegative approximate controllability is rather general and will be satisfied by any operator (including uniformly second-order elliptic operator of the form (7)) for which such an element p_f can be built.*

5 Further comments

5.1 Properties of the value function in the general case

For general linear operators generating a C_0 semigroup, fixing Ω , L , ε , y_0 and y_f , we analyse the dependence with respect to the final time T , for the optimal control problem (19) studied in Section 3 for system (5).

By Lemma 3.4 and Proposition 3.5, the optimal control problem (19) is well-posed, *i.e.*, optimal controls exist (see also Remark 3.3), hence we may consider

$$\Pi(T) := \frac{1}{2}(\overline{M}(T))^2 := \inf\{F_T(u), \quad u \in E, \quad \|L_T u - (y_f - S_T y_0)\|_{L^2} \leq \varepsilon\}, \quad T > 0. \quad (31)$$

When A^* satisfies the **(GUC)** property and $\partial_t - A^*$ is analytic-hypoelliptic, $\overline{M}(T)$ is the amplitude of the unique optimal control in Proposition 3.7.

Recall that by strong duality, we have

$$\Pi(T) = \frac{1}{2}(\overline{M}(T))^2 = -J_{T,\varepsilon}(p_T^*), \quad \forall T \geq 0, \quad (32)$$

where p_T^* is the unique minimiser of $J_{T,\varepsilon}$. This is exactly the identity obtained for the HUM method where the cost functional F_T is just $\frac{1}{2}\|\cdot\|_E^2$.

We first establish the continuity of $T \mapsto \overline{M}(T)$.

Proposition 5.1. \overline{M} (and thus Π) are continuous on $(0, +\infty)$.

Proof. Using (32), we prove the continuity by showing that $(p_f, T) \mapsto J_{T,\varepsilon}(p_f)$ (given by (23)) satisfies the assumptions of Lemma A.9 with $H = L^2(\Omega)$ and $Z = (0, +\infty)$. Clearly, the first, second and fourth assumptions are satisfied, hence we are left with proving that $(p_f, T) \mapsto J_{T,\varepsilon}(p_f)$ is weak-strong lower semicontinuous over $L^2(\Omega) \times (0, +\infty)$. The last two terms of (23) are easily seen to be weak-strong lower semicontinuous over $L^2(\Omega) \times (0, +\infty)$, hence we investigate the property for the remaining term $F_T^*(L_T^*p_f)$.

Given $p_f \in L^2(\Omega)$ and $T > 0$, let (p_f^n) and (T_n) be two sequences such that $p_f^n \rightharpoonup p_f$, $T_n \rightarrow T$. We decompose

$$F_{T_n}^*(L_{T_n}^*p_f^n) = F_T^*(L_T^*p_f^n) + (F_{T_n}^*(L_{T_n}^*p_f^n) - F_T^*(L_T^*p_f^n)).$$

By weak (sequential) lower semicontinuity of F_T^* over $L^2(0, T; L^2(\Omega))$, we find that the first term satisfies

$$F_T^*(L_T^*p_f) \leq \liminf_{n \rightarrow +\infty} F_T^*(L_T^*p_f^n).$$

To conclude, we only need to prove that the second term tends to 0 as $n \rightarrow +\infty$.

Using the notation q_n for the solution to the forward adjoint problem such that $q_n(0) = p_f^n$, i.e., $q_n(t) = S_t^*p_f^n$, we have

$$\begin{aligned} F_{T_n}^*(L_{T_n}^*p_f^n) - F_T^*(L_T^*p_f^n) &= \frac{1}{2} \left(\int_0^{T_n} \sigma_{\overline{u}_L}(q_n(T_n - t)) dt \right)^2 - \frac{1}{2} \left(\int_0^T \sigma_{\overline{u}_L}(q_n(T - t)) dt \right)^2 \\ &= \frac{1}{2} \left(\int_T^{T_n} \sigma_{\overline{u}_L}(q_n(t)) dt \right) \left(\int_0^{T_n} \sigma_{\overline{u}_L}(q_n(t)) dt + \int_0^T \sigma_{\overline{u}_L}(q_n(t)) dt \right) \end{aligned}$$

Using the bound $0 \leq \sigma_{\overline{u}_L}(p) \leq |\Omega|^{1/2} \|p\|_{L^2}$ (see the proof of Lemma 3.4) and the estimate $\|S_t\|_{L(L^2(\Omega))} \leq C$ valid for all $t \in [0, T+1]$ with $C > 0$ some constant independent of n , we have

$$\left| \int_0^{T_n} \sigma_{\overline{u}_L}(q_n(t)) dt + \int_0^T \sigma_{\overline{u}_L}(q_n(t)) dt \right| \leq C|\Omega|^{1/2}(T + T_n) \|p_f^n\|_{L^2},$$

a bounded quantity, and

$$\left| \int_T^{T_n} \sigma_{\overline{u}_L}(q_n(t)) dt \right| \leq C|\Omega|^{1/2}|T - T_n| \|p_f^n\|_{L^2},$$

which tends to 0 as $n \rightarrow +\infty$. \square

We now study the behaviour of $\overline{M}(T)$ near $T = 0$ and $T = +\infty$. We recall that $\overline{M}(T)$ also depends on all other parameters y_0, y_f, ε and L .

We now recall (see [35]) that there exist $C_s > 0, \alpha \in \mathbb{R}$ such that for all $t \geq 0$, $\|S_t\|_{L(L^2(\Omega))} \leq C_s e^{\alpha t}$, and the semi-group generated by $(A, D(A))$ is said to be *exponentially stable* if $\alpha < 0$.

Proposition 5.2. *We have*

$$\forall T > 0, \quad \overline{M}(T) \geq |\alpha| \frac{\|y_f - S_T y_0\|_{L^2} - \varepsilon}{\sqrt{L|\Omega|}(1 - e^{\alpha T})}. \quad (33)$$

Proof. Let u_T^* be an optimal control in time T for the optimal control problem (31), then

$$\begin{aligned} \|L_T u_T^*\|_{L^2} &= \left\| \int_0^T S_{T-t} u_T^*(t, \cdot) dt \right\|_{L^2} \leq \int_0^T \|S_{T-t} u_T^*(t, \cdot)\|_{L^2} dt \\ &\leq \int_0^T e^{\alpha(T-t)} \|u_T^*(t, \cdot)\|_{L^2} dt \leq \frac{1}{|\alpha|} (1 - e^{\alpha T}) \overline{M}(T) \sqrt{L|\Omega|}. \end{aligned}$$

Now, by definition of our control problem, for all $T > 0$, $\|y_f - S_T y_0\|_{L^2} - \varepsilon \leq \|L_T u_T^*\|_{L^2}$, and the result follows. \square

Corollary 5.3. *Assume that $y_f \notin \overline{B}(y_0, \varepsilon)$. Then:*

$$\frac{1}{T} = O_{T \rightarrow 0}(\overline{M}(T)). \quad (34)$$

In particular, $\overline{M}(T) \xrightarrow{T \rightarrow 0} +\infty$.

Assume that $y_f \notin \overline{B}(0, \varepsilon)$. If, additionally, $(S_t)_{t \geq 0}$ is exponentially stable, then

$$\liminf_{T \rightarrow +\infty} \overline{M}(T) > 0. \quad (35)$$

Proof. The estimate (34) is obtained by passing to the limit in (33), using that $S_T y_0 \xrightarrow{T \rightarrow 0} y_0$: the lower bound behaves as $\frac{\|y_f - y_0\|_{L^2} - \varepsilon}{\sqrt{L|\Omega|}} \frac{1}{T}$. The inequality (35) is obtained by passing to the limit $T \rightarrow +\infty$ in (33), using that $S_T y_0 \xrightarrow{T \rightarrow \infty} 0$:

$$\liminf_{T \rightarrow +\infty} \overline{M}(T) \geq |\alpha| \frac{\|y_f\|_{L^2} - \varepsilon}{\sqrt{L|\Omega|}} > 0. \quad \square$$

5.2 Obstructions

We further investigate the behaviour of \overline{M} , and establish results on the corresponding minimal time problem (37). The comparison principle formulated in (6) will be a key ingredient in our study.

5.2.1 Obstruction to reachability and small-time controllability

Given the controllability result of Theorem 3.1, in order to study possible obstructions, we introduce a new bound on the amplitude of the control, of the form:

$$M(u) := 2\sqrt{F_T(u)} \leq M_{\max}, \quad u \in E, \quad (36)$$

for some $M_{\max} > 0$. Note that such a constraint imposes nonnegativity of the control. With this new constraint on the controls, we illustrate a general property that is well known for finite-dimensional systems: exponential stability prevents reachability.

In particular, the result below holds for uniformly elliptic operators of the form (7) with 0th order coefficient satisfying $c \leq 0$.

Proposition 5.4. *Assume that $(S_t)_{t \geq 0}$ is exponentially stable. Let (y_0, y_f) be such that for all $T \geq 0$, $y_f \geq S_T y_0$ and $\|S_T y_0 - y_f\|_{L^2} \geq \delta$ for some $\delta > 0$. Then, for all $0 < \varepsilon < \delta$ there exists $M_{\max} M(y_0, y_f, \varepsilon) > 0$ satisfying*

- if $M_{\max} > M_{\max}(y_0, y_f, \varepsilon)$, there exists a time $T > 0$ and a control $u \in E$ satisfying (36), steering y_0 to $\overline{B}(y_f, \varepsilon)$ in time T . If A^* satisfies the (GUC) property and $\partial_t - A^*$ is analytic-hypoelliptic, the control may be chosen to be in $\mathcal{U}_{\text{shape}}^L$.
- if $M_{\max} < M_{\max}(y_0, y_f, \varepsilon)$, no such control exists.

Moreover, for all $M_{\max} > 0$, the control system (5) is not nonnegatively approximately controllable with controls in $\{M(u) \leq M_{\max}\}$ in any time $T > 0$.

Proof. Given Corollary 5.3, the function $\overline{M}(T)$ goes to $+\infty$ as $T \rightarrow 0$, is bounded away from 0 at infinity, and does not vanish over the interval $(0, +\infty)$. Since it is continuous, we define

$$M_{\max}(y_0, y_f, \varepsilon) := \inf_{T > 0} \overline{M}(T) > 0,$$

and the first two claims follow. When A^* satisfies the (GUC) property and $\partial_t - A^*$ is analytic-hypoelliptic, the control may be chosen to be in $\mathcal{U}_{\text{shape}}^L$ by Theorem 3.1.

Then, let $M_{\max} > 0$. Taking $y_f \in L^2(\Omega)$ such that $\|y_f\|_{L^2} > \frac{\sqrt{L|\Omega|}}{|\alpha|} M_{\max} + \varepsilon$ and $y_0 \in L^2(\Omega)$ such that $y_f \geq S_T y_0$ and $\|S_T y_0 - y_f\|_{L^2} \geq \delta > 0$. Thanks to the proof of Corollary 5.3, we infer $M_{\max}(y_0, y_f, \varepsilon) \geq |\alpha| \frac{\|y_f\|_{L^2} - \varepsilon}{\sqrt{L|\Omega|}} > M_{\max}$. It follows from the second claim that y_0 cannot be steered to y_f in any time $T > 0$ with a control u such that $M(u) \leq M_{\max}$. Thus, system (5) is not nonnegatively approximately controllable with such controls in any time $T > 0$. \square

5.2.2 Characterisation of minimal time controls

Throughout this section, we let $\varepsilon > 0$, $y_f \in L^2(\Omega)$, we assume that (21) holds, and let $y_0 = 0$. Hence we must have $\|y_f\|_{L^2} > \varepsilon$ and the condition (21) is independent of T . Finally, $y_f \geq S_T y_0$ here simply amounts to $y_f \geq 0$.

Given the obstruction result of Proposition 5.4, we consider the *minimal time* control problem:

$$T^*(\lambda) = \inf\{T > 0, \exists u \in E, \|L_T u - y_f\|_{L^2} \leq \varepsilon, F_T(u) \leq \lambda\}, \quad \lambda > 0. \quad (37)$$

From our study of the optimal control problem (19), we know that this minimal time is well defined for $\lambda \in \overline{M}((0, +\infty))$. Under appropriate assumptions, we will show that it is reached, and characterise the minimal time controls, by establishing a form of equivalence between the optimal control problem and the corresponding minimal time problem. This is now a well-known feature for parabolic equations (see [19, 39, 46]).

Further study of the value function \overline{M} . Using strong duality again, we will establish that \overline{M} is a non-increasing function under the assumption that A^* satisfies the comparison principle (6). We start with the following general lemma:

Lemma 5.5. *Given any $0 < T_1 < T_2$, and $y_0 = 0$, for a general unbounded linear operator A , the dual functional defined by (23) satisfies:*

$$J_{T_1, \varepsilon}(p_f) \leq J_{T_2, \varepsilon}(p_f), \quad \forall p_f \in L^2(\Omega), \quad (38)$$

with equality if and only if

$$L_{T_2}^* p_f(t) \leq 0, \quad \forall t \in [0, T_2 - T_1]. \quad (39)$$

Proof. Since $y_0 = 0$, inequality (38) follows immediately from the comparison of the integral terms in the expression of the $J_{T_i, \varepsilon}$, $i \in \{1, 2\}$. Moreover, for $p_f \in L^2(\Omega)$, one has $J_{T_1, \varepsilon}(p_f) = J_{T_2, \varepsilon}(p_f)$ if and only if

$$\int_0^{T_1} \sigma_{\overline{u}_L} (L_{T_1}^* p_f(t)) dt = \int_0^{T_2} \sigma_{\overline{u}_L} (L_{T_2}^* p_f(t)) dt,$$

that is, by definition of the operators $L_{T_i}^*$ (see (9) which are obviously related by $L_{T_1, \varepsilon}^* p_f(t) = L_{T_2, \varepsilon}^* p_f(T_2 - T_1 + t)$ for all $t \in (0, T_1)$,

$$\int_0^{T_2 - T_1} \sigma_{\bar{u}_L} (L_{T_2}^* p_f(t)) dt = 0.$$

Using the definition of the support function $\sigma_{\bar{u}_L}$ (see the proof of Lemma 2.7), this is equivalent to (39). \square

Corollary 5.6. *The function \bar{M} (and hence Π) are non-increasing on $(0, +\infty)$.*

We now denote $\mu_- = \mu_-(y_f) := \lim_{T \rightarrow +\infty} \Pi(T) = \lim_{T \rightarrow +\infty} \frac{1}{2} \bar{M}(T)^2$. Note that $\mu_- \in [0, +\infty)$, and if the semi-group generated by A is exponentially stable, $\mu_- > 0$ as established by (35) in Corollary 5.3.

Proposition 5.7. *Assume A^* satisfies the comparison principle (6). Then, there exists $T_\ell = T_\ell(y_f) \in (0, +\infty]$ such that \bar{M} is decreasing on $[0, T_\ell)$, and constant on $[T_\ell, +\infty)$.*

Remark 5.8. *The proposition above implies in particular that \bar{M} either decreases on the whole of $(0, +\infty)$ to its limit μ_- (if $T_\ell = +\infty$), or reaches it at $T_\ell < +\infty$ and then remains constant.*

Proof. By strong duality, Lemma 5.5 implies that \bar{M} is non-increasing. Let $T_2 > T_1 > 0$, and denote $p_{T_1}^*, p_{T_2}^*$ the associated dual minimisers. Assume that

$$\bar{M}(T_1) = \bar{M}(T_2). \quad (40)$$

From Lemma 5.5, and by definition of $p_{T_1}^*$, we know that

$$J_{T_1, \varepsilon}(p_{T_1}^*) \leq J_{T_1, \varepsilon}(p_{T_2}^*) \leq J_{T_2, \varepsilon}(p_{T_2}^*). \quad (41)$$

From (32), (40) implies that $J_{T_1, \varepsilon}(p_{T_1}^*) = J_{T_2, \varepsilon}(p_{T_2}^*)$, so that all the inequalities in (41) actually are equalities.

By uniqueness of the dual optimal variable (Proposition 3.10), the first equality implies that

$$p_{T_1}^* = p_{T_2}^* =: p_f^*. \quad (42)$$

From Lemma 5.5, the second equality implies that

$$L_{T_2}^* p_f^*(t) \leq 0, \quad \forall t \in [0, T_2 - T_1]. \quad (43)$$

From (42) and (43), we get $p_T^* = p_f^*$ for all $T \in [T_1, T_2]$. Now, for $T > T_2$, the comparison principle (6) and inequality (43) imply that $L_T^* p_f^*(t) \leq 0$ for all $t \in [0, T - T_1]$. From Lemma 5.5, we then get $J_{T, \varepsilon}(p_f^*) = J_{T_1, \varepsilon}(p_f^*)$, which implies $J_{T, \varepsilon}(p_f^*) = J_{T_1, \varepsilon}(p_f^*) \leq J_{T, \varepsilon}(p_T^*)$. By definition of the dual minimiser p_T^* of $J_{T, \varepsilon}$, we also have $J_{T, \varepsilon}(p_T^*) \leq J_{T, \varepsilon}(p_f^*)$, and then finally, $J_{T, \varepsilon}(p_T^*) = J_{T, \varepsilon}(p_f^*)$, i.e., $p_T^* = p_f^*$. This implies, thanks to (32), that $\bar{M}(T) = \bar{M}(T_1) = \bar{M}(T_2)$, which proves the proposition. \square

Remark 5.9. *It follows from all the above and (43) that, when A^* satisfies the comparison principle (6), if $T_\ell < +\infty$, then*

$$L_T^* p_{T_\ell}^*(t) \leq 0, \quad \forall T \geq T_\ell, \quad \forall t \in [0, T - T_\ell],$$

and

$$u_T^*(t) = \begin{cases} 0 & \text{if } t \in (0, T - T_\ell), \\ u_{T_\ell}^*(t - T + T_\ell) & \text{if } t \in (T - T_\ell, T), \end{cases}, \quad \forall T \geq T_\ell$$

is an optimal control on $[0, T]$ whenever $u_{T_\ell}^*$ is an optimal control on $[0, T_\ell]$.

We now establish the relationship between the optimal control problem (31) and the minimal time control problem.

Proposition 5.10. *Assume that A^* satisfies the comparison principle (6). Then, for all $T \in (0, T_\ell)$, any optimal control for (31) on $[0, T]$ is a minimal time control, that is,*

$$T^*(\Pi(T)) = T.$$

Moreover, for any $\lambda > \mu_-$,

$$\Pi(T^*(\lambda)) = \lambda.$$

Proof. We proceed by contradiction. Assume that $T^*(\Pi(T)) < T$. Then, there exists $\delta > 0$ and a control $u_\delta \in L^2(0, T - \delta; L^2(\Omega))$ such that $F_T(u_\delta) \leq \Pi(T)$. Now, any optimal control u_δ^* (in the sense of optimal control problem (31) in time $T - \delta$) satisfies $F_T(u_\delta^*) \leq F_T(u_\delta)$ (the inequality is not necessarily strict, as u_δ could be an optimal control), i.e., $\Pi(T - \delta) = F_T(u_\delta^*) \leq F_T(u_\delta) \leq \Pi(T)$, which contradicts the fact that $T \mapsto \Pi(T)$ is a decreasing function on $(0, T_\ell)$. Thus, (5.10) holds.

Now, let $\lambda > \mu_-$. From Corollaries 5.3, 5.6 and Proposition 5.1, there exists $T \in (0, T_\ell)$ such that $\Pi(T) = \lambda$. Applying T^* to the above and using (5.10), we get $T^*(\lambda) = T^*(\Pi(T)) = T$. Then, applying Π to the above yields $\Pi(T^*(\lambda)) = \Pi(T) = \lambda$. \square

We can also formulate the above result in the following way: for all $\lambda > \mu_-$,

$$T^*(\lambda) = \inf\{T > 0, \quad \Pi(T) \leq \lambda\},$$

that is, T^* is the pseudo-inverse of Π on $(\mu_-, +\infty)$.

In terms of the time optimal control problem, we now have a complete characterisation of time optimal controls for (37):

Theorem 5.11. *Assume that A^* satisfies the comparison principle (6). For any $\lambda > \mu_-$, $T^*(\lambda) < +\infty$, and $T^*(\lambda) \xrightarrow{\lambda \rightarrow \infty} 0$, $T^*(\lambda) \xrightarrow{\lambda \rightarrow \mu_-} +\infty$. As a consequence, the domain of definition of T^* is $(\mu_-, +\infty)$, and on its domain of definition, T^* is continuous and decreasing.*

Moreover, if A^* satisfies the (GUC) property and $\partial_t - A^*$ is analytic-hypoelliptic, there exists a unique minimal time control for (37), given by the optimal control problem (19), and it lies in $\mathcal{U}_{\text{shape}}^L$.

Acknowledgments. The authors are grateful to Rémy Abergel for enlightening discussions about Fenchel duality. All three authors acknowledge the support of the ANR project TRECOS, grant number ANR-20-CE40-0009.

A Convex analysis

A.1 Core properties of Fenchel conjugation

A fundamental property of conjugation is involution (over $\Gamma_0(H)$):

Theorem A.1 (Fenchel-Moreau). *Given any $f \in \Gamma_0(H)$, there holds $f^* \in \Gamma_0(H)$ and $f^{**} = f$.*

Analogously to the classical gradient, the subdifferential can be used to study optimality:

Proposition A.2 (Fermat's rule). *Let $f \in \Gamma_0(H)$. f attains a finite global minimum over H in x^* if and only if*

$$0 \in \partial f(x^*).$$

We now list further useful properties of the Fenchel conjugate:

- multiplication by a real number: for $\alpha \in \mathbb{R}$,

$$(\alpha f)^*(y) = \begin{cases} \alpha f^*\left(\frac{y}{\alpha}\right) & \text{if } \alpha \neq 0, \\ \sigma_{\text{dom}(f)}(y) & \text{if } \alpha = 0. \end{cases} \quad (44)$$

- the (suitably normalised) squared norm is its own conjugate:

$$\left(\frac{1}{2}\|\cdot\|_H^2\right)^* = \frac{1}{2}\|\cdot\|_H^2. \quad (45)$$

Let us also mention a result about composition [17]. First, let $f \in \Gamma_0(H)$ and $g \in \Gamma_0(\mathbb{R})$ be non-decreasing. Then,

$$(g \circ f)^*(y) = \min_{\alpha \geq 0} \left(g^*(\alpha) + \alpha f^*\left(\frac{y}{\alpha}\right) \right).$$

Following (44), the convention for $\alpha = 0$ is $0 f^*\left(\frac{y}{0}\right) = \sigma_{\text{dom}(f)}(y)$.

Link with the subdifferential. We now give another characterisation of the subdifferential set, which illustrates the link with convex conjugation: for $f \in \Gamma_0(H)$,

$$\partial f(x) = \{p \in H, \langle p, x \rangle_H - f(x) = f^*(p)\} = \{p \in H, \langle x, p \rangle_H - f^*(p) = f(x)\} \quad (46)$$

Essentially, the subdifferential is the set of linear forms on which the convex conjugate is attained.

Using this characterisation, we then get the Legendre-Fenchel identity, which allows us to “flip” subdifferentials:

$$p \in \partial f(x) \iff x \in \partial f^*(p), \quad f \in \Gamma_0(H), \quad \forall x, p \in H.$$

A.2 Some properties of indicator and support functions

Indicator functions are a crucial tool to encode constraints in convex optimisation problems. Their properties are closely linked to topological properties of their indicated sets:

Proposition A.3. *We have $\delta_C, \sigma_C \in \Gamma_0(H)$ as soon as C is non-empty, convex and closed.*

The characterisation (46) of the subdifferential yields a useful result on indicator functions:

Proposition A.4. *Let $C \subset H$ be a closed convex set with nonempty interior. Then, for $x \in H$ we have the following:*

$$x \in \partial C \iff \partial \delta_C(x) \text{ is a nontrivial cone.}$$

Equivalently, by convex conjugation,

$$\exists p \neq 0, x \in \arg \max_{v \in C} \langle v, p \rangle \iff \exists p \neq 0, x \in \partial \sigma_C(p) \iff x \in \partial C.$$

Indicator function of a ball in a Hilbert space. Consider the closed unit ball $\overline{B}(0, 1)$ of H . We have seen before that

$$\sigma_{\overline{B}(0,1)}(y) = (\delta_{\overline{B}(0,1)})^*(y) = \|y\|_H.$$

Using (46), we get the following: for $x \in \overline{B}(0, 1)$,

$$\partial \delta_{\overline{B}(0,1)}(x) = \{p \in H, \langle p, x \rangle_H = \sigma_{\overline{B}(0,1)}(p)\} = \{p \in H, \langle p, x \rangle_H = \|p\|_H\}.$$

From the Cauchy-Schwarz inequality we know that $\langle p, x \rangle_H \leq \|p\|_H \|x\|_H$, it follows that $\langle p, x \rangle_H = \|p\|_H$ if and only if $x = \frac{p}{\|p\|_H}$. This implies that

$$\partial \delta_{\overline{B}(0,1)}(x) = \begin{cases} \{0\} & \text{if } \|x\|_H < 1, \\ \{\lambda x, \lambda \geq 0\} & \text{if } \|x\|_H = 1. \end{cases} \quad (47)$$

A.3 Technical lemmas

Lemma A.5. *Let $f \in \Gamma_0(H)$ be such that*

$$F : u \in L^2(0, T; H) \longmapsto \int_0^T f(u(t)) dt,$$

is well-defined and proper. Then $F \in \Gamma_0(L^2(0, T; H))$, and its Fenchel conjugate and subdifferential are given by

$$\forall p \in L^2(0, T; H), \quad F^*(p) = \int_0^T f^*(p(t)) dt,$$

$$\partial F(u) = \{p \in L^2(0, T; H), \quad p(t) \in \partial f(u(t)), \text{ for a.e. } t \in (0, T)\}, \quad \forall u \in L^2(0, T; H).$$

Proof. Since F is obviously convex, we only need to justify that F is lsc to infer $F \in \Gamma_0(L^2(0, T; H))$. We let $u_n \rightarrow u$ be in $L^2(0, T; H)$ and must show that $F(u) \leq \liminf F(u_n)$. Upon extraction of a subsequence, we may assume that $F(u_n) \rightarrow \liminf F(u_n)$, and that $u_n(t) \rightarrow u(t)$ in H for a.e. $t \in (0, T)$. Then, using successively the lsc of f and Fatou's lemma, we find

$$F(u) = \int_0^T f(u(t)) dt \leq \int_0^T \liminf f(u_n(t)) dt \leq \liminf \int_0^T f(u_n(t)) dt = \liminf F(u_n).$$

For $p \in L^2(0, T; H)$, we compute

$$F^*(p) = \sup_{u \in L^2(0, T; H)} \langle p, u \rangle_{L^2(0, T; H)} - \int_0^T f(u(t)) dt = \int_0^T \left(\sup_{u \in H} \langle p(t), u \rangle_H - f(u(t)) \right) dt = \int_0^T f^*(p(t)) dt.$$

Using the characterisation given in (46), and Lemma A.5, we have the following:

$$\begin{aligned} \partial F(u) &= \arg \max_{p \in L^2(0, T; H)} \{ \langle p, u \rangle - F^*(p) \} \\ &= \arg \max_{p \in L^2(0, T; H)} \left\{ \int_0^T \langle p(t), u(t) \rangle dt - \int_0^T f^*(p(t)) dt \right\} \\ &= \arg \max_{p \in L^2(0, T; H)} \left\{ \int_0^T (\langle p(t), u(t) \rangle - f^*(p(t))) dt \right\} \\ &= \left\{ p \in L^2(0, T; H), \quad p(t) \in \arg \max_{p \in H} \{ \langle p, u(t) \rangle - f^*(p) \} \right\}, \end{aligned}$$

and the result follows by the same characterisation of the subdifferential set $\partial f(u(t))$. \square

A.4 Fenchel-Rockafellar duality

Let E and F be two Hilbert spaces. Let f and g be functions in $\Gamma_0(E)$ and $\Gamma_0(F)$, respectively, and $A : E \rightarrow F$ be a bounded operator. Consider the (primal) optimisation problem

$$\pi = \inf_{x \in E} (f(x) + g(Ax)). \quad (\mathcal{C})$$

and its dual problem

$$d = \sup_{z \in F} (-f^*(A^*z) - g^*(-z)) = - \inf_{z \in F} (f^*(A^*z) + g^*(-z)) \quad (\mathcal{D})$$

With the above notations, weak duality always holds, *i.e.*, we always have $\pi \geq d$. The Fenchel-Rockafellar theorem states when and how strong duality holds, *i.e.*, when $d = \pi$ [40].

Theorem A.6. *If there exists $\bar{x} \in E$ such that g is continuous at $A\bar{x}$ and $f(\bar{x}) < +\infty$, then*

$$\pi = d \quad \text{and} \quad d \text{ is attained if finite.}$$

Symmetrically, if there exists $\bar{z} \in F$ such that f^ is continuous at $A^*\bar{z}$ and $g^*(-\bar{z}) < +\infty$, then*

$$d = \pi \quad \text{and} \quad \pi \text{ is attained if finite.}$$

The second part of the theorem is obtained by applying the first part to (\mathcal{D}) , $\inf_{z \in F} (f^*(A^*z) + g^*(-z))$, seen as a primal problem, and (\mathcal{C}) , rewritten as $\sup_{x \in E} (-f(x) - g(Ax))$, seen as its dual problem. This yields $-d \geq -\pi$, with equality under the corresponding assumptions.

Lagrangian and saddle-point interpretation. Let us now define the Lagrangian for $(x, y) \in E \times F$ by

$$\mathcal{L}(x, y) := \langle y, Ax \rangle + f(x) - g^*(y).$$

If (x^*, y^*) is a saddle point of the Lagrangian, *i.e.*,

$$x^* \in \arg \min_{x \in E} \mathcal{L}(x, y^*) \quad \text{and} \quad y^* \in \arg \max_{y \in F} \mathcal{L}(x^*, y),$$

then (x^*, z^*) (with $z^* = -y^*$) is a pair of primal and dual optimal variables, and strong duality holds.

What matters is the converse: if (x^*, z^*) is a pair of primal and dual optimal variables and if strong duality holds, then (x^*, y^*) (with $y^* = -z^*$) is a saddle point of \mathcal{L} .

Whenever (x^*, y^*) is a primal-dual optimal pair, Fermat's rule and the Legendre-Fenchel identity yield

$$x^* \in \arg \min_{x \in E} \mathcal{L}(x, y^*) \quad \iff \quad -A^*y^* \in \partial f(x^*) \quad \iff \quad x^* \in \partial f^*(-A^*y^*),$$

as well as

$$y^* \in \arg \max_{y \in F} \mathcal{L}(x^*, y) \quad \iff \quad Ax^* \in \partial g^*(y^*) \quad \iff \quad y^* \in \partial g(Ax^*),$$

Summing up, we have the following proposition:

Proposition A.7. *Let (x^*, z^*) be a pair of primal and dual optimal variables. If strong duality holds, then*

$$\begin{aligned} x^* \in \partial f^*(A^*z^*), \quad Ax^* \in \partial g^*(-z^*), \\ z^* \in -\partial g(Ax^*), \quad A^*z^* \in \partial f(x^*). \end{aligned} \tag{48}$$

Reinterpreting the Fenchel-Rockafellar theorem with the above and in a way that is useful for control-lability issues, we end up with

Proposition A.8. *Under the assumption that there exists $\bar{x} \in E$ such that g is continuous at $A\bar{x}$ and $f(\bar{x}) < +\infty$, if π is finite, and attained at $x^* \in E$, then d is attained at $z^* \in F$ satisfying*

$$z^* \in -\partial g(Ax^*), \quad A^*z^* \in \partial f(x^*). \tag{49}$$

Conversely, if (x^, z^*) satisfies (49), (x^*, z^*) is a pair of primal and dual optimal variables.*

Similarly, under the assumption that there exists $\bar{z} \in F$ such that f^ is continuous at $A^*\bar{z}$ and $g^*(-\bar{z}) < +\infty$, if d is finite, and attained at $z^* \in F$, then π is attained at $x^* \in E$ satisfying*

$$x^* \in \partial f^*(A^*z^*), \quad Ax^* \in \partial g^*(-z^*). \tag{50}$$

Conversely, if (x^, z^*) satisfies (50), (x^*, z^*) is a pair of primal and dual optimal variables.*

A.5 Parametric convex optimisation

Lemma A.9. Let H be a Hilbert space, Z be a metric space, $f : H \times Z \rightarrow \mathbb{R} \cup \{+\infty\}$. Assume that

- $\forall \alpha \in Z$, $f(\cdot, \alpha)$ is convex on H ,
- $\forall x \in H$, $f(x, \cdot)$ is continuous on Z ,
- f is sequentially weak-strong lower semicontinuous on $H \times Z$, i.e.,

$$\forall x_n \rightharpoonup x, \forall \alpha_n \rightarrow \alpha, \quad f(x, \alpha) \leq \liminf_{n \rightarrow +\infty} f(x_n, \alpha_n),$$

- there exists a unique $x_\alpha \in H$ such that $\inf_{x \in H} f(x, \alpha) = f(x_\alpha, \alpha)$.

Then the mapping

$$\alpha \in Z \mapsto \inf_{x \in H} f(x, \alpha)$$

is continuous on Z .

Proof. Let $\alpha_n \rightarrow \alpha$. Denoting $m(\alpha) = \inf_{x \in H} f(x, \alpha) = f(x_\alpha, \alpha)$, let us show that $m(\alpha_n)$ converges to $m(\alpha)$.

Upper semicontinuity. For $x \in H$ fixed, thanks to the continuity of $f(x, \cdot)$, we pass to the limit in $f(x, \alpha_n) \geq m(\alpha_n)$ and find

$$m(\alpha) = \inf_{x \in H} f(x, \alpha) \geq \limsup_{n \rightarrow +\infty} m(\alpha_n).$$

Lower semicontinuity. We denote $x_n = x_{\alpha_n}$. Let us for the moment admit that (x_n) is bounded. Upon extraction, we may assume that $x_n \rightharpoonup \bar{x}$ for some $x \in H$. By sequential weak-strong lower semicontinuity,

$$f(\bar{x}, \alpha) \leq \liminf_{n \rightarrow +\infty} f(x_n, \alpha_n) = \liminf_{n \rightarrow +\infty} m(\alpha_n).$$

Since the left-hand side is bounded from below by $m(\alpha)$, we have proved lower semicontinuity (and in fact $\bar{x} = x_\alpha$).

We are left to proving the boundedness of (x_n) to conclude the proof. Assume that (x_n) is not bounded. Upon extraction, we may assume that

$$y_n := \frac{x_n}{\|x_n\|_H} \rightharpoonup y,$$

for some $y \in H$. For any fixed $\lambda > 0$, we shall prove that $x_\alpha + \lambda y$ minimises $f(\cdot, \alpha)$, which contradicts the fourth assumption that there exists a single minimum point.

Indeed, we notice that

$$\left(1 - \frac{\lambda}{\|x_n\|_H}\right)x_\alpha + \frac{\lambda}{\|x_n\|_H}x_n \rightharpoonup x_\alpha + \lambda y.$$

Hence, by weak-strong lower semicontinuity, convexity, the fact x_n minimises $f(\cdot, \alpha_n)$ and continuity,

$$\begin{aligned} f(x_\alpha + \lambda y, \alpha) &\leq \liminf_{n \rightarrow +\infty} f\left(\left(1 - \frac{\lambda}{\|x_n\|_H}\right)x_\alpha + \frac{\lambda}{\|x_n\|_H}x_n, \alpha_n\right) \\ &\leq \liminf_{n \rightarrow +\infty} \left(1 - \frac{\lambda}{\|x_n\|_H}\right)f(x_\alpha, \alpha_n) + \frac{\lambda}{\|x_n\|_H}f(x_n, \alpha_n) \\ &\leq \liminf_{n \rightarrow +\infty} \left(1 - \frac{\lambda}{\|x_n\|_H}\right)f(x_\alpha, \alpha_n) + \frac{\lambda}{\|x_n\|_H}f(x_\alpha, \alpha_n) \\ &= \liminf_{n \rightarrow +\infty} f(x_\alpha, \alpha_n) = f(x_\alpha, \alpha). \end{aligned}$$

□

B The classical bathtub principle

The classical bathtub principle characterises the maximisers, and gives the maximum value, of the constrained scalar product maximisation:

$$\sup_{u \in \tilde{\mathcal{U}}_L^*} \int_{\Omega} u(x)v(x) dx, \quad (51)$$

where $v \in L^2(\Omega)$ is arbitrary, and

$$\tilde{\mathcal{U}}_L^* := \left\{ u \in L^2(\Omega), 0 \leq u \leq 1 \text{ and } \int_{\Omega} u = L|\Omega| \right\},$$

is the convex hull (and L^∞ weak-* closure) of the set of characteristic functions whose support has the corresponding fixed measure:

$$\tilde{\mathcal{U}}_L := \{ \chi_{\omega}, \quad \omega \subset \Omega, \quad |\omega| = L|\Omega| \}.$$

Recalling the notations (14) and (15) introduced in Section 2.3, the classical bathtub principle reads:

Lemma B.1 (classical bathtub principle). *Let $v \in L^2(\Omega)$. Denote $\rho(v) := \Phi_v^{-1}(L|\Omega|)$. The maximum in (51) equals*

$$\left(\int_{v > \rho(v)} v \right) + \rho(v)(L|\Omega| - |\{v > \rho(v)\}|) = \int_0^{L|\Omega|} \Phi_v^{-1},$$

and the maximisers are given by

$$u^* := \chi_{\{v > \rho(v)\}} + c(x)\chi_{\{v = \rho(v)\}},$$

where c is any measurable function such that $0 \leq c \leq 1$ and

$$\int_{\{v = \rho(v)\}} c = L|\Omega| - |\{v > \rho(v)\}|.$$

In particular, if all the level sets of the function v have zero measure, then the maximum is uniquely attained by

$$u^* := \chi_{\{v > \rho(v)\}},$$

and the maximum hence equals $\int_{\{v > \rho(v)\}} v$.

Now, recall that we defined \mathcal{U}_L by (11) and its convex hull $\bar{\mathcal{U}}_L$ by (12) in Section 2.3. They are respective relaxations of $\tilde{\mathcal{U}}_L$, and its convex hull $\tilde{\mathcal{U}}_L^*$.

For $v \in L^2(\Omega)$, we consider the relaxed version of (51):

$$\sup_{u \in \bar{\mathcal{U}}_L} \int_{\Omega} u(x)v(x) dx. \quad (52)$$

Then, the complete solution of Lemma 2.3 is given by the following:

Lemma B.2 (relaxed bathtub principle). *Let $v \in L^2(\Omega)$. Denote $h(v) = \max(0, \Phi_v^{-1}(L|\Omega|)) = \max(0, \rho(v))$. Then, the maximum in (52) equals*

$$\int_0^{\min(\Phi_v(0), L|\Omega|)} \Phi_v^{-1},$$

and the maximisers are given by

$$u^* := \chi_{\{v > h(v)\}} + c(x)\chi_{\{v = h(v)\}},$$

where c is any measurable function such that $0 \leq c \leq 1$ and

$$\begin{cases} \int_{\{v=h(v)\}} c = L|\Omega| - |\{v > h(v)\}| & \text{if } h(v) > 0 \\ \int_{\{v=h(v)\}} c \leq L|\Omega| - |\{v > h(v)\}| & \text{if } h(v) = 0 \end{cases}$$

Remark B.3. In particular, if $h(v) > 0$, there is a unique maximiser in (52) if and only if

$$|\{v > h(v)\}| + |\{v = h(v)\}| = L|\Omega|.$$

Indeed, when the above does not hold, c can be chosen to have values in $(0, 1)$ on a set of nonzero measure, and the maximisers is no longer unique.

On the other hand, if $h(v) = 0$, as soon as $|\{v = h(v)\}| > 0$ the maximisers are no longer unique.

References

- [1] N. U. Ahmed. Finite-time null controllability for a class of linear evolution equations on a Banach space with control constraints. *Journal of optimization Theory and Applications*, 47(2):129–158, 1985. Publisher: Springer.
- [2] Nachman Aronszajn. A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order. Technical report, KANSAS UNIV LAWRENCE, 1956.
- [3] Larbi Berrahmoune. A variational approach to constrained controllability for distributed systems. *Journal of Mathematical Analysis and Applications*, 416(2):805–823, 2014.
- [4] Larbi Berrahmoune. Constrained null controllability for distributed systems and applications to hyperbolic-like equations. *ESAIM: Control, Optimisation and Calculus of Variations*, 25:32, 2019.
- [5] Larbi Berrahmoune. A variational approach to constrained null controllability for the heat equation. *European Journal of Control*, 52:42 – 48, 2020.
- [6] Jean-Michel Bony. Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés. *Annales de l’Institut Fourier*, 19(1):277–304, 1969.
- [7] Franck Boyer. On the penalised HUM approach and its applications to the numerical approximation of null-controls for parabolic problems. *ESAIM: Proc.*, 41:15–58, 2013.
- [8] Robert F Brammer. Controllability in linear autonomous systems with positive controllers. *SIAM Journal on Control*, 10(2):339–353, 1972.
- [9] Haïm Brézis. *Functional analysis, Sobolev spaces and partial differential equations*, volume 2. Springer, 2011.
- [10] Antonin Chambolle and Thomas Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of Mathematical Imaging and Vision*, 40, 05 2011.
- [11] Frank H Clarke. *Optimization and nonsmooth analysis*. SIAM, 1990.
- [12] J.-M. Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [13] O Yu Emanuilov. Controllability of parabolic equations. *Sbornik: Mathematics*, 186(6):879–900, jun 1995.

- [14] Klaus-Jochen Engel and Rainer Nagel. One-parameter semigroups for linear evolution equations. *Semigroup Forum*, 63:278–280, 06 2001.
- [15] Sylvain Ervedoza. Control issues and linear projection constraints on the control and on the controlled trajectory. *North-W. Eur. J. of Math.*, 6:165–197, 2020.
- [16] Lawrence C Evans. Partial differential equations. *Graduate studies in mathematics*, 19(4):7, 1998.
- [17] Jean-Baptiste Hiriart-Urruty. A note on the Legendre-Fenchel transform of convex composite functions. In *Nonsmooth Mechanics and Analysis*, pages 35–46. Springer, 2006.
- [18] Bernhard Kawohl. *Rearrangements and convexity of level sets in PDE*, volume 1150. Springer, 2006.
- [19] Karl Kunisch and Lijuan Wang. Time optimal control of the heat equation with pointwise control constraints. *ESAIM: Control, Optimisation and Calculus of Variations*, 19(2):460–485, 2013.
- [20] Stéphane Labbé and Emmanuel Trélat. Uniform controllability of semidiscrete approximations of parabolic control systems. *Systems & Control Letters*, 55(7):597–609, 2006.
- [21] Gontran Lance, Emmanuel Trélat, and Enrique Zuazua. Shape turnpike for linear parabolic PDE models. *Systems & Control Letters*, 142:104733, 2020.
- [22] Gilles Lebeau and Luc Robbiano. Contrôle exact de l'équation de la chaleur. *Communications in Partial Differential Equations*, 20(1-2):335–356, 1995.
- [23] Elliott H. Lieb and Michael Loss. *Analysis*. Crm Proceedings & Lecture Notes. American Mathematical Society, 2001.
- [24] Jacques-Louis Lions. Remarks on approximate controllability. *Journal d'Analyse Mathématique*, 59(1):103, 1992.
- [25] Jérôme Lohéac, Emmanuel Trélat, and Enrique Zuazua. Minimal controllability time for the heat equation under unilateral state or control constraints. *Mathematical Models and Methods in Applied Sciences*, 27(09):1587–1644, 2017.
- [26] Jérôme Lohéac, Emmanuel Trélat, and Enrique Zuazua. Minimal controllability time for finite-dimensional control systems under state constraints. *Automatica*, 96:380–392, 2018.
- [27] Jérôme Lohéac, Emmanuel Trélat, and Enrique Zuazua. Nonnegative control of finite-dimensional linear systems. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 38(2):301–346, 2021.
- [28] Idriss Mazari. Quantitative estimates for parabolic optimal control problems under L^∞ and L^1 constraints in the ball: Quantifying parabolic isoperimetric inequalities. *Nonlinear Analysis*, 215:112649, 2022.
- [29] Sorin Micu, Ionel Roventa, and Marius Tucsnak. Time optimal boundary controls for the heat equation. *Journal of Functional Analysis*, 263(1):25–49, 2012. Publisher: Elsevier.
- [30] Sorin Micu and Enrique Zuazua. An introduction to the controllability of partial differential equations. *Quelques questions de théorie du contrôle. Sari, T., ed., Collection Travaux en Cours Hermann, to appear*, 2004.
- [31] Victor J Mizel and Thomas I Seidman. An abstract bang-bang principle and time-optimal boundary control of the heat equation. *SIAM Journal on Control and Optimization*, 35(4):1204–1216, 1997.
- [32] G. Métivier. Hypocoellipticité d'opérateurs à caractéristiques multiples. *Séminaire Équations aux dérivées partielles (Polytechnique)*, pages 1–13, 1980-1981.
- [33] Edward Nelson. Analytic vectors. *Annals of Mathematics*, pages 572–615, 1959.

- [34] Khoa Nguyen Son. A unified approach to constrained approximate controllability for the heat equations and the retarded equations. *Journal of Mathematical Analysis and Applications*, 150(1):1–19, 1990. Publisher: Elsevier.
- [35] Amnon Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44. Springer Science & Business Media, 2012.
- [36] Dario Pighin and Enrique Zuazua. Controllability under positivity constraints of semilinear heat equations. *Mathematical Control & Related Fields*, 8(3&4):935–964, 2018.
- [37] Camille Pouchol, Emmanuel Trélat, and Enrique Zuazua. Phase portrait control for 1D monostable and bistable reaction–diffusion equations. *Nonlinearity*, 32(3):884, 2019.
- [38] Yannick Privat, Emmanuel Trélat, and Enrique Zuazua. Optimal shape and location of sensors for parabolic equations with random initial data. *Archive for Rational Mechanics and Analysis*, 216(3):921–981, 2015. Publisher: Springer.
- [39] Shulin Qin and Gengsheng Wang. Equivalence between minimal time and minimal norm control problems for the heat equation. *SIAM Journal on Control and Optimization*, 56(2):981–1010, 2018.
- [40] Ralph Rockafellar. Duality and stability in extremum problems involving convex functions. *Pacific Journal of Mathematics*, 21(1):167–187, 1967.
- [41] Domènec Ruiz-Balet and Enrique Zuazua. Control under constraints for multi-dimensional reaction-diffusion monostable and bistable equations. *Journal de Mathématiques Pures et Appliquées*, 143:345–375, 2020.
- [42] David L. Russell. Controllability and stabilizability theory for linear partial differential equations: Recent progress and open questions. *SIAM Review*, 20(4):639–739, 1978.
- [43] E.J.P. Georg Schmidt. The “bang-bang” principle for the time-optimal problem in boundary control of the heat equation. *SIAM Journal on Control and Optimization*, 18(2):101–107, 1980.
- [44] Emmanuel Trélat, Gengsheng Wang, and Yashan Xu. Characterization by observability inequalities of controllability and stabilization properties. *Pure and Applied Analysis*, 2(1):93–122, 2020.
- [45] Gengsheng Wang, Yashan Xu, and Yubiao Zhang. Attainable Subspaces and the Bang-Bang Property of Time Optimal Controls for Heat Equations. *SIAM Journal on Control and Optimization*, 53(2):592–621, 2015.
- [46] Gengsheng Wang and Enrique Zuazua. On the Equivalence of Minimal Time and Minimal Norm Controls for Internally Controlled Heat Equations. *SIAM Journal on Control and Optimization*, 50(5):2938–2958, 2012.
- [47] Dong-Hui Yang, Bao-Zhu Guo, Weihua Gui, and Chunhua Yang. The Bang–Bang Property of Time-Varying Optimal Time Control for Null Controllable Heat Equation. *Journal of Optimization Theory and Applications*, 182(2):588–605, 2019. Publisher: Springer.