



HAL
open science

Mediterranean red coral as a spawning ground for cuttlefish

Federica Maggioni, Bramanti Lorenzo

► **To cite this version:**

Federica Maggioni, Bramanti Lorenzo. Mediterranean red coral as a spawning ground for cuttlefish. Global Journal of Ecology, 2022, 7 (2), pp.120-121. 10.17352/gje.000071 . hal-03892998

HAL Id: hal-03892998

<https://hal.science/hal-03892998v1>

Submitted on 6 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Opinion

Mediterranean red coral as a spawning ground for cuttlefish

Federica Maggioni^{1,2} and Lorenzo Bramanti^{2*}

¹ENTROPIE, IRD, University of Reunion, CNRS, IFREMER, University of New Caledonia, Nouméa 98800, New Caledonia

²CNRS-Sorbonne University, Laboratory of Ecogeochemistry of Benthic Environments, LECOB, Oceanological Observatory of Banyuls sur Mer, 1 avenue Pierre Fabre – 66650, Banyuls sur Mer, France

Received: 21 November, 2022
Accepted: 24 November, 2022
Published: 25 November, 2022

*Corresponding author: Lorenzo Bramanti, CNRS-Sorbonne University, Laboratory of Ecogeochemistry of Benthic Environments, LECOB, Oceanological Observatory of Banyuls sur Mer, 1 avenue Pierre Fabre – 66650, Banyuls sur Mer, France, Tel: +33781927635; E-mail: bramanti@obs-banyuls.fr

ORCID: <https://orcid.org/0000-0002-4872-840X>

Copyright License: © 2022 Maggioni F, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<https://www.peertechzpublications.com>



The Mediterranean red coral (*Corallium rubrum*) is a long-lived and slow-growing gorgonian, which suffered intense exploitation due to its precious calcium carbonate skeleton used in jewelry and traded worldwide [1].

Due to its commercial and ecological value, knowledge of the biology of the species has increased noticeably during the last decades as a support to management; however, evidence of its ecological role is scarce [2]. Demographic studies revealed a shift of *C. rubrum* population structures towards small-sized colonies and raised concern about the risk of the loss of the ecological function of the species [2].

Similarly to other gorgonians, *C. rubrum* can form dense forests [3] which increase structural complexity and serve as feeding, shelter, and foraging grounds for many associated organisms. Furthermore, coral forests provide favorable conditions for the egg deposition of several species [4].

In the framework of a transplanting experiment near the Marine Protected Area of Cérbere-Banyuls (North Western Mediterranean Sea, France), we observed, at the end of May 2019, one egg mass deposition of the European cuttlefish (*Sepia officinalis*) on a transplanted healthy *C. rubrum* colony at 30 m depth (Figure 1). We report some characteristics of the eggs: the diameter of the 9 eggs was comprised between 1.5 and 2 cm, and they presented a swollen and fuller appearance, suggesting that they were at a late stage of embryological development. Moreover, the egg mass contained empty egg capsules indicating that some cuttlefish already hatched (Figure 1) and then suggesting that the *C. rubrum* colony was a viable structure for their embryological development.




Figure 1: Cuttlefish egg deposition on a *C. rubrum* colony transplanted at 30 m depth in Banyuls sur mer, France. a: distant view. b: close up on the eggs (photo by B. Hesse).

Cephalopods play a key role in marine food webs, and most of them, such as *S. officinalis* in the Mediterranean and particularly in the area in which the study was performed, are important fishery resources. Cuttlefish egg depositions have been observed on different structures including gorgonians and are not common at depths >25 m [5]. Our observation increased our knowledge of this complex ecological interaction that is difficult to test and demonstrate. In addition, functional linkage among relevant species is a crucial step toward the development of conservation and management strategies. Our observation highlights the importance of Mediterranean red coral habitats as a spawning ground and functional repository for recruits of commercially important species and suggests that small-sized *C. rubrum* colonies may still fulfill some ecological functions. Those results could help re-evaluate the ecological ecosystem services provided by this precious



octocoral, which have been underestimated with respect to the economical ones.

Ethical approval

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed by the authors.

Sampling and field studies

All necessary permits for sampling and observational field studies have been obtained by the authors from the competent authorities.

Data availability

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

1. Tsounis G, Grigg R, Gili J. The exploitation and conservation of precious corals. *Oceanogr Mar Biol An Annu Rev* 48: 161–212.
2. Bramanti L, Iannelli M, Santangelo G. Mathematical modeling for conservation and management of gorgonians corals: young and olds, could they coexist? *Ecol. Modell.* 2009; 220: 2851–2856.
3. Rossi S, Bramanti L, Gori A, Orejas C. An Overview of the Animal Forests of the World. In book: *Marine animal forests*. 1-26.
4. Cau A, Follesa MC, Moccia D, Bellodi A, Mulas A, Bo M, Canese S, Angiolillo M, Cannas R. *Leiopathes glaberrima* millennial forest from SW Sardinia as a nursery ground for the small-spotted catshark *Scyliorhinus canicula*. *Aquatic conservation-Marine freshwater ecosystems*. 2017; 27:731–735.
5. Guerra A, Hernandez-Urcera J, Garci ME, Sestelo M, Regueira M, Gilcoto M, Gonzalez AF. Spawning habitat selection by the common cuttlefish *Sepia officinalis* in the Cíes Islands (Northwest Spain). *Fishery Research*. 2016; 183: 44-54.

Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

Highlights

- ❖ Signatory publisher of ORCID
- ❖ Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- ❖ Articles archived in world's renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- ❖ Journals indexed in ICMJE, SHERPA/ROMEO, Google Scholar etc.
- ❖ OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- ❖ Dedicated Editorial Board for every journal
- ❖ Accurate and rapid peer-review process
- ❖ Increased citations of published articles through promotions
- ❖ Reduced timeline for article publication

Submit your articles and experience a new surge in publication services (<https://www.peertechz.com/submission>).

Peertechz journals wishes everlasting success in your every endeavours.