



HAL
open science

Minimizing Movements for Anisotropic and Inhomogeneous Mean Curvature Flows

Antonin Chambolle, Daniele de Gennaro, Massimiliano Morini

► **To cite this version:**

Antonin Chambolle, Daniele de Gennaro, Massimiliano Morini. Minimizing Movements for Anisotropic and Inhomogeneous Mean Curvature Flows. *Advances in Calculus of Variation*, 2023, 10.1515/acv-2022-0102 . hal-03894146v1

HAL Id: hal-03894146

<https://hal.science/hal-03894146v1>

Submitted on 12 Dec 2022 (v1), last revised 6 Dec 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Minimizing Movements for Anisotropic and Inhomogeneous Mean Curvature Flows

Antonin Chambolle, Daniele De Gennaro, Massimiliano Morini

Abstract

In this paper we address anisotropic and inhomogeneous mean curvature flows with forcing and mobility, and show that the minimizing movements scheme converges to level set/viscosity solutions and to distributional solutions *à la* Luckhaus-Sturzenhecker to such flows, the latter result holding in low dimension and conditionally to the convergence of the energies. By doing so we generalize recent works concerning the evolution by mean curvature by removing the hypothesis of translation invariance, which in the classical theory allows one to simplify many arguments.

1 Introduction

In this paper we deal with the anisotropic, inhomogeneous mean curvature flow with forcing and mobility. By inhomogeneous we mean that the flow is driven by surface tensions depending on the position in addition to the orientation of the surface. The evolution of sets $t \mapsto E_t \subseteq \mathbb{R}^N$ considered is (formally) governed by the law

$$V(x, t) = \psi(x, \nu_{E_t}(x)) \left(-H_{E_t}^\phi(x) + f(x, t) \right), \quad x \in \partial E_t, \quad t \in (0, T), \quad (1.1)$$

smooth law

where $V(x, t)$ is the (outer) normal velocity of the boundary ∂E_t at x , $\phi(x, p)$ is a given anisotropy representing the *surface tension*, H^ϕ is the *anisotropic mean curvature* of ∂E_t associated to ϕ , $\psi(x, p)$ is an anisotropy evaluated at the outer unit normal $\nu_{E_t}(x)$ to ∂E_t which represents a *velocity* modifier (also called the *mobility* term), and f is the *forcing* term. We will be mainly concerned with smooth anisotropies (and the regularity assumptions will be made precise later on): in this case, the curvature H^ϕ is the first variation of the anisotropic and inhomogeneous perimeter associated to the anisotropy ϕ (in short, ϕ -perimeter) defined as

$$P_\phi(E) := \int_{\partial^* E} \phi(x, \nu_E(x)) \, d\mathcal{H}^{N-1}(x) \quad (1.2)$$

for any set E of finite perimeter (where $\partial^* E$ denotes the reduced boundary of E) and, if E is sufficiently smooth, it takes the form

$$H_E^\phi(x) = \operatorname{div}(\nabla_p \phi(x, \nu_E(x))),$$

where with ∇_p we denote the gradient made with respect to the second variable. Note that evolution (1.1) can be read as the motion of sets in \mathbb{R}^N , when the latter is endowed with the Finsler

metric induced by the anisotropy (see Remark 4.14). Equation (1.1) is relevant in Material Sciences, Crystal Growth, Image Segmentation, Geometry Processing and other fields see e.g. [1, 20, 26, 37, 38].

The mathematical literature for inhomogeneous mean curvature flows is not as extensive as in the homogeneous case, mainly due to the difficulties arising from the lack of translational invariance. Indeed, assuming that the evolution is invariant under translations allows to simplify many arguments used in the classical proofs of, for example, comparison results and estimates on the speed of evolution. In the homogeneous case the well-posedness theory is nowadays well established and quite satisfactory, both in the local and nonlocal case, and even in the much more challenging crystalline case (that is, when the anisotropy ϕ is piecewise affine) see [2, 3, 9, 12, 13, 14, 17, 25, 32, 33, 35] to cite a few. Concerning the inhomogeneous mean curvature flow, we cite [27, 28] where the short time existence of smooth solutions on manifolds is shown, and [24, 30], where the viscosity level set approach (introduced for the homogeneous evolution in [17, 22]) is extended, respectively, to the equation (1.1) and to the Riemannian setting.

In the present work we implement the minimising movement approach à la Almgren-Taylor-Wang [3] to prove existence via approximation of a level set solution to the generalized anisotropic and inhomogeneous motion (1.1). To carry on this scheme (which has only been sketched in [9], but lacks a formal proof) we gain insights from [14]. We also show that, under the additional hypothesis of convergence of the energies (1.6) and low dimension (2.6) (which are nowadays classical for this approach), the same approximate solutions provide in the limit a suitable notion of “BV-solutions”, also termed distributional solutions, see [33, 35].

The main limitations of this work are the following. To begin with, the new arguments which are used to compensate the lack of translation invariance are based on the locality of the anisotropic curvature H^ϕ associated with a smooth anisotropy ϕ . This implies that the proofs are not straightforwardly adaptable to the so-called “variational curvatures” considered in [14], which are non-local in nature. On the other hand, since the crystalline curvatures are highly nonlocal and degenerate operators (see e.g. [12, 15]), they do not fall in the theory constructed in the present work. In principle, it would be possible to follow the same perturbative study conducted in [12] in order to prove at least existence for an inhomogeneous and crystalline mean curvature flow. However, a satisfactory characterization of the limiting motion equation bearing a comparison principle is lacking so far.

This work can be seen as a first step towards constructing a general theory of motions driven by non translational invariant and possibly nonlocal curvatures, in the spirit of [14].

1.1 Main results

Now briefly recall the minimizing movements procedure in order to state the main results of the paper. Given an initial bounded set E_0 and a parameter $h > 0$, we define the *discrete flow* $E_t^{(h)} := T_{h,t-h} E_{t-h}^{(h)}$ for any $t \geq h$ and $E_t^{(h)} = E_0$ for $t \in [0, h)$, where the functional $T_{h,t}$ is defined for $t \geq 0$ as follows: for any bounded set E we set $T_{h,t} E$ (or, sometimes, $T_{h,t}^- E$) as the minimal solution to the problem

$$\min \left\{ P_\phi(F) + \int_F \left(\frac{\text{sd}_E^\psi(x)}{h} + \int_{[\frac{t}{h}]h}^{[\frac{t}{h}]h+h} f(x, s) ds \right) d\mathcal{H}^{N-1}(x) : F \text{ is measurable} \right\}, \quad (1.3)$$

where $\text{sd}_E^\psi(x)$ is the signed geodesic distance between x and E induced by the anisotropy ψ (see (2.2) for the precise definition) and $[s] = \max\{n \leq s, n \in \mathbb{N} \cup \{0\}\}$ denotes the integer part of

a non-negative real number $s \in [0, +\infty)$. We will then define $T_{h,t}^+$ as the maximal solution to the problem above. Any L^1 -limit point as $h \rightarrow 0$ of the family $\{E_t^{(h)}\}_{t \geq 0}$ will be called a *flat flow*. In the whole paper we will assume that it holds

$$\begin{aligned} &\phi \in \mathcal{L} \text{ (see Definition 2.2) and } \psi \text{ is an anisotropy as in Definition 2.1,} \\ &\forall x \in \mathbb{R}^N, t \in [0, +\infty) \text{ it holds } f(\cdot, t) \in C^0(\mathbb{R}^N), \|f\|_{L^\infty(\mathbb{R}^N \times [0, +\infty))} < \infty. \end{aligned} \quad (\text{H})$$

standing hp

With more effort one could weaken the hypothesis and require $\int_0^t f(\cdot, s) ds$ to be continuous (see [16]). For the sake of simplicity we will require the global-in-time boundedness. We prove existence and Holder regularity for flat flows.

istence flat flow

Theorem 1.1 (Existence of flat flows). *Assume (H). Let E_0 be a bounded set of finite perimeter and ϕ, ψ, f satisfy the hypothesis above. For any $h > 0$, let $\{E_t^{(h)}\}_{t \geq 0}$ be a discrete flow with initial datum E_0 . Then, there exists a family of sets of finite perimeter $\{E_t\}_{t \geq 0}$ and a subsequence $h_k \searrow 0$ such that*

$$E_t^{(h)} \rightarrow E_t \quad \text{in } L^1,$$

for a.e. $t \in [0, +\infty)$. Such flow satisfies the following regularity property: for any $T > 0$ and for every $0 \leq s \leq t < T$,

$$\begin{aligned} |E_s \Delta E_t| &\leq c|t - s|^{1/2}, \\ P_\phi(E_t) &\leq P_\phi(E_0) + c, \end{aligned}$$

where the constant c depends on T also.

Subsequently, we will show that the flat flow are distributional solutions, as defined in [33]. We will require additional hypothesis: firstly, low dimension (2.6) (linked to the complete regularity of the ϕ -perimeter minimizer, compare [33, 35]), moreover

$$\exists c_\psi > 0 \text{ s.t. } |\psi(x, v) - \psi(y, v)| \leq c_\psi |x - y|, \quad \forall x, y \in \mathbb{R}^N, v \in S^N, \quad (1.4)$$

hp lip psi 1

$$f \in C^0(\mathbb{R}^N \times [0, \infty)). \quad (1.5)$$

hp f 1

ditional solutions

Theorem 1.2 (Existence of distributional solutions). *Assume (H), (1.4), (1.5) and (2.6). For any $T > 0$, if*

$$\lim_{k \rightarrow \infty} \int_0^T P_\phi(E_t^{(h_k)}) = \int_0^T P_\phi(E_t), \quad (1.6)$$

hp per

then $\{E_t\}_{t \geq 0}$ is a distributional solution (1.1) with initial datum E_0 in the following sense:

(1) for a.e. $t \in [0, T)$ the set E_t has weak ϕ -curvature $H_{E_t}^\phi$ (see (2.11) for details) satisfying

$$\int_0^T \int_{\partial^* E_t} |H_{E_t}^\phi|^2 < \infty;$$

(2) there exist $v : \mathbb{R}^N \times (0, T) \rightarrow \mathbb{R}$ with $\int_0^T \int_{\partial^* E_t} v^2 d\mathcal{H}^{N-1} dt < \infty$ and $v(\cdot, t)|_{\partial E_t} \in L^2(\partial E_t)$ for a.e. $t \in [0, T)$, such that

$$-\int_0^T \int_{\partial^* E_t} v \eta d\mathcal{H}^{N-1} dt = \int_0^T \int_{\partial^* E_t} (H_{E_t}^\phi - f) \eta d\mathcal{H}^{N-1} dt \quad (1.7)$$

legge curvatura

$$\int_0^T \int_{E_t} \partial_t \eta \, dx \, dt + \int_{E_0} \eta(\cdot, 0) \, dx = - \int_0^T \int_{\partial^* E_t} \psi(\nu_{E_t}) v \eta \, d\mathcal{H}^{N-1} \, dt. \quad (1.8)$$

legge velocita

for every $\eta \in C_c^1(\mathbb{R}^N \times [0, +\infty))$.

The definitions 1), 2) extend to our case the the definition of *BV*-solutions of [33] and the distributional solutions of [35]. We recall that hypotheses (1.6) ensures that the evolving sets avoid the so-called ‘‘fattening’’ phenomenon. It is known that this hypothesis is satisfied in the case of evolution of convex or mean-convex sets, see e.g. [15, 18], but in general is not known under which general hypothesis it is valid. We also remark that the proof of the theorem above provides a rigorous proof of [15, Theorem 3.2], which had only been sketched in that paper and was invoking a Bernstein type argument which might not be known in that setting, and which we bypass using regularity results and a double blow-up argument.

In the second part of the work we will focus on the level set approach. Briefly, given an initial compact set E_0 , we set u_0 such that $\{u_0 \geq 0\} = E_0$ and we look for a solution u in the viscosity sense (in a sense made precise in Definition 4.6) to

$$\begin{cases} \partial_t u + \psi(x, -\nabla u) (\operatorname{div} \nabla_p \phi(x, \nabla u(x)) - f(x, t)) = 0 \\ u(\cdot, t) = u_0. \end{cases} \quad (1.9)$$

intro cauchy

Classical remarks ensure that any level set $\{u \geq s\}$ is evolving following the mean curvature flow (1.1). To prove existence for (1.9) we use an approximating procedure. For $h > 0$ and $t \in (0, +\infty)$ we set iteratively $u_h^\pm(\cdot, t) = u_0$ for $t \in [0, h)$ and for $t \geq h$

$$\begin{aligned} u_h^+(x, t) &:= \sup \left\{ s \in \mathbb{R} : x \in T_{h, t-h}^+ \{u_h^+(\cdot, t-h) \geq s\} \right\} \\ u_h^-(x, t) &:= \sup \left\{ s \in \mathbb{R} : x \in T_{h, t-h}^- \{u_h^-(\cdot, t-h) > s\} \right\}, \end{aligned}$$

where the operator $T_{h,t}^\pm$ has been previously introduced. We remark that these are maps piecewise constant in time, since $T_{h,t}^\pm = T_{h, \lfloor t/h \rfloor h}^\pm$, which are only upper and lower semicontinuous in space respectively. Then, we will pass to the limit $h \rightarrow 0$ on the families $\{u_h^\pm\}_h$ to find functions u^+, u^- which are viscosity sub and supersolution respectively of equation (1.9). Passing to the limit as $h \rightarrow 0$ in our case is not straightforward. The main issue is that we do not have an uniform estimate on the modulus of continuity of the functions u_h (compare [14]) and thus we can not pass to the (locally) uniform limit of the sequence. (More precisely, our best estimate contained in Lemma 4.9 decays too fast as $h \rightarrow 0$ to provide any useful information). Nonetheless, motivated by [6, 7, 8] we can define the half-relaxed limits

$$\begin{aligned} u^+(x, t) &:= \sup_{(x_h, t_h) \rightarrow (x, t)} \limsup_{h \rightarrow 0} u_h^+(x_h, t_h) \\ u^-(x, t) &:= \inf_{(x_h, t_h) \rightarrow (x, t)} \liminf_{h \rightarrow 0} u_h^-(x_h, t_h), \end{aligned} \quad (1.10)$$

def sottosoluzion

and prove that the functions defined above are sub and supersolutions, respectively, to (1.9). The main difficulty in this regard is that we need to work with just semicontinuous functions in space, as in the translationally invariant setting one can easily prove the uniform equicontinuity of the approximating sequence. We prove the following.

teo sol viscosa

Theorem 1.3. Assume (H), (1.4) and $f \in C^0(\mathbb{R}^n \times [0, +\infty))$. The function u^+ (respectively u^-) defined in (1.10) is a viscosity subsolution (respectively a viscosity supersolution) of (1.9).

Thanks to the results of [17] we then prove that, under the additional hypothesis

$$\begin{aligned} \nabla_x \nabla_p \phi(\cdot, p) \text{ and } \nabla_p^2 \phi(\cdot, p) \text{ are Lipschitz, uniformly for } p \in S^N \\ \nabla_p^2 \phi^2 \text{ is uniformly elliptic (apart from the degenerate direction } p) \\ \psi(\cdot, p) \text{ Lipschitz continuous, uniformly in } p \\ f(\cdot, t) \text{ Lipschitz continuous, uniformly in } t, \end{aligned} \tag{1.11} \quad \text{hp 2}$$

the following uniqueness result holds.

viscosity solutions

Theorem 1.4. Assume (H) and (1.11). If u_0 is a continuous function which is spatially constant outside a compact set, equation (1.9) with initial condition u_0 admits a unique continuous viscosity solution u given by (1.10). In particular, $u^+ = u^- = u$ is the unique continuous viscosity solution to (1.9) and $u_h^\pm \rightarrow u$ as $h \rightarrow 0$, locally uniformly.

The previous result yields a proof of consistency between the level set approach and the minimizing movements one to study the evolution (1.1). We recall that it has been established for the classical mean curvature flow in [11], in the anisotropic but homogeneous case in [21] and in a very general nonlocal setting in [14].

2 Preliminaries

We start introducing some notations. We consider $0 \in \mathbb{N}$. We will use both $B_r(x)$ and $B(x, r)$ to denote the Euclidean ball in \mathbb{R}^N centered in x and of radius r ; with $B_r^{N-1}(x)$ we denote the Euclidean ball in \mathbb{R}^{N-1} centered in x and of radius r ; with S^N we denote the sphere $\partial B_1(0) \subseteq \mathbb{R}^N$. In the following, we will always speak about measurable sets and refer to a set as the union of all the points of density 1 of that set i.e. $E = E^{(1)}$. Moreover, if not otherwise stated, we implicitly assume that the function spaces considered are defined on \mathbb{R}^N , e.g. $L^\infty = L^\infty(\mathbb{R}^N)$. Moreover, we often drop the measure with respect to which we are integrating, if clear from the context.

def anisotropy

Definition 2.1. We define *anisotropy* (sometimes defined as an *elliptic integrand*) a function ψ with the following properties: $\psi(x, p) : \mathbb{R}^N \times \mathbb{R}^N \rightarrow [0, +\infty)$ is a continuous function, which is convex and positively 1-homogeneous in the second variable, such that

$$\frac{1}{c_\psi} |p| \leq \psi(x, p) \leq c_\psi |p|$$

for any point $x \in \mathbb{R}^N$ and vector $p \in \mathbb{R}^N$.

We remark that, as standard, we define a real function f positively 1-homogeneous if for any $\lambda \geq 0$, it holds $f(\lambda x) = \lambda f(x)$. In particular, the anisotropies that we will consider are not symmetric. In the following, we will always denote the gradient of an anisotropy with respect to the first (respectively second) variable as $\nabla_x \psi$ (respectively $\nabla_p \psi$). We then recall the definition of some well-known quantities (see [9]). Define the polar function of an anisotropy ψ , denoted with ψ° , as

$$\psi^\circ(\cdot, \xi) := \sup_{p \in \mathbb{R}^N} \{\xi \cdot p : \psi(\cdot, p) \leq 1\}. \tag{2.1} \quad \text{CS type ineq}$$

Using the definition it is easy to see that for all $p, \xi \in \mathbb{R}^N$ it holds

$$\psi(\cdot, p)\psi^\circ(\cdot, \xi) \geq p \cdot \xi, \quad -\psi(\cdot, -p)\psi^\circ(\cdot, \xi) \leq p \cdot \xi.$$

Furthermore, one can prove that (see [9])

$$\psi^\circ(\nabla_p \psi) = 1, \quad \psi(\nabla_p \psi^\circ) = 1, \quad (\psi^\circ)^\circ = \psi.$$

We define for any $x, y \in \mathbb{R}^N$ the geodesic distance induced by ψ , or ψ -distance in short, as

$$\text{dist}^\psi(x, y) := \inf \left\{ \int_0^1 \psi^\circ(\gamma(t), \dot{\gamma}(t)) dt : \gamma \in W^{1,1}([0, 1]; \mathbb{R}^N), \gamma(0) = x, \gamma(1) = y \right\}.$$

We remark that this function is not symmetric in general. We define the *signed distance function* from a closed set $E \subseteq \mathbb{R}^N$ as

$$\text{sd}_E^\psi(x) := \inf_{y \in E} \text{dist}^\psi(y, x) - \inf_{y \notin E} \text{dist}^\psi(x, y), \quad (2.2)$$

def sd

so that $\text{sd}_E^\psi \geq 0$ on E^c and $\text{sd}_E^\psi \leq 0$ in E . We remark that the bounds stated in Definition 2.1 imply

$$\frac{1}{c_\psi} \text{dist} \leq \text{dist}^\psi \leq c_\psi \text{dist}, \quad (2.3)$$

inclusione psi p

where here and in the following we will denote with dist, sd the Euclidean distance and signed distance function respectively. We define the ψ -balls as the balls associated to the ψ -distance, that is

$$B_\rho^\psi(x) := \{y \in \mathbb{R}^N : \text{dist}^\psi(y, x) < \rho\},$$

which in general are not convex nor symmetric.

def reg ell integr

Definition 2.2. Given $\lambda \geq 1, l \geq 0$ we say that an anisotropy ϕ is a *regular elliptic integrand* if $\phi(x, \cdot)|_{S^N} \in C^{2,1}(S^N)$ and for every $x, y, e \in \mathbb{R}^N, \nu, \nu' \in S^N$ one has:

$$\begin{aligned} \frac{1}{\lambda} &\leq \phi(x, \nu) \leq \lambda, \\ |\phi(x, \nu) - \phi(y, \nu)| + |\nabla_p \phi(x, \nu) - \nabla_p \phi(y, \nu)| &\leq l|x - y| \\ |\nabla_p \phi(x, \nu)| + \|\nabla_p^2 \phi(x, \nu)\| + \frac{\|\nabla_p^2 \phi(x, \nu) - \nabla_p^2 \phi(x, \nu')\|}{|\nu - \nu'|} &\leq \lambda \\ e \cdot \nabla_p^2 \phi(x, \nu)[e] &\geq \frac{|e - (e \cdot \nu)\nu|^2}{\lambda}. \end{aligned}$$

We will denote a regular parametric elliptic integrand as $\phi \in \mathcal{E}$. Given any set of finite perimeter E , one can define the ϕ -perimeter P_ϕ as follows

$$P_\phi(E) := \int_{\partial^* E} \phi(x, \nu_E(x)) d\mathcal{H}^{N-1}(x),$$

where $\partial^* E$ is the reduced boundary of E and ν_E is the measure-theoretic outer normal, see [34] for further references on sets of finite perimeter. The perimeter of a set of finite perimeter E in an open set A is defined as

$$P_\phi(E; A) := \int_{\partial^* E \cap A} \phi(x, \nu_E(x)) d\mathcal{H}^{N-1}(x).$$

We remark that, by definition of regular elliptic integrand, for any C^2 set E it holds

$$\frac{1}{\lambda}P(E) \leq P_\phi(E) \leq \lambda P(E).$$

Some additional remarks on this definition can be found in [19]. We just recall the submodularity property of the ϕ -perimeter, whose proof can be found in [19, Remark 2.4].

Proposition 2.3 (Submodularity property). *For any two sets $E, F \subseteq \mathbb{R}^N$ of finite perimeter in an open set $A \subseteq \mathbb{R}^N$, one has*

$$P_\phi(E \cup F) + P_\phi(E \cap F) \leq P_\phi(E) + P_\phi(F). \quad (2.4)$$

submod

Moreover, by homogeneity, (2.1) and recalling that for any set E of finite perimeter it holds $D\chi_E = -\nu_E \, d\mathcal{H}^{N-1}|_{\partial^* E}$ we have the following equivalent definitions

$$\begin{aligned} P_\phi(E) &= \sup \left\{ \int_{\mathbb{R}^N} -D\chi_E \cdot \xi \ : \ \xi \in C_c^1(\mathbb{R}^N; \mathbb{R}^N), \phi^\circ(\cdot, \xi) \leq 1 \right\} \\ &= \sup \left\{ \int_E \operatorname{div} \xi \, d\mathcal{H}^{N-1} \ : \ \xi \in C_c^1(\mathbb{R}^N; \mathbb{R}^N), \phi^\circ(\cdot, \xi) \leq 1 \right\}. \end{aligned} \quad (2.5)$$

P via calibration

Concerning the regularity property of the ϕ -perimeter minimizers, we refer to [36]. We just recall the following results. Given two anisotropies $\phi, \psi \in \mathcal{E}$, we define the “distance” between them as

$$\begin{aligned} \operatorname{dist}_{\mathcal{E}}(\phi, \psi) &:= \sup \{ |\phi(x, p) - \psi(x, p)| \\ &\quad + |\nabla_p \phi(x, p) - \psi(x, p)| + |\nabla_p^2 \phi(x, p) - \nabla_p^2 \psi(x, p)| : x \in \mathbb{R}^N, p \in S^N \}, \end{aligned}$$

where $|\cdot|$ denotes the Euclidian norm. Then, some regularity properties of minimizers of ϕ -perimeter can be found in the corollary in part II 6 and the theorem in part II 7 in [36], which are recalled below.

Theorem 2.4. *Assume $\phi \in \mathcal{E}$. Then, for any Λ -minimizer E of the ϕ -perimeter, the reduced boundary $\partial^* E$ of the set E is of class $C^{1,1/2}$ and the singular set $\Sigma := \partial E \setminus \partial^* E$ satisfies*

$$\mathcal{H}^{N-3}(\Sigma) = 0.$$

Moreover, if $\operatorname{dist}_{\mathcal{E}}(\phi, |\cdot|)$ is small enough, then

$$\mathcal{H}^{N-7}(\Sigma) = 0.$$

We sum up these hypotheses that yield the complete regularity of Λ -minimizers of parametric elliptic integrands:

$$\begin{aligned} &\text{either } \phi \in \mathcal{E} \text{ and } N \leq 3, \\ &\text{or } \operatorname{dist}_{\mathcal{E}}(\phi, |\cdot|) \text{ is small enough and } N \leq 7. \end{aligned} \quad (2.6)$$

hp dim

2.1 The first variation of the ϕ -perimeter

In this section we compute the first variation of the ϕ -perimeter and define some additional operators associated to it.

Assume E is of class C^2 . Let X be a vector field and assume $\Psi(x, t) =: \Psi_t(x)$ is the associated flow. To simplify the notation, we write

$$\nu(x, t) = \nabla_x \text{sd}_{\Psi(E, t)}(x).$$

By classical formulae (see e.g. [10]) we can compute the following. For the sake of brevity, we avoid writing the evaluation $\phi = \phi(x, \nu_E(x))$, if not otherwise specified, and assume that all the integrals are made with respect to the Hausdorff $(N - 1)$ -dimensional measure \mathcal{H}^{N-1} .

$$\begin{aligned}
\frac{d}{dt} \Big|_{t=0} P_\phi(E_t) &= \frac{d}{dt} \Big|_{t=0} \int_{\partial E} \phi(\Psi_t(x), \nu(\Psi_t(x), t)) J\Psi_t \, d\mathcal{H}^{N-1}(x) \\
&= \int_{\partial E} \nabla_x \phi \cdot X + \nabla_p \phi \cdot (-\nabla_\tau(X \cdot \nu) + D\nu[X]) + \phi \operatorname{div}_\tau X \\
&= \int_{\partial E} \nabla_x \phi \cdot X + \nabla_p \phi \cdot (-\nabla_\tau(X \cdot \nu) + D\nu[X]) + \operatorname{div}_\tau(\phi X) - \nabla \phi \cdot X + (\nabla \phi \cdot \nu)(X \cdot \nu) \\
&= \int_{\partial E} \nabla_x \phi \cdot X + \nabla_p \phi \cdot (-\nabla_\tau(X \cdot \nu) + D\nu[X]) - \nabla_x \phi \cdot X - D\nu[\nabla_p \phi] \cdot X \\
&\quad + \operatorname{div}_\tau(\phi X) + (\nabla \phi \cdot \nu)(X \cdot \nu) \\
&= \int_{\partial E} -\nabla_p \phi \cdot \nabla_\tau(X \cdot \nu) + (\nabla_x \phi \cdot \nu)(X \cdot \nu) + (D\nu[\nabla_p \phi] \cdot \nu)(X \cdot \nu) + \operatorname{div}_\tau(\phi X) \\
&= \int_{\partial E} \operatorname{div}_\tau(\nabla_p \phi(X \cdot \nu)) - \nabla_p \phi \cdot \nabla_\tau(X \cdot \nu) + (X \cdot \nu)(\nabla_x \phi \cdot \nu) \\
&= \int_{\partial E} (\operatorname{div}_\tau \nabla_p \phi)(X \cdot \nu) + \nabla_p \phi \cdot \nabla_\tau(X \cdot \nu) - \nabla_p \phi \cdot \nabla_\tau(X \cdot \nu) + (\nabla_x \phi \cdot \nu)(X \cdot \nu) \\
&= \int_{\partial E} (X \cdot \nu) (\operatorname{div}_\tau \nabla_p \phi + \nabla_x \phi \cdot \nu) = \int_{\partial E} (X \cdot \nu) \operatorname{div} \nabla_p \phi
\end{aligned} \tag{2.7} \quad \text{eq align}$$

where the last equality follows from the definition of div_τ and the fact that ϕ is 1-homogeneous with respect to the p variable, since

$$\begin{aligned}
\operatorname{div} \nabla_p \phi &= \operatorname{div}_\tau \nabla_p \phi + \sum_i \nu_i \left(\frac{d}{dx_i} \nabla_p \phi \right) [\nu] \\
&= \operatorname{div}_\tau \nabla_p \phi + \sum_i \nu_i \nabla_p (\partial_{x_i} \phi) \cdot \nu + \nu \cdot (\nabla_p^2 \phi D\nu) [\nu] \\
&= \operatorname{div}_\tau \nabla_p \phi + \nabla_x \phi \cdot \nu.
\end{aligned}$$

Therefore, we define the first variation of a C^2 -regular set E , induced by the vector field X , as

$$\delta P_\phi(E)[X \cdot \nu] := \int_{\partial E} (X(x) \cdot \nu(x)) \operatorname{div} \nabla_p \phi(x, \nu(x)) \, d\mathcal{H}^{N-1}(x) \tag{2.8} \quad \text{first variation p}$$

and the ϕ -curvature of the set E as

$$H_E^\phi(x) := \operatorname{div} \nabla_p \phi(x, \nu(x)). \tag{2.9} \quad \text{def curvature}$$

If we now consider equation (2.7), we develop the tangential gradient to find

$$\nabla_p \phi \cdot (-\nabla_\tau(X \cdot \nu) + D\nu[X]) = \nabla_p \phi \cdot (-\nabla_\tau X[\nu] - D\nu[X] + D\nu[X]) = 0.$$

This shows that for any set E of class C^2 it holds

$$\delta P_\phi(E)[X \cdot \nu] := \int_{\partial E} (\nabla_x \phi \cdot X + \phi \operatorname{div}_\tau X) \, d\mathcal{H}^{N-1},$$

where we dropped the evaluation of ϕ at $(x, \nu_E(x))$. We remark that the expression on the right hand side makes sense even if the set E is just of finite perimeter. Defining the ϕ -divergence operator div_ϕ as

$$\operatorname{div}_\phi X := \nabla_x \phi \cdot X + \phi \operatorname{div}_\tau X, \quad (2.10)$$

we are led to define the distributional ϕ -curvature of a set E of finite perimeter as an operator $H_E^\phi \in L^1(\partial E)$ (if it exists) such that the following representation formula holds

$$\int_{\partial E} \operatorname{div}_\phi X \, d\mathcal{H}^{N-1} = \int_{\partial E} H_E^\phi \nu_E \cdot X \, d\mathcal{H}^{N-1}, \quad \forall X \in C_c^\infty(\mathbb{R}^N; \mathbb{R}^N). \quad (2.11)$$

def curvature var

The previous computations allow to say that the distributional ϕ -curvature can be expressed as (2.9) if the set is of class C^2 . Finally, since ϕ is a regular elliptic integrand, one can prove the following monotonicity result.

Lemma 2.5. *Let E, F be two C^2 sets of finite ϕ -perimeter with $E \subseteq F$, and assume that $x \in \partial F \cap \partial E$: then $H_F^\phi(x) \leq H_E^\phi(x)$.*

Proof. Since the anisotropy is smooth, we can expand the curvature formula (2.9) as

$$H^\phi = \operatorname{tr} (\nabla_x \nabla_p \phi(x, \nu) + \nabla_p^2 \phi(x, \nu) D\nu) \quad (2.12)$$

curvature expande

and compare H_E^ϕ with H_F^ϕ . We consider separately the two terms appearing in (2.12). The first one depends on ν just by the value it has at the point x . Therefore, since $\nu_E(x) = \nu_F(x)$ we have the equality. The second one falls in the classical framework of smooth anisotropies that do not depend on the space variable. Indeed, the second term in (2.12) is the same as the curvature in x associated to the anisotropy ϕ with the first variable freezed at x . Then, the result follows from classical results on curvatures invariant under translations (see e.g. [14, Lemma 4.3]). \square

3 The minimizing movements approach

In this section we follow the work of [35] (see also [3, 33]) to prove the existence for the mean curvature flow via the *minimizing movements* approach. We recall that in the whole paper we will assume the hypothesis (H).

3.1 The discrete scheme

In this subsection we will define the discrete scheme approximating the weak solution of the mean curvature flow, and we shall study some of its properties.

We define the following iterative scheme. Given $h > 0$, $f \in L^\infty(\mathbb{R}^N \times [0, \infty))$ and $t \geq h$, and given a bounded set of finite perimeter F , we minimize the energy functional

$$\mathcal{F}_{h,t}^F(E) = P_\phi(E) + \frac{1}{h} \int_E \text{sd}_F^\psi(x) \, dx - \int_E F_h(x, t) \, dx \quad (3.1) \quad \boxed{\text{problema discret}}$$

in the class of all measurable sets $E \subseteq \mathbb{R}^N$, and where we have set

$$F_h(x, t) := \int_t^{t+h} f(x, s) \, ds.$$

Equivalently, we could define the energy functional as

$$\mathcal{F}_{h,t}^F(E) = P_\phi(E) + \frac{1}{h} \int_{E \Delta F} |\text{sd}_F^\psi| - \int_E F_h(x, t) \, dx,$$

which agrees with (3.1) up to a constant. We will call the *incremental problem* this minimizing procedure. Then, we denote

$$T_{h,t}F = E \in \operatorname{argmin} \mathcal{F}_{h,t}^F.$$

It is well-known (compare (2.8) and [34, Proposition 17.8]) that a minimum of (3.1) of class C^2 satisfies the Euler-Lagrange equation

$$\int_{\partial E_t} H_{E_t}^\phi X \cdot \nu_{E_t} \, d\mathcal{H}^{N-1} = - \int_{\partial E} \left(\frac{1}{h} \text{sd}_{E_{t-h}}^\psi(x) - F_h(x, t) \right) X(x) \cdot \nu_{E_t}(x) \, d\mathcal{H}^{N-1}(x) = 0 \quad (3.2) \quad \boxed{\text{EL equation}}$$

for all $X \in C_c^\infty(\mathbb{R}^N; \mathbb{R}^N)$. We can then define the *discrete flow*, which can be seen as a discrete-in-time approximation of the mean curvature flow starting from the initial set E_0 . We first iteratively define the sequence $(E_n^{(h)})_{n \in \mathbb{N}}$ by setting $E_0^{(h)} = E_0$,

$$E_n^{(h)} = T_{h, (n-1)h} E_{n-1}^{(h)} \quad \forall n \geq 1.$$

Then, we define the *discrete flow* simply as

$$E_t^{(h)} = E_{[t/h]}^{(h)}, \quad t \in [0, +\infty). \quad (3.3) \quad \boxed{\text{def discrete flow}}$$

This section is devoted to recall and prove some estimates on the discrete flow. The first one is a well-known existence result.

Lemma. *For any measurable function $g : \mathbb{R}^N \rightarrow \mathbb{R}$ such that $\min\{g, 0\} \in L_{loc}^1$, the problem*

$$\min \left\{ P(E) + \int_E g : E \text{ is of finite perimeter} \right\}$$

admits a solution.

Consider now F as a bounded set of finite perimeter. Then, the function $g = \text{sd}_F^\psi/h - F_h$ is coercive, thus $\min\{g, 0\} \in L^1$. Therefore, by the previous result and by classical arguments, one can prove the following result.

Lemma 3.1. *For any given set F of finite perimeter, the problem (3.1) admits a solution E , which satisfies the discrete dissipation inequality*

$$P_\phi(E) + \frac{1}{h} \int_{E\Delta F} |\text{sd}_F^\psi| \leq P_\phi(F) + \int_{E\setminus F} F_h(x, t) \, dx - \int_{F\setminus E} F_h(x, t) \, dx.$$

Moreover, the problem (3.1) admits a minimal and a maximal solution.

We define $T_{h,t}^+ F$ (respectively $T_{h,t}^- F$) as the maximal (respectively minimal) solution to (3.1) having as initial datum F . In the following, whenever no confusion is possible, we shall write $T_{h,t}$ instead of $T_{h,t}^-$.

A comparison result holds. We will consider just bounded sets as datum for the problem (3.1), but the same result holds in general for unbounded sets (see also Section 4.1 for the case of unbounded sets with bounded boundary). The proof of this result is classical (see e.g. [14]) and it is based on the submodularity of the perimeter (2.4). We will omit it.

Lemma 3.2 (Weak comparison principle). *Assume that F_1, F_2 are bounded sets with $F_1 \subseteq\subseteq F_2$ and consider $g_1, g_2 \in L^\infty$ with $g_1 \geq g_2$. Then, for any two solutions E_i , $i = 1, 2$ of the problems*

$$\min \left\{ P_\phi(E) + \int_E \frac{\text{sd}_{F_i}^\psi}{h} + g_i : E \text{ is of finite perimeter} \right\},$$

we have $E_1 \subseteq E_2$. If, instead, $F_1 \supseteq F_2$, then we have that the minimal (respectively maximal) solution to (3.1) for $i = 1$ is contained in the minimal (respectively maximal) solution to (3.1) for $i = 2$.

We now prove the volume-density estimates for minimizers of problem (3.1). This result is based on the minimality properties of almost-minimizers for perimeters induced by regular elliptic integrands (see [19, Remark 1.9] for further results). These estimates have the disadvantage that the smallness condition on the radius depends on the parameter h . Subsequently, we will recall a finer result in the spirit of [33], where we can drop this dependence by making some restrictions on the balls considered.

Lemma 3.3. *Let $g \in L^\infty$ and assume E minimizes the functional*

$$\mathcal{F}(F) = P_\phi(F) + \int_F g$$

among all measurable subsets of \mathbb{R}^N . Then the density estimate

$$\begin{aligned} \sigma \rho^N &\leq |B_\rho(x) \cap E| \leq (1 - \sigma) \rho^N \\ \sigma \rho^{N-1} &\leq P_\phi(E; B_\rho(x)) \leq (1 - \sigma) \rho^{N-1} \end{aligned} \tag{3.4}$$

holds for all $x \in \partial^* E$, $0 < \rho < (2\lambda \|g\|_\infty)^{-1} := \rho_0$, for a suitable $\sigma = \sigma(N, c_\psi, \lambda)$.

Proof. By minimality,

$$P_\phi(E) \leq P_\phi(F) + \|g\|_\infty |E\Delta F| \quad \forall F \subseteq \mathbb{R}^N,$$

thus [19, Lemma 2.8] implies the thesis. \square

Remark 3.4. We remark that the previous result allows us to choose the minimal solution to (3.1) to be an open set, and the maximal one to be a closed set. This follows from the fact that the density estimates imply that the boundary of any minimizer has zero measure.

We now recall [12, Lemma 3.7], which is an anisotropic version of [33, Remark 1.4]. It provides volume-density estimates for minimizers of (3.1) starting from E , uniform in ψ and h , holding in the exterior of E . We remark that, even if in the reference the anisotropy ϕ considered did not depend on x , all the arguments hold with minor modifications also in our case. We recall the proof of this result, as similar techniques will be used later on.

Lemma 3.5. *Let E be a bounded, closed set, $h > 0$, and $g \in L^\infty(\mathbb{R}^N)$. Let E' be a minimizer of*

$$P_\phi(F) + \int_F \frac{\text{sd}_E^\psi}{h} + g.$$

Then, there exists $\sigma > 0$, depending on λ , and $r_0 \in (0, 1)$, depending only on $N, \lambda, G := \|g\|_{L^\infty(F)}$, with the following property: if \bar{x} is such that $|E' \cap B_s(\bar{x})| > 0$ for all $s > 0$ and $B_r(\bar{x}) \cap E = \emptyset$ with $r \leq r_0$, then

$$|E' \cap B_r(\bar{x})| \geq \sigma r^N. \tag{3.5}$$

Analogously, if \bar{x} is such that $|B_s(\bar{x}) \setminus E'| > 0$ for all $s > 0$ and $B_r(\bar{x}) \subseteq E$ with $r \leq r_0$, then

$$|B_r(\bar{x}) \setminus E'| \geq \sigma r^N.$$

Proof. For all $s \in (0, r)$, set $E'(s) := E' \setminus B_s(\bar{x})$. Note that, for a.e. s we have

$$P_\phi(E'(s)) = P_\phi(E') - P_\phi(E' \cap B_s(\bar{x})) + \int_{E' \cap \partial B_s(\bar{x})} (\phi(x, \nu(x)) + \phi(x, -\nu(x))) \, d\mathcal{H}^{N-1}(x),$$

where ν denotes the outer normal vector of the set $E' \cap \partial B_s(\bar{x})$. Since $\text{sd}_E^\psi \geq 0$ in E^c , one has $\int_{E'(s)} \text{sd}_E^\psi \leq \int_{E'} \text{sd}_E^\psi$, and the minimality of E' implies

$$P_\phi(E' \cap B_s(\bar{x})) + \int_{E' \cap B_s(\bar{x})} g \leq \int_{E' \cap \partial B_s(\bar{x})} (\phi(x, \nu(x)) + \phi(x, -\nu(x))) \, d\mathcal{H}^{N-1}(x).$$

By the bound on the ϕ -perimeter and using the classical isoperimetric inequality (whose constant is denoted C_N) we obtain

$$\begin{aligned} 2\lambda \mathcal{H}^{N-1}(E' \cap \partial B_s(\bar{x})) &\geq \frac{1}{\lambda} P(E' \cap B_s(\bar{x})) + \int_{E' \cap B_s(\bar{x})} g \\ &\geq \frac{1}{\lambda} C_N |E' \cap B_s(\bar{x})|^{\frac{N-1}{N}} - \|g\|_\infty |E' \cap B_s(\bar{x})| \geq \frac{C_N}{2\lambda} |E' \cap B_s(\bar{x})|^{\frac{N-1}{N}}, \end{aligned}$$

provided $|E' \cap B_s(\bar{x})|^{1/N} \leq C_N / (2\lambda \|g\|_\infty)$, which is true if r_0 is small enough. Since the rhs is positive for every s , we conclude

$$\frac{d}{ds} |E' \cap B_s(\bar{x})|^{\frac{1}{N}} \geq \frac{C_N}{4\lambda^2 N} \quad \text{for a.e. } s \in (0, r). \tag{3.6}$$

The thesis follows by integrating the above differential inequality. The other case is analogous. \square

Remark 3.6. Requiring that the anisotropy ψ is bounded uniformly from above and below ensures that the results of the previous Lemmas 3.3 and 3.5 can be read in terms of the ψ -balls. For example, for any $r \geq 0$ and $x \in \mathbb{R}^N$, equation (3.5) could be read as $|E' \cap B_r^\psi(\bar{x})| \geq \sigma c_\psi^{-N} r^N$, provided \bar{x} is such that $|E' \cap B_s^\psi(\bar{x})| > 0$ for all $s > 0$ and $B_r^\psi(\bar{x}) \cap E = \emptyset$, and holds for all $r \leq r_0/c_\psi$. Here, σ is as in Lemma 3.5 and depends only on λ . Analogous statements holds for Lemma 3.9.

We now provide some estimates on the evolution of balls under the discrete flow. We start by a simple remark concerning the boundedness of the evolving sets.

Remark 3.7. A simple estimate on the energies implies that the minimizers of (3.1) are bounded whenever F is bounded. Indeed, assume $F \subseteq B_R$ and consider $B_\rho(x) \cap (E \setminus B_R) \neq \emptyset$: testing the minimality of E against F we easily deduce

$$\frac{R}{2h} |B_\rho(x) \cap E| \leq \int_{E \cap B_\rho(x)} \frac{\text{sd}_F^\psi}{h} \leq P_\phi(F) + \|F_h(\cdot, t)\|_\infty |E \Delta F| \leq P_\phi(F) + \|f\|_\infty (|F| + |E|).$$

Employing the density estimates of Proposition 3.5 and sending $R \rightarrow \infty$, we get a contradiction, as the isoperimetric inequality implies that $|E|$ is bounded since $\mathcal{F}_{h,t}^F(F) < \infty$.

We now want to prove finer estimates on the speed of evolution of balls. These estimates are classically a crucial step in order to prove existence of the flow. In the case under study, the main difficulties come from the inhomogeneity of the functionals considered, as in the homogeneous case convexity arguments easily yield the boundedness result, for example. We will use a ‘‘variational’’ approach in the spirit of [14] (but see also [35, Lemma 3.8] for a different proof relying more on the smoothness of the evolving set).

Lemma 3.8. *For every $R_0 > 0$ there exist $h_0(R_0) > 0$ and $C(R_0, \phi, \psi, f) > 0$ with the following property: For all $R \geq R_0$, $h \in (0, h_0)$, $t > 0$ and $x \in \mathbb{R}^N$ one has*

$$T_{h,t}(B_R(x)) \supset B_{R-Ch}(x). \quad (3.7)$$

Proof. We divide the proof into three steps. In the following, the constants σ, r_0 are those of Lemma 3.5. We will assume $x = 0$ for simplicity. We fix $R \geq R_0$ and denote $E := T_{h,t}B_R$.

Step 1. We prove that, given $a \in (0, \sigma)$, $\varepsilon \in (0, 1)$, we can ensure $|B_{R(1-\varepsilon)} \setminus E| < a R^N (1-\varepsilon)^N$ for h small enough. Indeed, assume by contradiction $|B_{R(1-\varepsilon)} \setminus E| \geq a R^N (1-\varepsilon)^N$. Testing the minimality of E against B_R , we obtain

$$\int_{(B_{R(1-\varepsilon)} \setminus E) \cup (E \setminus B_R)} \frac{|\text{sd}_{B_R}^\psi|}{h} \leq \frac{1}{h} \int_{B_R \Delta E} |\text{sd}_{B_R}^\psi| \leq P_\phi(B_R) + \int_{B_R \setminus E} - \int_{E \setminus B_R} F_h(x, t) \, dx,$$

and estimating $|\text{sd}_{B_R}^\psi| \geq \varepsilon/c_\psi$ on $B_{R(1-\varepsilon)} \setminus E$, we get

$$\frac{\varepsilon}{hc_\psi} |B_{R(1-\varepsilon)} \setminus E| \leq P_\phi(B_R) + \|f\|_\infty (\omega_N R^N + |B_{R(1+\varepsilon)} \setminus B_R|) + \int_{E \setminus B_{R(1+\varepsilon)}} \left(F_h - \frac{|\text{sd}_{B_R}^\psi|}{h} \right) \, dx.$$

Taking $h \leq \varepsilon/(c_\psi \|f\|_\infty)$, the last term on the rhs is negative, thus we obtain for ε small

$$\frac{\varepsilon}{hc_\psi} |B_{R(1-\varepsilon)} \setminus E| \leq P_\phi(B_R) + \|f\|_\infty R^N (\omega_N + 2^{N+1} \varepsilon).$$

We employ the hypothesis to obtain

$$\frac{a}{hc_\psi} \varepsilon(1-\varepsilon)^N R^N \leq c_\psi N \omega_N R^{N-1} + cR^N,$$

a contradiction for $h \leq ca\varepsilon(1-\varepsilon)^N \min\{1, R\}$, where c is a constant depending on $N, \phi, \psi, \|f\|_\infty$. **Step 2.** We now prove that, for h small enough, if $|B_{3R/4} \setminus E| < \sigma(3R/4)^N$, then $B_{R/2} \subseteq E$. We first assume that $R \leq r_0$. In this case, if $|B_s \setminus E| > 0$ for all $s \leq R \leq r_0$, we can follow the second part of the proof of Lemma 3.5 (since obviously $B_s \subseteq R_R$) to obtain equation (3.6), which reads

$$\frac{d}{ds} |B_s \setminus E|^{1/N} \geq \frac{C_N}{4\lambda^2 N} = \sigma^{1/N} \quad \text{for a.e } s \in (0, R).$$

Using the condition

$$|B_{3R/4} \setminus E|^{1/N} < \sigma^{1/N} (3R/4),$$

we deduce the existence of a positive extinction radius

$$R^* = \frac{3R}{4} - \frac{|B_{3R/4} \setminus E|^{1/N}}{\sigma^{1/N}} \tag{3.8}$$

extinction radius

such that $|B_{R^*} \setminus E| = 0$. Applying Step 1 with the values $a = \sigma/3^N$, $\varepsilon = 1/4$, we get

$$|B_{3R/4} \setminus E| \leq \frac{\sigma}{4^N} R^N,$$

holding for $h \leq c(N, \phi, \psi, f)R$, which combined with (3.8) gives $R^* > R/2$. Clearly, taking $h \leq cR_0$ the smallness assumption on h is uniform for $R \geq R_0$.

If $R \geq r_0$ one simply uses a covering argument. For any $x \in B_{R-r_0}$, applying the previous result to the ball $B_{r_0}(x)$ and using the comparison principle of Lemma 3.2, we conclude that $\forall h \leq cr_0$ it holds

$$\bigcup_{x \in B_{R-r_0}} B_{r_0/2}(x) \subseteq E.$$

Step 3. We conclude the proof. By the previous two steps, taking h small enough we see that

$$\rho = \sup\{r > 0 : |B_r \setminus E| = 0\} \in (R/2, \tilde{R}],$$

where \tilde{R} is the constant of Remark 3.7. We can assume $\rho \leq R$, otherwise the result of the lemma is trivial. Consider the vector field $\nabla_p \phi \left(x, \frac{x}{|x|} \right) \in C^1(\mathbb{R}^N, \mathbb{R}^N)$. Then, recalling (2.5), we get $P_\phi(F) \geq -\int_{\mathbb{R}^N} D\chi_G \cdot \nabla_p \phi(x, x/|x|)$ for all G set of finite perimeter and

$$P_\phi((1+\varepsilon)B_\rho) = \int_{\mathbb{R}^N} D\chi_{(1+\varepsilon)B_\rho} \cdot \left(-\nabla_p \phi \left(x, \frac{x}{|x|} \right) \right).$$

Setting $W_\varepsilon = (1+\varepsilon)B_\rho \setminus E$, by submodularity on $(1+\varepsilon)B_\rho, E$ and exploiting the minimality of E , we obtain

$$\begin{aligned} \int_{\mathbb{R}^N} \nabla_p \phi \left(x, \frac{x}{|x|} \right) \cdot D\chi_{W_\varepsilon} &= \int_{\mathbb{R}^N} \nabla_p \phi \left(x, \frac{x}{|x|} \right) \cdot (D\chi_{(1+\varepsilon)B_\rho} - D\chi_{(1+\varepsilon)B_\rho \cap E}) \\ &\leq P_\phi((1+\varepsilon)B_\rho \cap E) - P_\phi((1+\varepsilon)B_\rho) \end{aligned}$$

$$\begin{aligned}
&\leq P_\phi(E) - P_\phi((1+\varepsilon)B_\rho \cup E) \\
&\leq \frac{1}{h} \int_{W^\varepsilon} \text{sd}_{B_R}^\psi + \int_{W^\varepsilon} F_h(x,t) \, dx.
\end{aligned}$$

We conclude, using the divergence theorem ,

$$\int_{W^\varepsilon} -\text{div} \nabla_p \phi \left(x, \frac{x}{|x|} \right) \leq \frac{1}{h} \int_{W^\varepsilon} \text{sd}_{B_R}^\psi + \|f\|_\infty |W^\varepsilon|.$$

Dividing by $|W^\varepsilon|$ and sending $\varepsilon \rightarrow 0$ we obtain

$$\int_{\partial B_\rho \cap \partial E} -\text{div} \nabla_p \phi \left(x, \frac{x}{|x|} \right) \leq \frac{1}{c_\psi} \frac{\rho - R}{h} + \|f\|_\infty.$$

Exploiting the regularity assumptions on ϕ , we remark that

$$|\text{div} \nabla_p \phi| = |\text{tr} (\nabla_x \nabla_p \phi + \nabla_p^2 \phi \nabla(x/|x|))| \leq C \left(1 + \frac{1}{|x|} \right).$$

Thus, we obtain

$$-C \left(1 + \frac{1}{\rho} \right) \leq \frac{\rho - R}{h},$$

which implies that $\rho \in (0, \rho_1) \cup (\rho_2, R)$ for $\rho_{1,2} = (R - Ch \mp \sqrt{(R - Ch)^2 - 4Ch})/2$, as long as $h \leq R_0^2/(4C)$. Since the choice $\rho \leq \rho_1 < R/2$ is not admissible, we conclude the proof by estimating

$$\rho_2 = R - Ch + \frac{R - Ch}{2} \left(\sqrt{1 - \frac{4Ch}{(R - Ch)^2}} - 1 \right) \geq R - Ch - \frac{Ch}{R - Ch},$$

from which the thesis follows. \square

The proof of the previous result can be employed to prove an estimate from above of the evolution speed of the flow, as the following result shows. Since the proof follows the same lines and is easier in this case, we only sketch it.

a priori estimate

Lemma 3.9. *Fix $T > 0$ and $R_0 > 0$. Then, there exists positive constants $C = C(\phi, \psi, f, R_0)$ and $h_0 = h_0(R_0)$ such that, for every $R \geq R_0$ and $h \leq h_0$, if $E_0 \subseteq B_R$, then $E_t^{(h)} \subseteq B_{R+CT}$ for all $t \in (0, T)$.*

Proof. Choose h small as in the previous result and set

$$\rho = \inf\{r > 0 : |E \setminus B_r| = 0\} \in (R/2, \tilde{R}],$$

for h small enough, where \tilde{R} is the constant of Remark 3.7. We can assume $\rho \geq R$, otherwise the result is trivial. Defining $W_\varepsilon = E \setminus (1 - \varepsilon)B_\rho$ and reasoning as before we obtain

$$\begin{aligned}
\int_{\mathbb{R}^N} \nabla_p \phi \left(x, \frac{x}{|x|} \right) \cdot D\chi_{W^\varepsilon} &= \int_{\mathbb{R}^N} \nabla_p \phi \left(x, \frac{x}{|x|} \right) \cdot (D\chi_{(1-\varepsilon)B_\rho \cup E} - D\chi_{(1-\varepsilon)B_\rho}) \\
&\geq -P_\phi((1-\varepsilon)B_\rho \cup E) + P_\phi((1-\varepsilon)B_\rho)
\end{aligned}$$

$$\begin{aligned}
&\geq -P_\phi(E) + P_\phi((1-\varepsilon)B_\rho \cap E) \\
&\geq \frac{1}{h} \int_{W^\varepsilon} \text{sd}_{B_R}^\psi + \int_{W^\varepsilon} F_h(x, t) \, dx.
\end{aligned}$$

As in the previous proof, we arrive at

$$\frac{\rho - R}{h} \leq C \left(1 + \frac{1}{\rho} \right),$$

which implies that $\rho \leq \rho_2 = \left(R + Ch + \sqrt{(R + Ch)^2 + 4Ch} \right) / 2 \leq R + Ch$, up to changing C . \square

3.2 Existence of flat flows

In the following, we will prove that the discrete flow (defined in (3.3)) defines a discrete-in-time approximation of a weak solution to the mean curvature flow, which is usually known as a “flat” flows (because the approximating surfaces $\partial^* E_t^{(h)}$ converge in the “flat” distance of Whitney to the limit $\partial^* E_t$, see [3]).

We start by proving uniform bounds on the distance between two consecutive sets of the discrete flow and on the symmetric difference between them. We introduce the time-discrete normal velocity: for all $t \geq 0$ and $x \in \mathbb{R}^N$, we set

$$v_h(x, t) := \begin{cases} \frac{1}{h} \text{sd}_{E_{t-h}^{(h)}}^\psi(x) & \text{for } t \in [h, +\infty) \\ 0 & \text{for } t \in [0, h). \end{cases}$$

The following result provides a bound on the L^∞ -norm of the discrete velocity. Since the proof is essentially the same of [33, Lemma 2.1], we will omit it. The only difference is that we use the upper and lower bounds of (2.3) to work with Euclidean balls.

L infy estimate

Lemma 3.10. *There exists a dimensional constant c with the following property. Let E_0 be a bounded set of finite perimeter and let $\{E_t^{(h)}\}_{t \in (0, T)}$ be a discrete flow starting from E_0 . Then,*

$$\sup_{E_t^{(h)} \Delta E_{t-h}^{(h)}} |v_h(\cdot, t)| \leq c_\infty h^{-1/2},$$

where c_∞ is a constant depending on N, ψ only.

The following result can be found in [35, Proposition 3.4] (see also [23, Lemma 2.2]): it provides an estimate on the volume of the symmetric difference of two consecutive sets of the discrete flow. The proof is analogous to the one in the reference.

Lemma. *Let F be a bounded set of finite perimeter and let E be a minimizer of (3.1). Then, there exists a constant C depending on ϕ, ψ such that*

$$|E \Delta F| \leq C \left(l P_\phi(E) + \frac{1}{l} \int_{E \Delta F} |\text{sd}_F^\psi| \right) \quad \forall l \leq c_\infty c_\psi \sqrt{h}. \quad (3.9)$$

stima L1

We are now able to prove an uniform bound on the perimeter of the evolving sets. The proof follows [23, Proposition 2.3].

Lemma 3.11. *For any initial bounded set E_0 of finite ϕ -perimeter and h small enough, the discrete flow $\{E_t^{(h)}\}$ satisfies*

$$P_\phi(E_t^{(h)}) \leq C_T \quad \forall t \in (0, T),$$

for a suitable constant $C_T = C_T(T, E_0, f, \phi, \psi)$.

Proof. Set $k = [T/h]$. By testing the minimality of $E_t^{(h)}$ against $E_{t-h}^{(h)}$ we obtain $\forall t \in [h, T]$

$$P_\phi(E_t^{(h)}) + \int_{E_t^{(h)} \Delta E_{t-h}^{(h)}} |\text{sd}_{E_{t-h}^{(h)}}^\psi| \leq P_\phi(E_{t-h}^{(h)}) + \|f\|_\infty |E_t^{(h)} \Delta E_{t-h}^{(h)}|. \quad (3.10) \quad \text{eq 2.4 FJM}$$

Combining this estimate with (3.9) for $l = 2ch\|f\|_\infty \ll c_\infty c_\psi \sqrt{h}$, with c being a suitably large constant, we obtain

$$P_\phi(E_t^{(h)}) + \frac{1}{2h} \int_{E_t^{(h)} \Delta E_{t-h}^{(h)}} |\text{sd}_{E_{t-h}^{(h)}}^\psi| \leq (1 + 2ch\|f\|_\infty^2) P_\phi(E_{t-h}^{(h)}) \quad (3.11) \quad \text{eq iterativa}$$

Iterating the previous estimate, we find

$$P_\phi(E_t^{(h)}) \leq (1 + ch)^{[\frac{t}{h}]-1} P_\phi(E_h^{(h)}) \leq c^{h[\frac{t}{h}]-h} P_\phi(E_h^{(h)})$$

For the first iterate, we start by observing that by Remark 3.7, for $h = h(E_0)$ small enough, we have $E_h^{(h)} \subseteq B_{2r}$, where $E_0 \subseteq B_r$. Therefore, by (3.10) for $t = h$ we obtain $P_\phi(E_h^{(h)}) \leq P_\phi(E_0) + c$ and we conclude $P_\phi(E_t^{(h)}) \leq C_T(P_\phi(E_0) + 1)$. \square

Following the previous proof, employing again (3.11) we find

$$\begin{aligned} P_\phi(E_{2h}) + \int_{E_2^{(h)} \Delta E_1^{(h)}} |v_h| + \int_{E_1^{(h)} \Delta E_0^{(h)}} |v_h| &\leq (1 + ch)P_\phi(E_1^{(h)}) + \int_{E_1^{(h)} \Delta E_0^{(h)}} |v_h| \\ &\leq (1 + ch) \left(P_\phi(E_1^{(h)}) + \int_{E_1^{(h)} \Delta E_0^{(h)}} |v_h| \right) \leq (1 + ch)^2 P_\phi(E_0). \end{aligned}$$

Iterating, we conclude as before

$$\sum_{k=1}^{[T/h]} \int_{E_{kh}^{(h)} \Delta E_{(k-1)h}^{(h)}} |v_h| \leq C_T(P_\phi(E_0) + 1). \quad (3.12) \quad \text{eq 3.18 MugSeiSp}$$

Therefore, combining the previous results and applying (3.9) with $l = h \ll \sqrt{h}$, we obtain

$$\int_h^T |E_t^{(h)} \Delta E_{t-h}^{(h)}| \leq c \sum_{k=1}^{[T/h]} \left(hP_\phi(E_{kh}^{(h)}) + \int_{E_{kh}^{(h)} \Delta E_{(k-1)h}^{(h)}} |v_h| \right) \leq C_T(P_\phi(E_0) + 1). \quad (3.13) \quad \text{eq 3.19 MugSeiSp}$$

With the previous estimates, the local Hölder continuity in time of the discrete flow, uniformly in h , is now easily deduced as in [23, Proposition 2.3]. We omit the proof as it is similar.

Proposition 3.12. *Let E_0 be an initial bounded set of finite ϕ -perimeter and $T > 0$. Then, for h small enough, for a discrete flow $\{E_t^{(h)}\}$ starting from E_0 it holds*

$$|E_t^{(h)} \Delta E_s^{(h)}| \leq C_T |t - s|^{1/2} \quad \forall h \leq t \leq s < T,$$

for a suitable constant $C_T = C_T(T, E_0, f, \phi, \psi)$.

We finally prove the main result of this section, the existence of flat flows.

Proof of Theorem 1.1. The proof is classical and we only sketch it. By the uniform equicontinuity of the approximating sequence of Proposition 3.12 and compactness of sets of finite perimeter (by Lemma 3.9 and 3.11) we can use the Ascoli-Arzelà theorem to prove that the sequence $(E_t^{(h_k)})_{k \in \mathbb{N}}$ converges in L^1 to sets E_t for all times $t \geq 0$ and that the family $\{E_t\}_{t \geq 0}$ satisfies the $1/2$ -Hölder continuity property, locally uniformly in time. The other property is then easily deduced. \square

3.3 Existence of distributional solutions

From Theorem 1.1 we deduce the existence of a subsequence $(h_k)_{k \geq 0}$ such that

$$D\chi_{E_t^{(h_k)}} \xrightarrow{*} D\chi_{E_t} \quad \forall t \geq 0. \quad (3.14)$$

convergenza senso

We will also assume (1.6), remarking that it implies

$$\lim_{k \rightarrow \infty} P_\phi(E_t^{(h_k)}) = P_\phi(E_t) \quad \text{for a.e. } t \in [0, +\infty). \quad (3.15)$$

convergenza perim

Our aim is to derive (1.7) and (1.8) from the Euler-Lagrange equation (3.2) and passing to the limit $h \rightarrow 0$. To achieve so, we will prove that the discrete velocity is a good approximation (up to multiplicative factors) of the discrete evolution speed of the sets. Notice that (1.7) is a weak formulation of (1.1), while (1.8) establishes the link between v and the velocity of the boundaries of E_t . Indeed, law (1.1) can be interpreted as looking for a family $\{E_t\}_{t \geq 0}$ of sets, whose normal vector ν_{E_t} and ϕ -curvature $H_{E_t}^\phi$ are well-defined objects and a function $v : [0, \infty) \times \mathbb{R}^N \rightarrow \mathbb{R}$ such that

$$\begin{cases} v &= -H_{E_t}^\phi + f \\ V &= \psi(x, \nu_{E_t})v, \end{cases} \quad (3.16)$$

where V represents the normal velocity of evolution, obtained as limit as $h \rightarrow 0$ (in a suitable sense) of the ratio

$$\frac{\chi_{E_t} - \chi_{E_{t-h}}}{h}.$$

In this whole section we will assume that hypothesis (2.6) holds. In particular, the sets defining the discrete flow are smooth hypersurfaces in \mathbb{R}^N . Moreover, we require hypotheses (1.4) to hold.

We start by estimating in time the L^2 -norm of the discrete velocity. The proof is the same of [35, Lemma 3.6], up to using the density estimates on the ϕ -perimeter of Lemma 3.3 and considering the ψ -balls instead of the Euclidean one.

Proposition 3.13. *Let $\{E_t^{(h)}\}_{t \geq 0}$ be a discrete flow starting from an initial bounded set E_0 of finite ϕ -perimeter. Then, for any $T > 0$ and for h small enough, it holds*

$$\int_0^T \int_{\partial E_t^{(h)}} v_h^2 d\mathcal{H}^{N-1} dt \leq C_T,$$

for a suitable constant $C_T = C_T(T, E_0, \phi, \psi, f)$.

Recalling now the Euler-Lagrange equation (3.2) and Lemma 3.11 we conclude

$$\int_0^T \int_{\partial E_t^{(h)}} \left(H_{E_t^{(h)}}^\phi \right)^2 = \int_0^T \int_{\partial E_t^{(h)}} (v_h - F_h)^2 \leq C_T, \quad (3.17)$$

We now prove an estimate on the error between the discrete velocity $\psi(\cdot, \nu_{E_t})v_h(\cdot, t)$ and the discrete time derivative of χ_h . The proof of this result is based on a double blow-up argument, and the smoothness of sets (locally) minimizing the ϕ -perimeter is essential. We will split the proof in various lemmas: the first one concerns the composition of blow-ups.

Lemma 3.14 (Composition of blow-ups). *Consider $0 < \beta < \beta' < 1$. Assume that A is a set such that the following blow-ups converge as $h \rightarrow 0$*

$$\begin{aligned} \frac{A - x_0}{h^\beta} &\rightarrow A_1 \quad \text{in } L_{loc}^1 \\ h^{-(\beta' - \beta)} A &\rightarrow A_2 \quad \text{in } L_{loc}^1, \end{aligned}$$

where $x_0 \in \partial A$. Then, if the composition of the blow-ups $h^{-\beta'}(A - x_0)$ converges in L_{loc}^1 , the limit coincides with A_2 .

Proof. We can assume wlog $x_0 = 0$. Denote with $A_3 = L_{loc}^1 - \lim_{h \rightarrow 0} h^{-\beta'} A$. We fix a ball B_M and $\varepsilon > 0$. There exists h^* such that $\forall h \leq h^*$ it holds

$$|((h^{-\beta'} A) \Delta A_3) \cap B_M| \leq \varepsilon, \quad |((h^{-\beta' + \beta} A_1) \Delta A_2) \cap B_M| \leq \varepsilon.$$

We fix h and wlog assume $Mh^{\beta' - \beta} \leq 1$. Taking $\tilde{h} < h$ suitably small (depending on h, ε), we can ensure

$$|((\tilde{h}^{-\beta} A) \Delta A_1) \cap B_1| \leq \varepsilon h^{N(\beta' - \beta)}.$$

Since $\tilde{h}^{-\beta} h^{-(\beta' - \beta)} > h^{-\beta'}$, there exists $\bar{h} < h$ such that $\bar{h}^{-\beta'} = \tilde{h}^{-\beta} h^{-(\beta' - \beta)}$. We can then estimate

$$\begin{aligned} |(A_3 \Delta A_2) \cap B_M| &\leq |(A_3 \Delta \bar{h}^{-\beta'} A) \cap B_M| + |((h^{-\beta' + \beta} A_1) \Delta (\bar{h}^{-\beta'} A)) \cap B_M| \\ &\quad + |((h^{-\beta' + \beta} A_1) \Delta A_2) \cap B_M| \\ &\leq 2\varepsilon + h^{-N(\beta' - \beta)} |(A_1 \Delta (\tilde{h}^{-\beta} A)) \cap B_{Mh^{\beta' - \beta}}| \\ &\leq 2\varepsilon + h^{-N(\beta' - \beta)} |(A_1 \Delta (\tilde{h}^{-\beta} A)) \cap B_1| \leq 3\varepsilon. \end{aligned}$$

□

We now compute some estimates on the normal vector on the boundary of the evolving sets. We fix c_∞ as the constant appearing in Lemma 3.10.

Lemma 3.15. *Assume (H) and (1.4). For given constants $1/2 < \beta' < \alpha < 1$ and $T > 2$, there exists a continuous increasing function $\omega : [0, 1] \rightarrow \mathbb{R}$ with $\omega(0) = 0$ and with the following property. Consider $t \in [2h, T]$ and $x_0 \in \partial E_t^{(h)}$ such that*

$$|v_h(t, y)| \leq h^{\alpha-1} \quad \forall y \in B_{c_\infty \sqrt{h}}(x_0) \cap (E_t^{(h)} \Delta E_{t-h}^{(h)}). \quad (3.18)$$

eq 4.6 MugSeiSpa

Then, there exists $\nu \in S^N$ such that

$$\begin{aligned} |\nu_{E_t^{(h)}}(\cdot) - \nu| &\leq \omega(h) \quad \text{in } B_{h^{\beta'}}(x_0) \cap \partial E_t^{(h)} \\ |\nu_{E_{t-h}^{(h)}}(\cdot) - \nu| &\leq \omega(h) \quad \text{in } B_{h^{\beta'}}(x_0) \cap \partial E_{t-h}^{(h)}. \end{aligned} \quad (3.19)$$

eq 4.8 MugSeiSpa

Proof. We fix $\frac{1}{2} < \beta < \beta' < \alpha$ and $0 < R < h^{\frac{1}{2}-\beta}/c_\psi$. Testing the minimality of $E_s^{(h)}$, $s = t, t-h$, we find

$$P_\phi(E_s^{(h)}, B_{Rh^\beta}(x_0)) \leq P_\phi(G, B_{Rh^\beta}(x_0)) + \frac{1}{h} \int_{G \Delta E_s^{(h)}} |\text{sd}_{E_{s-h}^{(h)}}^\psi| + \int_{G \Delta E_s^{(h)}} |F_h|, \quad (3.20)$$

eq 4.9 MugSeiSpa

for any set G of finite perimeter such that $G \Delta E_s^{(h)} \subseteq B_{Rh^\beta}(x_0)$. Using Lemma 3.10, the 1-Lipschitz regularity of sd^ψ and (3.18), we deduce $|v_h(s, y)| \leq c_\psi R h^{\beta-1} + c_\infty h^{-1/2} \leq (1+c_\infty)h^{-1/2}$ for any $y \in B_{Rh^\beta}(x_0) \cap (E_s^{(h)} \Delta F)$. Plugging this inequality in (3.20), we find

$$P_\phi(E_s^{(h)}, B_{Rh^\beta}(x_0)) \leq P_\phi(G, B_{Rh^\beta}(x_0)) + \frac{1+c}{\sqrt{h}} |F \Delta E_s^{(h)}| + \|f\|_\infty |G \Delta E_s^{(h)}|. \quad (3.21)$$

eq 4.10 MugSeiSpa

We then introduce the blown-up sets for $s = t, t-h$, defined as

$$E_s^{(h),\beta} := h^{-\beta} (E_s^{(h)} - x_0).$$

Rescaling equation (3.21), we easily find that $E_s^{(h),\beta}$ is a (Λ_h, r_h) -minimizer of the $\phi(x_0 + h^\beta \cdot, \cdot)$ -perimeter, with $\Lambda_h = (1+c)h^{\beta-1/2}$, $r_h = h^{1/2-\beta}$. Moreover, scaling the density estimates (3.4) we have a uniform bound on the perimeters of the sets $E_s^{(h),\beta}$ in each ball B_R . By compactness, there exist two sets E_1^β, E_2^β such that

$$E_t^{(h),\beta} \rightarrow E_1^\beta, \quad E_{t-h}^{(h),\beta} \rightarrow E_2^\beta \quad \text{in } L_{loc}^1.$$

Then, by scaling and (3.18) we find

$$|\text{sd}_{E_{t-h}^{(h),\beta}}^\psi(\cdot)| \leq h^{\alpha-\beta} \quad \text{on } B_{h^{1/2-\beta}}(0) \cap (E_t^{(h),\beta} \Delta E_{t-h}^{(h),\beta}),$$

thus we easily conclude that $E^\beta := E_1^\beta = E_2^\beta$. Moreover, by closeness of Λ_h -minimizers under L_{loc}^1 -convergence (see e.g. [19, Theorem 2.9]), one can see that E^β is a local $\phi(x_0, \cdot)$ -perimeter minimizer. Thus, by complete regularity, it is a smooth C^2 set. We can then employ the classic blow-up theorem to deduce that, for a fixed $\beta' \in (\beta, \alpha)$, the blow-up $h^{-(\beta'-\beta)} E^\beta$ converges to a half-space $\mathbb{H} = \{x \cdot \nu \leq 0\}$ as $h \rightarrow 0$. Moreover, the blow-ups

$$E_s^{(h),\beta'} := \frac{E_s^{(h)} - x_0}{h^{\beta'}}$$

admit a converging subsequence by compactness of sets of finite perimeter and by rescaling equation (3.21). Thus, the previous Lemma 3.14 implies

$$E_s^{(h),\beta'} \rightarrow \mathbb{H} \quad \text{in } L_{loc}^1$$

as $h \rightarrow 0$. To conclude, the ε -regularity Theorem for Λ -minimizers (see e.g. [19, Theorem 3.1]) ensures that $E_s^{(h),\beta'}$ are uniformly $C^{1,\frac{1}{2}}$ sets in $B_1(0)$ for $s = t, t - h$ as $h \rightarrow 0$. \square

We recall here an approximation result proved in [33] (see also [35] for a more detailed proof). We remark that the proof of this result is purely geometric and does not rely on the variational problem satisfied by the sets $E_t^{(h)}, E_{t-h}^{(h)}$.

Corollary (Corollary 4.3 in [35]). *Under the hypotheses of Lemma 3.15, fix $0 < \beta < \alpha$ and let $\mathbf{C}_{h,\beta}$ be the open cylinder defined as*

$$\mathbf{C}_{h,\beta}(x_0, \nu) := \left\{ x \in \mathbb{R}^N : |(x - x_0) \cdot \nu| < \frac{h^\beta}{2}, \left| (x - x_0) - ((x - x_0) \cdot \nu) \nu \right| < \frac{h^\beta}{2} \right\}.$$

Then, it holds

$$\begin{aligned} & \left| \int_{\mathbf{C}_{h,\beta/2}(x_0, \nu)} (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) dx - \int_{\partial E_t^{(h)} \cap \mathbf{C}_{h,\beta/2}(x_0, \nu)} \text{sd}_{E_{t-h}^{(h)}} d\mathcal{H}^{N-1} \right| \\ & \leq \omega(h) \int_{\mathbf{C}_{h,\beta/2}(x_0, \nu)} |\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}|. \end{aligned}$$

Carefully inspecting the proof, one indeed proves that there exists a geometric constant C such that for any $y \in B_{h^\beta/2}^{N-1}(x_0)$

$$|\text{sd}_{E_{t-h}^{(h)}}(y, f_t^{(h)}(y)) \sqrt{1 + |\nabla f_t^{(h)}(y)|^2} - (f_t^{(h)}(y) - f_{t-h}^{(h)}(y))| \leq C\omega(h)|f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|, \quad (3.22)$$

computation sd

where we set

$$\partial E_s^{(h)} \cap \mathbf{C} = \{(y, f_s^{(h)}(y)) \in \mathbb{R}^{N-1} \times \mathbb{R}, |y| \leq h^\beta/2\},$$

for $s = t, t - h$.

We briefly recall some classical results. Consider an anisotropy ψ , independent of the position. It is well-known that, for any bounded, C^2 set $G \subseteq \mathbb{R}^N$, setting sd^ψ as the distance induced by ψ° , then the gradient of sd^ψ exists almost everywhere (and everywhere in a neighborhood of the set G) and satisfies the following relations: the first one is known as the eikonal equation

$$\psi(\nabla \text{sd}^\psi) = 1 \quad \text{a.e.},$$

the second one is a useful representation formula, holding at every point $x \notin G$ having a unique point on $x' \in G$ of minimal distance, which reads

$$\nabla \text{sd}_G^\psi(x) = \frac{\nu_G(x')}{\psi(\nu_G(x'))}.$$

Moreover, in this particular case, in the definition of dist^ψ we can consider just straight lines as follows from a simple application of Jensen's inequality: for any curve γ as in the definition of dist^ψ , we have

$$\int_0^1 \psi^\circ(\dot{\gamma}(t)) dt \geq \psi^\circ\left(\int_0^1 \dot{\gamma}\right) = \psi^\circ(y - x).$$

Proposition 3.16 (Estimate on almost flat sets). *Under the hypotheses of Lemma 3.15, fix $\beta \in (0, \alpha)$ and let \mathbf{C}_{h^β} be the open cylinder defined as*

$$\mathbf{C}_{h^\beta}(x_0, \nu) := \left\{ x \in \mathbb{R}^N : |(x - x_0) \cdot \nu| < \frac{h^\beta}{2}, \left| (x - x_0) - ((x - x_0) \cdot \nu) \nu \right| < \frac{h^\beta}{2} \right\}.$$

Then, it holds

$$\begin{aligned} & \left| \int_{\mathbf{C}_{h^\beta/2}(x_0, \nu)} (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) dx - \int_{\partial E_t^{(h)} \cap \mathbf{C}_{h^\beta/2}(x_0, \nu)} \psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_{t-h}^{(h)}}^\psi d\mathcal{H}^{N-1} \right| \\ & \leq \omega(h) \int_{\mathbf{C}_{h^\beta/2}(x_0, \nu)} |\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}|. \end{aligned}$$

Proof. From the previous Lemma 3.15 we know that, for h suitably small, both $\partial E_t^{(h)}$ and $\partial E_{t-h}^{(h)}$ in $\mathbf{C}_{h^\beta/2}(x_0, \nu)$ can be written as graphs of functions of class $C^{1, \frac{1}{2}}$. Up to a change of coordinates, we can assume wlog that $x_0 = 0, \nu = e_N$. For simplicity, we set $\mathbf{C} = \mathbf{C}_{h^\beta/2}(0, e_N)$. We thus find

$$\partial E_s^{(h)} \cap \mathbf{C} = \{(y, f_s^{(h)}(y)) \in \mathbb{R}^{N-1} \times \mathbb{R}, |y| \leq h^\beta/2\}$$

for $s = t, t - h$, where $f_s^{(h)} : B_{h^\beta/2}^{N-1} \rightarrow \mathbb{R}$ are $C^{1, \frac{1}{2}}$ functions with

$$\|\nabla f_s^{(h)}\|_{L^\infty(B_{h^\beta/2})} \leq \omega(h).$$

We want to prove the following slightly stronger pointwise inequality: namely, that for any point $x = (y, f_t^{(h)}(y)) \in \partial E_t^{(h)} \cap \mathbf{C}$, it holds

$$\left| \text{sd}_{E_{t-h}^{(h)}}^\psi(x) \psi(x, \nu_{E_t^{(h)}}(x)) \sqrt{1 + |\nabla f_t^{(h)}(y)|} - (f_t^{(h)}(y) - f_{t-h}^{(h)}(y)) \right| \leq \omega(h) |f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|. \quad (3.23)$$

estimate flatness

Integrating the previous inequality over \mathbf{C} yields the thesis. Clearly, it is enough to prove (3.23) at each point x such that $|\text{sd}_{E_{t-h}^{(h)}}^\psi(x)| > 0$. We thus fix $x = (y, f_t^{(h)}(y)) \in \partial E_t^{(h)} \cap \mathbf{C}$ and denote by $x' := (y, f_{t-h}^{(h)}(y))$. It can be assumed without loss of generality that $x \notin E_{t-h}^{(h)}$, as the other case is analogous. Considering the anisotropy $\psi(x', \cdot)$, we denote by sd'_G the geodesic distance function from the set G induced by the anisotropy above. We proceed estimating $|\text{sd}'_{E_{t-h}^{(h)}}(x) - \text{sd}_{E_{t-h}^{(h)}}^\psi(x)|$.

Let γ be a smooth curve, with $\gamma(0) = x, \gamma(1) \in \partial E_{t-h}^{(h)}$ to be used in the definition of the geodesic distance $\text{sd}_{E_{t-h}^{(h)}}^\psi$. Firstly, we remark that one could assume

$$\gamma([0, 1]) \subseteq B(x, 2c_\psi^2 |f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|) \quad (3.24)$$

bound p

Indeed, if it were not the case, the lower bounds contained in (2.3) and (3.22) allow us to estimate

$$\int_0^1 \psi^\circ(\gamma, \dot{\gamma}) dt \geq \frac{1}{c_\psi} \int_0^1 |\dot{\gamma}| dt \geq 2c_\psi |f_t^{(h)}(y) - f_{t-h}^{(h)}(y)| \geq 2c_\psi \text{sd}_{E_{t-h}^{(h)}}(x) \geq 2 \text{sd}_{E_{t-h}^{(h)}}^\psi(x), \quad (3.25)$$

estimate curve

a contradiction for h small. We can reason analogously for $\text{sd}'_{E_{t-h}^{(h)}}$. In particular, we can consider just curves having length $\int_0^1 |\dot{\gamma}| \leq c|f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|$. Therefore, we obtain (by homogeneity)

$$\begin{aligned} \text{sd}_{E_{t-h}^{(h)}}^\psi(x) &\leq \int_0^1 \psi^\circ(\gamma, \dot{\gamma}) dt \leq \int_0^1 \psi^\circ(x', \dot{\gamma}) dt + \sup_{\nu \in S^N, t \in [0,1]} |\psi(\gamma(t), \nu) - \psi(x', \nu)| \int_0^1 |\dot{\gamma}| \\ &\leq \int_0^1 \psi^\circ(x', \dot{\gamma}) dt + c\omega(h)|f_t^{(h)}(y) - f_{t-h}^{(h)}(h)|, \end{aligned}$$

and, taking the \inf_γ , we obtain $\text{sd}_{E_{t-h}^{(h)}}^\psi(x) \leq \text{sd}'_{E_{t-h}^{(h)}}(x) + \omega(h)|f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|$. The converse inequality can be proved analogously, yielding

$$|\text{sd}'_{E_{t-h}^{(h)}}(x) - \text{sd}_{E_{t-h}^{(h)}}^\psi(x)| \leq \omega(h)|f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|.$$

Therefore, in what follows we will consider always the anisotropy frozen in x' , and use sd' instead of sd^ψ . Finally, define $p \in \partial E_{t-h}^{(h)}$ as a point of minimal $\text{sd}'_{E_{t-h}^{(h)}}$ distance from x . In the following, with $\Pi_{\mathbb{H}}^v z$ we denote the projection on the hyperplane \mathbb{H} of the point z along the direction v .

Step 1. We assume that $\partial E_{t-h}^{(h)} \cap \mathbf{C}$ coincides with the hyperplane $\mathbb{H} = p + \{z \cdot \nu = 0\}$. Assume that ψ° is strictly convex. The gradient $\nabla \text{sd}'_{\mathbb{H}}$ exists for every point $z \notin \mathbb{H}$ as it has a unique point of minimal distance from \mathbb{H} (by strict convexity). By the eikonal equation we have

$$\frac{\nu}{\psi(x', \nu)} = \nabla \text{sd}'_{\mathbb{H}}(z). \quad (3.26) \quad \boxed{\text{approx 1}}$$

Since $\psi^\circ(x', \cdot)$ is positively 1-homogeneous, setting $\tilde{z} \in \mathbb{H}$ the point of minimal $\psi(x', \cdot)$ -distance from z , we obtain also

$$\nabla \text{sd}'_{\mathbb{H}}(z) \cdot (z - \tilde{z}) = \text{sd}'_{\mathbb{H}}(z). \quad (3.27) \quad \boxed{\text{approx 2}}$$

Combining (3.26), (3.27) and choosing $z = x$, we have

$$\text{sd}'_{\mathbb{H}}(x)\psi(x', \nu) = (x - \tilde{x}) \cdot \nu = |x - \Pi_{\mathbb{H}}^{\nu} x| = \text{sd}_{\mathbb{H}}(x). \quad (3.28) \quad \boxed{\text{id dis}}$$

We remark that $\text{sd}'_{\mathbb{H}}(x) = \text{sd}'_{E_{t-h}^{(h)}}(x)$ by convexity, thus we conclude (3.23) by combining (3.28) with (3.22).

The same formula holds even if ψ° is not strictly convex, as an approximation result shows. It is indeed enough to approximate ψ° with strictly convex anisotropies. One way of doing this is the following: define W^ψ as the unit Wulff shape associated to ψ and set $W_n^\psi := W^\psi + \frac{1}{n}B_1$. The anisotropies ψ_n associated to the Wulff shapes W_n^ψ are now strictly convex and converging to ψ as $n \rightarrow \infty$. Remarking that the distance functions induced by these anisotropies converge to $\text{sd}'_{\mathbb{H}}$, we pass to the limit in (3.28) to conclude.

Step 2. Set $\nu = \nu_{E_{t-h}^{(h)}}$ and consider the hyperplane $\mathbb{H} = p + \{z \cdot \nu = 0\}$. Define $w := x' - \Pi_{\mathbb{H}}(x')$. We shall prove that $|w| \leq c\omega(h)|f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|$. To see this, we start by remarking that (3.19) implies

$$|e_N - e_N(e_N \cdot \nu_{E_{t-h}^{(h)}})| \leq \omega(h) \quad \text{in } \partial E_{t-h}^{(h)} \cap \mathbf{C},$$

implying $e_N \cdot \nu_{E_{t-h}^{(h)}} \geq 1 - \omega(h)$, and thus, for any versor v tangent to $\partial E_{t-h}^{(h)} \cap \mathbf{C}$ one has $|v \cdot e_N| \leq \omega(h)$. Therefore, we have $(x' - p) \cdot e_N \leq \omega(h)|x' - p|$ and also

$$\begin{aligned} \frac{x' - p}{|x' - p|} \cdot \nu &= \frac{x' - p}{|x' - p|} \cdot (e_N(\nu \cdot e_N) + \nu - e_N(\nu \cdot e_N)) \\ &\leq \omega(h) + |\nu - e_N(\nu \cdot e_N)| = \omega(h) + (1 - |\nu \cdot e_N|^2)^{1/2} \\ &\leq 3\sqrt{\omega(h)}, \end{aligned}$$




Figure 1: The situation in the proof of the lemma.

by choosing h small. Up to defining $\sqrt{\omega}$ as ω , using the previous estimate and the bounds (3.24) we see that

$$|w| = |x' - p| \left(\frac{x' - p}{|x' - p|} \cdot \nu \right) \leq c\omega(h)|x' - p| \leq c\omega(h)|f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|. \quad (3.29) \quad \boxed{\text{bound planes}}$$

We now remark that $\text{sd}'_{E_{t-h}^{(h)}}(x) = \text{sd}'_{\mathbb{H}}(x)$ (by convexity of the anisotropy $\psi(x', \cdot)$) and so, applying the previous step to \mathbb{H} we get

$$\left| \text{sd}_{E_{t-h}^{(h)}}^\psi(x) \psi(x, \nu_{E_{t-h}^{(h)}}(x)) \sqrt{1 + |\nabla f_t^{(h)}(y)|} - |x - \Pi_{\mathbb{H}}^{e_N} x| \right| \leq \omega(h)|x - \Pi_{\mathbb{H}}^{e_N} x|.$$

We conclude (3.23) by estimating

$$\left| |x - \Pi_{\mathbb{H}}^{e_N} x| - |x - x'| \right| \leq |x' - \Pi_{\mathbb{H}}^{e_N} x| = |w|/|\nu \cdot e_N| \leq c \frac{\omega(h)}{1 - \omega(h)} |f_t^{(h)}(y) - f_{t-h}^{(h)}(y)|,$$

where we used (3.29). We conclude the proof by a simple change of coordinates and using (3.23) to find

$$\begin{aligned} &\left| \int_{\partial E_t^{(h)} \cap \mathbf{C}} \psi(x, \nu_{E_t^{(h)}}(x)) \text{sd}_{E_{t-h}^{(h)}}^\psi(x) \, d\mathcal{H}^{N-1} - \int_{B_{h,\beta/2}} f_t^{(h)}(y) - f_{t-h}^{(h)}(y) \, dy \right| \\ &= \left| \int_{B_{h,\beta/2}} \psi((y, f_t^{(h)}(y)), \nu_{E_t^{(h)}}(y, f_t^{(h)}(y))) \text{sd}_{E_{t-h}^{(h)}}^\psi(y, f_t^{(h)}(y)) \sqrt{1 + |\nabla f_t^{(h)}(y)|^2} - (f_t^{(h)}(y) - f_{t-h}^{(h)}(y)) \, dy \right| \\ &\leq c\omega(h) \int_{B_{h,\beta/2}} |f_t^{(h)} - f_{t-h}^{(h)}| \, dy. \end{aligned}$$

□

Finally, we are able to prove that the error generated approximating the discrete velocity with v_h goes to zero as $h \rightarrow 0$.

osition 2.2 LucStu

Proposition 3.17 (Error estimate). *Under the hypothesis of Lemma 3.15, the error in the discrete curvature equation vanishes in the limit $h \rightarrow 0$, namely*

$$\lim_{h \rightarrow 0} \left| \frac{1}{h} \int_0^T \left(\int_{E_t^{(h)}} \eta \, dx - \int_{E_{t-h}^{(h)}} \eta \, dx \right) dt - \int_0^T \int_{\partial E_t^{(h)}} \psi(x, \nu_{E_t^{(h)}}) v_h \eta \, d\mathcal{H}^{N-1}(x) \, dt \right| = 0 \quad (3.30)$$

error estimate on

for all $\eta \in C_c^1(\mathbb{R}^N \times [0, T])$.

Proof. We fix $t \in [2h, \infty)$ and $\alpha \in (\frac{1}{2}, \frac{N+2}{2N+2})$. For any point $x_0 \in \partial E_t^{(h)}$ we define the open set A_{x_0} defined as follows:

- (i) if (3.18) holds, we set $A_{x_0} = \mathbf{C}_{h^\beta/2}(x_0, \nu)$, with the notations of Corollary 3.16;
- (ii) otherwise we set $A_{x_0} = B(x_0, c_\infty \sqrt{h})$, where c_∞ is the constant of Lemma 3.10.

By Lemma 3.10, the family $\{A_{x_0} : x_0 \in \partial E_t^{(h)}\}$ is a covering of $E_t^{(h)} \triangle E_{t-h}^{(h)}$. By a simple application of Besicovitch's theorem (see e.g. [34]), we find a finite collection of points $I \subseteq \partial E_t^{(h)}$ such that $\{A_{x_0}\}_{x_0 \in I}$ is a covering of $E_t^{(h)} \triangle E_{t-h}^{(h)}$ with the finite intersection property. We proceed to estimate (3.30) on each A_{x_0} belonging to this family.

Estimate in case (i) We use Proposition 3.16 to deduce

$$\begin{aligned} & \left| \int_{A_{x_0}} (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) \eta \, dx - \int_{\partial E_t^{(h)} \cap A_{x_0}} \psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_{t-h}^{(h)}}^\psi \eta \, d\mathcal{H}^{N-1} \right| \\ & \leq |\eta(x_0, t)| \left| \int_{A_{x_0}} (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) - \int_{\partial E_t^{(h)} \cap A_{x_0}} \psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_{t-h}^{(h)}}^\psi \, d\mathcal{H}^{N-1} \right| \\ & + \left| \int_{A_{x_0}} (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) (\eta - \eta(x_0, t)) - \int_{\partial E_t^{(h)} \cap A_{x_0}} (\eta - \eta(x_0, t)) \psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_{t-h}^{(h)}}^\psi \, d\mathcal{H}^{N-1} \right| \\ & \leq C(\omega(h) \|\eta\|_\infty + h^\beta \|\nabla \eta\|_\infty) \int_{A_{x_0}} |\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}| \, d\mathcal{H}^{N-1} + ch^\beta \|\nabla \eta\|_\infty P(E_t^{(h)}, A_{x_0}). \end{aligned} \quad (3.31)$$

eq 4.13 MugSeiSp

Estimate in case (ii) By assumption $\exists y \in B_{c_\infty \sqrt{h}}(x_0) \cap (E_t^{(h)} \triangle E_{t-h}^{(h)})$ such that $|v_h(t, y)| > h^{\alpha-1}$. We can assume wlog $y \in E_t^{(h)}$. We then have $B(y, h^\alpha/(2c_\psi)) \subseteq \mathbb{R}^N \setminus E_{t-h}^{(h)}$ and $\text{sd}_{E_{t-h}^{(h)}}^\psi > h^\alpha/(2c_\psi^2)$ on $B(y, h^\alpha/(2c_\psi))$. Since $h^\alpha \ll h^{1/2}$, we can use the density estimates of Lemma 3.3 to deduce

$$ch^{(N+1)\alpha-1} \leq \int_{B(y, h^\alpha/(2c_\psi)) \cap (E_t^{(h)} \triangle E_{t-h}^{(h)})} |v_h| \, dx.$$

Analogously, recalling also Lemma 3.10, we deduce

$$\int_{B(x_0, c_\infty \sqrt{h}) \cap \partial E_t^{(h)}} |\psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_{t-h}^{(h)}}^\psi| \, d\mathcal{H}^{N-1}(x) \leq ch^{\frac{N}{2}}.$$

Combining the two previous equations and $B(y, h^\alpha/(2c_\psi)) \subseteq B(y, c\sqrt{h})$, we infer

$$\begin{aligned} & \int_{A_{x_0}} |\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}| + \int_{A_{x_0} \cap \partial E_t^{(h)}} |\psi(x, \nu_{E_{t-h}^{(h)}}) \text{sd}_{E_{t-h}^{(h)}}^\psi| d\mathcal{H}^{N-1} \\ & \leq ch^{\frac{N}{2} - (N+1)\alpha + 1} \int_{A_{x_0} \cap (E_t^{(h)} \Delta E_{t-h}^{(h)})} |\psi(x, \nu_{E_{t-h}^{(h)}}) v_h|. \end{aligned} \quad (3.32)$$

eq 4.16 MugSeiSpa

Summing over $x_0 \in I$ both (3.31) and (3.32), and using the local finiteness of the covering, we get

$$\begin{aligned} & \left| \int (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) \eta dx - \int_{\partial E_t^{(h)}} \psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_t^{(h)}}^\psi \eta d\mathcal{H}^{N-1} \right| \\ & \leq \sum_{x_0 \in I} \left| \int_{A_{x_0}} (\chi_{E_t^{(h)}} - \chi_{E_{t-h}^{(h)}}) \eta dx - \int_{\partial E_t^{(h)} \cap A_{x_0}} \psi(x, \nu_{E_t^{(h)}}) \text{sd}_{E_t^{(h)}}^\psi \eta d\mathcal{H}^{N-1} \right| \\ & \leq c \left(\omega(h) \|\eta\|_\infty + h^\beta \|\nabla \eta\|_\infty + h^{\frac{N}{2} - (n+1)\alpha + 1} \|\eta\|_\infty \right) \\ & \quad \cdot \left(P(E_t^{(h)}) + |E_t^{(h)} \Delta E_{t-h}^{(h)}| + \int_{E_t^{(h)} \Delta E_{t-h}^{(h)}} |v_h| \right) \end{aligned}$$

where the last constant c depends on N, ψ . We then use Lemma 3.11, (3.12) and (3.13) to conclude

$$\begin{aligned} & \left| \int_{2h}^T \frac{1}{h} \left(\int_{E_t^{(h)}} \eta dx - \int_{E_{t-h}^{(h)}} \eta dx \right) - \int_h^T \int_{\partial E_t^{(h)}} \psi(x, \nu_{E_t^{(h)}}) v_h \eta d\mathcal{H}^{N-1} \right| \\ & \leq c \left(\omega(h) \|\eta\|_\infty + h^\beta \|\nabla \eta\|_\infty + h^{\frac{N}{2} - (n+1)\alpha + 1} \|\eta\|_\infty \right), \end{aligned}$$

where $c = c(E_0, f, T, \psi)$ and T is chosen such that $\text{spt } \eta \subseteq \mathbb{R}^N \times [0, T]$. The conclusion follows using the definition of α and taking the limit $h \rightarrow 0$. \square

The proof of our main theorem of this section is now a consequence of the previous results. In particular, hypothesis (3.14) and (3.15) imply that the discrete flow converges to the flat flow in the sense of varifolds and this allows to prove (1.7), while (1.8) is a consequence of Proposition 3.17. In order to prove the convergence of the approximations in time of the forcing term, we need to require additionally that (1.5) holds.

Proof of Theorem 1.2. Firstly, combining [29, Theorem 4.4.2] with the bounds contained in (3.17) and in Proposition 3.13, we conclude the existence of functions $v, H^\phi, \tilde{f} : \mathbb{R}^N \times [0, \infty) \rightarrow \mathbb{R}$ satisfying

$$\int_0^T \int_{\partial E_t} |v|^2 + |H^\phi|^2 + |\tilde{f}|^2 d\mathcal{H}^{N-1} dt \leq C_T$$

and the following properties

$$\lim_k \int_0^T \int_{\partial E_t^{(h_k)}} v_{h_k} \eta d\mathcal{H}^{N-1} dt = \int_0^T \int_{\partial E_t} \eta v d\mathcal{H}^{N-1} dt$$

$$\begin{aligned}
\lim_k \int_0^T \int_{\partial E_t^{(h_k)}} F_{h_k}(x, t) \eta \, d\mathcal{H}^{N-1} \, dt &= \int_0^T \int_{\partial E_t} \eta \tilde{f} \, d\mathcal{H}^{N-1} \, dt \\
\lim_k \int_0^T \int_{\partial E_t^{(h_k)}} H_{E_t^{(h_k)}}^\phi \eta \, d\mathcal{H}^{N-1} \, dt &= \int_0^T \int_{\partial E_t} \eta H^\phi \, d\mathcal{H}^{N-1} \, dt,
\end{aligned} \tag{3.33}$$

for any $\eta \in C_c^0(\mathbb{R}^N \times [0, T])$. We now employ an approximation procedure to prove that $H^\phi(\cdot, t)$ is the ϕ -mean curvature of E_t for a.e. $t \in [0, \infty)$, following the lines of [33, 35]. Fixed $t \in [0, +\infty)$ and $\varepsilon > 0$, set ν_ε a continuous function such that $\int_{\partial E_t} (\nu_{E_t} - \nu_\varepsilon)^2 \, d\mathcal{H}^{N-1} < \varepsilon$. Then, by (3.14) one could prove that $\lim_{k \rightarrow \infty} \int_{\partial E_t^{(h_k)}} (\nu_{E_t^{(h_k)}} - \nu_\varepsilon)^2 \, d\mathcal{H}^{N-1} < \varepsilon$. Considering test functions in (3.33) of the form $\eta(x, t) = a(t)g(x)$, one has for a.e. $t \in [0, +\infty)$

$$\lim_k \int_{\partial E_t^{(h_k)}} H_{E_t^{(h_k)}}^\phi g \, d\mathcal{H}^{N-1} = \int_{\partial E_t} H^\phi g \, d\mathcal{H}^{N-1}.$$

Thus, for a.e. $t \in [0, +\infty)$ and for any $X \in C_c^0(\mathbb{R}^N; \mathbb{R}^N)$ it holds

$$\lim_k \int_{\partial E_t^{(h_k)}} H_{E_t^{(h_k)}}^\phi \nu_{E_t^{(h_k)}} \cdot X \, d\mathcal{H}^{N-1} = \int_{\partial E_t} H^\phi \nu_{E_t} \cdot X \, d\mathcal{H}^{N-1}$$

by approximating the normal vectors of $E_t^{(h_k)}$ with ν_ε . Furthermore, by the convergence (3.14) and the hypothesis (3.15) we can use the Reshetnyak's continuity theorem (see e.g. [4, Theorem 2.39]), ensuring

$$\int_{\partial E_t^{(h_k)}} L(x, \nu_{E_t^{(h_k)}}) \, d\mathcal{H}^{N-1} \rightarrow \int_{\partial E_t} L(x, \nu_{E_t}) \, d\mathcal{H}^{N-1}$$

as $k \rightarrow \infty$, for any $L \in C_c^0(\mathbb{R}^N \times \mathbb{R}^N)$. We choose $L(x, \nu) = \operatorname{div}_\phi X$ for some $X \in C_c^1(\mathbb{R}^N; \mathbb{R}^N)$ to obtain

$$\begin{aligned}
\int_{\partial E_t} \operatorname{div}_\phi X \, d\mathcal{H}^{N-1} &= \lim_k \int_{\partial E_t^{(h_k)}} \operatorname{div}_\phi X \, d\mathcal{H}^{N-1} \\
&= \lim_k \int_{\partial E_t^{(h_k)}} X \cdot \nu_{E_t^{(h_k)}} H_{E_t^{(h_k)}}^\phi \, d\mathcal{H}^{N-1} \\
&= \int_{\partial E_t} X \cdot \nu_{E_t} H^\phi \, d\mathcal{H}^{N-1},
\end{aligned}$$

which shows that $H^\phi(\cdot, t)$ is the ϕ -mean curvature of the set E_t for a.e. $t \in [0, +\infty)$. Moreover, we remark that $F_{h_k}(x, t) \rightarrow f(x, t)$ for every (x, t) , thus for any test function $\eta \in C_c^0(\mathbb{R}^N \times [0, +\infty))$ and $t \in [0, +\infty)$ we have

$$\begin{aligned}
\left| \int_{\partial E_t^{(h_k)}} F_{h_k}(x, t) \eta(x, t) \, d\mathcal{H}_x^{N-1} - \int_{\partial E_t} f \eta \, d\mathcal{H}_x^{N-1} \right| &\leq \left| \int_{\partial E_t^{(h_k)}} F_{h_k} \eta - \int_{\partial E_t} F_{h_k} \eta \right| + \int_{\partial E_t} |F_{h_k} - f| \eta \\
&\leq \|f\|_\infty \|\eta\|_\infty \left(P(E_t^{(h_k)}) - P(E_t) \right) + \int_{\partial E_t} |F_{h_k} - f| \eta \rightarrow 0
\end{aligned}$$

applying the dominated convergence theorem and recalling Lemma 3.9. Thus, $\tilde{f} = f$. We then prove (1.7) by passing to the limit in the Euler-Lagrange equation (3.2).

To prove (1.8) we employ Proposition 3.17: for every $\eta \in C_c^0(\mathbb{R}^N \times [0, T])$, by a change of variables we have that

$$\int_h^T \left[\int_{E_t^{(h)}} \eta \, dx - \int_{E_{t-h}^{(h)}} \eta \, dx \right] dt = \int_h^T \int_{E_t^{(h)}} (\eta(x, t) - \eta(x, t-h)) \, dx \, dt - h \int_{E_0} \eta \, dx$$

where we have used that $E_t^{(h)} = E_0$ for $t \in [0, h)$. Therefore, a simple convergence argument yields

$$\lim_{h \rightarrow 0} \frac{1}{h} \int_h^T \left[\int_{E_t^{(h)}} \eta \, dx - \int_{E_{t-h}^{(h)}} \eta \, dx \right] dt = - \int_h^T \partial_t \eta(x, t) \, dx \, dt - \int_{E_0} \eta.$$

Combining the previous estimate with Proposition 3.17 and passing to the limit, we obtain (1.8). \square

4 Viscosity solutions

In this section we will prove the existence of another weak notion of solution for the mean curvature flow starting from a compact set. We will follow the so-called level set approach based on the theory of viscosity solution. we recall that in the first part we work with the standing assumptions of the paper (H). Additionally, we require (1.4).

4.1 The discrete scheme for unbounded sets

In this short subsection we will define the discrete evolution scheme for unbounded sets having compact boundary. The idea would be to define this evolution simply as the complement of the evolution of the complementary set, but since the anisotropies we are considering are not symmetric, we need additional care.

We recall that, given an anisotropy ϕ , we define $\tilde{\phi}(x, \nu) := \phi(x, -\nu)$. This anisotropy has all the properties of the original one, concerning regularity and bounds. We start remarking the following simple fact. One can see that $\text{dist}^\psi(x, y) = \text{dist}^{\tilde{\psi}}(y, x)$, since for any curve $\gamma \in W^{1,1}([0, 1]; \mathbb{R}^N)$, $\gamma(0) = x, \gamma(1) = y$, a simple change of variable yields

$$\int_0^1 \psi^\circ(\gamma(t), \dot{\gamma}(t)) \, dt = \int_0^1 \psi^\circ \left(\gamma(1-t), -\frac{d}{dt}(\gamma(1-t)) \right) \, dt = \int_0^1 \widetilde{(\psi^\circ)}(\eta(t), \dot{\eta}(t)) \, dt,$$

for $\eta(t) = \gamma(1-t)$, once one sees that

$$\widetilde{(\psi^\circ)}(\cdot, \nu) = \sup_{\psi(\cdot, \xi) \leq 1} \xi \cdot (-\nu) = \sup_{\tilde{\psi}(\cdot, -\xi) \leq 1} (-\xi) \cdot \nu = (\tilde{\psi})^\circ(\cdot, \nu).$$

Therefore, by definition of signed distance we have

$$\text{sd}_E^\psi(x) = -\text{sd}_{E^c}^{\tilde{\psi}}(x). \quad (4.1)$$

invers dist

For every compact set F and $h > 0, t \geq 0$, we will denote by $\tilde{T}_{h,t}^\pm F$ the maximal and the minimal solution to problem (3.1), according to Lemma 3.1 with P_ϕ and sd^ψ , respectively, replaced by P_ϕ^- and $\text{sd}^{\tilde{\psi}}$. Finally, for every set E with compact boundary we define

$$T_{h,t}^\pm E := \left(\tilde{T}_{h,t}^\mp E^c \right)^c. \quad (4.2)$$

pb discr compl

As in the case for compact sets, we set $T_{h,t}E := T_{h,t}^-E$. Given an open, unbounded set E_0 having compact boundary, we can then define the discrete flow $\{E_t^{(h)}\}_{t \geq 0}$ as follows: $E_t^{(h)} := E_0$ for $t \in [0, h)$ and

$$E_t^{(h)} = T_{h,t}E_{t-h}^{(h)}, \quad \forall t \in [h, +\infty).$$

One easily checks that analogous results to Lemmas 3.2, 3.9 and 3.8 hold also for this problem. We state the corresponding results.

principle, unbounded

Lemma 4.1. *Let $F_1 \subseteq F_2$ be open, unbounded sets with compact boundary and fix $h > 0, t \geq 0$. Then, $T_{h,t}F_1 \subseteq T_{h,t}F_2$.*

Lemma 4.2. *For any $T > 0$ there exists a constant $C_T(\phi, \psi, f, T)$ such that for every $R > 0$ the following holds. If the initial open set $E \supset B_R^c$, then $E_t^{(h)} \supset B_{C_T R}^c$ for all $t \in [0, T]$.*

Lemma 4.3. *For every $R_0 > 0$ there exist $h_0(R_0) > 0$ and $C(R_0, \phi, \psi, f) > 0$ with the following property: For all $R \geq R_0, h \in (0, h_0), t > 0$ and $x \in \mathbb{R}^N$ one has*

$$T_{h,t}((B_R(x))^c) \subseteq (B_{R-Ch}(x))^c.$$

We now state a comparison principle between bounded and unbounded sets, following the line of [14, Lemma 6.10].

bounded-unbounded

Lemma 4.4. *Let E_1 be a compact set and let E_2 be an open, unbounded set, with compact boundary, and such that $E_1 \subseteq E_2$. Then, for every $h \in (0, 1), t \geq 0$ it holds $T_{h,t}^\pm E_1 \subseteq T_{h,t}^\pm E_2$.*

Proof. We fix $h \in (0, 1), t \in [0, T]$ for $T > 0$. Set $R > 0$ such that $E_1, E_2^c \subseteq B_R$ and note that by Lemmas 3.2 and 3.9 (applied to P_ϕ^- instead of P_ϕ) we get

$$(T_{h,t}^+ E_2)^c \subseteq \tilde{T}_{h,t}^- E_2^c \subseteq T_{h,t}^- B_R \subseteq B_{C_T R}, \quad (4.3)$$

eq 6.9 Nonlocal

for some $C_T(\phi, \psi, f, T)$. Since $\tilde{T}_{h,t}^- E_2^c$ is the minimal solution of

$$\min \left\{ P_\phi^-(E) + \frac{1}{h} \int_E \text{sd}_{E_2^c}^\psi(x) dx - \int_E F_h(x, t) dx \right\},$$

considering the change of variables $\tilde{E} = E^c$ and using (4.1), we easily conclude that $T_{h,t}^+ E_2 = (\tilde{T}_{h,t}^- E_2^c)^c$ is the maximal solution of

$$\min \left\{ P_\phi(\tilde{E}) + \frac{1}{h} \int_{B_{C_T R}} \text{sd}_{E_2}^\psi - \frac{1}{h} \int_{\tilde{E}^c} \text{sd}_{E_2}^\psi - \int_{\tilde{E}^c} F_h(x, t) dx \right\} - \frac{1}{h} \int_{B_{C_T R}} \text{sd}_{E_2}^\psi.$$

we then note that

$$\int_{B_{C_T R}} \text{sd}_{E_2}^\psi = \int_{\tilde{E}} \text{sd}_{E_2}^\psi \chi_{B_{C_T R}} + \int_{\tilde{E}^c} \text{sd}_{E_2}^\psi,$$

for every \tilde{E} such that $\tilde{E}^c \subseteq B_{C_T R}$. By (4.3), we conclude that $T_{h,t}^+ E_2$ is the maximal solution of

$$\min \left\{ P_\phi(\tilde{E}) + \frac{1}{h} \int_{\tilde{E}} \text{sd}_{E_2}^\psi \chi_{B_{C_T R}} - \int_{\tilde{E}^c} F_h(x, t) dx : \tilde{E} \subseteq B_{C_T R} \right\}. \quad (4.4)$$

eq 6.10 Nonlocal

Analogously, one proves that $T_{h,t}^- E_2$ is the minimal solution of (4.4). Finally, we remark that $\text{sd}_{E_2}^\psi \chi_{B_{C_T R}} \leq \text{sd}_{E_1}^\psi$ and that $T_{h,t}^\pm E_1 \cup T_{h,t}^\pm E_2, T_{h,t}^\pm E_1 \cap T_{h,t}^\pm E_2$ are both admissible competitors for (4.4), one argues exactly as in the proof of Lemma 3.2 to conclude $T_{h,t}^\pm E_1 \subseteq T_{h,t}^\pm E_2$. \square

4.2 The level set approach

Consider a function $u : \mathbb{R}^N \times [0, +\infty) \rightarrow \mathbb{R}$ whose superlevel set evolve according to the mean curvature equation

$$V(x, t) = -\psi(x, \nu_{\{u(\cdot, t) \geq s\}}) \left(H_{\{u(\cdot, t) \geq s\}}^\phi(x) - f(x, t) \right).$$

The function u then satisfies (recalling that $-\nabla u/|\nabla u|$ is the *outer normal vector* to the superlevel set $\{u(\cdot, t) \geq u(x, t)\}$) the equation

$$\begin{aligned} \partial_t u &= |\nabla u| V(x) = -\psi(x, -\nabla u) \left(H_{\{u(\cdot, t) \geq u(x, t)\}}^\phi(x) - f(x, t) \right) \\ &= -\psi(x, -\nabla u) (\operatorname{div} \nabla_p \phi(x, -\nabla u) - f(x, t)) \\ &= -\psi(x, -\nabla u) \left(\sum_i \partial_{x_i} \partial_p \phi(x, -\nabla u) - \nabla_p^2 \phi(x, -\nabla u) : \nabla^2 u - f(x, t) \right) \\ &:= -\psi(x, -\nabla u) (H(x, \nabla u, \nabla^2 u) - f(x, t)), \end{aligned}$$

where we defined the Hamiltonian $H : \mathbb{R}^N \times \mathbb{R}^N \setminus \{0\} \times S^N \rightarrow \mathbb{R}$ as

$$H(x, p, X) := \sum_i \partial_{x_i} \partial_p \phi(x, -p) - \nabla_p^2 \phi(x, -p) : X, \quad (4.5)$$

def hamiltonian

and S^N denotes the space of the symmetric matrices of dimension $N \times N$. We therefore focus on solving the parabolic Cauchy problem

$$\begin{cases} \partial_t u + \psi(x, -\nabla u) (H(x, \nabla u, \nabla^2 u) - f(x, t)) = 0 \\ u(\cdot, t) = u_0. \end{cases} \quad (4.6)$$

Hamilton-Jacobi e

The appropriate setting for this type of geometric evolution equations is the one of viscosity solutions, in the framework of [24, 31] (see also [14]). Before giving the definition of viscosity solutions in this setting, we need to introduce an appropriate class of test functions. To do so, we introduce an auxiliary class of functions whose properties can be found in [31] (see also [14, Section 2.2]).

family F

Definition 4.5. The family \mathcal{F} is composed of smooth functions $g \in C_c^\infty([0, +\infty))$ satisfying $g(0) = g'(0) = g''(0) = 0$, $g''(r) > 0$ in a neighborhood of 0, g constant in $(0, M)^c$ for some $M > 0$ (depending on g), and

$$\lim_{r \rightarrow 0} \frac{g'(r)}{r} = 0.$$

We will focus on the evolution of sets with compact boundary on compact time intervals of the form $[0, T]$. We now define the notion of admissible test function. With a slight abuse of notation, in the following we will say that a function is spatially constant outside a compact set even if the value of such constant is time-dependent.

def visco sol

Definition 4.6. Let $\hat{z} = (\hat{x}, \hat{t}) \in \mathbb{R}^N \times (0, T)$ and let $A \subseteq (0, T)$ be any open interval containing \hat{t} . We will say that $\eta \in C^0(\mathbb{R}^N \times \bar{A})$ is admissible at the point \hat{z} if it is of class C^2 in a neighborhood of \hat{z} , if it is constant out of a compact set, and, in case $\nabla \eta(\hat{z}) = 0$, the following holds: there exists $f \in \mathcal{F}$ and $\omega \in C^\infty([0, \infty))$ with $\omega'(0) = 0, \omega(r) > 0$ for $r \neq 0$ and such that

$$|\eta(x, t) - \eta(\hat{z}) - \eta_t(\hat{z})(t - \hat{t})| \leq f(|x - \hat{x}|) + \omega(|t - \hat{t}|),$$

for all $(x, t) \in \mathbb{R}^N \times A$.

One can then define the following notion of viscosity solution.

Definition 4.7. An upper semicontinuous function $u : \mathbb{R}^N \times [0, T] \rightarrow \mathbb{R}$ (in short, $u \in usc(\mathbb{R}^N \times [0, T])$), constant outside a compact set, is a viscosity subsolution of the Cauchy problem (4.6) if $u(\cdot, 0) \leq u_0$ and for all $z := (x, t) \in \mathbb{R}^N \times [0, T]$ and all C^∞ -test functions η such that η is admissible at z and $u - \eta$ has a maximum at z (in the domain of definition of η) the following holds:

i) If $\nabla\eta(z) = 0$, then it holds

$$\eta_t(z) \leq 0 \tag{4.7}$$

eq viscosa degen

ii) If $\nabla\eta(z) \neq 0$, then

$$\partial_t\eta(z) + \psi(z, -\nabla\eta(z)) (H(z, \nabla\eta(z), \nabla^2\eta(z)) - f(z, t)) \leq 0. \tag{4.8}$$

eq viscosa

A lower semicontinuous function $u : \mathbb{R}^N \times [0, T] \rightarrow \mathbb{R}$ (in short, $u \in lsc(\mathbb{R}^N \times [0, T])$), constant outside a compact set, is a viscosity supersolution of the Cauchy problem (4.6) if $u(\cdot, 0) \geq u_0$ and for all $z := (x, t) \in \mathbb{R}^N \times [0, T]$ and all C^∞ -test functions η such that η is admissible at z and $u - \eta$ has a minimum at z (in the domain of definition of η) the following holds:

i) If $\nabla\eta(z) = 0$, then $\eta_t(z) \geq 0$;

ii) If $\nabla\eta \neq 0$ then

$$\partial_t\eta(z) + \psi(z, -\nabla\eta(z)) (H(z, \nabla\eta(z), \nabla^2\eta(z)) - f(z, t)) \leq 0.$$

Finally, a function u is a viscosity solution for the Cauchy problem (4.6) if it is both a subsolution and a supersolution of (4.6).

Remark. By classical arguments, one could assume that the maximum of $u - \eta$ is strict in the definition of subsolution above (an analogous remark holds for supersolutions).

Remark. We remark that, if $-u$ is a subsolution to (4.6) with initial datum $-u_0$, then u is a supersolution for (4.6) for the initial datum u_0 and where ϕ, ψ are replaced by $\tilde{\phi}, \tilde{\psi}$ respectively, as defined in Section 4.1.

We will first prove existence for viscosity solutions of (4.6) via an approximation-in-time technique, and then prove uniqueness of solutions to (4.6) to link the approximate solution to the mean curvature flow equation. We would like to proceed with the classical construction of e.g. [11, 14, 21], but in our case the lack of continuity of the evolving functions forces us to be particularly careful with the procedure.

We use the shorthand notation of *lsc* for lower semicontinuous and *usc* for upper semicontinuous. Given a bounded, *usc* function v which is constant outside a compact set, we define the transformation

$$T_{h,t}^+ v(x) = \sup \left\{ s : x \in T_{h,t}^+ \{v \geq s\} \right\}. \tag{4.9}$$

def increm supers

Firstly, we see that $T_{h,t}^+ v(x) \in \mathbb{R}$, as v is bounded. Moreover, it turns out that the function $T_{h,t}^+ v$ is *usc*, bounded and constant outside a compact set. Indeed, definition (4.9) is equivalent to

$$T_{h,t}^+ v(x) = \inf \left\{ s : x \notin T_{h,t}^+ \{v \geq s\} \right\} = \inf_{s \in \mathbb{R}} \left(s + \mathbf{1}_{(T_{h,t}^+ \{v \geq s\})^c}(x) \right),$$

where $\mathbb{1}_A(x)$ is the indicatrix function of a set A , being 0 on the set and $+\infty$ outside. By definition, $\mathbb{1}_A$ is an *usc* function for any open set A . Thus, recalling Remark 3.4, in the equation above we are taking the infimum of a family of *usc* functions, which is then a *usc* function. The other two properties follows from the previous study of the discrete evolution. Analogously, given a bounded *lsc* function g , we define

$$T_{h,t}^- g(x) = \sup \left\{ s : x \in T_{h,t}^- \{g > s\} \right\} = \sup_{s \in \mathbb{R}} \left(s - \mathbb{1}_{T_{h,t}^- \{g > s\}} \right), \quad (4.10)$$

which is now a bounded *lsc* function (as sup of *lsc* functions), constant outside a compact set.

We are now ready to give the definition of the discrete-in-time approximations of sub and super solution to (4.6). Given an initial compact set E_0 , set u_0 as a (uniformly) continuous function, spatially constant outside a compact set, such that $\{u_0 \geq 0\} = E_0$. We remark that for every $s \in \mathbb{R}$, the superlevel set $\{u_0 \geq s\}$ is either compact or it is unbounded with compact boundary. Then, for $h > 0$ we introduce the following family of maps as $u^\pm(\cdot, t) = u_0$ for $t \in [0, h)$ and

$$u_h^\pm(\cdot, t) := T_{h,t-h}^\pm u_h^\pm(\cdot, t-h) \quad \text{for } t \geq h. \quad (4.11)$$

def subsol iter

We easily see that the maps above are functions (as implied by the comparison principle contained in Lemmas 3.2, 4.1 and 4.4) piecewise constant in time (as $T_{h,t}^\pm = T_{h,[t/h]h}^\pm$). Moreover, by the previous remarks, we have that $u_h^+(\cdot, t)$ is an *usc* function, while $u_h^-(\cdot, t)$ is a *lsc* function, for every $t \in [0, +\infty)$. Some further properties of the approximating scheme are listed below.

Lemma 4.8. *For any $h > 0$, $t \geq 0$ we have the following. It holds*

$$u_h^-(\cdot, t) \leq u_h^+(\cdot, t). \quad (4.12)$$

ineq subsol

Furthermore, given any $\lambda \in \mathbb{R}$ and $t \geq h$ it holds

$$\begin{aligned} \{u_h^+(\cdot, t) > \lambda\} &\subseteq T_{h,t-h}^+ \{u_h^+(\cdot, t-h) \geq \lambda\} \subseteq \{u_h^+(\cdot, t) \geq \lambda\} \\ \{u_h^-(\cdot, t) > \lambda\} &\subseteq T_{h,t-h}^- \{u_h^-(\cdot, t-h) > \lambda\} \subseteq \{u_h^-(\cdot, t) \geq \lambda\}. \end{aligned} \quad (4.13)$$

inclusion level s

Proof. Fix $x \in \mathbb{R}^N$, $t \in [0, h)$. For any given $\sigma < u_h^-(x, h)$ we have that there exists a sequence $(s_n) \nearrow \sigma$ so that $x \in T_{h,t-h}^- \{u_0 > s_n\} \subseteq T_{h,t-h}^+ \{u_0 \geq s_n\}$. Thus, $u_h^+(x, t) \geq \sigma$. We then conclude by induction. Then, (4.13) follows easily by the definition (4.11). \square

We then prove that the half-relaxed limits (in the spirit of [6], see also the references therein) of the families of functions u_h^\pm

$$\begin{aligned} u^+(x, t) &:= \sup_{(x_h, t_h) \rightarrow (x, t)} \limsup_{h \rightarrow 0} u_h^+(x_h, t_h) \\ u^-(x, t) &:= \inf_{(x_h, t_h) \rightarrow (x, t)} \liminf_{h \rightarrow 0} u_h^-(x_h, t_h), \end{aligned} \quad (4.14)$$

def subsol supers

are (respectively) sub and supersolutions in the viscosity sense of (4.6), see Theorem 1.2 (note that, by definition, u^+ is *usc*, while u^- is *lsc*). The proof of this result is the subject of the following section and we recall that the hypothesis required are (H), (1.4) and $f \in C^0(\mathbb{R}^N \times [0, \infty))$ only. Once the existence of sub and super-solutions to the equation is settled, we need to properly define the notion of level-set solution to the mean curvature flow. To do so, we first prove uniqueness for

(4.6) via a comparison principle and under additional hypothesis. Then, we show that the evolution of the zero superlevel set of the solution does not depend on the choice of the initial function u_0 .

We start with a comparison result between u^+, u^- and u_0 at the initial time: it will ensure that the classical hypothesis for the comparison principle are satisfied. We first prove an estimate for the speed of decay of the level sets of the evolving functions. While it will only be needed in the following section, in the proof of the forthcoming Lemma 4.10 we will use similar techniques, so we preferred to state it here.

imate decay balls

Lemma 4.9. *Let $u^+(x, t)$ be the function defined in (4.14), let $\sigma \in \mathbb{R}$. Assume that, for a suitable x_0 and $R > 0$, it holds $B(x_0, R) \subseteq \{u^+(\cdot, t_0) \geq \sigma\}$. Then, there exists $C = C(R, \phi, \psi, f)$ such that $B(x_0, R - C(t - t_0)) \subseteq \{u^+(\cdot, t) \geq \sigma\}$ for every $t \leq t_0 + R/(2C)$. An analogous statement holds for u^- by considering its open sublevel sets.*

Proof. We focus on the case $\{u^+(\cdot, t_0) \geq \sigma\}$ bounded, the other case being analogous. By assumption, for any $R_0 < R$, if h is small enough, we have $B(x_0, R_0) \subseteq \{u_h^+(\cdot, t_0) \geq \sigma\}$. Set $C = C(R_0/2, \phi, \psi, f)$ as the constant of Lemma 3.8. Let R_n be defined recursively following law (3.7), that is $R_{n+1} = R_n - Ch$, as long as $R_n \geq R_0/2$. By simple iteration we find that $R_n = R_0 - nCh$, as long as $R_n \geq R_0/2$, which can be ensured enforcing $hn \leq R_0/(2C)$. Therefore, for any $t \geq t_0$ such that $t - t_0 \leq R_0/(2C)$, we set $n = \lfloor (t - t_0)/h \rfloor$ and send $h \rightarrow 0$ to deduce (recalling also Lemma 3.2)

$$\{u^+(\cdot, t) \geq \sigma\} \supset B(x_0, R_0 - C(t - t_0)).$$

Since the choice of R_0 is arbitrary, we conclude. \square

We are now ready to prove a comparison result for the functions u^\pm and a continuity estimate at the initial time $t = 0$.

ma initial datum

Lemma 4.10. *For any $(x, t) \in \mathbb{R}^N \times [0, +\infty)$ it holds*

$$u^-(x, t) \leq u^+(x, t).$$

Moreover $u^-(\cdot, 0) = u^+(\cdot, 0) = u_0$, so that there exists a modulus of continuity ω such that $\forall x, y \in \mathbb{R}^N$

$$u^+(x, 0) - u^-(y, 0) \leq \omega(|x - y|).$$

Proof. The proof of the first inequality essentially follows from (4.12) and the definition of u^\pm . To prove the equality at the initial time $t = 0$, we start by remarking that $u^+(\cdot, 0) \geq u_0$ as can be seen taking sequences of the form $(x_h, 0)$ in (4.14). Then, consider ω as a continuous, strictly increasing modulus of continuity for u_0 . We can also see that $\forall \varepsilon > 0 \{u_0 \leq u_0(x) + \varepsilon\} \supseteq B(x, \omega^{-1}(\varepsilon))$ by uniform continuity. Thus, reasoning iteratively as in Lemma 4.9 and using (4.13), we obtain that there exists $h_0(\varepsilon)$ such that $\forall h \leq h_0$ it holds

$$\{u_h^+(\cdot, t) \leq u_h^+(x, 0) + \varepsilon\} \supseteq \left(T_{h, t-h}^+ \{u_0 > u_0(x) + \varepsilon\}\right)^c = T_{h, t-h}^- \{u_0 \leq u_0(x) + \varepsilon\} \supseteq B(x, \omega^{-1}(\varepsilon/2)),$$

as long as $t \leq (\omega^{-1}(\varepsilon) - \omega^{-1}(\varepsilon/2))/(2C) =: t_\varepsilon$, and where we recalled that $u_h^\pm(\cdot, 0) = u_0$. Now, fix $\sigma > 0, x \in \mathbb{R}^N$ such that $u(x, 0) > \sigma$ and a sequence $(x_{h_k}, t_{h_k}) \rightarrow (x, 0)$ such that $\lim_k u_{h_k}^+(x_{h_k}, t_{h_k}) > \sigma$. Then, for k large enough $(x_{h_k}, t_{h_k}) \in B(x, \omega^{-1}(\varepsilon/2)) \times [0, t_\varepsilon]$ and so we conclude

$$\sigma < \lim_k u_{h_k}^+(x_{h_k}, t_{h_k}) \leq u_0(x, 0) + \varepsilon.$$

Letting $\varepsilon \rightarrow 0$ we conclude $u(\cdot, 0)^+ \leq u_0$. The proof for u^- is essentially the same. The last claim follows from the previous one, recalling that ω is a modulus of uniform continuity for u_0 . \square

In order to prove a comparison principle for (4.6), we will need to assume (1.11). Under these additional hypotheses, we are able to prove uniqueness for the parabolic Cauchy problem (4.6). The proof of this result follows from [24, Theorem 4.2]: we will just show in detail that the assumption of the aforementioned theorem hold in our case, following [9, Proposition 6.1] and [24, pag. 465].

Proof of Theorem 1.4. The proof of this result essentially follows from [24, Theorem 4.2], combined with the existence result of Theorem 1.3. Referring to the notation of [24], we firstly remark that in our case $\Omega = \mathbb{R}^N$, thus the parabolic boundary of $U = \Omega \times [0, T]$ is simply $\partial_p U = \mathbb{R}^N \times \{0\}$. Therefore, the initial conditions (A1) – (A3) are all verified by Lemma 4.10. We then define the continuous Hamiltonian $F : [0, T] \times \mathbb{R}^N \times (\mathbb{R}^N \setminus \{0\}) \times M^{N \times N} \rightarrow \mathbb{R}$ as follows

$$F(t, x, p, X) := \psi(x, -p) \left(- \sum_i \partial_{x_i} \partial_p \phi(x, -p) + \nabla_p^2 \phi(x, -p) : X + f(x, t) \right), \quad (4.15)$$

Hamiltonian Giga

and focus on the conditions (F1), (F3) – (F5), (F6'), (F7), (F9), (F10) that F must satisfy. The assumptions (F1), (F3) – (F5), (F9) are easily checked. (F6') follows from the Lipschitz regularity of ϕ and ψ , as $\forall t \in [0, T], x \in \mathbb{R}^N, |p| \geq \rho, |q| + |X| \leq R$ one has

$$\begin{aligned} |F(t, x, p, X) - F(t, x, q, X)| &\leq c_\psi |p - q| \left| - \sum_i \partial_{x_i} \partial_p \phi(x, -p) + \nabla_p^2 \phi(x, -p) : X \right| \\ &+ \psi(x, -q) \left| - \sum_i (\partial_{x_i} \partial_p \phi(x, -p) - \partial_{x_i} \partial_p \phi(x, -q)) + (\nabla_p^2 \phi(x, -p) - \nabla_p^2 \phi(x, -q)) : X \right| \\ &\leq c_R |p - q| \left(1 + \frac{1}{|p|} \right) + c_R |p - q| \leq c_{R, \rho} |p - q|. \end{aligned}$$

For (F7), we remark that the first term in the parenthesis in (4.15) is 0-homogeneous in p , while the second one is (-1) -homogeneous in p but 1-homogeneous in X . Lastly, we sketch how to prove (F10). Since it concerns the X -terms, we focus simply on

$$\nabla_p^2 \phi(x, -p) : X = \text{tr} (\nabla_p^2 \phi(x, -p) X^T).$$

Multiplying by $\phi(x, -p)$, we rewrite $\phi(x, -p) \text{tr} (\nabla_p^2 \phi(x, -p) X^T) = \text{tr} (A(x, -p) X^T)$, where $A = B - (\nabla_p \phi \otimes \nabla_p \phi)$, with B being the uniformly elliptic operator $\frac{1}{2} \nabla_p^2 \phi^2$. We can then factorize $B = \tilde{L} \tilde{L}^T$, with \tilde{L} being a nondegenerate, lower triangular matrix. Then, following the proof of [9, Proposition 6.1] and [24, pg. 465], we obtain (F10). \square

Once uniqueness is settled, one can finally define the notion of *level set* solution to the mean curvature flow as follows.

Definition 4.11. Let E_0 be a compact initial set. Define a uniformly continuous, bounded function $u_0 : \mathbb{R}^N \rightarrow \mathbb{R}$ such that $\{u_0 \geq 0\} = E_0$. Then, let $u : \mathbb{R}^N \times [0, +\infty) \rightarrow \mathbb{R}$ be the unique continuous viscosity solution to (4.6) given by Theorem 1.4. Then, the family $E_t := \{u^+(\cdot, t) \geq 0\}_{t \geq 0}$ will be called the level set solution to the mean curvature flow.

This definition is well posed since the Hamiltonian defined in (4.5) satisfies the so-called *geometricity condition*. Namely, one can easily check that for any $\lambda \neq 0, p \in \mathbb{R}^N \setminus 0, q \in \mathbb{R}^N$ and any symmetric $N \times N$ matrix X one has

$$H(x, \lambda p, \lambda X + p \otimes q + q \otimes p) = \frac{\lambda}{|\lambda|} H(x, p, X).$$

Thus, one can prove by classical arguments (see e.g. [14, Remark 3.9]) the following result.

Lemma 4.12. *Let u_0, \tilde{u}_0 two initial data for (4.6) such that $\{u_0 \geq 0\} = \{\tilde{u}_0 \geq 0\}$. Then, denoting by u, \tilde{u} the corresponding solutions to (4.6), one has*

$$\{u(\cdot, t) \geq 0\} = \{\tilde{u}(\cdot, t) \geq 0\} \quad \text{for all } t \in [0, T],$$

and the same identity holds for the open superlevel sets.

4.3 Proof of Theorem 1.3

In this section we will prove that the limiting functions u^\pm are respectively a viscosity sub and supersolutions to (4.6). We remark that we work assuming (H), (1.4) and that $f \in C^0(\mathbb{R}^N \times [0, +\infty))$. We will be following the structure of the proof of [14, Theorem 6.16], but taking into account the weaker definition of u holding in our case. We will be using the O, o notations with respect to $h \rightarrow 0$ and focus on proving that u is a subsolution. The proof for u^- is analogous.

Proof of Theorem 1.3. Consider u^+ as defined in (1.10): we need to prove that it is a subsolution. In the following, we will denote $u := u^+$ and $u_h := u_h^+$. Let $\eta(x, t)$ be an admissible test function in $\bar{z} := (\bar{x}, \bar{t})$ and assume that (\bar{x}, \bar{t}) is a strict maximum point for $u - \eta$. Assume furthermore that $u - \eta = 0$ in such point. We need to show that either (4.7) or (4.8) holds at \bar{z} .

Step 1. Let us first assume that $\nabla \eta(\bar{z}) \neq 0$. By classical arguments, we can assume that \bar{z} is a strict maximum point with $\bar{t} > 0$ and that η is smooth. By the definition of u , there exist a sequence $\tilde{z}_k := (\tilde{x}_{h_k}, \tilde{t}_{h_k}) \rightarrow \bar{z}$ such that $\lim_k u_{h_k}(\tilde{z}_k) = u(\bar{z})$. By standard arguments (compare e.g. [5, Lemma 6.1]), there exists a radius $\rho > 0$ such that all functions $u_{h_k} - \eta$ achieve a local supremum in $B_\rho(\bar{z})$ at points $z_k = (x_k, t_k)$. We remark that we could modify slightly the definition of $u_{h_k}(x, t)$ to be

$$\sum_{l \in \mathbb{N}} u_{h_k}(x, t) \chi_{[lh, (l+1)h]}(t) \quad \text{or} \quad \sum_{l \in \mathbb{N}} u_{h_k}(x, t) \chi_{(lh, (l+1)h]}(t), \quad (4.16) \quad \boxed{\text{def equi}}$$

as this does not change the value of $u(x, t)$. Recalling that u_{h_k} is *usc* in space, we can thus ensure that z_k is a local maximum point for $u_{h_k} - \eta$ by choosing the definition in (4.16) that makes u_{h_k} *usc* in time at z_k . Then, passing to a further subsequence we can ensure that $z_k \rightarrow w \in B_\rho(\bar{z})$, and we use the definition of u to obtain

$$(u - \eta)(w) \geq \limsup_k (u_{h_k} - \eta)(z_k) \geq \limsup_k (u_{h_k} - \eta)(\tilde{z}_k) = (u - \eta)(\bar{z}).$$

Therefore, $w = \bar{z}$ by maximality. Thus we can assume that each function $u_{h_k} - \eta$ achieves a local supremum in $B_\rho(\bar{z})$ at a point $z_{h_k} =: (x_k, t_k)$ and that $u_{h_k}(z_{h_k}) \rightarrow u(\bar{z})$ as $k \rightarrow \infty$. Finally, we can assume also that $\nabla \eta(x_k, t_k) \neq 0$ for k large enough.

In particular, one has

$$u_h(x, t) \leq \eta(x, t) + c_k \quad (4.17)$$

eq 6.19 Nonlocal

where $c_k := u_{h_k}(x_k, t_k) - \eta(x_k, t_k)$, with equality if $(x, t) = (x_k, t_k)$. Let $\sigma > 0$ and set

$$\eta_{h_k}^\sigma(x) := \eta(x, t_k) + c_k + \frac{\sigma}{2}|x - x_k|^2.$$

Then, for all $x \in \mathbb{R}^N$,

$$u_{h_k}(x, t_k) \leq \eta_{h_k}^\sigma(x)$$

with equality if and only if $x = x_k$. We set $l_k = u_{h_k}(x_k, t_k) = \eta_{h_k}^\sigma(x_k)$. We fix $\varepsilon > 0$, to be chosen later, and write $E_{\varepsilon, k} := \{u_{h_k}(\cdot, t_k - h_k) \geq l_k - \varepsilon\}$. We define¹

$$W_\varepsilon := \left(T_{h, t_k - h_k}^+ E_{\varepsilon, k}\right) \setminus \{\eta_{h_k}^\sigma(\cdot) > l_k + \varepsilon\}. \quad (4.18)$$

We immediately see that $W_\varepsilon \rightarrow \{x_k\}$ in the Kuratowski sense as $\varepsilon \rightarrow 0$ since by (4.13)

$$\{u_{h_k}(\cdot, t_k) > l_k - \varepsilon\} \setminus \{\eta_{h_k}^\sigma(\cdot) > l_k + \varepsilon\} \subseteq W_\varepsilon \subseteq \{u_{h_k}(\cdot, t_k) \geq l_k - \varepsilon\} \setminus \{\eta_{h_k}^\sigma(\cdot) > l_k + \varepsilon\}, \quad (4.19)$$

inclusions W_e

see also (4.24) below. Then, we check that $|W_\varepsilon| > 0$ for all ε small enough. By the continuity of η^σ and $|\nabla \eta(\bar{z})| \neq 0$, for any ε there exist a radius r_ε such that $W_\varepsilon \supseteq B(x_k, r_\varepsilon) \cap T_{h, t_k - h_k}^+ E_{\varepsilon, k}$. Furthermore, for any $\varepsilon > 0$, using (4.13) again yields $x_k \in T_{h_k, t_k - h_k}^+ \{u_{h_k}(\cdot, t_k - h_k) \geq l_k - \varepsilon\}$, and the latter set coincides with the closure of its points of density 1 by Lemma 3.3. Thus, x_k satisfies lower density estimates and so we conclude that $|W_\varepsilon| > 0$. Now, assume $E_{\varepsilon, k}$ is bounded. By minimality we have

$$\begin{aligned} & P_\phi(T_{h, t_k - h_k}^+ E_{\varepsilon, k}) + \frac{1}{h_k} \int_{T_{h, t_k - h_k}^+ E_{\varepsilon, k}} \text{sd}_{E_{\varepsilon, k}}^\psi(x) \, dx + \int_{W_\varepsilon} F_{h_k}(x, t_k - h_k) \, dx \\ & \leq P_\phi\left(\left(T_{h, t_k - h_k}^+ E_{\varepsilon, k}\right) \cap \{\eta_{h_k}^\sigma > l_k + \varepsilon\}\right) + \frac{1}{h_k} \int_{\left(T_{h, t_k - h_k}^+ E_{\varepsilon, k}\right) \cap \{\eta_{h_k}^\sigma > l_k\}} \text{sd}_{E_{\varepsilon, k}}^\psi. \end{aligned} \quad (4.20)$$

eq 6.21 Nonlocal

Adding to both sides the term $P_\phi\left(\{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cup T_{h, t_k - h_k}^+ E_{\varepsilon, k}\right)$ and using the submodularity (2.4), we obtain

$$\begin{aligned} & P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cup W_\varepsilon) - P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\}) + \frac{1}{h_k} \int_{W_\varepsilon} \text{sd}_{E_{\varepsilon, k}}^\psi(x) \, dx \\ & \quad + \int_{W_\varepsilon} F_{h_k}(x, t_k - h_k) \, dx \leq 0. \end{aligned}$$

By (4.17), $\{u_{h_k}(\cdot, t_k - h_k) \geq l_k - \varepsilon\} \subseteq \{\eta(\cdot, t_k - h_k) \geq l_k - c_k - \varepsilon\}$, therefore it holds

$$\begin{aligned} & P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cup W_\varepsilon) - P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\}) + \frac{1}{h_k} \int_{W_\varepsilon} \text{sd}_{\{\eta(\cdot, t_k - h_k) \geq l_k - c_k - \varepsilon\}}^\psi(x) \, dx \\ & \quad + \int_{W_\varepsilon} F_{h_k}(x, t_k - h_k) \, dx \leq 0. \end{aligned} \quad (4.21)$$

eq 6.22 Nonlocal

¹We need to define the sets W_ε in this way (compare the different definition in [14]) since firstly, we can not rule out that the inclusions in (4.19) are strict, and secondly it is not clear if otherwise $|W_\varepsilon| > 0$.

If instead $E_{\varepsilon,k}$ is an unbounded set with compact boundary, we replace inequality (4.20) by

$$\begin{aligned} & P_\phi(T_{h,t_k-h_k} E_{\varepsilon,k}) + \frac{1}{h_k} \int_{(T_{h,t_k-h_k}^+ E_{\varepsilon,k}) \cap B_R} \text{sd}_{E_{\varepsilon,k}}^\psi(x) \, dx + \int_{W_\varepsilon} F_{h_k}(x, t_k - h_k) \, dx \\ & \leq P_\phi((T_{h,t_k-h_k}^+ E_{\varepsilon,k}) \cap \{\eta_{h_k}^\sigma > l_k + \varepsilon\}) + \frac{1}{h_k} \int_{(T_{h,t_k-h_k}^+ E_{\varepsilon,k}) \cap \{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cap B_R} \text{sd}_{E_{\varepsilon,k}}^\psi, \end{aligned}$$

for $R > 0$ sufficiently large, see (4.4). Then, one can argue as before to obtain (4.21).

The first two terms $P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cup W_\varepsilon) - P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\})$ can be estimated as done in Lemma 3.8. Indeed, we consider the vector field $v = \nabla_p \phi(x, \nabla \eta_{h_k}^\sigma)$ in (2.5) and we use the divergence theorem to get

$$\begin{aligned} P_\phi(\{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cup W_\varepsilon) - P_\phi(\{\eta_{h_k}^\sigma \geq l_k + \varepsilon\}) & \geq \int_{\partial(\{\eta_{h_k}^\sigma > l_k + \varepsilon\} \cup W_\varepsilon)} v \cdot \nu - \int_{\partial\{\eta_{h_k}^\sigma > l_k + \varepsilon\}} v \cdot \nu \\ & = |W_\varepsilon| \int_{W_\varepsilon} \text{div } v, \end{aligned} \tag{4.22}$$

eq 6.30 Nonlocal

where ν denotes the unit outer vector to the set we are integrating on. We then remark that $\int_{W_\varepsilon} \text{div } v \rightarrow H_{\{\eta_{h_k}^\sigma > l_k\}}^\phi(x_k)$ and $\int_{W_\varepsilon} F_{h_k}(x, t_k - h_k) \, dx \rightarrow F_{h_k}(x_k, t_k - h_k)$ as $\varepsilon \rightarrow 0$ by continuity.

The other term in (4.21), $\int_{W_\varepsilon} \text{sd}_{\{\eta(\cdot, t_k - h_k) > l_k - c_k - \varepsilon\}}^\psi$, can be treated as follows. For any $z \in W_\varepsilon$, we have

$$\eta(z, t_k) + c_k + \frac{\sigma}{2} |z - x_k|^2 \leq l_k + \varepsilon. \tag{4.23}$$

eq 6.23 Nonlocal

Since, in turn, $\eta(z, t_k) + c_k > l_k - \varepsilon$ it follows that $\sigma |z - x_k|^2 < 4\varepsilon$ and thus, for ε small enough,

$$W_\varepsilon \subseteq B_{c\sqrt{\varepsilon}}(x_k). \tag{4.24}$$

eq 6.24 Nonlocal

By a Taylor expansion, for every $z \in W_\varepsilon$ we have

$$\eta(z, t_k - h_k) = \eta(z, t_k) - h_k \partial_t \eta(z, t_k) + h_k^2 \int_0^1 (1-s) \partial_{tt}^2 \eta(z, t_k - sh_k) \, ds. \tag{4.25}$$

eq 6.25 Nonlocal

Then, we consider $y, y_e \in \{\eta(\cdot, t_k - h_k)(y) = l_k - c_k - \varepsilon\}$ being respectively, a point of minimal ψ -distance and Euclidean distance from z . We shall prove that

$$|z - y| = O(h_k). \tag{4.26}$$

decay y

In order to prove this result, we start remarking that for $k \rightarrow \infty$ and choosing $\varepsilon \ll h_k$, one has $\text{sd}_{\{\eta(\cdot, t_k - h_k) \geq l_k - c_k - \varepsilon\}}^\psi(z) \rightarrow 0$ (as $z \rightarrow x_k$ for $\varepsilon \rightarrow 0$ and $x_k \in \{\eta(\cdot, t_k) \geq l_k - c_k\}$). In particular, recalling the bounds (2.3) one has

$$|z - y_e| \leq c_\psi^2 |z - y| \leq c_\psi^3 |\text{sd}_{\{\eta(\cdot, t_k - h_k) \geq l_k - c_k - \varepsilon\}}^\psi(z)| \rightarrow 0$$

as $k \rightarrow \infty$. By (4.23) we deduce in particular $\eta(z, t_k) + c_k < l_k + \varepsilon$, that is,

$$0 \leq \eta(z, t_k) - \eta(y, t_k - h_k) \leq 2\varepsilon, \tag{4.27}$$

eq 6.28 Nonlocal

and the same inequality substituting y_e to y . Thus, one has

$$\eta(z, t_k) - \eta(y_e, t_k - h_k) = \nabla\eta(y, t_k - h_k) \cdot (z - y_e) - h_k \partial_t \eta(y, t_k - h_k) + O(|z - y_e|^2 + h_k^2)$$

which we combine with $\nabla\eta(y, t_k - h_k) \cdot (z - y_e) = \pm |\nabla\eta(y, t_k - h_k)| |z - y_e|$ (see [14] for details) and (4.27) to get

$$|z - y_e| |\nabla\eta(y, t_k - h_k)| \leq 2\varepsilon + O(h_k) + O(|z - y_e|^2).$$

Recalling that $|\nabla\eta(y, t_k - h_k)| \geq c > 0$ for h_k small enough, we divide by $|\nabla\eta(y, t_k - h_k)|$ to conclude $|z - y_e| = O(h_k)$ as $\varepsilon \ll h_k$. Finally, employing again (2.3), we conclude (4.26).

Then, we consider a geodesic curve for the definition of $\text{sd}_{\{\eta(\cdot, t_k - h_k) \geq l_k - c_k - \varepsilon\}}^\psi(z)$: if this distance is positive, we choose $\gamma : [0, 1] \rightarrow \mathbb{R}^N$ with $\gamma(0) = z, \gamma(1) = y$, with y as before, otherwise we take γ such that $\gamma(0) = y, \gamma(1) = z$. In the following, we will assume $\text{sd}_{\{\eta(\cdot, t_k - h_k) \geq l_k - c_k - \varepsilon\}}^\psi(z) > 0$, the other case being analogous. Recalling (2.1), we have

$$\begin{aligned} \eta(z, t_k - h_k) &= \eta(y, t_k - h_k) + \int_0^1 \nabla\eta(\gamma, t_k - h_k) \cdot \dot{\gamma} \, dt \\ &\geq \eta(y, t_k - h_k) - \int_0^1 \psi(\gamma, -\nabla\eta(\gamma, t_k - h_k)) \psi^\circ(\gamma, \dot{\gamma}) \, dt \\ &\geq \eta(y, t_k - h_k) - \psi(y, -\nabla\eta(y, t_k - h_k)) \text{sd}_{\{\eta(\cdot, t_k - h_k) = l_k - c_k - \varepsilon\}}^\psi(z) \\ &\quad - \int_0^1 (\psi(\gamma, -\nabla\eta(\gamma, t_k - h_k)) - \psi(y, -\nabla\eta(y, t_k - h_k))) \psi^\circ(\gamma, \dot{\gamma}) \, dt \\ &\geq \eta(y, t_k - h_k) - (\psi(y, -\nabla\eta(y, t_k - h_k)) + c|z - y|) \text{sd}_{\{\eta(\cdot, t_k - h_k) = l_k - c_k - \varepsilon\}}^\psi(z), \end{aligned}$$

where in the last line we reasoned as in (3.25) to obtain the bound $\sup_t |\gamma(t) - y| \leq c|z - y|$. Recalling (4.26) one has

$$\eta(z, t_k - h_k) \geq \eta(y, t_k - h_k) - \psi(y, -\nabla\eta(y, t_k - h_k)) \text{sd}_{\{\eta(\cdot, t_k - h_k) = l_k - c_k - \varepsilon\}}^\psi(z) + o(h_k). \quad (4.28)$$

eq 6.26 Nonlocal

Combining (4.25) with (4.28) and using (4.27), we deduce

$$\begin{aligned} &\text{sd}_{\{\eta(\cdot, t_k - h_k) = l_k - c_k - \varepsilon\}}^\psi(z) \psi(y, -\nabla\eta(y, t_k - h_k)) + o(h_k) \\ &\geq -2\varepsilon + h_k \partial_t \eta(z, t_k) - h_k^2 \int_0^1 (1-s) \partial_{tt}^2 \eta(z, t_k - sh_k) \, ds. \end{aligned}$$

Note that, in view of (4.23) and (2.3), $|\eta(z, t_k) - \eta(y, t_k)| \leq c\varepsilon + ch_k = O(h_k)$, provided $\varepsilon \ll h_k$ and small enough. Thus, using also (4.24), (4.26) we deduce

$$\begin{aligned} \frac{1}{h_k} \text{sd}_{\{\eta(\cdot, t_k - h_k) = l_k - c_k - \varepsilon\}}^\psi(z) &\geq \frac{\partial_t \eta(z, t_k) - \frac{2\varepsilon}{h_k} - O(h_k) - O_{h_k}(1)}{\psi(y, -\nabla\eta(y, t_k - h_k))} \\ &= \frac{\partial_t \eta(x_k, t_k) + O(\sqrt{\varepsilon}) - \frac{2\varepsilon}{h_k} - O(h_k) - O_{h_k}(1)}{\psi(x_k, -\nabla\eta(x_k, t_k - h_k)) + O(\sqrt{\varepsilon}) + O(h_k)}. \end{aligned} \quad (4.29)$$

eq 6.29 Nonlocal

We conclude by employing (4.21), (4.22) and (4.29), dividing by $|W_\varepsilon|$ and sending $\varepsilon \rightarrow 0$ to obtain

$$\frac{\partial_t \eta(x_k, t_k) - O_{h_k}(1)}{\psi(x_k, -\nabla\eta(x_k, t_k)) + O(h_k)} + H_{\{\eta_{h_k}^\sigma \geq \eta_{h_k}^\sigma(x_k)\}}^\phi(x_k) - F_{h_k}(x_k, t_k - h_k) \leq 0.$$

Letting simultaneously $\sigma \rightarrow 0$ and $k \rightarrow \infty$, recalling the continuity properties of H^ϕ , we deduce (4.8). Indeed the sets $\{\eta_{\tilde{h}_k}^\sigma > \eta_{\tilde{h}_k}^\sigma(x_k)\}$ are converging in C^2 to the set $\{\eta > \eta(x)\}$, $x_k \rightarrow x$ and thus

$$H_{\{\eta_{\tilde{h}_k}^\sigma > \eta_{\tilde{h}_k}^\sigma(x_k)\}}^\phi(x_k) \rightarrow H_{\{\eta > \eta(x)\}}^\phi(x),$$

and we conclude the proof of this step.

Step 2. Now we consider the case $\nabla\eta(\bar{x}, \bar{t}) = 0$ and we show that $\partial_t\eta(\bar{x}, \bar{t}) \leq 0$. The proof follows the line of the one in [14], we just highlight the differences.

Since $\nabla\eta(\bar{z}) = 0$, there exist $g \in \mathcal{F}$ and $\omega \in C^\infty(\mathbb{R})$ with $\omega'(0) = 0$ such that

$$|\eta(x, t) - \eta(\bar{z}) - \partial_t\eta(\bar{z})(t - \bar{t})| \leq g(|x - \bar{x}|) + \omega(|t - \bar{t}|),$$

thus, we can define

$$\begin{aligned} \tilde{\eta}(x, t) &= \partial_t\eta(\bar{z})(t - \bar{t}) + 2g(|x - \bar{x}|) + 2\omega(|t - \bar{t}|) \\ \tilde{\eta}_k(x, t) &= \tilde{\eta}(x, t) + \frac{1}{k(\bar{t} - t)}. \end{aligned}$$

We remark that $u - \tilde{\eta}$ achieves a strict maximum in \bar{z} and the local maxima of $u - \tilde{\eta}_k$ in $\mathbb{R}^N \times [0, \bar{t}]$ are in points $(x_k, t_k) \rightarrow \bar{z}$ as $k \rightarrow \infty$, with $t_n \leq \bar{t}$. From now on, the only difference from [14] is in the case $x_k = \bar{x}$ for an (unrelabeled) subsequence. We assume $x_k = \bar{x} \forall k > 0$ and define $b_k = \bar{t} - t_k > 0$ and the radii

$$r_k := g^{-1}(a_k b_k),$$

where $a_k \rightarrow 0$ must be chosen such that the extinction time for $B(\cdot, r_k)$ is greater than $\bar{t} - t_k$, for k large enough. To show that such a choice for a_k is possible, recalling Lemma 4.9 we set

$$r_k = 2\sqrt{C(\bar{t} - t_k)} = 2\sqrt{C b_k}$$

(which has extinction time greater than $2(\bar{t} - t_k)$) and it is enough to prove that

$$\frac{g(2\sqrt{C b_k})}{b_k} \rightarrow 0, \quad \text{as } k \rightarrow \infty,$$

which follows from the definition of the family \mathcal{F} given in Definition 4.5. We then have

$$\begin{aligned} B(\bar{x}, r_k) &\subseteq \{\tilde{\eta}_k(\cdot, t_k) \leq \tilde{\eta}_k(\bar{x}, t_k) + 2g(r_k)\} \\ &\subseteq \{u(\cdot, t_k) \leq u(\bar{x}, t_k) + 2g(r_k)\}, \end{aligned}$$

by maximality of $u - \tilde{\eta}_k$ at z_k . Since Lemma 4.9 ensures

$$\bar{x} \in \{u(\cdot, \bar{t}) \leq u(\bar{x}, t_k) + 2g(r_k)\},$$

we use again the maximality of $u - \eta$ at \bar{z} and the choice of r_k to obtain

$$\frac{\eta(\bar{x}, t_k) - \eta(\bar{z})}{t_k - \bar{t}} = \frac{\eta(\bar{x}, t_k) - \eta(\bar{z})}{-b_k} \leq \frac{u(\bar{x}, t_k) - u(\bar{x}, \bar{t})}{-b_k} \leq \frac{-2g(r_k)}{-b_k} = -2a_k.$$

Passing to the limit $k \rightarrow \infty$, we conclude that $\partial_t\eta(\bar{z}) \leq 0$. \square

We conclude with two remarks concerning some possible generalizations of the results presented.

Remark 4.13. The results presented in this work can be immediately extended to unbounded initial open sets E_0 , whose boundary is compact. Indeed, defining the discrete flow as $E_t^{(h)} = E_0$ if $t \in [0, h)$, otherwise by induction $E_t^{(h)} = T_{h,t}^- E_{t-h}^{(h)}$, where the operator $T_{h,n}^-$ is the one defined in (4.2), this evolution is uniquely characterized by the one of the complement. Thus, all the results presented in this paper can be extended to this particular unbounded case.

rmk rel geom

Remark 4.14. Following the lines of [9] (in the spirit of [3]) one can see that the results of this paper may be extended to prove existence of flat flows and level set solutions to the mean curvature flow on \mathbb{R}^N endowed with the geometric structure induced by a Finsler metric ϕ° . For example, the perimeter functional in this setting is defined as follows. Given a set E of finite perimeter, its (intrinsic) perimeter is

$$\mathcal{P}_{\phi^\circ}(E) = \int_{\partial^* E} \phi(x, \nu_E(x)) d\mathcal{H}_{\phi^\circ}^{N-1}(x),$$

where the Hausdorff measure $\mathcal{H}_{\phi^\circ}^{N-1}$ is the one induced by the metric ϕ° . In particular, one can compute $d\mathcal{H}_{\phi^\circ}^{N-1}(x) = \omega_N |B^{\phi^\circ}(x)|^{-1} d\mathcal{H}^{N-1}(x)$ (see [9]), thus this approach is equivalent to consider in our framework a slightly different (but still regular) anisotropy, namely $\phi^*(x, \nu) := \omega_N |B^{\phi^\circ}(x)|^{-1} \phi(x, \nu)$. In particular, this approach leads to considering the evolution of hypersurfaces E_t moving according to the evolution law

$$V_{\phi^\circ}(x, t) = -\mathcal{H}_{E_t}(x) + f(x, t) \quad x \in \partial E_t, \quad t \in (0, T)$$

where now V_{ϕ° represents the speed of evolution along the anisotropic normal outer vector $n_{\phi^\circ}(x) = \nabla_p \phi(x, \nu_E(x))$ and \mathcal{H} is the “intrinsic” mean curvature, thus the first variation of the perimeter \mathcal{P}_{ϕ° . Recalling that $n_{\phi^\circ}(x) \cdot \nu_E(x) = \phi(x, \nu_E(x))$, we see that the hypersurfaces are evolving with a normal (in the Euclidean sense) velocity given by the law

$$V(x, t) = \phi(x, \nu_{E_t}(x)) \left(-H_{E_t}^{\phi^*}(x) + f(x, t) \right).$$

After this transformation, we can apply the results previously proved.

Acknowledgements

D. De Gennaro has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 94532. .

References

AllCah

- [1] S. M. Allen and J. W. Cahn. “A macroscopic theory for antiphase boundary motion and its application to antiphase domain coarsening”. In: *Acta Metall.* 27.5 (1979), pp. 1085–1095.

AlmChaNov

- [2] L. Almeida, A. Chambolle, and M. Novaga. “Mean curvature flow with obstacles”. In: *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 29.5 (2012), pp. 667–681. ISSN: 0294-1449. DOI: [10.1016/j.anihpc.2012.03.002](https://doi.org/10.1016/j.anihpc.2012.03.002).

AlmTayWan

- [3] F. Almgren, J. E. Taylor, and L. Wang. “Curvature-driven flows: a variational approach”. In: *SIAM J. Control Optim.* 31.2 (1993), pp. 387–438. ISSN: 0363-0129. DOI: [10.1137/0331020](https://doi.org/10.1137/0331020).

- AmbFusPal** [4] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000, pp. xviii+434. ISBN: 0-19-850245-1.
- Bar-book** [5] G. Barles. “An introduction to the theory of viscosity solutions for first-order Hamilton-Jacobi equations and applications”. In: *Hamilton-Jacobi equations: approximations, numerical analysis and applications*. Vol. 2074. Lecture Notes in Math. Springer, Heidelberg, 2013, pp. 49–109. DOI: [10.1007/978-3-642-36433-4_2](https://doi.org/10.1007/978-3-642-36433-4_2). URL: https://doi.org/10.1007/978-3-642-36433-4_2.
- BarSonSou** [6] G. Barles, H. M. Soner, and P. E. Souganidis. “Front propagation and phase field theory”. In: *SIAM J. Control Optim.* 31.2 (1993), pp. 439–469. ISSN: 0363-0129. DOI: [10.1137/0331021](https://doi.org/10.1137/0331021).
- Bar-rev** [7] Guy Barles. “An introduction to the theory of viscosity solutions for first-order Hamilton-Jacobi equations and applications”. In: *Hamilton-Jacobi equations: approximations, numerical analysis and applications*. Vol. 2074. Lecture Notes in Math. Springer, Heidelberg, 2013, pp. 49–109. DOI: [10.1007/978-3-642-36433-4_2](https://doi.org/10.1007/978-3-642-36433-4_2).
- BarSou** [8] Guy Barles and Panagiotis E. Souganidis. “A new approach to front propagation problems: theory and applications”. In: *Arch. Rational Mech. Anal.* 141.3 (1998), pp. 237–296. ISSN: 0003-9527. DOI: [10.1007/s002050050077](https://doi.org/10.1007/s002050050077).
- BelPao** [9] G. Bellettini and M. Paolini. “Anisotropic motion by mean curvature in the context of Finsler geometry”. In: *Hokkaido Math. J.* 25.3 (1996), pp. 537–566. ISSN: 0385-4035. DOI: [10.14492/hokmj/1351516749](https://doi.org/10.14492/hokmj/1351516749).
- CagMorMor** [10] F. Cagnetti, M. G. Mora, and M. Morini. “A second order minimality condition for the Mumford-Shah functional”. In: *Calc. Var. Partial Differential Equations* 33.1 (2008), pp. 37–74. ISSN: 0944-2669. DOI: [10.1007/s00526-007-0152-3](https://doi.org/10.1007/s00526-007-0152-3).
- Cha** [11] A. Chambolle. “An algorithm for mean curvature motion”. In: *Interfaces Free Bound.* 6.2 (2004), pp. 195–218. ISSN: 1463-9963. DOI: [10.4171/IFB/97](https://doi.org/10.4171/IFB/97). URL: <https://doi.org/10.4171/IFB/97>.
- ChaMorNovPon** [12] A. Chambolle, M. Morini, M. Novaga, and Marcello Ponsiglione. “Existence and uniqueness for anisotropic and crystalline mean curvature flows”. In: *J. Amer. Math. Soc.* 32.3 (2019), pp. 779–824. ISSN: 0894-0347. DOI: [10.1090/jams/919](https://doi.org/10.1090/jams/919).
- ChaMorPon12** [13] A. Chambolle, M. Morini, and M. Ponsiglione. “A nonlocal mean curvature flow and its semi-implicit time-discrete approximation”. In: *SIAM J. Math. Anal.* 44.6 (2012), pp. 4048–4077. DOI: [10.1137/120863587](https://doi.org/10.1137/120863587).
- ChaMorPon15** [14] A. Chambolle, M. Morini, and M. Ponsiglione. “Nonlocal curvature flows”. In: *Arch. Ration. Mech. Anal.* 218.3 (2015), pp. 1263–1329. DOI: [10.1007/s00205-015-0880-z](https://doi.org/10.1007/s00205-015-0880-z).
- ChaNov22** [15] A. Chambolle and M. Novaga. “Anisotropic and crystalline mean curvature flow of mean-convex sets”. In: *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 23.2 (2022), pp. 623–643. ISSN: 0391-173X.
- ChaNov08** [16] A. Chambolle and M. Novaga. “Implicit time discretization of the mean curvature flow with a discontinuous forcing term”. In: *Interfaces Free Bound.* 10.3 (2008), pp. 283–300. ISSN: 1463-9963. DOI: [10.4171/ifb/190](https://doi.org/10.4171/ifb/190). URL: <https://doi.org/10.4171/ifb/190>.
- CheGigGot** [17] Y. G. Chen, Y. Giga, and S. Goto. “Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations”. In: *J. Differential Geom.* 33.3 (1991), pp. 749–786.

- [18] G. De Philippis and T. Laux. “Implicit time discretization for the mean curvature flow of mean convex sets”. In: *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 21 (2020), pp. 911–930.
- [19] G. De Philippis and F. Maggi. “Regularity of free boundaries in anisotropic capillarity problems and the validity of Young’s law”. In: *Arch. Ration. Mech. Anal.* 216.2 (2015), pp. 473–568. ISSN: 0003-9527. DOI: [10.1007/s00205-014-0813-2](https://doi.org/10.1007/s00205-014-0813-2).
- [20] M. Desbrun, M. Meyer, P. Schröder, and A. H. Barr. “Implicit fairing of irregular meshes using diffusion and curvature flow”. In: *SIGGRAPH ’99*. 1999.
- [21] T. Eto, Y. Giga, and K. Ishii. “An area-minimizing scheme for anisotropic mean-curvature flow”. In: *Adv. Differential Equations* 17.11-12 (2012), pp. 1031–1084. ISSN: 1079-9389.
- [22] L. C. Evans and J. Spruck. “Motion of level sets by mean curvature. I”. In: *J. Differential Geom.* 33.3 (1991), pp. 635–681. URL: <http://projecteuclid.org/euclid.jdg/1214446559>.
- [23] N. Fusco, V. Julin, and M. Morini. “Stationary sets and asymptotic behavior of the mean curvature flow with forcing in the plane”. In: *J. Geom. Anal.* 32.2 (2022), Paper No. 53, 29. ISSN: 1050-6926. DOI: [10.1007/s12220-021-00806-x](https://doi.org/10.1007/s12220-021-00806-x).
- [24] Y. Giga, S. Goto, H. Ishii, and M.-H. Sato. “Comparison principle and convexity preserving properties for singular degenerate parabolic equations on unbounded domains”. In: *Indiana Univ. Math. J.* 40.2 (1991), pp. 443–470. ISSN: 0022-2518. DOI: [10.1512/iumj.1991.40.40023](https://doi.org/10.1512/iumj.1991.40.40023).
- [25] Y. Giga and N. Požár. “Motion by crystalline-like mean curvature: a survey”. In: *Bull. Math. Sci.* 12.2 (2022), Paper No. 2230004, 68. ISSN: 1664-3607. DOI: [10.1142/S1664360722300043](https://doi.org/10.1142/S1664360722300043).
- [26] M. E. Gurtin. “Toward a nonequilibrium thermodynamics of two-phase materials”. In: *Arch. Rational Mech. Anal.* 100.3 (1988), pp. 275–312. ISSN: 0003-9527. DOI: [10.1007/BF00251518](https://doi.org/10.1007/BF00251518). URL: <https://doi.org/10.1007/BF00251518>.
- [27] G. Huisken. “Contracting convex hypersurfaces in Riemannian manifolds by their mean curvature”. In: *Invent. Math.* 84.3 (1986), pp. 463–480. ISSN: 0020-9910. DOI: [10.1007/BF01388742](https://doi.org/10.1007/BF01388742).
- [28] G. Huisken and A. Polden. “Geometric evolution equations for hypersurfaces”. In: *Calculus of variations and geometric evolution problems (Cetraro, 1996)*. Vol. 1713. Lecture Notes in Math. Springer, Berlin, 1999, pp. 45–84. DOI: [10.1007/BFb0092669](https://doi.org/10.1007/BFb0092669).
- [29] J. E. Hutchinson. “Second fundamental form for varifolds and the existence of surfaces minimizing curvature”. In: *Indiana Univ. Math. J.* 35.1 (1986), pp. 45–71. ISSN: 0022-2518. DOI: [10.1512/iumj.1986.35.35003](https://doi.org/10.1512/iumj.1986.35.35003).
- [30] T. Ilmanen. “The level-set flow on a manifold”. In: *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990)*. Vol. 54. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1993, pp. 193–204. DOI: [10.1090/pspum/054.1/1216585](https://doi.org/10.1090/pspum/054.1/1216585).
- [31] H. Ishii and P. Souganidis. “Generalized motion of noncompact hypersurfaces with velocity having arbitrary growth on the curvature tensor”. In: *Tohoku Math. J. (2)* 47.2 (1995), pp. 227–250. ISSN: 0040-8735. DOI: [10.2748/tmj/1178225593](https://doi.org/10.2748/tmj/1178225593).
- [32] T. Laux and F. Otto. “Convergence of the thresholding scheme for multi-phase mean-curvature flow”. In: *Calc. Var. Partial Differential Equations* 55.5 (2016), Art. 129, 74. ISSN: 0944-2669. DOI: [10.1007/s00526-016-1053-0](https://doi.org/10.1007/s00526-016-1053-0).

- LucStu** [33] S. Luckhaus and T. Sturzenhecker. “Implicit time discretization for the mean curvature flow equation”. In: *Calc. Var. Partial Differential Equations* 3.2 (1995), pp. 253–271. ISSN: 0944-2669. DOI: [10.1007/BF01205007](https://doi.org/10.1007/BF01205007).
- Mag-book** [34] F. Maggi. *Sets of finite perimeter and geometric variational problems*. Vol. 135. Cambridge Studies in Advanced Mathematics. An introduction to geometric measure theory. Cambridge University Press, Cambridge, 2012, pp. xx+454. ISBN: 978-1-107-02103-7. DOI: [10.1017/CB09781139108133](https://doi.org/10.1017/CB09781139108133).
- MugSeiSpa** [35] L. Mugnai, C. Seis, and E. Spadaro. “Global solutions to the volume-preserving mean-curvature flow”. In: *Calc. Var. Partial Differential Equations* 55.1 (2016), Art. 18, 23. ISSN: 0944-2669. DOI: [10.1007/s00526-015-0943-x](https://doi.org/10.1007/s00526-015-0943-x).
- SchSimAlm** [36] R. Schoen, L. Simon, and F. J. Almgren Jr. “Regularity and singularity estimates on hypersurfaces minimizing parametric elliptic variational integrals. I, II”. In: *Acta Math.* 139.3-4 (1977), pp. 217–265. ISSN: 0001-5962. DOI: [10.1007/BF02392238](https://doi.org/10.1007/BF02392238).
- Set** [37] J. A. Sethian. *Level set methods and fast marching methods*. Second Edition. Vol. 3. Cambridge Monographs on Applied and Computational Mathematics. Evolving interfaces in computational geometry, fluid mechanics, computer vision, and materials science. Cambridge University Press, Cambridge, 1999, pp. xx+378.
- Tau** [38] G. Taubin. “A signal processing approach to fair surface design”. In: *Proceedings of the 22nd annual conference on Computer graphics and interactive techniques* (1995).