



HAL
open science

Certain Paracontact Metrics Satisfying the Critical Point Equation

Dhriti Patra

► **To cite this version:**

Dhriti Patra. Certain Paracontact Metrics Satisfying the Critical Point Equation. 2022. hal-03904450v1

HAL Id: hal-03904450

<https://hal.science/hal-03904450v1>

Preprint submitted on 16 Dec 2022 (v1), last revised 14 Feb 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Certain Paracontact Metrics Satisfying the Critical Point Equation

Dhriti Sundar Patra

Abstract: The aim of this paper is to study the CPE (Critical Point Equation) on some paracontact metric manifolds. First, we prove that if a para-Sasakian metric satisfies the CPE, then it is Einstein with constant scalar curvature $-2n(2n+1)$. Next, we prove that if (κ, μ) -paracontact metric satisfies the CPE, then it is locally isometric to the product of a flat $(n+1)$ -dimensional manifold and n -dimensional manifold of negative constant curvature -4 .

Mathematics Subject Classification 2010: 53C15, 53C25, 53D10

Keywords: The critical point equation, paracontact metric manifold, para-Sasakian manifold, (κ, μ) -paracontact manifold

1 Introduction

In [8], Kaneyuki and Williams introduce a new structure on pseudo-Riemannian geometry called paracontact structure, as a natural odd-dimensional counterpart to para-Hermitian structures, just like contact metric structures correspond to the Hermitian one. The importance of paracontact geometry comes from the theory of para-Kähler manifolds. A systematic study of paracontact metric manifolds and their subclasses was started by Zamkovoy [19]. Since then, many authors investigated paracontact geometry using various meaningful geometric conditions. We refer the reader to ([7, 9–18]) for some related results on paracontact geometry.

On the other hand, in [1] (for details, see Chapter 2), A. Besse study the the Hilbert-Einstein functional and proved that the critical points of this functional are the Einstein metrics. The Hilbert-Einstein functional has the following Euler-Lagrange equation

$$\mathcal{L}_g^*(\lambda) = -(\Delta_g \lambda)g + \text{Hess}_g \lambda - \lambda \text{Ric}_g = \text{Ric}_g^o, \quad (1.1)$$

for a critical point g , where Δ_g , Ric_g , Ric_g^o and $\text{Hess} \lambda$ are respectively the Laplacian, the Ricci tensor, the traceless Ricci tensor and the Hessian of the smooth function λ on M . Here $\mathcal{L}_g^*(\lambda)$ is the formal L^2 -adjoint of the linearized scalar curvature operator $\mathcal{L}_g(\lambda)$. The Eq. (1.1) is called the Critical Point Equation (shortly, CPE). The function λ is known as the potential function. From now, we consider a metric g with a non-trivial potential function λ as a solution of the

CPE and is denoted by (g, λ) . Also, we can express the equation (1.1) in the form

$$\left(\text{Ric}_g - \frac{r}{n-1}g\right)\lambda - \text{Hess}_g\lambda = \frac{r}{n}g - \text{Ric}_g. \quad (1.2)$$

First, we note that if λ is constant in the equation (1.2), then $\lambda = 0$ and g becomes Einstein. Further, tracing (1.2) we deduce $\Delta_g\lambda = -\frac{r}{n-1}\lambda$. From which it follows that λ is an eigenfunction of the Laplacian. Since the Laplacian has non-positive spectrum, the scalar curvature must be positive. In [1], A. Besse first conjectured that the solution of the CPE must be Einstein (known as the CPE conjecture). In my knowledge many authors study the CPE satisfying either, some curvature conditions or, some conditions on the potential functions on Riemannian manifolds. In [2], Barros and Ribeiro proved that the CPE conjecture is also true for half conformally flat. Further in [3], Hwang proved that the CPE conjecture is valid under certain conditions on the bounds of the potential function λ . Recently, Nato [4] obtained a necessary and sufficient condition on the norm of the gradient of the potential function for a CPE metric to be Einstein. Recently, the author considered the CPE on contact metric manifolds (see [5, 6]) and proved that the CPE conjecture is true for K -contact manifold. However, the CPE has not yet been considered on pseudo-Riemannian manifolds, for instance paracontact metric manifolds. Hence it deserves special attention to consider the CPE on certain class of paracontact metric manifolds. Here we characterize the solution of the CPE on certain classes of paracontact metric manifolds and prove that the CPE conjecture is true for para-Sasakian manifold.

The paper is structured as follows: in Section 2, very brief review of paracontact geometry is given. Next, we consider the CPE on para-Sasakian manifold in Section 3 and we prove that a if a para-Sasakian metric satisfies the CPE, then it is Einstein with Einstein constant $-2n$ and has constant scalar curvature. Finally, in Section 4, we consider the CPE on (κ, μ) -paracontact manifold and prove that if (κ, μ) -paracontact metric satisfies the CPE, then it is locally isometric to the product of a flat $(n+1)$ -dimensional manifold and n -dimensional manifold of negative constant curvature.

2 Preliminaries

In this section, we shall collect some fundamental results regarding paracontact metric manifolds (for more details see [17, 19]). A $(2n+1)$ -dimensional smooth manifold M^{2n+1} has an almost paracontact structure (φ, ξ, η) if it admits a $(1, 1)$ -tensor field φ , a vector field ξ and a 1-form η such that

$$\varphi^2 = I - \eta\circ\xi, \quad \varphi(\xi) = 0, \quad \eta\circ\varphi = 0, \quad \eta(\xi) = 1, \quad (2.1)$$

and there exists a distribution $\mathcal{D} : p \in M \rightarrow \mathcal{D}_p \subset T_p M : \mathcal{D}_p = \text{Ker}(\eta) = \{x \in T_p M : \eta(x) = 0\}$, called paracontact distribution generated by η . Let $\mathfrak{X}(M)$ is the Lie algebra of all vector fields on M^{2n+1} . If an almost paracontact manifold admits a pseudo-Riemannian metric g such that

$$g(\varphi X, \varphi Y) = -g(X, Y) + \eta(X)\eta(Y), \quad X, Y \in \mathfrak{X}(M), \quad (2.2)$$

then we say that M has an almost paracontact metric structure (φ, ξ, η, g) and g is called compatible metric. Any compatible metric g with a given almost paracontact structure is necessarily of signature $(n+1, n)$. The fundamental 2-form Φ of an almost paracontact metric structure (φ, ξ, η, g) defined by

$$\Phi(X, Y) = g(X, \varphi Y), \quad X, Y \in \mathfrak{X}(M).$$

If $\Phi = d\eta$, then the manifold $M^{2n+1}(\varphi, \xi, \eta, g)$ is called a paracontact metric manifold. On a paracontact metric manifold $M^{2n+1}(\varphi, \xi, \eta, g)$, we consider a self-adjoint operator $h = \frac{1}{2}\mathcal{L}_\xi\varphi$, where \mathcal{L}_ξ is the Lie-derivative along ξ . The operator h satisfy [19]:

$$\text{Tr}_g h = 0, \quad h\xi = 0, \quad h\varphi = -\varphi h.$$

On a paracontact metric manifold [19]:

$$\nabla_X \xi = -\varphi X + \varphi hX, \quad X \in \mathfrak{X}(M), \quad (2.3)$$

where ∇ is the operator of covariant differentiation of g . If the vector field ξ is a Killing (equivalently $h = 0$) then M is said to be a K -paracontact manifold. Moreover, on any K -paracontact manifold [19]:

$$\nabla_X \xi = -\varphi X, \quad (2.4)$$

$$R(X, Y)\xi = \eta(X)Y - \eta(Y)X, \quad (2.5)$$

$$Q\xi = -2n\xi, \quad (2.6)$$

for all vector fields X, Y on M , where R is the Riemann curvature tensor of g and Q denotes the Ricci operator associated with the Ricci tensor given by $\text{Ric}_g(X, Y) = g(QX, Y)$ for all vector fields X, Y on M .

Moreover, a paracontact metric manifold $M^{2n+1}(\varphi, \xi, \eta, g)$ is said to be para-Sasakian if the following integrability condition is satisfied $[\varphi, \varphi] = -2d\eta \otimes \xi$, where $[\varphi, \varphi]$ denotes the Nijenhuis

tensor of φ . Equivalently, a paracontact metric manifold is said to para-Sasakian if

$$(\nabla_X \varphi)Y = -g(X, Y)\xi + \eta(Y)X,$$

for all vector fields X, Y on M , see [19]. A para-Sasakian manifold is K -paracontact [13] but the converse is true only in dimension 3, see [11].

3 On K -paracontact and para-Sasakian manifolds

In this section, we study the CPE on K -paracontact and para-Sasakian manifolds. Before entering into our main results we prove the following.

Lemma 3.1 *On a K -paracontact manifold $M^{2n+1}(\varphi, \xi, \eta, g)$, we have*

$$(i) (\nabla_X Q)\xi = Q\varphi X + 2n\varphi X,$$

$$(ii) (\nabla_\xi Q)X = Q\varphi X - \varphi QX,$$

for all vector fields X on M .

Proof: First, taking the covariant derivative of (2.6) along an arbitrary vector field X on M and then using the result (2.4) we have the first one. Further, since ξ is Killing, we have $\mathcal{L}_\xi \text{Ric}_g = 0$. This implies $(\mathcal{L}_\xi Q)X = 0$ for any vector field X on M . From which it follows

$$\begin{aligned} 0 &= \mathcal{L}_\xi(QX) - Q(\mathcal{L}_\xi X) \\ &= \nabla_\xi QX - \nabla_{QX}\xi - Q(\nabla_\xi X) + Q(\nabla_X \xi) \\ &= (\nabla_\xi Q)X - \nabla_{QX}\xi + Q(\nabla_X \xi), \end{aligned}$$

for any vector field X on M . Using (2.4) the last equation gives the second result. This completes the proof. \square

Lemma 3.2 *Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a K -paracontact manifold. If (g, λ) is a non-constant solution of the CPE, then we have*

$$(Q\varphi + \varphi Q)X = -4n\varphi X \tag{3.1}$$

for all vector fields X on M .

Proof: First, we note that (1.1) implies $\Delta_g \lambda = -\frac{r\lambda}{2n}$. Thus, equation (1.1) can be exhibited as

$$\nabla_X D\lambda = (\lambda + 1)QX + fX, \tag{3.2}$$

for all vector fields X on M ; where $f = -r(\frac{\lambda}{2n} + \frac{1}{2n+1})$. Using this in the well known expression of the curvature tensor $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$, we obtain

$$\begin{aligned} R(X, Y)D\lambda &= (X\lambda)QY - (Y\lambda)QX + (\lambda + 1)\{(\nabla_X Q)Y - (\nabla_Y Q)X\} \\ &\quad + (Xf)Y - (Yf)X, \end{aligned} \quad (3.3)$$

for all vector fields X, Y on M . Now, substituting X by ξ in (3.3) and using the Lemma 3.1 we can compute

$$\begin{aligned} R(\xi, Y)D\lambda &= (\xi\lambda)QY + 2n(Y\lambda)\xi - (\lambda + 1)\{\varphi QY + 2n\varphi Y\} \\ &\quad + (\xi f)Y - (Yf)\xi, \end{aligned} \quad (3.4)$$

for any vector fields Y on M . On the other hand, we obtain from (2.4) that

$$g(R(X, Y)\xi, Z) = g(\nabla_X \varphi)Y, Z) - g(\nabla_Y \varphi)X, Z),$$

for all vector fields X, Y on M . Applying Bianchi's first identity and the last equation we achieve $R(\xi, X)Y = (\nabla_\varphi X)Y$ for all vector fields X, Y on M . Making use of this in the scalar product of (3.4) with an arbitrary vector field X on M provides

$$\begin{aligned} g((\nabla_Y \varphi)X, D\lambda) &= (\lambda + 1)\{g(Q\varphi Y, X) + 2ng(\varphi Y, X)\} - (\xi\lambda)g(QY, X) \\ &\quad - (\xi f)g(X, Y) - \{2n(Y\lambda) - (Yf)\}\eta(X). \end{aligned} \quad (3.5)$$

Taking into account of (2.1), (2.4) and putting $X = \varphi X, Y = \varphi Y$ in (3.5) yields

$$\begin{aligned} g((\nabla_{\varphi X} \varphi)\varphi X, D\lambda) &= (\lambda + 1)\{2ng(Y, \varphi X) - g(Q\varphi Y, X)\} \\ &\quad - (\xi\lambda)g(Q\varphi Y, \varphi X) + (\xi f)\{g(X, Y) - \eta(X)\eta(Y)\}, \end{aligned} \quad (3.6)$$

Now, adding (3.5) with (3.6) and using (2.1), (2.6) and the well known formula (see [19])

$$(\nabla_{\varphi X} \varphi)\varphi Y - (\nabla_X \varphi)Y = 2g(X, Y)\xi - \eta(Y)(X - hX + \eta(X)\xi),$$

one can compute

$$\begin{aligned} 2\xi(\lambda - f)g(X, Y) &= Y\{(2n + 1)\lambda - f\}\eta(X) \\ &\quad + \xi(\lambda - f)\eta(X)\eta(Y) - (\lambda + 1)g(Q\varphi Y + \varphi QY, X) \\ &\quad + 4n(\lambda + 1)g(\varphi X, Y) - (\xi\lambda)\{g(Q\varphi Y, \varphi X) - g(QY, X)\}. \end{aligned}$$

Anti-symmetrizing the foregoing equation yields

$$Y\{(2n+1)\lambda - f\}\eta(X) - X\{(2n+1)\lambda - f\}\eta(Y) + 8n(\lambda+1)g(\varphi X, Y) + 2(\lambda+1)g(Q\varphi X + \varphi QX, Y) = 0. \quad (3.7)$$

At this point, we replace X by φX and Y by φY in (3.7) to achieve

$$(\lambda+1)\left\{g(Q\varphi X, Y) + g(\varphi QX, Y) + 4ng(\varphi X, Y)\right\} = 0.$$

Since λ is non-constant in the interior of M , the last equation gives us the required result. \square

Thus, from last Lemma, one can easily conclude the following result.

Theorem 3.1 *Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a K -paracontact manifold and the Ricci operator Q commutes with paracontact structure φ . If (g, λ) is a non-constant solution of the CPE, then g is Einstein with Einstein constant $-2n$ and has constant scalar curvature $-2n(2n+1)$.*

Proof: By hypothesis, the Ricci operator Q commutes with paracontact structure φ , i.e., $Q\varphi = \varphi Q$. Applying this in (3.1) and then substituting X by φX , and using (2.1) gives that $QX = -2nX$ for any vector fields X . This shows that M is Einstein with Einstein constant $-2n$. This completes that proof. \square

On para-Sasakian manifold, the Ricci operator satisfies (see [19], Lemma 3.15):

$$QX - \varphi Q\varphi X = -2n\eta(X)\xi, \quad (3.8)$$

for all vector fields X on M . Operating (3.8) by φ and using (2.1) we see that the Ricci operator Q commutes with paracontact structure φ . Hence, from Theorem 3.1, we can deduce the following result.

Theorem 3.2 *Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a para-Sasakian manifold. If (g, λ) is a non-constant solution of the CPE, then g is Einstein with Einstein constant $-2n$ and has constant scalar curvature $-2n(2n+1)$.*

Remark 3.1 *There are some partial answers to the CPE Conjecture. For example, Lafontaine proved that the CPE conjecture is true under conformally flat assumption with $\text{Ker } \mathcal{L}_g^*(\lambda) \neq 0$. Further, Barros and Ribeiro [2] proved that the CPE conjecture is also true for half conformally flat. Therefore, in this section we prove the CPE conjecture in a subclass paracontact metric manifolds, for instance, para-Sasakian manifolds.*

4 On (κ, μ) -paracontact manifolds

In [12], Cappelletti-Montano et al introduce the notion of nullity conditions in paracontact geometry. According to them a (κ, μ) -paracontact manifold is a paracontact metric manifold $M^{2n+1}(\varphi, \xi, \eta, g)$ whose curvature tensor satisfies

$$R(X, Y)\xi = \kappa\{\eta(Y)X - \eta(X)Y\} + \mu\{\eta(Y)hX - \eta(X)hY\}, \quad (4.1)$$

for all vector fields X, Y on M and for some real numbers (κ, μ) . The class of (κ, μ) -paracontact metric manifold contains para-Sasakian manifolds. Since then, many geometers have studied (κ, μ) -paracontact manifold and obtained various important properties of these manifolds (see, for instance, [20]). On (κ, μ) -paracontact manifold the following formulas are valid (e.g., [12]):

$$h^2 = (\kappa + 1)\varphi^2, \quad (4.2)$$

$$Q\xi = 2n\kappa\xi. \quad (4.3)$$

First of all, we recall a lemma for our main proof.

Lemma 4.1 (See [12]) *Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a (κ, μ) -paracontact manifold with $\kappa > -1$, then the Ricci operator Q of M can be expressed as*

$$\begin{aligned} QX &= [2(1-n) + n\mu]X + [2(n-1) + \mu]hX \\ &\quad + [2(n-1) + n(2\kappa - \mu)]\eta(X)\xi, \end{aligned} \quad (4.4)$$

for any vector field X on M . In this case, the scalar curvature of M is $2n(2(1-n) + \kappa + n\mu)$.

Now, we prove a lemma for later use.

Lemma 4.2 *On a (κ, μ) -paracontact manifold $M^{2n+1}(\varphi, \xi, \eta, g)$ with $\kappa > -1$, we have*

$$(\nabla_\xi Q)X = \mu(2(n-1) + \mu)h\varphi X, \quad (4.5)$$

for any vector field X in M .

Proof: Replacing Y by ξ in (4.1) and using $\varphi^2 = I - \eta \otimes \xi$, we can easily deduce that

$$R(X, \xi)\xi = \kappa\{X - \eta(X)\xi\} + \mu hX, \quad (4.6)$$

for any vector field X in M . Recalling a formula on paracontact metric manifold [19]:

$$(\nabla_\xi h)X = -\varphi X + \varphi h^2 X - \varphi R(X, \xi)\xi, \quad (4.7)$$

for any vector field X in M . By virtue of (2.1), (4.2), (4.6) and $\varphi h = -h\varphi$, the Eq. (4.7) yields

$$(\nabla_\xi h)X = -\mu\varphi hX, \quad (4.8)$$

for any vector field X in M . Covariant derivative of (4.4) along ξ , we have

$$\begin{aligned} (\nabla_\xi Q)X + Q(\nabla_\xi X) = & \{2(1-n) + n\mu\}\nabla_\xi X + \{2(n-1) + \mu\}\{(\nabla_\xi h)X \\ & + h(\nabla_\xi X)\} + \{2(n-1) + n(2\kappa - \mu)\}g(\nabla_\xi X, \xi)\xi, \end{aligned} \quad (4.9)$$

for any vector field X on M . Next, putting $X = \nabla_\xi X$ in (4.4), we get

$$\begin{aligned} Q(\nabla_\xi X) = & \{2(1-n) + n\mu\}\nabla_\xi X + \{2(n-1) + \mu\}h(\nabla_\xi X) \\ & + \{2(n-1) + n(2\kappa - \mu)\}g(\nabla_\xi X, \xi)\xi. \end{aligned}$$

Using this and (4.8) in (4.9) we get the required result. \square

Lemma 4.3 *Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a (κ, μ) -paracontact manifold with $\kappa > -1$. If (g, λ) is a non-constant solution of the CPE, then we have*

$$\kappa(2 - \mu) = \mu(n + 1). \quad (4.10)$$

Proof: First, differentiating (4.3) along an arbitrary vector field X on M and using (2.3) it follows that

$$(\nabla_X Q)\xi = Q(\varphi - \varphi h)X - 2n\kappa(\varphi - \varphi h)X. \quad (4.11)$$

Thus, the scalar product of (3.3) with ξ and making use of (4.3), (4.11) gives that

$$\begin{aligned} g(R(X, Y)D\lambda, \xi) = & 2n\kappa[(X\lambda)\eta(Y) - (Y\lambda)\eta(X)] + (\lambda + 1)g(Q\varphi X + \varphi QX, Y) \\ & - (\lambda + 1)g(Q\varphi hX + h\varphi QX, Y) + 4n\kappa(\lambda + 1)g(\varphi X, Y) \\ & + (Xf)\eta(Y) - (Yf)\eta(X). \end{aligned} \quad (4.12)$$

Replacing X by φX , Y by φY in (4.12) and noting that $R(\varphi X, \varphi Y)\xi = 0$ (follows from (4.1)), we obtain

$$(\lambda + 1)[Q\varphi X + \varphi QX - \varphi QhX - hQ\varphi X - 4n\kappa\varphi X] = 0,$$

for any vector field X on M . Since λ is non-constant in the interior M , the foregoing equation provides

$$Q\varphi X + \varphi QX + \varphi QhX + hQ\varphi X - 4n\kappa\varphi X = 0, \quad (4.13)$$

for any vector field X on M . Now, substituting X by φX in the relation (4.4) gives that

$$Q\varphi X = [2(1-n) + n\mu] \varphi X + [2(n-1) + \mu] h\varphi X.$$

On the other hand, by acting h on the previous equation and making use of (4.2) implies that

$$hQ\varphi X = [2(1-n) + n\mu] h\varphi X + (\kappa + 1)[2(n-1) + \mu] \varphi X.$$

Also, operating φ on (4.4) gives

$$\varphi QX = [2(1-n) + n\mu] \varphi X + [2(n-1) + \mu] \varphi hX.$$

Taking hX instead of X and using (4.2), the last equation reduces to

$$\varphi QhX = [2(1-n) + n\mu] \varphi hX + (\kappa + 1)[2(n-1) + \mu] \varphi X.$$

By help of the last four equations in (4.13) and using (4.2) we obtain the required result. Hence the proof. \square

Theorem 4.1 *Let $M^{2n+1}(\varphi, \xi, \eta, g)$ be a (κ, μ) -paracontact manifold $\kappa > -1$. If (g, λ) is a non-constant solution of the CPE, then M is locally isometric to the product of a flat $(n+1)$ -dimensional manifold and n -dimensional manifold of negative constant curvature -4 .*

Proof: Taking into account of (4.3) and (4.6) and substituting ξ instead of X in (4.12) we obtain

$$(2n+1)\kappa\{D\lambda - (\xi\lambda)\xi\} + \mu hD\lambda - (\xi f)\xi + Df = 0. \quad (4.14)$$

Contracting (3.3) over X with respect to an orthonormal basis $\{e_i\}$, we have

$$\lambda \left\{ \sum_{i=1}^{2n+1} g((\nabla_{e_i} Q)Y, e_i) - g((\nabla_Y Q)e_i, e_i) \right\} - r(Y\lambda) - 2n(Yf) = 0. \quad (4.15)$$

As the scalar curvature $r = 2n\{2(1-n) + \kappa + n\mu\}$ (from Lemma 4.1) is constant, using the formula $\operatorname{div} Q = \frac{1}{2}dr$ (follows from contraction of Bianchi's second identity) one can compute

$$\begin{aligned} \sum_{i=1}^{2n+1} g((\nabla_{e_i} Q)Y, e_i) - g((\nabla_Y Q)e_i, e_i) &= (\operatorname{div} Q)Y - Y(\operatorname{Tr}_g Q) \\ &= \frac{1}{2}(Yr) - (Yr) = -\frac{1}{2}(Yr) = 0. \end{aligned}$$

Using this in (4.15) yields

$$r D\lambda + 2n Df = 0. \quad (4.16)$$

Combining this with (4.14), we get

$$2n(2n + 1)\kappa\{D\lambda - (\xi\lambda)\xi\} + 2n\mu hD\lambda - 2n(\xi f)\xi - r D\lambda = 0. \quad (4.17)$$

From (3.2) and (4.3), we have

$$\nabla_\xi D\lambda = [2n\kappa(\lambda + 1) + f]\xi. \quad (4.18)$$

Now, taking covariant derivative of (4.17) along ξ and using (4.5), (4.8), (4.18), we obtain

$$\begin{aligned} & \{2n(2n + 1)\kappa - r\}\{2n\kappa(\lambda + 1) + f\}\xi - 2n(2n + 1)\kappa\xi(\xi\lambda)\xi \\ & + 2n\mu^2 h\varphi D\lambda - 2n\xi(\xi f)\xi = 0. \end{aligned} \quad (4.19)$$

Next, operating (4.19) by φ and using (2.1) we have $\mu^2 hD\lambda = 0$. By the action of h and using (4.2), (2.1) gives

$$(\kappa + 1)\mu^2 (D\lambda - (\xi\lambda)\xi) = 0.$$

As $\kappa > -1$, we have either (i) $\mu = 0$, or (ii) $\mu \neq 0$.

Case (i): In this case, it follows from (4.10) that $\kappa = 0$. Hence $R(X, Y)\xi = 0$ for any vector field X, Y on M , and therefore M^{2n+1} is the product of a flat $(n + 1)$ -dimensional manifold and n -dimensional manifold of negative constant curvature -4 (see [9], Theorem 3.3).

Case (ii): This case yields $D\lambda = (\xi\lambda)\xi$. Differentiating this along an arbitrary vector field X together with (2.3) entails that $\nabla_X D\lambda = X(\xi\lambda)\xi - (\xi\lambda)(\varphi X - \varphi hX)$. Since $g(\nabla_X D\lambda, Y) = g(\nabla_Y D\lambda, X)$, the foregoing equation shows

$$X(\xi\lambda)\eta(Y) - Y(\xi\lambda)\eta(X) + (\xi\lambda) d\eta(X, Y) = 0.$$

Since $d\eta$ is non-zero for a paracontact metric structure, replacing X by φX and Y by φY in previous result, we get $\xi\lambda = 0$. Hence $D\lambda = 0$, and therefore, λ is constant, which is a contradiction. This completes the proof. \square

Acknowledgments: The author is thankful to the reviewer for some valuable comments that improves the paper.

References

- [1] Besse A., *Einstein manifolds*, Springer-Verlag, New York, 2008.
- [2] Barros A. and Ribeiro Jr. E., *Critical point equation on four-dimensional compact manifolds*, Math. Nachr., **287**(2014), 1618–1623.
- [3] Hwang S., *Critical points of the total scalar curvature functionals on the space of metrics of constant scalar curvature*, Manuscripta Math., **103**(2000), 135–142.
- [4] Nato B. L., *A note on critical point metrics of the total scalar curvature functional*, J. Math. Anal. Appl. **424**(2015), 1544–1548.
- [5] Ghosh A. and Patra D. S., *The critical point equation and contact geometry*, J. Geom., **108**(2017), 185–194.
- [6] Patra D. S., Ghosh A. and Bhattacharyya A., *The Critical Point Equation on Kenmotsu and almost Kenmotsu manifolds*, Publ. Math. Debrecen, **97**(2020), 85–99.
- [7] Patra D.S., Ali A. and Mofarreh F., *Characterizations of Ricci-Bourguignon almost solitons on pseudo-Riemannian manifolds*, Mediterranean Journal of Mathematics, **19**, (2022), 176.
- [8] Kaneyuki S. and Williams F. L., *Almost paracontact and parahodge structures on manifolds*, Nagoya Math. J., **99** (1985), 173–187.
- [9] Zamkovoy S. and Tzanov V., *Non-existence of flat paracontact metric structures in dimension greater than or equal to five*, Annuaire Univ. Sofia Fac. Math. Inform., **100**(2011), 27–34.
- [10] Bejan C. L. and Crasmareanu M., *Second order parallel tensors and Ricci solitons in 3-dimensional normal paracontact geometry*, Ann. Glob. Anal. Geom., **46**(2) (2014), 117–127.
- [11] Martin-Molina V., *Local classification and examples of an important class of paracontact metric manifolds*, Filomat, **29**(3) (2015), 507–515.
- [12] Cappelletti-Montano B., Erken I. K. and Murathan C., *Nullity conditions in paracontact geometry*, Diff. Geom. Appl., **30**(2012), 665–693.
- [13] Calvaruso G., *Homogeneous paracontact metric three-manifolds*, Ill. J. Math., **55**(2011), 697–718.
- [14] Montano B. C., Carriazo A. and Martin-Molina V., *Sasaki-Einstein and paraSasaki-Einstein metrics from (κ, μ) -structures*, J. Geom. Phys., **73**(2013), 20–36.

- [15] Perrone A., *Some results on almost paracontact metric manifolds*, Mediterr. J. Math., **13**(5) (2016), 3311–3326.
- [16] Venkatesha and Naik D. M., *Certain results on K -paracontact and para-Sasakian manifolds*, J. Geom., **108**(3) (2017), 939–952.
- [17] Calvaruso G. and Perrone D., *Geometry of H -paracontact metric manifolds*, Publ. Math. Debrecen, **86** (2015), 325–346.
- [18] Patra D.S., *Ricci soliton and paracontact geometry*, Mediterr. J. Math. **16**, 137 (2019)
- [19] Zamkovoy S., *Canonical connection on paracontact manifolds*, Ann. Glob. Anal. Geom., **36**(1) (2009), 37–60.
- [20] Kupeli Erken I. and Murathan C., *A study of three dimensional paracontact (κ, μ) -spaces*, Int. J. Geom. Methods Mod. Phys., **14** (2017), no. 7, 1750106.

Dhriti Sundar Patra,
Department of Mathematics,
Indian Institute of Technology - Hyderabad,
Sangareddy-502285, India,
Email: dhriti@math.iith.ac.in; dhritimath@gmail.com