



HAL
open science

Propagation of chaos for topological interactions by a coupling technique

Pierre Degond, Mario Pulvirenti, Stefano Rossi

► **To cite this version:**

Pierre Degond, Mario Pulvirenti, Stefano Rossi. Propagation of chaos for topological interactions by a coupling technique. *Rendiconti Lincei. Matematica e Applicazioni*, 2023, 34, pp.641-655. hal-03904962

HAL Id: hal-03904962

<https://hal.science/hal-03904962>

Submitted on 17 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROPAGATION OF CHAOS FOR TOPOLOGICAL INTERACTIONS BY A COUPLING TECHNIQUE

PIERRE DEGOND, MARIO PULVIRENTI, AND STEFANO ROSSI

ABSTRACT. We consider a system of particles which interact through a jump process. The jump intensities are functions of the proximity rank of the particles, a type of interaction referred to as topological in the literature. Such interactions have been shown relevant for the modelling of bird flocks. We show that, in the large number of particles limit and under minimal smoothness assumptions on the data, the model converges to a kinetic equation which was derived in earlier works both formally and rigorously under more stringent regularity assumptions. The proof relies on the coupling method which assigns to the particle and limiting processes a joint process posed on the cartesian product of the two configuration spaces of the former processes. By appropriate estimates in a suitable Wasserstein metric, we show that the distance between the two processes tends to zero as the number of particles tends to infinity, with an error typical of the law of large numbers.

1. INTRODUCTION

Systems of self-propelled agents undergoing local interactions are ubiquitous in nature, from migrating cells [16] to locust swarms [2] and fish schools [18]. They form intriguing patterns such as coherent motion, travelling bands, oscillations etc. encompassed in the generic term of collective dynamics (see a review in [24]). Most models of collective dynamics are based on mean-field interactions (such as the Cucker-Smale [12] or Vicsek [23] models) or binary contact interactions [4]. However, a third type of interaction has been suggested following observations of bird flocks [1, 9] and referred to as “topological interaction”. In this kind of interaction, the strength of the interaction of an agent with another one is a function of the proximity rank of the latter with respect to the former. The seminal paper [1] has been followed by a number of papers studying various aspects of this phenomenon see e.g. [7, 8, 15, 20, 21].

Mathematically, flocking of systems of topologically interacting particles have been investigated in [19, 22, 26]. In [17], in addition to studying flocking, the author proposes kinetic and fluid models derived from mean-field topological interactions. The present work is strongly aligned with [5, 6, 13] where kinetic models are derived for topological interaction models based on jump processes. More precisely, [13] proves propagation of chaos and provides a rigorous proof of the model formally derived in [5]. The proof of [13] makes the limiting assumption that the interaction strength is an analytic function

Date: December 16, 2022.

MSC2020 subject classifications. 35Q92, 70K55, 92D50.

Key words and phrases. Propagation of chaos, Rank-based interaction, Coupling method.

of the normalized rank (a concept precisely defined below) and is based on the BBGKY hierarchy. In the present work, we propose an alternative proof of the result of [13] based on the coupling method. The advantage of the coupling method over the BBGKY hierarchy is that it only requires the interaction strength to be Lipschitz continuous, a much more general and natural assumption than that of [13]. On the other hand, [6] formally derives a kinetic model for a more singular interaction. The mathematical validity of this formal result is still open. The literature on propagation of chaos and derivation of kinetic models from particle ones is huge and it is difficult to provide a fair account of all relevant contributions in a short introduction. We refer the interested reader to the reviews [10, 11] which provide a fairly detailed description of the subject.

The outline of this paper is as follows. In Section 2, we present the model and provide a formal derivation of the macroscopic model. We then state the theorem and comment it in view of the previous results. Section 3 is devoted to the proof.

2. PRESENTATION OF THE MODEL AND MAIN RESULTS

We recall the model and notations introduced in [5, 13] and state our result. We study a N -particle system in \mathbb{R}^d , $d = 1, 2, 3 \dots$ (or in \mathbb{T}^d the d -dimensional torus). Each particle, say particle i , has a position x_i and velocity v_i . The configuration of the system is denoted by

$$Z_N = \{z_i\}_{i=1}^N = \{(x_i, v_i)\}_{i=1}^N = (X_N, V_N).$$

Given the particle i , we order the remaining particles j_1, j_2, \dots, j_{N-1} according to their distance from i , namely by the following relation

$$|x_i - x_{j_h}| \leq |x_i - x_{j_{h+1}}|, \quad h = 1, 2, \dots, N-1.$$

The rank $R(i, k)$ of particle $k = j_h$ (with respect to i) is h . Note that, if $B_r(x)$ denotes the closed ball of center $x \in \mathbb{R}^d$ and radius $r > 0$, we have

$$R(i, k) = \sum_{\substack{1 \leq h \leq N \\ h \neq i}} \mathcal{X}_{B_{|x_i - x_k|}(x_i)}(x_h),$$

where \mathcal{X}_A is the characteristic function of the set A .

Given a non-increasing Lipschitz continuous function

$$K : [0, 1] \rightarrow \mathbb{R}^+ \quad \text{s.t.} \quad \int_0^1 K(r) dr = 1,$$

we introduce the transition probabilities

$$\pi_{i,j}^N = \frac{K(r(i, j))}{\sum_s K(\frac{s}{N-1})}, \quad (2.1)$$

where $r(i, j)$ is the normalized rank:

$$r(i, j) = \frac{R(i, j)}{N-1} \in \left\{ \frac{1}{N-1}, \frac{2}{N-1}, \dots \right\}.$$

Thanks to the normalization in (2.1), we have that $\sum_j \pi_{i,j}^N = 1$. We can also rewrite $\pi_{i,j}^N$ as

$$\pi_{i,j}^N = \alpha_N K(r(i, j)), \quad (2.2)$$

where

$$\alpha_N = \frac{1}{(N-1)(1-e_K(N))} \quad (2.3)$$

and $e_K(N)$ is the error given by the Riemann sums

$$e_K(N) = \int_0^1 K(x) dx - \frac{1}{N-1} \sum_s K\left(\frac{s}{N-1}\right). \quad (2.4)$$

We are now in position to introduce a stochastic process describing alignment via a topological interaction. The particles go freely: $x_i + v_i t$. At some random time dictated by a Poisson process of intensity N , choose a particle (say i) with probability $\frac{1}{N}$ and a partner particle, say j , with probability $\pi_{i,j}$. Then perform the transition

$$(v_i, v_j) \rightarrow (v_j, v_j).$$

After that the system goes freely with the new velocities and so on.

The process is described by the following Markov generator given, for any $\Phi \in C_b^1(\mathbb{R}^{2dN})$, by

$$\begin{aligned} L_N \Phi(X_N, V_N) &= \sum_{i=1}^N v_i \cdot \nabla_{x_i} \Phi(X_N, V_N) \\ &+ \sum_{i=1}^N \sum_{\substack{1 \leq j \leq N \\ i \neq j}} \pi_{i,j}^N [\Phi(X_N, V_N^i(v_j)) - \Phi(X_N, V_N)], \end{aligned} \quad (2.5)$$

where $V_N^i(v_j) = (v_1 \dots v_{i-1}, v_j, v_{i+1} \dots v_N)$ if $V_N = (v_1 \dots v_{i-1}, v_i, v_{i+1} \dots v_N)$.

Note that $\pi_{i,j}^N$ depends not only on N but also on the whole spatial configuration X_N . Therefore the law of the process $W^N(t) = W^N(Z_N; t)$ is driven by the following evolution equation

$$\begin{aligned} \partial_t \int W^N(t) \Phi &= \int W^N(t) \sum_{i=1}^N v_i \cdot \nabla_{x_i} \Phi \\ &+ \int W^N(Z_N; t) \sum_{i=1}^N \sum_{\substack{1 \leq j \leq N \\ i \neq j}} \pi_{i,j}^N [\Phi(X_N, V_N^i(v_j)) - \Phi(X_N, V_N)], \end{aligned} \quad (2.6)$$

for any test function Φ .

We assume that the initial measure $W^N(0)$ factorizes, namely $W^N(0) = f_0^{\otimes N}$ where f_0 is the initial datum for the limiting kinetic equation we are going to establish. Note also that $W^N(Z_N; t)$, for $t \geq 0$, is symmetric in the exchange of particles.

The strong form of equation (2.6) is

$$\left(\partial_t + \sum_{i=1}^N v_i \cdot \nabla_{x_i} \right) W^N(t) = -N W^N(t) + \mathcal{L}_N W^N(t)$$

where

$$\mathcal{L}_N W^N(X_N, V_N; t) = \sum_{i=1}^N \sum_{\substack{1 \leq j \leq N \\ i \neq j}} \int du \pi_{i,j}^N W^N(X_N, V_N^{(i)}(u); t) \delta(v_i - v_j).$$

2.1. Heuristic derivation. We now want to derive the kinetic equation we expect to be valid in the limit $N \rightarrow \infty$. Setting $\Phi(Z_N) = \varphi(z_1)$ in (2.6), we obtain

$$\partial_t \int f_1^N \varphi = \int f_1^N v \cdot \nabla_x \varphi - \int f_1^N \varphi + \int W^N \sum_{j \neq 1} \pi_{i,j}^N \varphi(x_1, v_j). \quad (2.7)$$

Here f_1^N denotes the one-particle marginal of the measure W^N . We recall that the s -particle marginals are defined by

$$f_s^N(Z_s) = \int W^N(Z_s, z_{s+1} \cdots z_N) dz_{s+1} \cdots dz_N, \quad s = 1, 2, \dots, N \quad (2.8)$$

and are the distribution of the first s particles (or of any group of s tagged particles).

In order to describe the system in terms of a single kinetic equation, we expect that chaos propagates. Actually since W^N is initially factorizing, although the dynamics creates correlations, we hope that, due to the weakness of the interaction, factorization still holds approximately also at any positive time t , namely

$$f_s^N \approx f_1^{\otimes s}.$$

In this case the law of large numbers does hold, that is

$$\frac{1}{N} \sum_j \delta(z - z_j) \approx f_1^N(z, t)$$

for W^N -almost all $Z_N = \{z_1 \cdots z_N\}$. Then

$$\begin{aligned} \pi_{i,j}^N &\approx \frac{1}{N-1} K \left(\frac{1}{N-1} \sum_k \mathcal{X}_{B_{|x_i - x_j|}(x_i)}(x_k) \right) \\ &\approx \frac{1}{N-1} K \left(M_\rho(B_{|x_1 - x_2|}(x_1)) \right) \end{aligned}$$

where

$$M_\rho(B_R(x)) = \int_{B_R(x)} \rho(y) dy, \quad (2.9)$$

and $\rho(x) = \int dv f_1^N(x, v)$ is the spatial density. Motivated by this remark, from now on we use the following notation

$$M_{X_N}(B_{|x_i - x_j|}(x_i)) = r(i, j) = \frac{1}{N-1} \sum_k \mathcal{X}_{B_{|x_i - x_j|}(x_i)}(x_k).$$

Here M stands for ‘mass’ and the notation introduced is justified by the law of large numbers.

In conclusion we expect that, by (2.7), in the limit $N \rightarrow \infty$, $f_1^N \rightarrow f$ and $f_2^N \rightarrow f^{\otimes 2}$, where f solves

$$\partial_t \int f \varphi = \int f v \cdot \nabla_x \varphi - \int f \varphi + \int f(z_1) f(z_2) \varphi(x_1, v_2) K\left(M_\rho(B_{|x_1-x_2|}(x_1))\right)$$

which is the weak form of the equation

$$\left(\partial_t + v \cdot \nabla_x\right) f(x, v, t) = -f(x, v, t) + \rho(x, t) \int K\left(M_\rho(B_{|x-y|}(x))\right) f(y, v, t) dy. \quad (2.10)$$

We remark that existence and uniqueness of global solutions in $L^1(\mathbb{R}^{2d})$ for the kinetic equation (2.10) can be proved by using a standard Banach fixed-point argument.

Once known f , we can construct the one-particle nonlinear process given by the generator

$$L_1^{(1)} \phi(x, v) = (v \cdot \nabla_x - 1) \phi(x, v) + \int f(y, w) \phi(x, w) K\left(M_\rho(B_{|x-y|}(x))\right) dy dw.$$

We also introduce the N -particle process given by N independent copies of the above process. Its generator is

$$\begin{aligned} L_N^{(1)} \Phi(Z_N) &= V_N \cdot \nabla_{X_N} \Phi(Z_N) \\ &+ \sum_i \left[\int \Phi(X_N, V_N^i(w_i)) K\left(M_\rho(B_{|x_i-y_i|}(x_i))\right) f(y_i, w_i) dy_i dw_i - \Phi(X_N, V_N) \right]. \end{aligned} \quad (2.11)$$

2.2. Motivations and main result. This work aims to prove propagation of chaos for the N -particle process described by (2.5). Propagation of chaos consists in preparing a system of N particles with initial configurations i.i.d with a given law f_0 and show that, considering any group of fixed s particles between the N ones, this independence (chaos) is also recovered for future times for the fixed s -group when $N \rightarrow \infty$. This is expressed mathematically by saying that the s -particle marginal $f_s^N(t)$ introduced in (2.8) approximates $f^{\otimes s}(t)$ for positive times, where $f(t)$ is the solution with initial datum f_0 of the limit equation (2.10).

As mentioned in the introduction, the propagation of chaos result for (2.5) was already obtained in [13] using hierarchical techniques. Indeed, the BBGKY hierarchies are a powerful approach but their structure is such that the equation for the s -marginal depends only on the $(s+1)$ -marginal. In this case the non-binary nature of the topological interaction does not allow to derive this hierarchical structure, unless the interaction function K is real analytic and therefore expandable in series, which is exactly the assumption made in [13].

The reason for this work is to provide a different derivation of the limit kinetic equation, using the classic probabilistic coupling technique. In general, given two stochastic processes X and Y , a coupling is a realization of a new process on a product probability space that has as marginal distributions those of X and Y . This approach brings a more natural and general proof, avoiding the analyticity assumption on K .

Theorem 1. *Let $f \in C([0, T]; L^1(\mathbb{R}^{2d}))$ solution of the limit equation (2.10) with initial datum $f_0 \in L^1(\mathbb{R}^{2d})$. Assume that the interaction function K is Lipschitz-continuous and consider the N -particle dynamics such that $W_N(0) = f_0^{\otimes N}$.*

If f_s^N denotes the s -marginal as defined in (2.8), for $t \in [0, T]$ and $s \in \{1, \dots, N\}$, it holds that

$$\|f_s^N(t) - f^{\otimes s}(t)\|_{L^1(\mathbb{R}^{2ds})} \leq s \frac{e^{C_K T}}{\sqrt{N-1}}, \quad (2.12)$$

where C_K is a constant depending only on the Lipschitz constant of K .

The topological character of the interaction bring us naturally to work with norms of strong type and in particular with the L_1 /Total variation distance (see also [3] where a distance similar to the Total Variation has been used to prove the validity of the mean-field limit for a deterministic Cucker-Smale model with topological interactions introduced in [17]).

Indeed, given two measures ρ_1 and ρ_2 , from (2.9) we have

$$|M_{\rho_1}(B_r(x)) - M_{\rho_2}(B_r(x))| \leq \|\rho_1 - \rho_2\|_{TV}$$

where, given (X, \mathcal{A}) a measurable space and two measures μ and ν over X , the total variation distance is defined as

$$\|\mu - \nu\|_{TV} = \sup_{A \in \mathcal{A}} |\mu(A) - \nu(A)|.$$

In the present work, we use the equivalence between the L^1 distance and the Total variation for regular measures and the characterization of the TV distance given by the Wasserstein distance

$$\|\mu - \nu\|_{TV} = \inf_{\pi \in \mathcal{C}(\mu, \nu)} \int_{X \times X} d(x, y) \, d\pi(x, y),$$

where $\mathcal{C}(\mu, \nu)$ is the set of all couplings, *i.e.* measures on the product space with marginals respectively μ and ν in the first and second variables, and $d(a, b) = 1 - \delta_{a,b}$ is the discrete distance (see [25]).

3. PROOF OF THE RESULT

3.1. Coupling and strategy of the proof. We introduce, as a coupling between (2.5) and (2.11), the process $t \rightarrow (Z_N(t); \Sigma_N(t))$ on the product space $\mathbb{R}^{2dN} \times \mathbb{R}^{2dN}$, where $\Sigma_N(t) = (Y_N(t), W_N(t))$. The generator of the new process is

$$Q_N = Q_0 + \tilde{Q}_N,$$

where

$$Q_0 \Phi(Z_N; \Sigma_N) = (V_N \cdot \nabla_{X_N} + W_N \cdot \nabla_{Y_N}) \Phi(Z_N; \Sigma_N) \quad (3.1)$$

is the free-stream operator, while

$$\tilde{Q}_N \Phi(Z_N; \Sigma_N) = \sum_{i=1}^N \sum_{j \neq i} \lambda_{i,j} [\Phi(X_N, V_N^i(v_j); Y_N, W_N^{(i)}(w_j)) - \Phi(Z_N; \Sigma_N)] \quad (3.2a)$$

$$+ \sum_{i=1}^N \sum_{j \neq i} [\pi_{i,j}^N(X_N) - \lambda_{i,j}] [\Phi(X_N, V_N^i(v_j); \Sigma_N) - \Phi(Z_N; \Sigma_N)] \quad (3.2b)$$

$$+ \sum_{i=1}^N \sum_{j \neq i} [\pi^\rho(y_i, y_j) - \lambda_{i,j}] [\Phi(Z_N; Y_N, W_N^{(i)}(w_j)) - \Phi(Z_N; \Sigma_N)] \quad (3.2c)$$

$$+ \sum_{i=1}^N \int du \mathcal{E}_i^N(u) [\Phi(Z_N; Y_N, W_N^{(i)}(u)) - \Phi(Z_N; \Sigma_N)] \quad (3.2d)$$

tends to penalize the discrepancies that can occur over time between Z_N and Σ_N .

Indeed, in (3.2a) the process jumps jointly on both variables with a rate given by

$$\lambda_{i,j}(X_N; y_i, y_j) := \min\{\pi_{i,j}^N(X_N), \pi^\rho(y_i, y_j)\}, \quad (3.3)$$

where

$$\pi^\rho(y_i, y_j) := \alpha_N K\left(M_\rho(B_{|y_i - y_j|}(y_i))\right). \quad (3.4)$$

In (3.2b) and (3.2c) the jumps occur only for one of the pair, with a transition probability given by the error between $\lambda_{i,j}$ and $\pi_{i,j}^N$ or π^ρ . Finally, in (3.2d),

$$\mathcal{E}_i^N(u) = \int K\left(M_\rho(B_{|y_i - y|}(y_i))\right) f(y, u) dy - \sum_{j \neq i} \pi^\rho(y_i, y_j) \delta(u - w_j)$$

is the last error due to the approximation of the limit kinetic equation by the N -particle dynamics with transition probabilities given by π^ρ and will be treated using the law of large numbers.

We remark that, since $\int K(x) dx = 1$, formally we have¹,

$$\begin{aligned} \int K\left(M_\rho(B_{|x-y|}(x))\right) \rho(y) dy &= \int_0^{+\infty} dr K(M_\rho(B_r(x))) \int_{|x-y|=r} \rho(y) d\mathcal{H}^{n-1}(dy) \\ &= \int_0^{+\infty} dr K(M_\rho(B_r(x))) \frac{d}{dr} [M_\rho(B_r(x))] = \int K(x) dx = 1. \end{aligned}$$

From this fact, it follows that Q_N is a coupling of the two previously described processes, i.e. we recover, considering test functions depending only Z_N and Σ_N respectively, the two processes as the two marginals.

We want to prove that f and f_1^N (defined as in (2.8)) agree asymptotically in the limit $N \rightarrow +\infty$. To do this we consider $R^N(t) = R^N(Z_N, \Sigma_N; t)$ the law at time t for the coupled process. As initial distribution at time 0 we assume

$$R^N(0) = f_0^{\otimes N}(Z_N) \delta(Z_N - \Sigma_N). \quad (3.5)$$

¹In general, the formula is true for $\rho \in L^1(\mathbb{R}^d)$ and it is a consequence of the coarea formula (see [14, Thm 3.12, p. 140]).

Let $D_N(t)$ be the average fraction of particles having different positions or velocities, *i.e.* using the symmetry of the law,

$$D_N(t) = \int dR^N(t) \frac{1}{N} \sum_{i=1}^N d(z_i, \sigma_i) = \int dR^N(t) d(z_1, \sigma_1), \quad (3.6)$$

where $z_i = (x_i, v_i)$, $\sigma_i = (y_i, w_i)$ and $d(a, b) = 1 - \delta_{a,b}$ is the discrete distance.

The aim is to show that $D_N(t) \rightarrow 0$. This means the following: initially the coupled system has all the pairs of particles overlapping. The dynamics creates discrepancies and the average number of separated pairs is exactly D_N which is also the Total Variation distance ($L^1(x, v)$ in our case) between f_1^N and f .

Notice that the convergence of the s -marginals f_s^N towards $f^{\otimes s}$ claimed in (2.12) is easily recovered by the fact that

$$\begin{aligned} \|f_s^N(t) - f^{\otimes s}(t)\|_{TV} &\leq \int \delta(Z_s, \Sigma_s) dR^N(Z_N, \Sigma_N; t) \\ &\leq \sum_{i=1}^s \int d(z_i, \sigma_i) dR^N(Z_N, \Sigma_N; t) = sD_N(t) \end{aligned}$$

where $\delta(a, b)$ denotes the discrete distance on the space $\mathbb{R}^{2ds} \times \mathbb{R}^{2ds}$.

3.2. Convergence estimates. Let S_t^N be the semigroup defined by the free-stream generator Q_0 in (3.1). To estimate $D_N(t)$ we apply the Duhamel formula in (3.6) and we get

$$\begin{aligned} \int dR^N(t) d(z_1, \sigma_1) &= \int dR^N(0) d(S_t^N(z_1, \sigma_1)) \\ &\quad + \int_0^t d\tau \int dR^N(\tau) \tilde{Q}_N d(S_{t-\tau}^N(z_1, \sigma_1)), \quad (3.7) \end{aligned}$$

where \tilde{Q}_N is defined in (3.2).

The first term in (3.7) is negligible: indeed, from (3.5), we have

$$\int dR^N(0) d(S_t^N(z_1, \sigma_1)) = \int df_0^{\otimes N}(Z_N) d(S_t^N(z_1, z_1)) \equiv 0.$$

Concerning the second term in (3.7), we define

$$\bar{z}_1 = (x_1 + v_1(t - \tau), v_1), \quad \bar{z}_1^{(j)} = (x_1 + v_1(t - \tau), v_j)$$

and $\bar{X}_N = (x_1 + v_1(t - \tau), \dots, x_N + v_N(t - \tau))$; similarly for $\bar{\sigma}$, $\bar{\sigma}^{(j)}$ and \bar{Y}_N .

By (3.2) we get

$$\int dR^N(\tau) \tilde{Q}_N d(S_{t-\tau}^N(z_1, \sigma_1)) = A_1(\tau) + A_2(\tau) + A_3(\tau),$$

where

$$A_1(\tau) = \sum_{j \neq 1} \int dR^N(\tau) \lambda_{1,j}(\bar{X}_N; \bar{y}_1, \bar{y}_j) [d(\bar{z}_1^{(j)}; \bar{\sigma}_1^{(j)}) - d(\bar{z}_1; \bar{\sigma}_1)]$$

is due to the term of the generator \tilde{Q}_N where the velocities of the particles jump simultaneously;

$$\begin{aligned} A_2(\tau) &= \sum_{j \neq 1} \int dR^N(\tau) (\pi_{1,j}^N(\bar{X}_N) - \lambda_{1,j}) [d(\bar{z}_1^{(j)}; \bar{\sigma}_1) - d(\bar{z}_1; \bar{\sigma}_1)] \\ &\quad + \sum_{j \neq 1} \int dR^N(\tau) (\pi_{1,j}^\rho(\bar{y}_1, \bar{y}_j) - \lambda_{1,j}) [d(\bar{z}_1; \bar{\sigma}_1^{(j)}) - d(\bar{z}_1; \bar{\sigma}_1)] \end{aligned}$$

is due to the terms of the generator where only one of the two coupled processes jump and

$$A_3(\tau) = \int dR^N(\tau) \int du \bar{\mathcal{E}}_1^N(u) [d(\bar{z}_1; \bar{\sigma}_1^{(u)}) - d(\bar{z}_1; \bar{\sigma}_1)]$$

is due to the remainder term. Here $\bar{\mathcal{E}}_1^N(u)$ is $\mathcal{E}_1^N(u)$ evaluated along the moving frame of the free transport.

Here, we have used that $d(z_1, \sigma_1)$ depends only on the configurations of the first particle; hence, the only non-zero contribution in the sum over i is given for $i = 1$.

Concerning $A_1(\tau)$, it follows from (2.3) and (2.4) that

$$|e_K(N)| \leq \frac{\text{Lip}(K)}{N-1}$$

and that, for $N > 2\text{Lip}(K) + 1$,

$$\alpha_N \leq \frac{4e^{\frac{\text{Lip}(K)}{N-1}}}{N-1},$$

using the inequality $1/(1-x) \leq 4e^x$ for $x \in (0, 1/2)$. Therefore, from (3.3) we get

$$\lambda_{1,j} \leq \alpha_N \|K\|_\infty \leq \frac{4\sqrt{e} \text{Lip}(K)}{N-1}.$$

By the symmetry of R^N and denoting $C_K := 8\sqrt{e} \text{Lip}(K)$,

$$A_1(\tau) \leq \frac{C_K}{2(N-1)} \sum_{j \neq 1} \int dR^N(\tau) [d(z_j, \sigma_j) + d(z_1, \sigma_1)] \leq C_K D_N(\tau), \quad (3.8)$$

since $d(\bar{z}_1^{(j)}; \bar{\sigma}_1^{(j)}) \leq d(z_j, \sigma_j) + d(z_1, \sigma_1)$. Indeed the right-hand side is vanishing iff $z_1 = \sigma_1$ and $z_j = \sigma_j$ and, in this case, also the left-hand side is clearly vanishing.

We now give a bound on $A_2(\tau)$. Since $\lambda_{1,j}$ is the minimum between $\pi_{1,j}^N$ and $\pi_{i,j}^\rho$, we have

$$|A_2(\tau)| \leq \sum_{j \neq 1} \int dR^N(\tau) |\pi_{1,j}^N(\bar{X}_N) - \pi_{1,j}^\rho(\bar{y}_1, \bar{y}_j)|. \quad (3.9)$$

From (2.2) and (3.4),

$$|\pi_{1,j}^N(\bar{X}_N) - \pi_{1,j}^\rho(\bar{y}_1, \bar{y}_j)| \leq \alpha_N \text{Lip}(K) |M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_\rho(\bar{B}_{1,j}^y)|,$$

where we are using the shorthand notation

$$\bar{B}_{1,j}^x = B_{|\bar{x}_1 - \bar{x}_j|}(\bar{x}_1) \quad \text{and} \quad \bar{B}_{1,j}^y = B_{|\bar{y}_1 - \bar{y}_j|}(\bar{y}_1).$$

By the triangular inequality

$$\begin{aligned} |M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_\rho(\bar{B}_{1,j}^y)| &\leq |M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_{\bar{X}_N}(\bar{B}_{1,j}^y)| \\ &\quad + |M_{\bar{X}_N}(\bar{B}_{1,j}^y) - M_{\bar{Y}_N}(\bar{B}_{1,j}^y)| + |M_{\bar{Y}_N}(\bar{B}_{1,j}^y) - M_\rho(\bar{B}_{1,j}^y)|. \end{aligned}$$

Hence we divide the estimate (3.9) respectively in three terms:

$$|A_2(\tau)| \leq T_1(\tau) + T_2(\tau) + T_3(\tau).$$

In $T_1(\tau)$ we are considering particles with spatial configuration given by X_N and we want to estimate the discrepancy of the configuration over two different balls $\bar{B}_{1,j}^x$ and $\bar{B}_{1,j}^y$. Since $\bar{B}_{1,j}^x = \bar{B}_{1,j}^y$ iff $z_1 = \sigma_1$ and $z_j = \sigma_j$, using that $M_{\bar{X}_N} \in [0, 1]$, we have

$$|M_{\bar{X}_N}(\bar{B}_{1,j}^x) - M_{\bar{X}_N}(\bar{B}_{1,j}^y)| \leq d(z_1, \sigma_1) + d(z_j, \sigma_j).$$

Therefore, by the symmetry of R^N ,

$$\begin{aligned} T_1(\tau) &\leq \alpha_N \text{Lip}(K) \sum_{j \neq 1} \int dR^N(\tau) [d(z_1, \sigma_1) + d(z_j, \sigma_j)] \\ &\leq C_K D_N(\tau). \end{aligned}$$

Regarding $T_2(\tau)$, we are considering the discrepancy of two different configurations over the same ball $\bar{B}_{1,j}^y$. Since

$$|M_{\bar{X}_N}(\bar{B}_{1,j}^y) - M_{\bar{Y}_N}(\bar{B}_{1,j}^y)| \leq \frac{1}{N} \sum_{i=1}^N d(z_i, \sigma_i),$$

using again the symmetry of the law, we get

$$T_2(\tau) \leq \alpha_N \text{Lip}(K) \sum_{j \neq 1} \int dR^N(\tau) d(z_1, \sigma_1) \leq C_K D_N(\tau).$$

The last estimate on $T_3(\tau)$ is a consequence of the law of large numbers. After a change of variable, using the symmetry of the law R^N and the fact that this last term depends only on the Y_N configuration, we have that

$$T_3(\tau) = \alpha_N \text{Lip}(K) \sum_{j \neq 1} \int d\rho^{\otimes N}(\tau) |M_{Y_N}(B_{1,j}^y) - M_\rho(B_{1,j}^y)|,$$

where $B_{1,j}^y = B_{|y_1 - y_j|}(y_1)$. By Cauchy-Schwartz,

$$\begin{aligned} &\left| \int d\rho^{\otimes N}(\tau) |M_{Y_N}(B_{1,j}^y) - M_\rho(B_{1,j}^y)| \right|^2 \\ &\leq \int d\rho^{\otimes N}(\tau) \left| \frac{1}{N-1} \sum_{h \neq 1} [\mathcal{X}_{B_{1,j}^y}(y_h) - M_\rho(B_{1,j}^y)] \right|^2 \\ &\leq \sum_{h_1, h_2 \neq 1} \int \frac{d\rho^{\otimes N}(\tau)}{(N-1)^2} [\mathcal{X}_{B_{1,j}^y}(y_{h_1}) - M_\rho(B_{1,j}^y)] [\mathcal{X}_{B_{1,j}^y}(y_{h_2}) - M_\rho(B_{1,j}^y)]. \end{aligned}$$

Thanks to the independence of the limit process, we get that the only non-zero contributions are given when $h_1 = h_2$ and this happens only for $N-1$ terms. Hence

$$T_3(\tau) \leq \frac{C_K}{\sqrt{N-1}}.$$

Collecting the estimates on T_1, T_2 and T_3 , we obtain that

$$A_2(\tau) \leq C_K \left(D_N(\tau) + \frac{1}{\sqrt{N-1}} \right). \quad (3.10)$$

We conclude the proof estimating $A_3(\tau)$. Since this term depends only on the independent Y_N configuration

$$\begin{aligned} |A_3(\tau)| &\leq \int \frac{df^{\otimes N}(\tau)}{N-1} \sum_{j \neq 1} \left| \int K \left(M_\rho(B_{|\bar{y}_1-y|}(\bar{y}_1)) \right) d\rho(y) - K(M_\rho(\bar{B}_{1,j}^y)) \right| \\ &\quad + \frac{1}{N-1} \int df^{\otimes N}(\tau) \frac{e_K(N)}{1-e_K(N)} \sum_{j \neq 1} K(M_\rho(\bar{B}_{1,j}^y)), \end{aligned}$$

where we added and subtracted the term $\sum_j K(M_\rho(\bar{B}_{1,j}^y))/(N-1)$.

Applying again the law of large numbers on the first term and estimating the second term thanks to

$$\frac{e_K(N)}{1-e_K(N)} \leq \frac{C_K}{N-1},$$

we arrive at

$$|A_3(\tau)| \leq \frac{C_K}{\sqrt{N-1}}. \quad (3.11)$$

Collecting the estimates in (3.8), (3.10) and (3.11) and using Gronwall's lemma, we conclude the proof of the theorem.

ACKNOWLEDGEMENTS

PD holds a visiting professor association with the Department of Mathematics, Imperial College London, UK.

REFERENCES

- [1] M. Ballerini, N. Cabibbo, R. Candelier, A. Cavagna, E. Cisbani, I. Giardina, V. Lecomte, A. Orlandi, G. Parisi, A. Procaccini, M. Viale, V. Zdravkovic *Interaction ruling animal collective behavior depends on topological rather than metric distance: evidence from a field study*. Proc. Natl. Acad. Sci. USA **105** (2008) 1232–1237.
- [2] S. Bazazi, J. Buhl, J. J. Hale, M. L. Anstey, G. A. Sword, S. J. Simpson, I. D. Couzin *Collective motion and cannibalism in locust migratory bands* Curr. Biol. **18** (2008) 735–739.
- [3] D. Benedetto, E. Caglioti, S. Rossi *Mean-field limit for particle systems with topological interactions* Math. Mech. Complex Syst. **9** (2021) 423–440.
- [4] E. Bertin, M. Droz, G. Grégoire *Boltzmann and hydrodynamic description for self-propelled particles* Phys. Rev. E **74** (2006) 022101.
- [5] A. Blanchet, P. Degond *Topological interactions in a Boltzmann-type framework* J. Stat. Phys. **163** (2016) 41–60.
- [6] A. Blanchet, P. Degond *Kinetic models for topological nearest-neighbor interactions* J. Stat. Phys. **169** (2017) 929–950.
- [7] N. W. Bode, D. W. Franks, A. J. Wood *Limited interactions in flocks: relating model simulations to empirical data* J. R. Soc. Interface. **8** (2011) 301–304.
- [8] M. Camperi, A. Cavagna, I. Giardina, G. Parisi, E. Silvestri *Spatially balanced topological interaction grants optimal cohesion in flocking models* Interface focus **2** (2012) 715–725.

- [9] A. Cavagna, A. Cimarelli, I. Giardina, G. Parisi, R. Santagati, F. Stefanini, R. Tavarone *From empirical data to inter-individual interactions: unveiling the rules of collective animal behavior* Math. Models Methods Appl. Sci. **20** Suppl. 01 (2010) 1491–1510.
- [10] L.P. Chaintron, A. Diez *Propagation of chaos: A review of models, methods and applications. I. Models and methods* Kinet. Relat. Models **15** (2022) 895–1015.
- [11] L.P. Chaintron, A. Diez *Propagation of chaos: A review of models, methods and applications. II. Applications* Kinet. Relat. Models **15** (2022) 1017–1173.
- [12] F. Cucker, S. Smale *Emergent behavior in flocks* IEEE Trans. Automat. Control **52** (2007) 852–862.
- [13] P. Degond, M. Pulvirenti *Propagation of chaos for topological interactions* Ann. Appl. Prob. **29** (2019) 2594–2612.
- [14] L.C. Evans, R.F. Gariepy *Measure theory and fine properties of functions* Chapman and Hall/CRC. (2015).
- [15] F. Ginelli, H. Chaté *Relevance of metric-free interactions in flocking phenomena* Phys. Rev. Lett. **105** (2010) 168103.
- [16] R. Giniūnaitė, R. E. Baker, P. M. Kulesa, P. K. Maini *Modelling collective cell migration: neural crest as a model paradigm* J. Math. Biol. **80** (2020) 481–504.
- [17] J. Haskovec *Flocking dynamics and mean-field limit in the Cucker-Smale-type model with topological interactions* Phys. D **261** (2013) 42–51.
- [18] U. Lopez, J. Gautrais, I. D. Couzin, G. Theraulaz *From behavioural analyses to models of collective motion in fish schools* Interface focus **2** (2012) 693–707.
- [19] S. Martin *Multi-agent flocking under topological interactions* Syst. Control. Lett. **69** (2014) 53–61.
- [20] T. Niizato, H. Murakami, Y. P. Gunji *Emergence of the scale-invariant proportion in a flock from the metric-topological interaction* BioSystems **119** (2014) 62–68.
- [21] Y. Shang, R. Bouffanais *Consensus reaching in swarms ruled by a hybrid metric-topological distance* Eur. Phys. J. B **87** (2014) 1–7.
- [22] Y. Shang, R. Bouffanais *Influence of the number of topologically interacting neighbors on swarm dynamics* Sci. Rep. **4** (2014) 4184.
- [23] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, O. Shochet *Novel type of phase transition in a system of self-driven particles* Phys. Rev. Lett. **75** (1995) 1226.
- [24] T. Vicsek, A. Zafeiris *Collective motion* Phys. Rep. **517** (2012) 71–140.
- [25] C. Villani *Topics in optimal transportation* Am. Math. Soc. (2016).
- [26] L. Wang, G. Chen *Synchronization of multi-agent systems with metric-topological interactions* Chaos **26** (2016) 094809.

PIERRE DEGOND
 INSTITUT DE MATHÉMATIQUES DE TOULOUSE ; UMR5219
 UNIVERSITÉ DE TOULOUSE ; CNRS
 UPS, F-31062 TOULOUSE CEDEX 9, FRANCE
 Email address: pierre.degond@math.univ-toulouse.fr

MARIO PULVIRENTI
 DIPARTIMENTO DI MATEMATICA,
 UNIVERSITÀ DI ROMA ‘LA SAPIENZA’
 PIAZZALE ALDO MORO 2, 00185 ROMA, ITALY

INTERNATIONAL RESEARCH CENTER M&MOCS,
 UNIVERSITÀ DEGLI STUDI DELL’AQUILA
 VIA GIOVANNI GRONCHI 18, 67100 L’AQUILA, ITALY

ACCADEMIA NAZIONALE DEI LINCEI
VIA DELLA LUNGARA 10, 00165 ROMA, ITALY

Email address: `pulviren@mat.uniroma1.it`

STEFANO ROSSI
DIPARTIMENTO DI MATEMATICA,
UNIVERSITÀ DI ROMA 'LA SAPIENZA'
PIAZZALE ALDO MORO 2, 00185 ROMA, ITALY

Email address: `stef.rossi@uniroma1.it`