



HAL
open science

Normal forms of z -graded q -manifolds

Alexei Kotov, Camille Laurent-Gengoux, Vladimir Salnikov

► **To cite this version:**

Alexei Kotov, Camille Laurent-Gengoux, Vladimir Salnikov. Normal forms of z -graded q -manifolds. 2022. hal-03906781

HAL Id: hal-03906781

<https://hal.science/hal-03906781>

Preprint submitted on 19 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NORMAL FORMS OF \mathbb{Z} -GRADED Q -MANIFOLDS

ALEXEI KOTOV, CAMILLE LAURENT-GENGOUX, AND VLADIMIR SALNIKOV

ABSTRACT. Following recent results of A.K. and V.S. on \mathbb{Z} -graded manifolds, we give several local and global normal forms results for Q -structures on those, i.e. for differential graded manifolds. In particular, we explain in which sense their relevant structures are concentrated along the zero-locus of their curvatures, especially when the negative-part is of Koszul-Tate type. We also give a local splitting theorem.

INTRODUCTION

This article is the sequel of [9], that studied normal forms of \mathbb{Z} -graded manifolds and where the analogue of the Batchelor’s theorem has been proven. We now equip a \mathbb{Z} -graded manifold with a degree +1 self-commuting vector field Q , thus making it a differential graded (DG) manifold, also called Q -manifold. The purpose of this paper is to provide several normal form type results in this setting.

The paper is organized as follows. In Section 1, we give some precise definitions and fix some usual notations related to graded manifolds. Then we proceed with the description of projective systems of algebras (recapitulated in Appendix A), which we specialize to the \mathbb{Z} -graded structure sheaves. Section 2 is devoted to the idea that “outside the zero locus of their curvatures, (\mathbb{Z}^* -graded) Q -manifolds can be made trivial”. A more precise statement is that on any open subset where the curvature is different from zero at all points, the dual \mathbb{Z}^* -graded Lie ∞ -algebroid can be chosen to have all k -ary bracket equal to zero, except for the 0-ary bracket, given by the nowhere vanishing curvature.

In section 3 we first recall the standard notion of Koszul–Tate resolution, which are examples of negatively graded Q -manifolds. Then we construct two structures on the zero locus $\{\kappa = 0\}$ of a Q -manifold that are independent of a choice of a splitting: a positively graded Q -structure on the zero locus $\{\kappa = 0\}$ and a negatively graded Q -manifold. We eventually show that Q -manifolds whose negative part is of Koszul-Tate type are entirely encoded by this positively graded Q -structure on the zero locus.

Last, in section 4 we choose a point in the zero locus, (on which leaves of the anchor map are well defined) and give a splitting theorem: near a leaf L in the zero locus, a Q -manifold is the direct product of the standard $T[1]L$ and a transverse Q -manifold. In the process, we also give some counter-examples to “naive beliefs” about the anchor maps of a Q -manifold. We conclude by mentioning some perspectives and potential applications.

Key words and phrases. \mathbb{Z} -graded manifolds, dg-manifolds, Q -structures, Lie ∞ -algebroids, normal forms, splitting theorems.

1. NOTATIONS AND PRELIMINARIES

1.1. \mathbb{Z}^* -graded manifolds. Let us give (recall) the definition of \mathbb{Z}^* -graded manifolds. We start with an important definition: a filtration which is used throughout the paper.

Definition 1.1. Let $\mathcal{O} = \bigoplus_{j \in \mathbb{Z}} \mathcal{O}_j$ be a \mathbb{Z} -graded commutative algebra. We call negative filtration the filtration

$$\mathcal{O} = F^0 \mathcal{O} \supset \mathcal{I}_- = F^1 \mathcal{O} \supset \dots \supset F^i \mathcal{O} \supset \dots$$

defined by $F^i \mathcal{O} = \mathcal{I}_-^i$ for all $i \in \mathbb{Z}_{\geq 0}$, where \mathcal{I}_-^i is generated by $\bigoplus_{j \leq -i} \mathcal{O}_j$.

This filtration allows to define the graded manifolds we are interested in.

Definition 1.2 ([9]). A \mathbb{Z}^* -graded manifold is a pair $M = (M_0, \mathcal{O})$, where M_0 is a smooth manifold (referred to as base manifold) and $\mathcal{O} = \bigoplus_{i \in \mathbb{Z}} \mathcal{O}_i$ is a sheaf of \mathbb{Z} -graded commutative algebras (whose sections are referred to as functions), such that each point of M_0 has a neighborhood $U \subset M_0$ over which $\mathcal{O}(U)$ is isomorphic to $\Gamma(\tilde{S}(\bigoplus_{i \in \mathbb{Z}^*} V_i))$, where each V_i is a graded vector bundle of degree i , and “ \tilde{S} ” stands for the completion of $\Gamma(S(\bigoplus_{i \in \mathbb{Z}^*} V_i))$ with respect to the negative filtration.

Remark 1.3. This definition is explained in details in [9]: for this paper to be self-consistent and for further use, we recollect some necessary facts about filtrations and their completions in what follows and in Appendix A.

Note that Definition 1.2 potentially allows a function to be a sum of infinitely many terms, which is also explained in [9]. \square

Remark 1.4. We write “ \mathbb{Z}^* -graded” instead of “ \mathbb{Z} -graded” to insist on the natural assumptions that there are no generators of \mathcal{O} of degree 0 which is not a coordinate function on M_0 . For instance, Kapranov dg-manifolds [13, 14] are not \mathbb{Z}^* -graded manifolds, the difference is in conventions though.

For \mathbb{Z}^* -graded manifolds, in contrast to the $\mathbb{Z}_{\geq 1}$ -graded or $(-\mathbb{Z}_{\geq 1})$ -graded cases, we do not have an isomorphism $C^\infty(M_0) \simeq \mathcal{O}_0$. For instance, the product of a function in \mathcal{O}_p with a function in \mathcal{O}_{-p} may very well produce a non-zero function: it then belongs to \mathcal{O}_0 but can not be considered as an element in $C^\infty(M_0)$. There is even no canonical inclusion $C^\infty(M_0) \hookrightarrow \mathcal{O}_0$, but there is a natural projection $\mathcal{O}_0 \rightarrow C^\infty(M_0)$, which corresponds to the inclusion $M_0 \hookrightarrow M$. \square

Remark 1.5. A \mathbb{Z}^* -graded manifold is complete with respect to the topology on \mathcal{O} given by the negative filtration, see [9]. \square

Remark 1.6. The negative filtration of \mathcal{O} is compatible with the negative filtration of the symmetric algebras that appear in Definition 1.2. Notice that elements in $F^i \mathcal{O}$ may be of any degree, although its generators have degree less or equal to $-i$. Also, notice that $\bigcap_{i \geq 0} F^i \mathcal{O} = \{0\}$. \square

According to [9], there are natural sheaves of graded ideals in \mathcal{O} :

- (1) the ideal \mathcal{I}_+ generated by $\bigoplus_{i \geq 1} \mathcal{O}_i$.
- (2) the ideal $\mathcal{I}_- = F^1 \mathcal{O}$ generated by $\bigoplus_{i \leq -1} \mathcal{O}_i$.
- (3) The ideal $\mathcal{I} = \mathcal{I}_+ + \mathcal{I}_-$.

Let us consider the quotient of \mathcal{O} by these three ideals:

- (1) The quotient $(M_0, \mathcal{O}/\mathcal{I}_+)$ is a graded manifold with grading now ranging from 0 to $-\infty$ that we call the *negative part of* (M_0, \mathcal{O}) .
- (2) The quotient $(M_0, \mathcal{O}/\mathcal{I}_-)$ is a graded manifold with grading now ranging from 0 to $+\infty$ that we call the *positive part of* (M_0, \mathcal{O}) .
- (3) The quotient $(M_0, \mathcal{O}/\mathcal{I})$ is simply the smooth manifold M_0 with its sheaf $C^\infty(M_0)$ of smooth functions (and, in particular, is concentrated in degree 0).

For $i \geq 0$ and $j \in \mathbb{Z}$, we denote by $(F^i \mathcal{O})_j$ elements of $F^i \mathcal{O}$ of degree j . Then, to a graded manifold $M = (M_0, \mathcal{O})$, one can associate (canonically) a family $(E_i)_{i \in \mathbb{Z}^*}$ of vector bundles over M_0 , as follows. The quotient space

$$(1.1) \quad \frac{\mathcal{I}}{\mathcal{I}^2} = \bigoplus_{i \in \mathbb{Z}^*} \left(\frac{\mathcal{I}}{\mathcal{I}^2} \right)_i$$

is a direct sum of projective $C^\infty(M_0)$ -modules, hence by Serre–Swan theorem, there exists for all $i \in \mathbb{Z}^*$ a vector bundle E_i such that $\Gamma(E^*)_i \simeq (\mathcal{I})_i / (\mathcal{I}^2)_i$. We call $E_\bullet := \bigoplus_{i \in \mathbb{Z}^*} E_i$ the *canonical graded vector bundle of* (M_0, \mathcal{O}) .

Theorem 1.7 (*Batchelor’s theorem*, [9] – Sections 3.3 and 4.2). *Let (M_0, \mathcal{O}) be a \mathbb{Z}^* -graded manifold with canonical \mathbb{Z}^* -graded bundle E_\bullet . There exists an isomorphism of sheaves (called *splitting*):*

$$\mathcal{O} \simeq \Gamma \left(\tilde{S}(\bigoplus_{i \in \mathbb{Z}^*} E_i^*) \right).$$

Here, \tilde{S} again refers to the completion of $\Gamma(S(\bigoplus_{i \in \mathbb{Z}^*} E_i^*))$ with respect to the its negative filtration as in Definition 1.1.

Remark 1.8. Notice that for every splitting, sections of $E_{-i}^* \equiv (E_{-i})^* = (E^*)_i$ become functions of degree $+i$ in \mathcal{O} . \square

Remark 1.9. Although Batchelor’s theorem claims that splitting exists, there is no canonical splitting in general. In contrast, the vector bundles $(E_i)_{i \in \mathbb{Z}^*}$ defined above are canonical. \square

Once a splitting is chosen, many different notions of “degree” can be defined, beside the degree that \mathcal{O} is equipped with by definition. More precisely, for a section $\alpha \in \Gamma(E^*)_i$, let us define three different degrees as follows:

$$\deg(\alpha) = i, \quad \text{pol}(\alpha) = 1, \quad \deg_+(\alpha) = \begin{cases} i & \text{for } i \geq 1 \\ 0 & \text{otherwise,} \end{cases}, \quad \deg_-(\alpha) = \begin{cases} -i & \text{for } i \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Then these degrees extend by multiplicativity to $\Gamma(S(\bigoplus_{i \in \mathbb{Z}^*} E_i^*))$. To avoid confusion, the degree \deg will be called the *total degree*, sometimes referred to as the *ghost degree*. It coincides with the degree that \mathcal{O} is initially equipped with. This degree is responsible¹ for all the commutation relations, i.e. the Koszul sign rule is defined by its reduction modulo 2. The degree \deg_- (resp. \deg_+) is called the *negative degree* (resp. *positive degree*) and plays an important role. Also,

$$\deg = \deg_+ - \deg_-.$$

¹We make this assumption for simplicity of the presentation in this paper, but the constructions work for a more general convention on the relation of the total degree and the (super) parity.

Last, pol is the *polynomial degree* (sometimes referred to as *arity*) that counts the number of sections in a product.

Example 1.10. Concretely, for a section of $E_{-5}^* \odot E_4^* \odot E_7^*$

- the total degree or ghost degree is $5 - 4 - 7 = -6$;
- the negative degree is $4 + 7 = +11$;
- the positive degree is $+5$;
- the polynomial degree is 3 (it is the product of three sections).

Remark 1.11. The negative degree is compatible with the filtration $F^i\mathcal{O}$ introduced above in the sense that $F^i\mathcal{O} = \{F \in \mathcal{O} \mid \deg_-(F) \geq i\}$. \square

1.2. Q -manifolds. Let us now define Q -manifolds, that is equip a \mathbb{Z}^* -graded manifold with a differential structure.

Definition 1.12. A vector field of degree k on a \mathbb{Z}^* -graded manifold (M, \mathcal{O}) is a degree k derivation of \mathcal{O} .

Vector fields of degree k shall be denoted as $\mathfrak{X}_k(\mathcal{O})$. The graded vector space of all vector fields:

$$\mathfrak{X}_\bullet(\mathcal{O}) = \bigoplus_{k \in \mathbb{Z}} \mathfrak{X}_k(\mathcal{O}),$$

form a graded Lie algebra when equipped with the graded commutator $[\cdot, \cdot]$.

Definition 1.13. A \mathbb{Z}^* -graded Q -manifold is a triple (M_0, \mathcal{O}, Q) , with $M = (M_0, \mathcal{O})$ a \mathbb{Z}^* -graded manifold and Q a degree $+1$ vector field which satisfies $[Q, Q] = 0$.

Since the degree of Q is $+1$, we have $Q[\mathcal{I}_+] \subset \mathcal{I}_+$, so that Q induces a degree $+1$ derivation Q^- of the quotient $\mathcal{O}/\mathcal{I}_+$ which is by definition the sheaf of functions of the negative part of (M_0, \mathcal{O}) . This allows the following definition.

Definition 1.14. We call the Q -manifold $(M_0, \mathcal{O}/\mathcal{I}_+, Q^-)$ the negative part of the Q -manifold (M_0, \mathcal{O}, Q) .

Remark 1.15. The vector field Q^- is $C^\infty(M_0)$ -linear, i.e. it is a vertical vector field. \square

1.3. An algebraic generalization: Q -varieties over a commutative algebra. Let \mathcal{A} be a unital commutative algebra (that may be thought as functions over an affine variety X_0 for instance). Definition 1.2 admits a generalization: a differential graded commutative algebra \mathcal{O} such that there exist finitely generated projective \mathcal{A} -modules $(\mathcal{V}_i)_{i \in \mathbb{N}}$ and a graded algebra isomorphism:

$$\mathcal{O} \simeq S_{\mathcal{A}}(\bigoplus_{i \geq 1} \mathcal{V}_i).$$

In particular, the following object will be important.

Definition 1.16. Let $I \subset C^\infty(M_0)$ be an ideal. A positively graded variety (resp. Q -variety) over $C^\infty(M_0)/I$ is a positively graded commutative algebra \mathcal{K}_+ (resp. positively graded commutative differential algebra (\mathcal{K}_+, Q_+)) that admits a splitting, i.e. an isomorphism

$$\mathcal{K}_+ \simeq \Gamma_I(S(\bigoplus_{i \geq 1} E_{-i}^*))$$

for a family of vector bundles $(E_{-i})_{i \geq 1}$ over M_0 . Here for any vector bundle $E \rightarrow M_0$,

$$\Gamma_I(E) := \Gamma(E) \otimes_{C^\infty(M_0)} C^\infty(M_0)/I.$$

Remark 1.17. There is no need to take completions in the definition above since every function of a given degree is necessarily polynomial with respect to non-zero degree variables.

1.4. Duality Q -manifolds \sim Lie ∞ -algebroids. Let (M_0, \mathcal{O}, Q) be a Q -manifold. Once a splitting $\mathcal{O} \simeq \Gamma(\oplus_{i \in \mathbb{Z}} E_i^*)$ is given, Q can be dualized to a Lie ∞ -algebroid, defined as follows.

Definition 1.18. [17, 3] *A \mathbb{Z}^* -graded Lie ∞ -algebroid of a \mathbb{Z}^* -graded vector bundle is the data of:*

- families indexed by $n \geq 1$ of vector bundle morphisms

$$\rho_n: S^n(\oplus_{i \in \mathbb{Z}^*} E_i)_{-1} \longrightarrow TM_0$$

called n -anchor maps,

- families of degree $+1$ maps:

$$\ell_n: S_{\mathbb{R}}^n(\Gamma(\oplus_{i \in \mathbb{Z}^*} E_i))_k \longrightarrow \Gamma(E_{k+1})$$

called n -bracket,

together with a section $\kappa \in \Gamma(E_{+1})$ called curvature that satisfy the higher Jacobi and higher Leibniz identities (see e.g. [16]).

Remark 1.19. It is not easy to attach a single name to the following proposition, based on a observation by Pavol Ševera [18], spelled out in the negative degree case in [2], and which can be proven using Theodore Voronov's derived brackets in [17]. \square

Proposition 1.20. *There is a one-to-one correspondence between \mathbb{Z}^* -graded Lie ∞ -algebroids structures on $\oplus_{i \in \mathbb{Z}} E_i \rightarrow M_0$ and Q -manifolds structures with sheaf of functions $\Gamma(\tilde{S}(\oplus_{i \in \mathbb{Z}^*} E_i^*))$.*

1.5. Projective systems associated to graded manifolds. In this section, we give a precise sense to the notion of the flow of a degree 0 vector field on a graded manifold. For the standard definitions of projective systems the reader is referred to Appendix A, while now we specialize the Proposition A.1 from there to the context we are interested in. Let (M_0, \mathcal{O}) be a \mathbb{Z}^* -graded Q -manifold over M_0 with the sheaf of functions \mathcal{O} . This sheaf of functions comes equipped with the (negative) filtration as in Definition 1.1, so that $A^i := \mathcal{O}/F^i \mathcal{O}$ is a projective system of algebras. Since $\bigcap_{i \in \mathbb{N}} F^i \mathcal{O} = \{0\}$, its projective limit A^∞ is canonically isomorphic to \mathcal{O} .

If a degree 0 vector field \mathfrak{v} such that $\mathfrak{v}[\mathcal{O}] \subset F^n \mathcal{O}$ for some $n \geq 1$ is given, then for every $i \in \mathbb{N}$, the family of endomorphisms

$$\begin{aligned} \mathcal{O}/F^i \mathcal{O} &\rightarrow \mathcal{O}/F^i \mathcal{O} \\ f &\mapsto \sum_{k \geq 0} \frac{t^k}{k!} \mathfrak{v}^k[f] \end{aligned}$$

is well-defined because the sum is finite, it is an algebra endomorphism for all $i \in \mathbb{N}$, and is a morphism of projective systems of algebras. We denote its projective limit by $e^{t\mathfrak{v}}$. By construction, for all $s, t \in \mathbb{R}$ we have $e^{s\mathfrak{v}} e^{t\mathfrak{v}} = e^{(s+t)\mathfrak{v}}$ and $e^{0\mathfrak{v}} = \text{Id}_{\mathcal{O}}$. As a consequence $e^{t\mathfrak{v}}$ is a diffeomorphism of the graded manifold (M_0, \mathcal{O}) .

Proposition 1.21. *Given a family $(v_n)_{n \in \mathbb{N}}$ of degree zero vector fields on a graded manifold (M_0, \mathcal{O}) such that*

$$v_n: \mathcal{O} \rightarrow F^n \mathcal{O}$$

the infinite composition $\bigcirc_{i \uparrow \in \mathbb{N}} e^{v_i}$ is a diffeomorphism of the graded manifold (M_0, \mathcal{O}) , well-defined in the sense of [9]

2. Q -MANIFOLDS WITH CURVATURE

2.1. Normal forms outside of the zero locus of the curvature. For (M_0, \mathcal{O}, Q) a \mathbb{Z}^* -graded Q -manifold², recall (Equation 1.1) that the vector bundle E_{+1} (in fact its dual) is defined by applying the Serre-Swan theorem:

$$\Gamma(E_{+1}^*) = \left(\frac{\mathcal{I}}{\mathcal{I}^2} \right)_{-1} = \frac{\mathcal{I}_{-1}}{\mathcal{I}_{-1}^2} = \frac{F^1 \mathcal{O}_{-1}}{F^2 \mathcal{O}_{-1}}.$$

Definition 2.1. *The composition*

$$F^1 \mathcal{O}_{-1} \xrightarrow{Q} \mathcal{O}_0 \longrightarrow \mathcal{O}(M_0) \simeq \frac{\mathcal{O}_0}{F^1 \mathcal{O}_0}$$

is $\mathcal{O}(M_0)$ -linear and admits $F^2 \mathcal{O}_{-1}$ in the kernel. It is therefore given by the contraction with a canonical section of E_{+1} that we call the curvature of the Q -manifold M and denote by κ .

Equivalently, the curvature is defined by the following commutative diagram, whose horizontal lines are exact:

$$\begin{array}{ccccc} F^2 \mathcal{O}_{-1} & \hookrightarrow & F^1 \mathcal{O}_{-1} & \twoheadrightarrow & \Gamma(E_{+1}^*) \\ \downarrow Q & & \downarrow Q & & \downarrow \mathbf{i}_\kappa \\ F^1 \mathcal{O}_0 & \hookrightarrow & \mathcal{O}_0 & \twoheadrightarrow & \mathcal{O}(M_0) \end{array}$$

Remark 2.2. The previous description of the curvature, although abstract, implies that it is a canonical notion, but it can be described in a more explicit manner, upon choosing a splitting. The polynomial degree is then well-defined, and \mathbf{i}_κ is the only component of Q of polynomial degree -1 .

$$Q = \mathbf{i}_\kappa + \sum_{i \geq 0} Q^{[i]}$$

where $Q^{[i]}$ is the component of polynomial degree i of Q . Also, after having chosen a splitting (which always exists in the smooth case) and local coordinates:

$$(2.2) \quad Q = \sum_{i=1}^{\text{rk}(E_{+1})} \tilde{\kappa}_i(x) \frac{\partial}{\partial \eta_i} + \sum_{j=1}^{\dim(M_0)} f_j \frac{\partial}{\partial x_j} + \sum_{i \in \mathbb{Z} \setminus \{0,1\}} \sum_{j=1}^{\text{rk}(E_i)} g_{i,j} \frac{\partial}{\partial \theta_{i,j}}.$$

Here the x_i 's are the variables in the base manifold, the η_i 's are the degree -1 variables, the $\theta_{i,j}$'s are the degree j variables for $j \neq 0, -1$, the functions $\tilde{\kappa}_i(x) \in \mathcal{O}_0$ are functions whose projection in $C^\infty(M_0)$ are the components of the section κ , $f_j \in \mathcal{O}_1$, and $g_{i,j} \in \mathcal{O}_{1-i}$. \square

²We present the results of this section for Q -manifolds with a smooth base. All results in section 2 extend to Q -manifolds over affine varieties or Q -manifolds over Stein varieties. This may no longer be true for the results of Section 3, where we will treat non-smooth cases separately.

It is well-known [15] that on a super-manifold of dimension (n, p) , every point where self-commuting odd vector field Q does not vanish on the zero section, there exist local coordinates $(x_1, \dots, x_n, \eta_1, \dots, \eta_p)$ such that

$$(2.3) \quad Q = \frac{\partial}{\partial \eta_1}.$$

Below is the equivalent of this statement for the \mathbb{Z}^* -graded case.

Proposition 2.3. *Let (M_0, \mathcal{O}, Q) be a \mathbb{Z}^* -graded Q -manifold with associated bundles $(E_i)_{i \in \mathbb{Z}^*}$ over M_0 . Over every open set $U \subset M_0$ over which the curvature $\kappa \in \Gamma(E_1)$ is different from zero at every point, there is a splitting $\mathcal{O}(U) \simeq \Gamma\left(\tilde{S}(\oplus_{i \in \mathbb{Z}^*} E_i^*)\right)$ under which*

$$Q = \mathbf{i}_\kappa,$$

i.e. the degree +1 vector field Q is given by the contraction with the curvature.

Remark 2.4. In the situation when there is duality (in the sense of Section 1.4), Proposition 2.3 may be restated as follows: every open subset on which the curvature $\kappa \in \Gamma(E_{+1})$ is different from zero at every point admits a dual \mathbb{Z}^* -graded Lie ∞ -algebroid for which all the brackets $(\ell_k)_{k \geq 1}$ are equal to zero except for the 0-ary bracket (which is κ). Also, it immediately implies the existence of local coordinates as in Equation (2.2) such that Q takes the form (2.3). \square

The proof of Proposition 2.3 goes through the next three lemmas (see Definition 1.14 for the negative part Q_- of the vector field Q).

Lemma 2.5. *There exists a degree -1 function $\alpha \in F^1\mathcal{O}$ such that $Q_-(\alpha) = 1 \in \mathcal{O}$.*

Proof. Take any splitting $\mathcal{O}(U) \simeq \Gamma\left(\tilde{S}(\oplus_{i \in \mathbb{Z}^*} E_i^*)\right)$. Since the curvature κ is a nowhere vanishing section of E_{+1} , there exists $\alpha \in \Gamma(E_{+1}^*) \subset F^1\mathcal{O}$ such that $\langle \kappa, \alpha \rangle = 1$. We then have $Q(\alpha) = \langle \kappa, \alpha \rangle + F = 1 + F$ for some function $F \in F^1\mathcal{O}_0 = \mathcal{O}_0 \cap \mathcal{I}_- = \mathcal{O}_0 \cap \mathcal{I}_+$. As a consequence, $Q_-(\alpha) = 1$. \blacksquare

Lemma 2.6. *There exists a splitting $\mathcal{O}(U) = \Gamma_U\left(\tilde{S}(\oplus_{i \neq 0} E_i^*)\right)$ such that $Q = Q_-$.*

Proof. The choice of a splitting $\mathcal{O}(U) \simeq \Gamma\left(\tilde{S}(\oplus E^*)\right)$ allows to decompose functions and vector fields according to their negative degree, and any function of given degree decomposes as a sum $f = \sum_{n \geq 0} f^{(n)}$ with $f^{(n)}$ a function of negative degree n ($\deg_-(f^{(n)}) = n$). For a degree +1 vector fields R , we have:

$$R = \sum_{i \geq -1} R^{(i)}$$

with $R^{(n)}$ a vector field of negative degree n . Notice that, for instance, $Q_- = Q^{(-1)}$.

We construct by induction a sequence $\Phi_n = e^{v_n}$ (starting at $n = 1$) of graded manifold isomorphisms that satisfy the following conditions:

- (1) v_n is a vector field such that $v_n : \mathcal{O} \rightarrow F^n\mathcal{O}$ for all $n \in \mathbb{N}$ (i.e. $v_n^{(i)} = 0$ for $i < n$).
- (2) the push-forward Q_n of the vector field Q by $\Phi_n \circ \dots \circ \Phi_1$ is of the form:

$$Q_{n+1} = Q^{(-1)} + Q_{n+1}^{(n+1)} + \dots$$

The sequence is constructed as follows: $Q_0 = Q$ and at each step we choose $\mathbf{v}_{n+1} = -\alpha Q_n^{(n)}$, with α as in Lemma 2.5. It follows from $[Q_n, Q_n] = 0$ that $[Q^{(-1)}, Q_n^{(n)}] = 0$. As a consequence, the push-forward vector of Q_n by $e^{\mathbf{v}_n}$, i.e. the derivation:

$$e^{-\mathbf{v}_n} Q_n e^{\mathbf{v}_n} = \sum_{k=0}^{\infty} \frac{1}{k!} \text{ad}_{\mathbf{v}_n}^k Q \quad (\text{all sums are finite for a given negative degree})$$

is given (up to components of negative degree $\geq n+1$) by

$$Q_n + [Q_n, \mathbf{v}_n] = Q^{(-1)} + Q_n^{(n)} - [Q^{(-1)}, \alpha Q_n^{(n)}] = Q^{(-1)} + Q_n^{(n)} - Q_n^{(n)} = Q^{(-1)}.$$

The henceforth constructed sequence satisfies the required assumption. We then apply Proposition 1.21 to construct the infinite composition $\Psi := \bigcirc_{i \uparrow \geq 1} e^{\mathbf{v}_i}$. By construction, the push-forward of Q through Ψ is Q_- , which completes the proof. ■

Lemma 2.7. *There exists a splitting $\mathcal{O}(U) = \Gamma(\tilde{S}(\bigoplus_{i \neq 0} E_i^*))$ such that $Q^{(-1)} = \mathbf{i}_\kappa$.*

Proof. The proof consists in repeating the steps of the proof of Lemma 2.6, by using now the polynomial degree, which is well-defined in the negative part. We write

$$Q^{(-1)} = \mathbf{i}_\kappa + Q^{[0]} + Q^{[1]} + \dots,$$

where $[i]$ now stands for the polynomial degree. We then transport $Q^{(-1)}$ through $e^{\alpha Q^{[0]}}$. Since $[\mathbf{i}_\kappa, Q^{[0]}] = 0$, the vector field obtained in such a way is now of the form:

$$Q_1^{(-1)} = \mathbf{i}_\kappa + Q_1^{[1]} + Q_1^{[2]} + \dots,$$

for new $(Q_1 - \mathbf{i}_\kappa)$ of polynomial degree ≥ 1 . We then construct recursively a collection of isomorphisms of the graded manifold M that satisfy the requirements of Proposition 1.21: since we only use negative variables at this point, the ideal of elements of polynomial degree k in negative variables is included in $F^k \mathcal{O}$ (cf. to be more precise [9]). Their infinite composition intertwines $Q^{(-1)}$ with \mathbf{i}_κ . ■

Proof. (of Proposition 2.3) The statement follows from Lemmas 2.6 and 2.7 above: Lemma 2.6 constructs an isomorphism of graded manifold under which Q becomes its negative part Q_- , and Lemma 2.7 constructs an isomorphism of graded manifold under which Q_- becomes \mathbf{i}_κ . ■

Corollary 2.8. *Let (M_0, \mathcal{O}, Q) be a \mathbb{Z}^* -graded Q -manifold. On every open set $U \subset M_0$ over which the curvature $\kappa \in \Gamma(E_1)$ is different from zero at every point, the cohomology of $(\mathcal{O}(U), Q)$ is zero in every degree.*

Proof. The statement follows from the easily-checked fact that multiplication by the function $\alpha \in \Gamma(E_{+1}^*)$ defined in Lemma 2.5 is a contracting homotopy for $Q = \mathbf{i}_\kappa$. ■

2.2. Geometry of the zero locus of the curvature of a Q -manifold.

Consider a Q -manifold (M_0, \mathcal{O}, Q) , with associated bundle $(E_i)_{i \in \mathbb{Z}^*}$ and curvature $\kappa \in \Gamma(E_{+1})$ (see Definition 2.1).

Definition 2.9. *We call the zero locus ideal of \mathcal{O} the image of*

$$\mathbf{i}_\kappa: \Gamma(E_{+1}^*) \rightarrow \mathcal{O}$$

and we denote it by $\langle \kappa \rangle$. We call functions on the zero locus the quotient algebra $\mathcal{O}/\langle \kappa \rangle$.

The space $\mathcal{I}_- + \mathcal{O}Q[\mathcal{I}_-] \subset \mathcal{O}$ is both an ideal of \mathcal{O} and stable by Q , so that the latter induces a derivation Q^+ of the quotient

$$\mathcal{K}_+ := \frac{\mathcal{O}}{\mathcal{I}_- + \mathcal{O}Q[\mathcal{I}_-]},$$

so that (\mathcal{K}_+, Q_+) is a differential graded algebra.

Definition 2.10. *The differential graded algebra (\mathcal{K}, Q_+) is called the zero locus DGA of a \mathbb{Z}^* -graded Q -manifold (M_0, \mathcal{O}, Q) .*

Here is an important result.

Proposition 2.11. *The zero locus DGA (\mathcal{K}_+, Q_+) of a \mathbb{Z}^* -graded Q -manifold (M_0, \mathcal{O}, Q) is a positively graded Q -variety over the algebra $C^\infty(M_0)/\langle \kappa \rangle$ of functions on the zero locus and there is a splitting*

$$(2.4) \quad \mathcal{K}_+ \simeq \Gamma_{\langle \kappa \rangle} (S(\oplus_{i \geq 1} E_{-i}^*)).$$

Here $\langle \kappa \rangle$ is the zero-locus ideal and $\Gamma_{\langle \kappa \rangle}(E) = \Gamma(E) \otimes_{C^\infty(M_0)} C^\infty(M_0)/\langle \kappa \rangle$ for every vector bundle $E \rightarrow M$.

We start with a lemma.

Lemma 2.12. *For any \mathbb{Z}^* -graded Q -manifold (M_0, \mathcal{O}, Q) :*

$$\mathcal{O}Q[\mathcal{I}_-] + \mathcal{I}_- = \langle \kappa \rangle \mathcal{O} + \mathcal{I}_-,$$

where κ stands for the curvature.

Proof. For any $\alpha \in \Gamma_{E_{+1}^*}$:

$$\langle \kappa, \alpha \rangle = Q[\alpha] + \sum_{i \geq 1} F_i G_i$$

where $F_i, G_i \in \mathcal{O}$ are functions of degree $-i$ and $+i$ respectively (the sum might be infinite). This proves the inclusion

$$\langle \kappa \rangle \mathcal{O} + \mathcal{I}_- \subset \mathcal{O}Q[\mathcal{I}_-] + \mathcal{I}_-.$$

The converse inclusion is straightforward. ■

Proof. (of Proposition 2.11) As a consequence of Lemma 2.12 above, the graded algebra morphism

$$\Gamma(S(\oplus_{i \geq 1} E_{-i}^*)) \rightarrow \mathcal{K}$$

is surjective, so that the following sequence is exact:

$$0 \rightarrow \langle \kappa \rangle \Gamma(S(\oplus_{i \geq 1} E_{-i}^*)) \rightarrow \Gamma(S(\oplus_{i \geq 1} E_{-i}^*)) \rightarrow \mathcal{K}_+ \rightarrow 0.$$

Consequently:

- (1) the degree of elements in \mathcal{K}_+ is non-negative by construction,
- (2) degree 0-elements can be identified with $\mathcal{O}(M_0)/\langle \kappa \rangle$,
- (3) for $k \geq 1$, degree $+k$ elements are elements of degree k in the symmetric algebra (over $\mathcal{O}(M_0)/\langle \kappa \rangle$) of $\oplus_{i \geq 1} \Gamma(E_{-i}^*) \otimes \mathcal{O}(M_0)/\langle \kappa \rangle$.

This yields the isomorphism of projective $\mathcal{O}(M_0)/\langle \kappa \rangle$ -module in Equation (2.4). ■

Definition 2.13. *Let (M_0, \mathcal{O}, Q) be a \mathbb{Z}^* -graded Q -manifold. We call zero locus NQ-variety the NQ-variety with sheaf of functions \mathcal{K}_+ and differential Q_+ .*

Remark 2.14. For (M_0, \mathcal{O}, Q) be a \mathbb{Z}^* -graded Q -manifold with splitting, Q can be decomposed by the negative degree as an infinite sum:

$$Q = q_{-1} + q_0 + \cdots + q_i + \dots$$

with q_i a degree +1 vector field of negative degree i for $i \geq -1$. Then, it is easy to see that q_{-1} induces the negative part of the Q -manifold and that q_0 (which commutes with q_{-1} , hence induces a derivation of \mathcal{K}_+) induces the differential Q_+ of the zero locus NQ-variety. \square

Remark 2.15. As explained in [12], when the ideal κ is the ideal of functions vanishing on a submanifold $X \subset M_0$, then the distribution $\mathcal{D} := \rho_1(\Gamma(E_{-1}))$ is made of vector fields tangent to X and its restriction to X is involutive on X . This singular foliation on the submanifold X is the basic singular foliation of the NQ-manifold (\mathcal{K}_+, Q_+) . The same conclusion holds when X is a singular subset, provided that vector fields on X can be defined in an appropriate manner (e.g.: an affine variety). \square

3. KOSZUL-TATE RESOLUTION AND VECTOR FIELDS ON THE ZERO LOCUS NQ-VARIETY

Recall that for any vector bundle $E \rightarrow M_0$ and any ideal $I \subset C^\infty(M_0)$, we use the following notation:

$$(3.5) \quad \Gamma_I(E) := \Gamma(E) \otimes_{C^\infty(M_0)} C^\infty(M_0)/I.$$

If I is the vanishing ideal of a submanifold $X_I \subset M_0$ (i.e. X_I is the zero locus of I), then $\Gamma_I(E)$ is simply the space of sections of the restriction of E to X_I .

3.1. Koszul-Tate resolutions. Let M_0 be a smooth manifold. We recall the usual definition of a Koszul-Tate resolution of an ideal.

Definition 3.1. A Koszul-Tate resolution³ of an ideal $I \subset C^\infty(M_0)$ is a \mathbb{Z}_- -graded Q -manifold $(M_0, \mathcal{O}_-, \delta)$ which

- (1) is concentrated in non-positive degree $\mathcal{O}_- = \bigoplus_{i \leq 0} \mathcal{O}_i$, with $\mathcal{O}_0 = C^\infty(M_0)$,
- (2) and satisfies that the cohomology of the (total) degree +1 vector field $\delta: \mathcal{O}_- \rightarrow \mathcal{O}_-$ is given by

$$(3.6) \quad H^i(\mathcal{O}_-, \delta) = \begin{cases} C^\infty(M_0)/I, & i = 0 \\ 0, & i < 0 \end{cases}$$

Example 3.2. For $M_0 = \mathbb{R}^n$ and I the ideal of functions vanishing at 0, the graded algebra exterior form $\Omega(M_0)$ equipped with $\delta = \mathbf{i}_E$ the contraction with the Euler vector field is a Koszul-Tate resolution of I . In that case, only $E_{+1} = TM_0$ is non-zero and the curvature is the Euler vector field.

We start with a few remarks that may help to understand the notion.

Remark 3.3. Item (1) in Definition 3.1 implies that the associated canonical graded vector bundle E_\bullet of a Koszul-Tate resolution is concentrated in positive degrees $E_\bullet = \bigoplus_{i \geq 1} E_{+i}$. \square

³We use the standard notations from [7].

Remark 3.4. Let $\kappa \in \Gamma(E_{+1})$ be the curvature of a Koszul-Tate resolution. The condition on $H^0(\mathcal{O}, \delta)$ in Definition 3.1 implies that the curvature ideal $\langle \kappa \rangle$ of $\kappa \in \Gamma(E_{+1})$ coincides with I , i.e. a function $F \in C^\infty(M_0)$ belongs to I if and only if there exists a section $\alpha \in \Gamma(E_{+1}^*)$ such that $F = \langle \kappa, \alpha \rangle = \delta(\alpha)$. \square

We will need a variation of Definition 3.1.

Definition 3.5. Let $I \subset C^\infty(M_0)$ be an ideal, and consider a positively-graded variety \mathcal{K}_+ on $C^\infty(M_0)/I$. A Koszul-Tate resolution of \mathcal{K}_+ is a pair made of

- (1) a splitting of \mathcal{K}_+ , i.e.

$$\mathcal{K}_+ \simeq \Gamma_I(S(\oplus_{i \geq 1} E_{-i}^*)),$$

- (2) a Koszul-Tate resolution of I with splitting

$$(\Gamma(S(\oplus_{i \geq 1} E_i^*)), \delta),$$

assembled into a Q -manifold $(M_0, \mathcal{O}, \tilde{\delta})$ with splitting

$$\mathcal{O} \simeq \tilde{S}(\Gamma(\oplus_{i \neq 0} E_i^*))$$

where $\tilde{\delta}$ is the extension of δ which is identically 0 on $\oplus_{i \geq 1} \Gamma(E_{-i}^*)$.

Here are a few comments about Definition 3.5.

Lemma 3.6. Let $(M_0, \mathcal{O}, \tilde{\delta})$ be a Koszul-Tate resolution of \mathcal{K}_+ as in Definition 3.5.

- (1) Its zero locus DGA $\frac{\mathcal{O}}{\mathcal{O}\tilde{\delta}(\mathcal{I}_-) + \mathcal{I}_-}$ is \mathcal{K}_+ .

- (2) The cohomology of the complex $(\mathcal{O}, \tilde{\delta})$ is given by:

$$(3.7) \quad H^i(\mathcal{O}, \tilde{\delta}) = \begin{cases} \mathcal{K}_+, & i = 0 \\ 0, & i > 0 \end{cases}$$

Here the degree considered is the negative degree \deg_- (compare with Equation (3.6) where the index was running over the total degree, i.e. the total degree of $\tilde{\delta}$ is still $+1$).

Proof. The first item is a direct consequence of the identification:

$$\mathcal{O}\tilde{\delta}(\mathcal{I}_-) + \mathcal{I}_- = I\mathcal{O} + \mathcal{I}_- = \langle I + \Gamma(\oplus_{i \geq 1} E_i^*) \rangle.$$

Let us prove the second item. The cohomology of the complex

$$(3.8) \quad (\Gamma(S(\oplus_{i \geq 1} E_{-i}^*)) \otimes_{C^\infty(M_0)} \Gamma(S(\oplus_{i \geq 1} E_{+i}^*)), \text{id} \otimes \delta)$$

is $\Gamma(S \oplus_{i \geq 1} E_{-i}^*) \otimes_{C^\infty(M_0)} C^\infty(M_0)/I \simeq \mathcal{K}_+$. Now, $(\mathcal{O}, \tilde{\delta})$ is the completion of the complex (3.8) with respect to the negative degree, but completion does not affect cohomology, and the result follows. \blacksquare

We conclude the section with an important definition. To any Q -manifold (M_0, \mathcal{O}, Q) was associated in Definition 1.14 another Q -manifold, called its negative part $(M_0, \mathcal{O}/\mathcal{I}_+, Q_-)$.

Definition 3.7. We say that a \mathbb{Z}^* -graded Q -manifold (M_0, \mathcal{O}, Q) with curvature κ has a Koszul-Tate negative part if its negative part is a Koszul-Tate resolution of the curvature ideal $\langle \kappa \rangle$.

3.2. Vector fields on Koszul-Tate resolutions I: the cohomology. The space $\mathfrak{X}(\mathcal{O}_-)$ of vector fields on a Koszul-Tate resolution $(M_0, \mathcal{O}_-, \delta)$ of an ideal $I \subset C^\infty(M_0)$ form a DGLA when equipped with the graded commutator and the differential ad_δ . In particular, $(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta)$ is a complex, whose cohomology we now compute. To start with, let us notice that $((\mathcal{O}_- \delta(\mathcal{I}_-) + \mathcal{I}_-) \mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta)$ is a subcomplex of $(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta)$. The quotient complex is canonically isomorphic to a complex of the form

$$(3.9) \quad \Gamma_I(TM) \mapsto \Gamma_I(E_{+1}) \mapsto \Gamma_I(E_{+2}) \mapsto \cdots,$$

recall Equation (3.5) for the notation Γ_I .

Definition 3.8. Let $(M_0, \mathcal{O}, \delta)$ be a Koszul-Tate resolution of an ideal $I \subset C^\infty(M_0)$. We call linearization of Koszul-Tate differential δ at the zero locus the complex (3.9), and denote it by $(\mathfrak{X}_{lin}, \delta_{lin})$.

Remark 3.9. The complex (3.9) can be understood as follows when I is the vanishing ideal of a subset $X_I \subset M_0$: the differential of the curvature $\kappa : M_0 \rightarrow E_{+1}$ is a vector bundle morphism:

$$T\kappa : TM_0 \rightarrow TE_{+1}$$

over $\kappa : M_0 \rightarrow E_{+1}$. Now, for any $m \in X_I$, since $\kappa(m) = 0_m$, there is a canonical decomposition $T_{0_m}E_{+1} = T_mM + E_{+1}|_m$, $pr_2 \circ T_m\kappa$ can be seen as a linear map $T_mM \rightarrow E_{+1}|_m$, where pr_2 being the projection onto the second component. This map easily checked to coincide with the first bundle morphism in (3.9). All remaining morphisms in (3.9) are simply the restriction to X_I of the component of polynomial degree 0 of δ (which is by construction a degree +1 vector bundle endomorphism of E_\bullet). \square

Let us now state the main result of this section. For a Koszul-Tate resolution, recall that the negative degree is simply the opposite of the total degree. Also, ad_δ is a negative degree -1 operator. Also, the following remark helps to understand the statement.

Remark 3.10. A ad_δ -cocycle q of degree 0 induces a vector field $\underline{q} \in \mathfrak{X}(M_0)$ which satisfies $\underline{q}[I] \subset I$, and therefore induces a derivation q^I of $C^\infty(M_0)/I$. If q is an ad_δ -coboundary, then $q^I = 0$. \square

Theorem 3.11. Let $(M_0, \mathcal{O}_-, \delta)$ be a Koszul-Tate resolution of an ideal $I \in C^\infty(M_0)$. With respect to the negative degree, we have:

$$(3.10) \quad H^{-i}(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta) = \begin{cases} H^{-i}(\mathfrak{X}_{lin}, \delta_{lin}), & i \leq 0 \\ 0, & i > 0 \end{cases}$$

In particular, an ad_δ -cocycle q of degree 0 is a coboundary if and only if its induced derivation q^I of $C^\infty(M_0)/I$ is zero.

Proof. Let us chose a splitting $\mathcal{O}_- \simeq \Gamma(S(\oplus_{i \geq 1} E_i^*))$, and a family of affine connections ∇^k on E_k^* (recall Remark 1.8 for notations). Consider the following bigrading (on the ‘‘North-West’’ quarter):

$$(3.11) \quad \mathfrak{X}(\mathcal{O}_-)_{a,b} = \begin{cases} \mathcal{O}_a \otimes \mathfrak{X}(M_0) & \text{for } a \geq 0 \text{ and } b = 0, \\ \mathcal{O}_a \otimes \Gamma(E_{-b}) & \text{for } a \geq 0 \text{ and } b \leq -1, \\ 0 & \text{otherwise.} \end{cases}$$

We adopt the following conventions:

- (1) All tensor products are over $C^\infty(M_0)$.
- (2) A section $e \in \Gamma(E_b)$ is seen as the vertical vector field given by the derivation \mathbf{i}_e of \mathcal{O}_- . Notice that for the negative degree, it is of degree $-b$.
- (3) A vector field $X \in \mathfrak{X}(M_0)$ is extended to a degree 0 derivation of \mathcal{O}_- by $X[\epsilon_k] = \nabla_X^k \epsilon_k$ for every section $\epsilon_k \in \Gamma(E_k^*)$.

It is indeed a bigrading, since:

$$\mathfrak{X}(\mathcal{O}_-)_i = \bigoplus_{a \geq 0} \mathfrak{X}(\mathcal{O}_-)_{a, i-a},$$

(infinite sums are allowed, since they converge with respect to the filtration $(F^i \mathcal{O})_{i \geq 0}$).

With respect to this bi-grading, ad_δ decomposes as follows:

$$\begin{array}{c}
 \mathfrak{X}(\mathcal{O}_-)_{a+2, b-3} \\
 \swarrow \\
 \mathfrak{X}(\mathcal{O}_-)_{a+1, b-2} \\
 \swarrow \quad \searrow \\
 \mathfrak{X}(\mathcal{O}_-)_{a, b-1} \xleftarrow{\text{id} \otimes D} \mathfrak{X}(\mathcal{O}_-)_{a, b} \\
 \downarrow \delta \otimes \text{id} \\
 \mathfrak{X}(\mathcal{O}_-)_{a-1, b}
 \end{array}$$

We can now use generic diagram chasing arguments: since all vertical lines are acyclic in degree $\neq 0$, the cohomology is concentrated in the 0-th cohomology of the line $a = 0$, which coincides with $\Gamma_I(TM_0)$ for $b = 0$ and $\Gamma_I(E_b)$ for $b \leq -1$. Equipped with the induced differential, a direct computation shows that it coincides (with opposite signs) with the differential of the complex (3.9). This proves (3.10).

Since all vertical lines are exact, a degree 0 ad_δ -cocycle is exact if and only if its bi-degree $(0, 0)$ component lies in the image of the vertical lines, i.e. belong to $I \otimes \mathfrak{X}(M_0)$. Equivalently, this means that this element induces the zero map on $C^\infty(M_0)/I$. This completes the proof. ■

Here is an immediate consequence of Theorem 3.11.

Corollary 3.12. *Let $(M_0, \mathcal{O}, \tilde{\delta})$ be a Koszul-Tate resolution of \mathcal{K}_+ as in Definition 3.5. With respect to the negative degree:*

$$(3.12) H^i(\mathfrak{X}(\mathcal{O}), \text{ad}_{\tilde{\delta}}) = \begin{cases} \mathcal{K}_+ \otimes_{C^\infty(M_0)/I} H^{-i}(\mathfrak{X}_{lin}, \delta_{lin}), & i < 0 \\ \mathcal{K}_+ \otimes_{C^\infty(M_0)/I} (\bigoplus_{i \geq 1} \Gamma_I(E_{-i}) \oplus H^0(\mathfrak{X}_{lin}, \delta_{lin})), & i = 0 \\ 0, & i > 0 \end{cases}$$

Proof. As in the proof of Theorem 3.11, one can use a family of connections on $(E_i)_{i \in \mathbb{Z}^*}$ to decompose the \mathcal{O} -module $\mathfrak{X}(\mathcal{O})$ as the sum of two submodules: one is

$$\mathcal{O} \otimes (\bigoplus_{i \geq 1} E_i \oplus TM_0) \text{ and } \mathcal{O} \otimes (\bigoplus_{i \geq 1} E_{-i})$$

Both modules are $\text{ad}_{\tilde{\delta}}$ -stable. On the second one, $\text{ad}_{\tilde{\delta}} = \tilde{\delta} \otimes \text{id}$, so that the cohomology is concentrated in negative degree 0 and coincides with

$$\frac{\Gamma(S(\bigoplus_{i \geq 1} E_{-i}^*))}{I} \otimes_{C^\infty(M_0)} \Gamma(\bigoplus_{i \geq 1} E_{-i}) = \mathcal{K}_+ \otimes_{C^\infty(M_0)/I} \Gamma_I(\bigoplus_{i \geq 1} E_{-i}).$$

The first one is the completion of the tensor product of $\Gamma(S(\oplus_{i \geq 1} E_{-i}^*))$ with the module $\mathfrak{X}(\mathcal{O}_-)$ of vector fields on a Koszul-Tate resolution of I , whose cohomology is given in Theorem 3.11. The differential being given by $\text{id} \otimes \text{ad}_{\bar{\delta}}$, the result then follows from Theorem 3.11 and the fact that \mathcal{K}_+ is a $C^\infty(M_0)/I$ -projective module, so that tensoring with \mathcal{K}_+ preserves cohomology. ■

3.3. Vector fields on Koszul-Tate resolutions II: the extension. We now consider another problem. As stated in Remark 3.10, for a Koszul-Tate resolution $(M_0, \mathcal{O}_-, \delta)$ of an ideal I , an $\text{ad}_{\bar{\delta}}$ -cocycle $q \in \mathfrak{X}(\mathcal{O}_-)$ induces a derivation q^I of $C^\infty(M_0)/I$, and an $\text{ad}_{\bar{\delta}}$ -coboundary induces a derivation equal to zero. In particular, there is a Lie algebra morphism

$$(3.13) \quad H^0(\mathfrak{X}(\mathcal{O}_-), \text{ad}_{\bar{\delta}}) \longrightarrow \text{Der}(C^\infty(M_0)/I).$$

The second part of Theorem 3.11 implies that this morphism is injective. The following statement shows that it is surjective.

Proposition 3.13. *Let $(M_0, \mathcal{O}_-, \delta)$ be a Koszul-Tate resolution of an ideal $I \subset C^\infty(M_0)$. Then the natural Lie algebra morphism 3.13 is an isomorphism*

$$H^0(\mathfrak{X}(\mathcal{O}_-), \text{ad}_{\bar{\delta}}) \simeq \text{Der}(C^\infty(M_0)/I).$$

In particular, every derivation q^I of $C^\infty(M_0)/I$ is induced by a degree 0 vector field $q \in \mathfrak{X}(\mathcal{O}_-)$ such that $[\delta, q] = 0$.

Proof. Denote the projection $C^\infty(M_0) \rightarrow C^\infty(M_0)/I$ by $F \mapsto \bar{F}$. Also, let us choose $(U_k, \chi_k)_{k \in K}$ a partition of unity of the manifold M_0 for which each U_k is a coordinate neighborhood on which each one of the vector bundles E_k admits a trivialization.

Let q^I be a derivation of $C^\infty(M_0)/I$. Let x_1, \dots, x_n be local coordinates on the open subset U_k for some $k \in K$. Consider any functions $F_1, \dots, F_r \in C^\infty(U_k)$ such that $q^I(\bar{x}_i) = \bar{F}_i$. The vector field

$$\nu_k^0 := \sum_{i=1}^r F_i \frac{\partial}{\partial x_i}$$

satisfies by construction that $\nu_k^0(I) \subset I$, since it induces the derivation of $C^\infty(U_k)/I$ which coincides with the restriction of q^I to U_k . These local vector fields ν_k^0 can be glued to a vector field ν^0 on M_0 :

$$\nu^0 = \sum_{k \in K} \chi_k \nu_k^0.$$

This vector field still satisfies $\nu^0[I] \subset I$ by construction.

Since $I = \delta(\Gamma(E_{+1}^*))$, for any local trivialization η_1, \dots, η_r of E_{+1}^* , defined on the open subset U_k , there exist functions $(\phi_{i,j})_{i,j=1}^r$ in $C^\infty(U_k)$ such that the collection of functions $\kappa_i := \delta(\eta_i)$, $i = 1, \dots, r$ locally generates the vanishing ideal of the zero locus and

$$\nu^0 \delta(\eta_i) = \nu^0(\kappa_i) = \sum_{j=1}^r \phi_{i,j} \kappa_j = \sum_{j=1}^r \phi_{i,j} \delta(\eta_j)$$

Consider the vector field

$$\nu_k^1 := \sum_{i,j=1}^r \phi_{j,i} \eta_i \frac{\partial}{\partial \eta_j}.$$

By construction, it satisfies

$$\nu^0 \circ \delta = \delta \circ \nu_k^1.$$

Since δ is $C^\infty(M_0)$ -linear, the vector field

$$\nu^1 := \sum_{k \in K} \chi_k \nu_k^1$$

also satisfies:

$$\nu^0 \circ \delta = \delta \circ \nu^1.$$

Now, ν^0, ν^1 extends to vector fields on \mathcal{O}_- , that we will denote by the same symbol. The proof then consists in constructing recursively $\nu^j \in \bigoplus_{b \leq -j} \mathfrak{X}(\mathcal{O}_-)_{a,b}$ such that

$$\nu^j \circ \delta = \delta \circ \nu^{j+1}.$$

Provided that ν_0, \dots, ν_j are constructed, the existence of ν_{j+1} follows from the fact that $\nu^j \circ \delta$ is valued in the kernel of δ :

$$\delta \circ \nu^j \circ \delta = \nu^{j-1} \circ \delta^2 = 0.$$

Since the cohomology of (\mathcal{O}_-, δ) is zero, $\nu^j \circ \delta$ is therefore valued in the image of δ , and since \mathcal{O}_- is a projective $C^\infty(M_0)$ -module, the existence of the vector field ν^{j+1} is granted.

Moreover, with respect to the bi-grading above (see Equation (3.11)),

$\nu^{j+1} \in \bigoplus_{b \leq -(j+1)} \mathfrak{X}(\mathcal{O}_-)_{a,b}$. As a consequence, the sequence

$$(3.14) \quad q^k := \sum_{j=0}^k \nu^j,$$

converges and the limit is the desired vector field q . ■

Consider now a Koszul-Tate resolution of a graded variety given by \mathcal{K}_+ as in Definition 3.5. Again, notice that a total degree k and negative degree 0 vector field q_0 such that $[\tilde{\delta}, q_0] = 0$ induces a degree k derivation of \mathcal{K}_+ . If q_0 is an $\text{ad}_{\tilde{\delta}}$ -cocycle, that derivation is zero. Proposition 3.13 extends easily to give the following result.

Corollary 3.14. *Let $(M_0, \mathcal{O}, \tilde{\delta})$ be a Koszul-Tate resolution of a graded variety \mathcal{K}_+ as in Definition 3.5. There is a natural isomorphism:*

$$H^{(0,k)}(\mathfrak{X}(\mathcal{O}), \text{ad}_{\tilde{\delta}}) = \text{Der}^k(\mathcal{K}_+)$$

where $H^{(0,k)}$ stands for the cohomology in negative degree 0 and total degree k and Der^k stands for derivations of degree k of \mathcal{K}_+ . In particular, for any degree k derivation Q_+ of \mathcal{K}_+ , there exists $q_0 \in \mathfrak{X}(\mathcal{O})$ of negative degree 0 and total degree k satisfying $[\tilde{\delta}, q_0] = 0$ and inducing the derivation Q_+ on \mathcal{K}_+ .

Remark 3.15 (Extension of derivations in the affine case). At the beginning of the proof of Proposition 3.13 it was shown that in the smooth category any derivation of the quotient algebra \mathcal{O}/I , considered as functions on the zero locus, can be extended to a derivation of the entire algebra of functions \mathcal{O} . This is also true in the affine case.

Let M_0 be an affine n -dimensional space over a field \mathbb{k} of characteristic 0 (we think of it as \mathbb{R} or \mathbb{C}) with affine coordinates $(z^i)_{i=1}^n$; and $I \subset \mathcal{O}(M_0) = \mathbb{k}[z^1, \dots, z^n]$ be an ideal, then every derivation of $\mathcal{K} = \mathcal{O}(M_0)/I$ admits an extension to a derivation of $\mathcal{O}(M_0)$. Indeed, let q^I be a derivation of \mathcal{K} . Define q such that $q(z^i)$ equals to the preimage of

$q^I[z^i] \in \mathcal{K}$ under the projection map $\mathcal{O}(M_0) \rightarrow \mathcal{K}$, where $[z^i] = z^i/I$. Let us extend q to the whole algebra of functions $\mathcal{O}(M_0)$ by the Leibniz rule. It is easy to see that $q(I) \subset I$ and $q/I = q^I$. Proposition 3.13 remains therefore valid in the context of affine varieties in algebraic geometry. \square

Remark 3.16. The above statements (Theorem 3.11 and Proposition 3.13) can be proved in a the following alternative way. Let $\mathfrak{X}_{\text{null}} = \mathfrak{X}_{\text{null}}(\mathcal{O}_-)$ be the Lie super subalgebra of derivations \mathfrak{v} of \mathcal{O}_- , satisfying $\mathfrak{v}(\mathcal{O}_-) \subset I_-$, where $I_- = I \oplus \bigoplus_{i < 0} \mathcal{O}_i$. In particular, $\mathfrak{X}_{\text{null}}$ contains all derivations of positive negative degree. Since $\mathcal{K} = \mathcal{O}_0/I \simeq \mathcal{O}_-/I_-$, $\mathfrak{X}_{\text{null}}$ can be identified with vector fields on the whole non-positively graded manifold $(M_{0,c}, \mathcal{O}_-)$, that vanish at the zero locus; therefore $\mathfrak{X}(\mathcal{O}_-)/\mathfrak{X}_{\text{null}} = \mathfrak{X}_I$, where $\mathfrak{X}_I = \mathfrak{X}(\mathcal{O}_-) \otimes_{\mathcal{O}_-} \mathcal{K}$. Notice that, since $\delta \in \mathfrak{X}_{\text{null}}$, the subalgebra $\mathfrak{X}_{\text{null}}$ is closed under the adjoint action of δ . Thus we have a short exact sequence of complexes

$$0 \rightarrow (\mathfrak{X}_{\text{null}}, \text{ad}_\delta) \rightarrow (\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta) \rightarrow (\mathfrak{X}_I, \text{ad}_\delta) \rightarrow 0$$

which leads to the long exact sequence in cohomology

$$\dots \rightarrow H^i(\mathfrak{X}_{\text{null}}, \text{ad}_\delta) \rightarrow H^i(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta) \rightarrow H^i(\mathfrak{X}_I, \text{ad}_\delta) \rightarrow H^{i+1}(\mathfrak{X}_{\text{null}}, \text{ad}_\delta) \rightarrow \dots$$

Notice that $(\mathfrak{X}_I, \text{ad}_\delta)$ coincides with (3.9). First we prove that the complex $(\mathfrak{X}_{\text{null}}, \text{ad}_\delta)$ is acyclic. Combining this statement with the fact that any derivation of \mathcal{K} extends to a derivation of \mathcal{O}_- (see the beginning of the Proposition 3.13), we prove Theorem 3.11 and the remaining part of Proposition 3.13 altogether. \square

Remark 3.17.

- In fact, Theorem 3.11 computed the cohomology of $(\mathfrak{X}(\mathcal{O}), \delta)$ by using the spectral sequence associated to the following filtration of $\mathfrak{X}(\mathcal{O}_-)$: $F^p \mathfrak{X}(\mathcal{O}_-)$ is the Lie subalgebra of vector fields on M which annihilate the subalgebra of functions generated by all elements of negative degree $0, \dots, p-1$. This spectral sequence converges in the second term. Its zero term gives sections of the graded vector bundle $\pi_-^*(TM_0 \oplus E_+)$, where $E_+ = \bigoplus_{k > 0} E_k$ and π_- is the projection of (M_0, \mathcal{O}) onto M_0 , while the first term of the spectral sequence - sections of the restriction of $TM_0 \oplus E_+$ on X together with the differential determined by the normal linearization of δ along X .
- The “restriction” of vector fields to the zero locus $\mathfrak{X}_I(\mathcal{O})$ is an \mathbb{N} -graded $C^\infty(M_0)/I$ -module, the homogeneous components of which of negative degree $i = 0$ and $i < 0$ are canonically isomorphic to $\mathfrak{X}_I(\mathcal{O}_-)$ and $\mathcal{K} \otimes_{\mathcal{O}_-} \Gamma(E_{-i})$, respectively.
- $\bigoplus_i H^i(\mathfrak{X}_{lin}, \delta_{lin})$ is a graded Lie-Rinehart $C^\infty(M_0)/I$ -algebra, which extends $\mathfrak{X}_I(\mathcal{O}_-)$; furthermore, the latter is embedded into the former as a Lie-Rinehart $C^\infty(M_0)/I$ -subalgebra consisting of all elements of negative degree.

\square

It follows from Theorem 3.11 that the ad_δ -cohomology of vector fields on a Koszul-Tate resolution $(M_0, \mathcal{O}_-, \delta)$ of I are zero near any point outside the zero locus of I . We also claim that non-trivial cohomologies of non-zero degree only appear on the singular part of the zero locus of I . The following examples illustrate this phenomenon. Example 3.18 is provided to show that the positive degree part in cohomology of $(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta)$, where (\mathcal{O}_0, δ) is the Koszul-Tate resolution of an ideal $I \subset \mathcal{O}(M_0)$, is related to singularities of

the zero locus of this ideal. Example 3.19 tells us that, even in the complete intersection case, the degree 1 cohomology of vector fields on a Koszul-Tate resolution can be non-trivial.

Example 3.18. Assume that $X \subset M_0$ is a smooth submanifold, I is the ideal of functions vanishing on X , δ is a Koszul-Tate differential which resolves I , such that X is the zero locus of δ regarded as a homological vector field on a non-positively graded M . It is possible to cover M by graded coordinate charts such that either such a chart does not intersect X , then the corresponding i_κ is non-vanishing at all points, so we can use Lemma 2.5 and technique from Corollary 2.8 to show that the ad_δ -cohomology of vector fields over this chart are vanishing, or there are adapted coordinates $(x^i, y^a, \eta^a, \xi^\alpha, \zeta^\alpha)$ such that⁴

$$\delta = \sum_a y^a \frac{\partial}{\partial \eta^a} + \sum_\alpha \zeta^\alpha \frac{\partial}{\partial \xi^\alpha}.$$

In such case the intersection of the above coordinate chart with X is given by equations $y^a = \eta^a = \xi^\alpha = \zeta^\alpha = 0$, therefore all sections of the restriction of TM onto X are of the form

$$v(x, \frac{\partial}{\partial x}) + \sum_a \left(f^a(x) \frac{\partial}{\partial y^a} + h^a(x) \frac{\partial}{\partial \eta^a} \right) + \sum_\alpha \left(\lambda^\alpha(x) \frac{\partial}{\partial \zeta^\alpha} + \mu^\alpha(x) \frac{\partial}{\partial \xi^\alpha} \right)$$

It is easy to see that $\left\{ \frac{\partial}{\partial y^a}, \frac{\partial}{\partial \eta^a}, \frac{\partial}{\partial \zeta^\alpha}, \frac{\partial}{\partial \xi^\alpha} \right\}$ generate an acyclic complex w.r.t. δ , therefore the cohomology of positive degree sections of $TM|_X$ are zero over this coordinate chart and thus on the whole X as δ is linear under the multiplication on functions on M_0 which allows us to apply the partition of unity technique.

Example 3.19. On $M_0 = \mathbb{R}^2$, equipped with affine coordinates (x, y) , let I be the ideal generated by xy , and X be an affine variety given by the equation $xy = 0$. A Koszul-tate resolution of I is determined by a homological vector field $\delta = xy \frac{\partial}{\partial \xi}$ on the non-positively graded affine manifold (M_0, \mathcal{O}_-) with graded coordinates (x, y, ξ) , where ξ has total degree -1 . It is routine to check directly that, as stated in Proposition 3.13:

$$H^0(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta) \simeq \left\{ xf(x) \frac{\partial}{\partial x} + yg(y) \frac{\partial}{\partial y} \mid f, g \in C^\infty(\mathbb{R}) \right\}.$$

Since a degree 1 vector field is a ad_δ -cocycle, since $[\delta, \frac{\partial}{\partial x}] = y \frac{\partial}{\partial \xi}$, $[\delta, \frac{\partial}{\partial y}] = x \frac{\partial}{\partial \xi}$, and $[\delta, \xi \frac{\partial}{\partial \xi}] = xy \frac{\partial}{\partial \xi}$, and since the quotient of $C^\infty(M_0)$ by the ideal generated by x, y, xy is \mathbb{R} , we also have

$$H^1(\mathfrak{X}(\mathcal{O}_-), \text{ad}_\delta) = \left\{ \lambda \frac{\partial}{\partial \xi} \mid \lambda \in \mathbb{R} \right\}.$$

In particular, the degree 1 cohomology is different from zero.

3.4. Koszul-Tate resolutions and singular locus NQ -variety. Here is the main result of this section.

Theorem 3.20. *Let M_0 be a manifold and $I \subset C^\infty(M_0)$ an ideal. Given*

⁴In mathematical physics $(y^a, \eta^a, \xi^\alpha, \zeta^\alpha)$ would be called contractible pairs.

(1) a Koszul-Tate resolution $(M_0, \mathcal{O}_-, \delta)$ of I with a splitting

$$\mathcal{O}_- = \Gamma(S(\oplus_{i \geq 1} E_i^*))$$

(2) a positively graded Q -variety (\mathcal{K}_+, Q_+) on $C^\infty(M_0)/I$ with a splitting

$$\mathcal{K}_+ = \Gamma_I(S(\oplus_{i \geq 1} E_{-i}^*)),$$

there exists a Q -manifold (M_0, \mathcal{O}, Q) with a splitting:

$$\mathcal{O} \simeq \Gamma(\tilde{S}(\oplus_{i \in \mathbb{Z}^*} E_i^*))$$

- (1) whose negative part is the Koszul-Tate resolution $(M_0, \mathcal{O}_-, \delta)$,
- (2) and whose positively graded NQ -variety is (\mathcal{K}_+, Q_+) .

Two such Q -vector fields are diffeomorphic through a diffeomorphism which is the composition of flows of degree zero vector fields as in Proposition 1.21, and that induce the identity maps of the base manifold M_0 , of the negative part \mathcal{O}_- , and of the positively graded NQ -variety \mathcal{K}_+ .

Proof. The idea of the proof consists in applying perturbation theory techniques, and construct Q (and the degree 0 vector fields defining Ψ) through a recursion by showing that the obstructions for the next step are cohomology classes that vanish.

Let $(M_0, \mathcal{O}, \tilde{\delta})$ be as in Definition 3.5. The first step consists in applying Corollary 3.14: there exists a vector field q_0 of negative degree 0 and total degree +1, such that $[q_0, \tilde{\delta}] = 0$ (which implies that Q_0 induces a derivation of \mathcal{K}_+) whose induced derivation of \mathcal{K}_+ is Q_+ . Now, the proof of the existence of the vector field Q consists in constructing recursively a family $(q_i)_{i \geq 1}$ such that

- (1) Each q_i is of negative degree i and total degree +1
- (2) $Q_i = \tilde{\delta} + q_0 + \cdots + q_i$ satisfies $[Q_i, Q_i] = 0$ up to vector fields of negative degree $\geq i$.

For instance $Q_0 = \tilde{\delta} + q_0$ satisfies the recursion for $i = 0$ since:

$$[\tilde{\delta} + Q_0, \tilde{\delta} + Q_0] = [Q_0, Q_0]$$

and since $[Q_0, Q_0]$ is of negative degree 0.

Now, by the graded Jacobi identity of the graded Lie algebra of vector fields $\mathfrak{X}(\mathcal{O})$,

$$\left[\tilde{\delta} + Q_0, [\tilde{\delta} + Q_0, \tilde{\delta} + Q_0] \right] = 0,$$

so that $\text{ad}_{\tilde{\delta}}([Q_0, Q_0]) = 0$ is an $\text{ad}_{\tilde{\delta}}$ -cocycle of negative degree 0. Now, since Q_0 induces Q_+ on \mathcal{K}_+ and since $Q_+^2 = 0$, the class of $[Q_0, Q_0]$ in $\mathcal{K}_+ \otimes H^0(\mathfrak{X}_{lin}, \delta_{lin})$ is zero, so that there exists a vector field q_1 of total degree +1 and negative degree +1 such that $[\tilde{\delta}, q_1] = [q_0, q_0]$. As a consequence

$$Q_1 = \tilde{\delta} + q_0 + q_1$$

satisfies the recursion condition for $i = 1$.

The proof then continues easily by noticing that if $Q_i := \tilde{\delta} + \sum_{k=1}^i q_k$ satisfies the recursion assumption at order i , then

$$[Q_i, Q_i] = \sum_{k=0}^i [q_k, q_{i-k}] + R_i$$

with R_i of negative degree $\geq i + 2$ and $\sum_{k=0}^i [q_k, q_{i-k}]$ being the component of negative degree $i + 1$. By the graded Jacobi identity, this implies that $\sum_{k=0}^i [q_k, q_{i-k}]$ is an $\text{ad}_{\bar{\delta}}$ -cocycle of negative degree $i + 1$. Since cohomology is zero in that degree by Corollary 3.12, there exists a vector field q_{i+1} of total degree 1 and negative degree $i + 1$ such that $-\sum_{k=0}^i [q_k, q_{i-k}] = \text{ad}_{\bar{\delta}} q_{i+1}$ which in turn implies that

$$Q_{i+1} = \tilde{\delta} + \sum_{k=0}^{i+1} q_k$$

satisfies the recursion relation for $i + 1$. Now, the series

$$\tilde{\delta} + \sum_{i=0}^{\infty} q_i$$

converges with respect to the negative degree filtration $(F^i \mathcal{O})_{i \geq 0}$. We denote by Q its limit. By construction, $[Q, Q] = 0$ (since $[Q, Q]$ is a derivation that takes values in $\cap_{i \geq 0} F^i \mathcal{O} = \{0\}$), and Q has total degree +1, so that (M_0, \mathcal{O}, Q) is a \mathbb{Z}^* -graded Q -manifold. By Remark 2.14, Q satisfies both requirements in the Theorem 3.20.

Now, let us show that any two such vector fields can be intertwined by a diffeomorphism for the desired form. Let Q and Q' be two vector fields as in Theorem 3.20. We will construct a family $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \dots$ of total degree 0 and of respective negative degrees 1, 2, 3, \dots such that the sequence of degree +1 vector fields defined by the recursion relation $Q_0 = Q$ and

$$(3.15) \quad Q_{i+1} = e^{\text{ad}_{\mathbf{u}_{i+1}}} Q_i$$

(which is well-defined, see Section 1.5) satisfies that Q_i coincides with Q' in negative degrees $-1, \dots, i - 1$. Proposition 1.21 implies then that the infinite composition of the exponentials of the vector fields \mathbf{u}_i intertwines Q and Q' through a diffeomorphism Ψ which is by construction of the desired form.

Let us first construct \mathbf{u}_1 . We have:

$$\begin{aligned} Q &= \tilde{\delta} + q_0 + \sum_{i \geq 1} q_i \\ Q' &= \tilde{\delta} + q'_0 + \sum_{i \geq 1} q'_i \end{aligned}$$

where q_i, q'_i are of negative degree i . Now, since both q_0, q'_0 are $\text{ad}_{\bar{\delta}}$ -cocycles, so is $q_0 - q'_0$. Since by construction, both q_0 and q'_0 induce the same derivation Q_+ on \mathcal{K}_+ , their difference induce the trivial derivation of \mathcal{K}_+ . This implies that $q_0 - q'_0$ is a $\text{ad}_{\bar{\delta}}$ -coboundary by Corollary 3.12, and there exists a vector field \mathbf{u}_{-1} of negative degree +1 and total degree 0 such that

$$q_0 - q'_0 = [\tilde{\delta}, \mathbf{u}_{-1}].$$

By construction,

$$Q_1 := e^{\text{ad}_{\mathbf{u}_{-1}}}(Q) = Q + \sum_{m=1}^{\infty} \frac{1}{m!} \text{ad}_{\mathbf{u}_{-1}}^m(Q)$$

is well-defined, squares to zero, and satisfies again the requirements of Theorem 3.20. Also, it coincides with Q' in negative degree -1 and 0.

Now, assume that $\mathbf{u}_1, \dots, \mathbf{u}_i$ are constructed. Consider the decompositions according to negative degrees:

$$\begin{aligned} Q' &= \tilde{\delta} + q'_0 + \dots + q'_i + q'_{i+1} + \dots \\ Q_i &= \tilde{\delta} + q'_0 + \dots + q'_i + q_{i+1} + \dots \end{aligned}$$

It follows from $[Q_i, Q_i] = 0$ and $[Q', Q'] = 0$ that

$$\text{ad}_{\tilde{\delta}} q_{i+1} = - \sum_{k=0}^{i+1} [q_k, q_{i+1-k}] \text{ and } \text{ad}_{\tilde{\delta}} q'_{i+1} = - \sum_{k=0}^{i+1} [q_k, q_{i+1-k}]$$

The difference $q_{i+1} - q'_{i+1}$ is therefore an $\text{ad}_{\tilde{\delta}}$ -cocycle. Since by Corollary 3.12, the cohomology is zero in degree $i+1$, there exists a vector field \mathbf{u}_{i+1} (of negative degree $i+1$ and total degree 0) such that $q'_{i+1} = q_{i+1} + \text{ad}_{\tilde{\delta}} \mathbf{u}_{i+1}$. The vector field Q_{i+1} defined as in (3.15) satisfies the recursion relation for $i+1$.

By Proposition 1.21, the infinite ordered product of automorphisms $\Psi = \lim_{k \rightarrow \infty} e^{\mathbf{u}^{-k}} \circ \dots \circ e^{\mathbf{u}^{-1}}$ exists and induces a diffeomorphism Ψ of the graded manifold M . Furthermore, one has $\Psi Q \Psi^{-1} = \lim_{k \rightarrow \infty} e^{\text{ad}_{\mathbf{u}^{-k}}} \circ \dots \circ e^{\text{ad}_{\mathbf{u}^{-1}}}(Q)$ and

$$\Psi Q \Psi^{-1} - Q' \in \bigcap_{j \geq 0} F^j \mathcal{O}(M) = \{0\},$$

therefore $\Psi Q \Psi^{-1} - Q' = 0$.

It is also clear that, since the total degree of each \mathbf{u}_i is zero, $\text{deg}_+(\mathbf{u}_{-i}) = \text{deg}_-(\mathbf{u}_{-i}) = i$ for each $i \geq 1$, so that the positive degree of $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \dots$ is $1, 2, 3, \dots$ respectively. This implies that $\Psi(F) - F \in \mathcal{I}_+$ for every $F \in \mathcal{O}$, and therefore that Ψ induces the identity on the negative part $\mathcal{O}_- = \mathcal{O}/\mathcal{I}_+$. These degree relations also imply that $\Psi(F) - F \in \mathcal{I}_-$, so that Ψ induces the identity of $S(\oplus_{i \geq 1} \Gamma(E_{-i}^*))$, and therefore of its quotient \mathcal{K}_+ . ■

Here is an immediate consequence of the second part of Theorem 3.20.

Corollary 3.21. *Any two \mathbb{Z}^* -graded manifolds (M_0, \mathcal{O}, Q) and (M_0, \mathcal{O}, Q') over the same graded manifold (M_0, \mathcal{O}) , whose negative parts coincide and are Koszul-Tate resolutions, and whose negatively graded NQ -varieties coincide, are diffeomorphic through a diffeomorphism as in Theorem 3.20.*

4. LOCAL STRUCTURES NEAR POINTS WHERE THE CURVATURE VANISHES

4.1. **Q -structure in local coordinates.** In the following, we consider local coordinates of a graded manifold (M_0, \mathcal{O}) of the form

$$(y_1, \dots, y_r, (x_i)_{i \in I}, \theta_1, \dots, \theta_r, (\eta_j)_{j \in J}),$$

where Latin letters x_i, y_k will be used for degree 0 variables, the Greek letters θ_k (resp. η_j) will be used for variables of degree +1 (resp. of degree different from 0). Last, we will also assume that the variables y_k and θ_k go “in pairs” and that there is the same number r of them. Last, an expression like

$$R \left(x, \eta, \frac{\partial}{\partial x_{\bullet}}, \frac{\partial}{\partial \eta_{\bullet}} \right)$$

stand for any local vector field of the form:

$$\sum_{i \in I} A_i(x, \eta) \frac{\partial}{\partial x_i} + \sum_{j \in J} B_j(x, \eta) \frac{\partial}{\partial \eta_j}$$

where $A_i(x, \eta), B_j(x, \eta)$ are functions that depend on the variables $(x_i)_{i \in I}, (\eta_j)_{j \in J}$ only. For any Q -manifold (M_0, \mathcal{O}, Q) , equipped with a splitting

$$\Phi: \mathcal{O} \simeq \Gamma \left(\hat{S} \oplus_{i \in \mathbb{Z}^*} E_i^* \right),$$

the anchor map $\rho: E_{-1} \rightarrow TM_0$ is the vector bundle morphism defined by:

$$\langle Q[f]^{(1)}, u \rangle = \rho(u)[f],$$

for every $u \in \Gamma(E_{-1})$ and $f \in C^\infty(M_0)$. Above, $Q[f]^{(1)}$ stands for the component of polynomial degree 1 of $Q[f]$: since $Q[f]$ is of degree 1, $Q[f]^{(1)}$ is a section of E_{-1}^* , so that the previous definition makes sense.

Remark 4.1. By construction, the anchor map of a Q -manifold is a vector bundle morphism $\rho: E_{-1} \rightarrow TM_0$ that depends on the choice of the splitting, although the vector bundles E_{-1} and TM_0 do not. For instance, in a splitting as in Proposition 2.3 for which $Q = \mathbf{i}_\kappa$, the anchor map is the zero map. But it may be non-zero in some other splitting. However, at every m that belongs to the zero locus of the curvature $\kappa \in \Gamma(E_{+1})$, the anchor map $\rho: E_{-1} \rightarrow M$ does not depend on the choice of a splitting, and is therefore canonical. \square

Remark 4.1 implies that the following theorem only makes sense when the point m is the zero locus of the curvature κ .

Theorem 4.2. *Let (M_0, \mathcal{O}, Q) be a Q -manifold. Let $\rho: E_{-1} \rightarrow TM_0$ be the anchor map corresponding to some splitting. Every point $m \in M_0$ on the zero locus of the curvature κ admits a coordinate neighborhood with variables $(y_1, \dots, y_r, (x_i)_{i \in I}, \theta_1, \dots, \theta_r, (\eta_j)_{j \in J})$ on which Q reads:*

$$Q = \sum_{k=1}^r \theta_k \frac{\partial}{\partial y_k} + R \left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet} \right)$$

where r is the rank of the anchor map $\rho: E_{-1} \rightarrow TM_0$ at m .

We start with a remark and two lemmas, before proving a proposition crucial for the proof of the above theorem.

Remark 4.3. Any degree 0 vector field \mathbf{v} on a \mathbb{Z}^* -graded manifold (M_0, \mathcal{O}) induces a vector field $\underline{\mathbf{v}}$ on M_0 : A degree 0 vector field being, by definition, a degree 0 derivation of \mathcal{O} , it preserves both negative and positive functions, so it preserves the maximal ideal \mathcal{I} , and induces a derivation of the quotient \mathcal{O}/\mathcal{I} , which is isomorphic to $C^\infty(M_0)$. This induced derivation is a vector field on M_0 . In coordinates, this assignment reads:

$$\begin{aligned} \mathfrak{X}(\mathcal{O})_0 &\rightarrow \mathfrak{X}(M_0) \\ \sum_i f_i(z, \zeta) \frac{\partial}{\partial z_i} + \sum_j g_j(z, \zeta) \frac{\partial}{\partial \zeta_j} &\mapsto \sum_i f_i(z, 0) \frac{\partial}{\partial z_i} \end{aligned}$$

where (z, ζ) are local coordinates of degree 0 and different from 0 respectively. \square

Lemma 4.4 extends to graded manifolds the well-known straightening theorem, also known as Hadamard Lemma.

Lemma 4.4. *Let \mathbf{v} be a vector field of degree 0 on a graded manifold (M_0, \mathcal{O}) with M_0 . Every point of the base manifold M_0 where the induced vector field $\underline{\mathbf{v}}$ is different from zero admits a coordinate neighborhood $(y, (x_i)_{i \in I}, (\eta_j)_{j \in J})$ on which $\mathbf{v} = \frac{\partial}{\partial y}$.*

Proof. The proof is rather straightforward: use the general form of the coordinate changes on graded manifolds (cf. [9]). ■

The following lemma is the result of an obvious computation.

Lemma 4.5. *Every vector field Q , defined on a coordinate neighborhood $(y, x_\bullet, \eta_\bullet)$, that satisfies $\left[Q, \frac{\partial}{\partial y}\right] = 0$ is of the form:*

$$Q = \tau(x_\bullet, \eta_\bullet) \frac{\partial}{\partial y} + R \left(x_\bullet, \eta_\bullet, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet} \right).$$

We can now prove the following statement:

Proposition 4.6. *Let (M_0, \mathcal{O}, Q) be a Q -manifold equipped with a splitting. Let $\rho: E_{-1} \rightarrow TM_0$ be the corresponding anchor map. Every point $m \in M_0$ in the zero locus of the curvature κ such that $\rho_m: E_{-1} \rightarrow TM_0$ is not the zero map admits a coordinate neighborhood with variables $((y, x_\bullet), (\theta, \eta_\bullet))$ on which Q reads:*

$$Q = \theta \frac{\partial}{\partial y} + R \left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet} \right).$$

Proof. The map $\rho: E_{-1} \rightarrow TM_0$ is different from zero at $m \in M_0$ if and only if there exists a section e in $\Gamma(E_{-1})$ such that the degree 0 vector field $\mathbf{v} := [Q, \mathbf{i}_e]$ (which is of degree 0) has a basic vector field $\underline{\mathbf{v}}$ (see Remark 4.3) different from 0 at m . By Lemma 4.4, there exists a coordinate neighborhood $(y, x_\bullet, \eta_\bullet)$ on which $\mathbf{v} = [Q, \mathbf{i}_e] = \frac{\partial}{\partial y}$. Since $[\mathbf{v}, Q] = 0$, Lemma 4.5 implies that in these coordinates:

$$Q = \tau(x, \eta) \frac{\partial}{\partial y} + R \left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet} \right).$$

Now, $\tau(x, \eta) = Q(y)$ is a degree +1 function whose component in $\Gamma((E_{-1})^*)$ cannot be zero in view of

$$\mathbf{i}_e \tau(x, \eta) = \mathbf{i}_e Q(y) = [\mathbf{i}_e, Q](y) + Q(\mathbf{i}_e[y]) = \frac{\partial}{\partial y}(y) + Q(\mathbf{i}_e[y]) = 1 + Q(\mathbf{i}_e[y])$$

and the fact that the projection of the degree 0 function $Q(\mathbf{i}_e[y])$ on $C^\infty(M_0)$ has to be an element of the zero locus ideal for degree reasons. We can therefore replace one of the degree -1 variables in the coordinates η_\bullet by $\tau(x, \eta)$: we denote by θ this new variable. Since $\theta = \tau(x, \eta)$ does not depend on the variable y , this change of coordinates does not affect $\theta \frac{\partial}{\partial y}$ and changes R in a vector field that again does not depend on y nor contains $\frac{\partial}{\partial y}$. But it may contain a component in $\frac{\partial}{\partial \theta}$. In conclusion:

$$Q = \theta \frac{\partial}{\partial y} + \tilde{R} \left(x, \eta, \theta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet} \right) + S(x_\bullet, \eta) \frac{\partial}{\partial \theta}.$$

Since $Q^2(y) = Q(\theta) = 0$, we have $S(x_\bullet, \eta) = 0$ and therefore:

$$Q = \theta \frac{\partial}{\partial y} + \tilde{R} \left(x, \eta, \theta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet} \right)$$

Since $\theta^2 = 0$, we have:

$$\tilde{R}\left(x, \eta, \theta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet}\right) = A\left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet}\right) + \theta B\left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet}\right)$$

so that

$$Q = \theta\left(\frac{\partial}{\partial y} + B\left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet}\right)\right) + A\left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet}\right)$$

There exists local coordinates (y', x'_\bullet, η') leaving θ untouched, where

$$\frac{\partial}{\partial y} + B\left(x, \eta, \frac{\partial}{\partial x_\bullet}, \frac{\partial}{\partial \eta_\bullet}\right) = \frac{\partial}{\partial y'}$$

We have in these coordinates:

$$Q = \theta \frac{\partial}{\partial y'} + A'\left(x', \eta', y', \frac{\partial}{\partial x'_\bullet}, \frac{\partial}{\partial \eta'_\bullet}, \frac{\partial}{\partial y'}\right).$$

Since $Q^2 = 0$, A' does not depend on y' , and:

$$Q = \theta \frac{\partial}{\partial y'} + A''\left(x', \eta', \frac{\partial}{\partial x'_\bullet}, \frac{\partial}{\partial \eta'_\bullet}\right) + T(x', \eta') \frac{\partial}{\partial y'}.$$

We now replace θ by $\theta' = \theta + T(x', \eta')$. Since $(\theta + T(x', \eta')) = Q(y') = \theta'$, we have $A''\left(x', \eta', \frac{\partial}{\partial x'_\bullet}, \frac{\partial}{\partial \eta'_\bullet}, \frac{\partial}{\partial \theta'}\right)\theta' = 0$, so that A'' has no component in $\frac{\partial}{\partial \theta'}$ and the vector field Q has the desired form in these coordinates. This completes the proof. ■

Proof. [Proof of Theorem 4.2] The theorem is now an immediate consequence of Proposition 4.6, upon making a finite recursion until the corresponding anchor map vanishes. ■

4.2. Examples and non-examples.

Example 4.7. Theorem 4.2, when applied to a Lie algebroids, gives back a classical result [5], which itself is similar to Weinstein splitting theorem for Poisson manifolds [19]. For Lie ∞ -algebroids, Theorem 4.2 gives back a similar statement in [4].

Example 4.8. For a Koszul-Tate resolution, Theorem 4.2 does not give any interesting result, since the anchor is zero at every point of the zero locus.

Example 4.9. For a positively graded Q -manifold over a manifold M_0 , the image of the anchor map

$$\rho: \Gamma(E_{-1}) \longrightarrow \mathfrak{X}(M_0)$$

is a singular foliation in the sense of [1], i.e. a locally finitely generated $C^\infty(M_0)$ -sub-module of $\mathfrak{X}(M_0)$ closed under Lie bracket. For \mathbb{Z}^* -graded Q -manifold with splitting, whose dual Lie ∞ -algebroid with anchor maps $(\rho_n)_{n \geq 1}$, it is natural to ask if

$$\bigoplus_{n \geq 1} \rho_n(\Gamma(S^n \oplus_{i \in \mathbb{Z}} E_i)_{-1})$$

is still a singular foliation. The answer is *no*: it is certainly a $C^\infty(M_0)$ -sub-module of $\mathfrak{X}(M_0)$, but, even when it is locally finitely generated, it may not be stable under Lie bracket. Here is a class of counter-examples: Let M_0 be a manifold, X_1, X_2 vector fields

such that $[X_1, X_2]$ is not in the C^∞ -module generated by X_1, X_2 , let θ_1, θ_2, η be additional variables of respective degrees 2, 2 and -1 , and consider

$$Q = \eta\theta_1 X_1 + \eta\theta_2 X_2.$$

It is straightforward to check that Q is a degree $+1$ vector field squaring to zero. The 2-ary anchor map is not zero and its image is the $C^\infty(M_0)$ module generated by X_1, X_2 , which generated a $C^\infty(M_0)$ -module; by assumption it is not stable under Lie bracket.

Example 4.10. Here is an example of a Q -manifold with a splitting, whose 2-ary anchor is not valued in vector fields tangent to the zero locus:

$$Q = (x - \epsilon\zeta) \frac{\partial}{\partial \eta} + \zeta\xi \frac{\partial}{\partial x} + \xi \frac{\partial}{\partial \epsilon},$$

where x is a degree 0 variables and $\eta, \zeta, \xi, \epsilon$ are variables of respective degrees $-1, 3, -2, -3$.

CONCLUSION / PERSPECTIVES

As mentioned in the introduction, the results of [9] on the \mathbb{Z} -graded manifolds and the technique of filtrations of functional spaces open a way to understanding the form of various geometric and algebraic structures on them. This permitted for example, to extend the results of [8] to the honest \mathbb{Z} -graded case and develop them in [10]. In the current paper we have added an important ingredient to the picture – a Q -structure – describing thus the normal form of differential \mathbb{Z}^* -graded manifolds. Our common thread is that ”for a \mathbb{Z}^* -graded Q -manifold, only the zero locus of the curvature matters”: Proposition 2.3 should be understood as meaning that outside the zero locus of their curvatures, \mathbb{Z}^* -graded Q -manifolds have a very trivial structure; then Theorem 3.20 makes more precise this general idea, by stating that positive part of a Q -manifold over its zero locus is the only piece that matters when its negative part is a Koszul-Tate resolution; and last, Theorem 4.2 adds an other layer to the same general idea, by stating that, at a point in the zero locus, the anchor map and its transverse Q -manifold are the only two non-trivial pieces of information.

On top of the pure mathematical significance of the above results we expect them to have straightforward consequences for gauge theories. According to [6], under rather natural assumptions one can read-off a Q -structure from the equations governing the theory. This language is also widely used for various quantization problems. Then, as explained in [11], a lot of information can be encoded in the language of mappings between Q -manifolds: the equations of motion (i.e. extrema of the functional describing the model) correspond to Q -morphisms, and gauge transformations (symmetries) to Q -homotopies. In this setting reducing a Q -structure to a (simple) canonical form by a homotopy would mean gauge fixing in an intelligent way.

APPENDIX A. PROJECTIVE SYSTEMS OF ALGEBRAS

We call *projective system of algebras* a pair made of a sequence $(A^i)_{i \in \mathbb{N}}$ of algebras, and a family of algebra morphisms $\pi^{[i \rightarrow j]}: A^i \rightarrow A^j$, defined for all integers $i \geq j$, subject to the two following conditions: $\pi^{[i \rightarrow i]} = \text{id}_{A^i}$ and

$$\pi^{[j \rightarrow k]} \circ \pi^{[i \rightarrow j]} = \pi^{[i \rightarrow k]}, \quad \forall i \geq j \geq k.$$

A *endomorphism of projective algebras* is a family $(\phi^{[i]})_{i \in \mathbb{N}}$ of algebra endomorphisms $\phi^{[i]} : A^i \rightarrow A^i$, defined for all $i \in \mathbb{N}$ such that $\phi^{[j]} \circ \pi^{[i \rightarrow j]} = \pi^{[i \rightarrow j]} \circ \phi^{[i]}$ for all $i \geq j$. The following diagram recapitulates the above commutativity properties for all $i \geq j \geq k$:

$$\begin{array}{ccccc}
 A^i & & \xrightarrow{\pi^{[i \rightarrow k]}} & & A^k \\
 \uparrow \phi^{[i]} & \searrow \pi^{[i \rightarrow j]} & & \nearrow \pi^{[j \rightarrow k]} & \uparrow \phi^{[k]} \\
 & & A^j & & \\
 & & \uparrow \phi^{[j]} & & \\
 A^i & & \xrightarrow{\pi^{[i \rightarrow k]}} & & A^k \\
 \searrow \pi^{[i \rightarrow j]} & & & & \nearrow \pi^{[j \rightarrow k]} \\
 & & A^j & &
 \end{array}$$

We define the *projective limit* A^∞ of a projective system of algebras to be the algebra of collections $i \mapsto a^i \in A^i$ such that $\pi^{[i \rightarrow j]}(a^i) = a^j$ for all $i \geq j$. By assigning to such a collection its i -th component, one defines, for all $i \in \mathbb{N}$, algebra morphisms $\pi^{[\infty \rightarrow i]} : A^\infty \rightarrow A^i$ that satisfy:

$$\pi^{[j \rightarrow k]} \circ \pi^{[\infty \rightarrow j]} = \pi^{[\infty \rightarrow k]}, \quad \forall j \geq k.$$

For any morphism of projective algebras $(\phi^{[i]})_{i \in \mathbb{N}}$, there exists a unique algebra endomorphism $\phi^{[\infty]} : A^\infty \rightarrow A^\infty$ such that $\phi^{[i]} \circ \pi^{[\infty \rightarrow i]} = \pi^{[\infty \rightarrow i]} \circ \phi^{[\infty]}$.

$$\begin{array}{ccc}
 A^\infty & \xrightarrow{\pi^{[\infty \rightarrow i]}} & A^i \\
 \uparrow \phi^{[\infty]} & & \uparrow \phi^{[i]} \\
 A^\infty & \xrightarrow{\pi^{[\infty \rightarrow i]}} & A^i
 \end{array}$$

We call $\phi^{[\infty]} : A^\infty \rightarrow A^\infty$ the *projective limit of $(\phi^{[i]})_{i \in \mathbb{N}}$* .

Proposition A.1. *Let $(A^i, \pi^{[i \rightarrow j]})$ be a projective system of algebras. For any family $(\phi_N)_{N \in \mathbb{N}}$ of endomorphisms of the latter such that $\phi_N^{[i]} = \text{id}_{A^i}$ for all $N \geq i$, the sequence of algebra endomorphisms defined for all $i \in \mathbb{N}$ by*

$$\begin{aligned}
 \psi^{[i]} : A^i &\rightarrow A^i \\
 a &\mapsto \cdots \circ \phi_3^{[i]} \circ \phi_2^{[i]} \circ \phi_1^{[i]}(a) \\
 &= \phi_i^{[i]} \circ \cdots \circ \phi_1^{[i]}(a) \quad (\text{by assumption})
 \end{aligned}$$

is an endomorphism of projective systems of algebras.

The projective limit $\psi^{[\infty]} : A^\infty \rightarrow A^\infty$ must be understood as the infinite composition of all the $(\phi_i)_{i \in \mathbb{N}}$, it will therefore be denoted by $\bigcirc_{i \uparrow \in \mathbb{N}} \phi_i$ or $\prod_{i \uparrow \in \mathbb{N}} \phi_i$, where by “ $i \uparrow \in \mathbb{N}$ ” we mean the ordered index i .

Acknowledgments. We are thankful to the “Research in Paris” program of the Institut Henri Poincaré, which hosted us in the beginning of our work on this paper. The work was also partially supported by the CNRS MITI Project “GraNum” and PHC Procope “GraNum 2.0”. A.K. also appreciates the support of the Faculty of Science of the University of Hradec Králové.

REFERENCES

1. Iakovos Androulidakis and Georges Skandalis, *The holonomy groupoid of a singular foliation*, J. Reine Angew. Math. **626** (2009), 1–37. MR 2492988
2. G. Bonavolontà and A. Kotov, *On the space of super maps between smooth supermanifolds*, arXiv:1304.0394 [math.DG] (2013), 16.
3. Giuseppe Bonavolontà and Norbert Poncin, *On the category of Lie n -algebroids*, J. Geom. Phys. **73** (2013), 70–90. MR 3090103
4. Henrique Bursztyn, Hudson Lima, and Eckhard Meinrenken, *Splitting theorems for Poisson and related structures*, J. Reine Angew. Math. **754** (2019), 281–312. MR 4000576
5. Jean-Paul Dufour, *Normal forms for Lie algebroids*, Lie algebroids and related topics in differential geometry (Warsaw, 2000), Banach Center Publ., vol. 54, Polish Acad. Sci. Inst. Math., Warsaw, 2001, pp. 35–41. MR 1881647
6. Melchior Grützmann and Thomas Strobl, *General Yang-Mills type gauge theories for p -form gauge fields: from physics-based ideas to a mathematical framework or from Bianchi identities to twisted Courant algebroids*, Int. J. Geom. Methods Mod. Phys. **12** (2015), no. 1, 1550009, 80.
7. Marc Henneaux and Claudio Teitelboim, *Quantization of gauge systems*, Princeton University Press, Princeton, NJ, 1992.
8. Benoit Jubin, Alexei Kotov, Norbert Poncin, and Vladimir Salnikov, *Differential graded Lie groups and their differential graded Lie algebras*, Transformation Groups **27** (2022), 497–523.
9. Alexei Kotov and Vladimir Salnikov, *The category of Z -graded manifolds: what happens if you do not stay positive*, arXiv:2108.13496 (2021), 26.
10. ———, *Various instances of Harish-Chandra pairs*, arXiv:2207.07083 (2022), 22.
11. Alexei Kotov and Thomas Strobl, *Characteristic classes associated to Q -bundles*, Int. J. Geom. Methods Mod. Phys. **12** (2015), no. 1, 1550006, 26.
12. Camille Laurent-Gengoux, Sylvain Lavau, and Thomas Strobl, *The universal Lie ∞ -algebroid of a singular foliation*, Doc. Math. **25** (2020), 1571–1652.
13. Camille Laurent-Gengoux, Mathieu Stiénon, and Ping Xu, *Exponential map and L_∞ algebra associated to a Lie pair*, C. R. Math. Acad. Sci. Paris **350** (2012), no. 17-18, 817–821.
14. ———, *Poincaré-Birkhoff-Witt isomorphisms and Kapranov dg -manifolds*, Adv. Math. **387** (2021), Paper No. 107792, 62.
15. Dimitri Leites, *Theory of Supermanifolds*, (rus), Moscow, 1983.
16. Leonid Ryvkin, *Observables and Symmetries of n -Plectic Manifolds*, Springer, 2016.
17. Th. Th. Voronov, *Q -manifolds and higher analogs of Lie algebroids*, XXIX Workshop on Geometric Methods in Physics, AIP Conf. Proc., vol. 1307, Amer. Inst. Phys., Melville, NY, 2010, pp. 191–202.
18. Pavol Ševera, *Some title containing the words “homotopy” and “symplectic”, e.g. this one*, Travaux mathématiques. Fasc. XVI, Trav. Math., vol. 16, Univ. Luxemb., Luxembourg, 2005, pp. 121–137.
19. Alan Weinstein, *The local structure of Poisson manifolds*, J. Differential Geom. **18** (1983), no. 3, 523–557. MR 723816

ALEXEI KOTOV: FACULTY OF SCIENCE, UNIVERSITY OF HRADEC KRALOVE, ROKITANSKEHO 62, HRADEC KRALOVE 50003, CZECH REPUBLIC
E-mail address: `oleksii.kotov@uhk.cz`

CAMILLE LAURENT-GENGOUX: INSTITUT ELIE CARTAN DE LORRAINE (IECL), UMR 7502 – 3 RUE AUGUSTIN FRESNEL, 57000 TECHNOPÔLE METZ, FRANCE
E-mail address: `camille.laurent-gengoux@univ-lorraine.fr`

VLADIMIR SALNIKOV: LASIE – CNRS & LA ROCHELLE UNIVERSITY, AV. MICHEL CRÉPEAU, 17042 LA ROCHELLE CEDEX 1, FRANCE
E-mail address: `vladimir.salnikov@univ-lr.fr`