



Interest of type I collagen as an additive in an alginate-based bioink for the production of cartilaginous substitutes by 3D printing

Océane Messaoudi, Christel Henrionnet, Laurent Grossin, Pierre Gillet, Didier Mainard, Laurent Galois, Damien Loeuille, Christophe Marquette, Astrid Pinzano

► To cite this version:

Océane Messaoudi, Christel Henrionnet, Laurent Grossin, Pierre Gillet, Didier Mainard, et al.. Interest of type I collagen as an additive in an alginate-based bioink for the production of cartilaginous substitutes by 3D printing. Journée scientifique de l'école doctorale BioSE, May 2022, Vandoeuvre-lès-Nancy, France. hal-03908036

HAL Id: hal-03908036

<https://hal.science/hal-03908036v1>

Submitted on 20 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Interest of type I collagen as an additive in an alginate-based bioink for the production of cartilaginous substitutes by 3D printing

Océane MESSAOUDI*, Christel HENRIONNET, Laurent GROSSIN, Pierre GILLET, Didier MAINARD, Laurent GALOIS, Damien LOEUILLE, Christophe MARQUETTE and Astrid PINZANO

Introduction

Hyaline cartilage




Stratified
Non vascularized
Non innervated
→ Limited self-repair properties

Trauma or Overload




Focal lesions :
Chondral
Osteochondral

Objectives



Methods

Cellularized bioink



Type I
collagen
0, 0.5, 1 or 5 mg

Maturation : 28 or 56 days

Conditions :
ITS 1%: Control
TGF-β1: Chondro-inductor

Analyses :


Cellular viability : LDH assay

Genes expressions : RT-qPCR


Matrix synthesis : Histological stainings

Results

1. Viability assay (LDH)




2. Gene expressions of (a) chondrogenic, (b) hypertrophic and (c) osteogenic markers (Day 28)



Mean ± SEM. Effect of TGF-β1: * p < 0.05, ** p < 0.01, and *** p < 0.001. Effect of the bioink composition : § p < 0.05, §§ p < 0.01, and §§§ p < 0.001. ; ns: not significant.

3. Matrix synthesis analyses: (i) Alcian blue and (ii) Alizarin red (Day 56)



Conclusions

Low type I collagen addition (0.5mg) :

- Proteoglycans
- Type II collagen
- Calcifications


→ Rich matrix adapted to recreate the superficial layers of the cartilage

High type I collagen addition (5mg) :

- Proteoglycans
- Type II collagen
- Calcifications

→ Calcified matrix mimicking the calcified layer of the cartilage

Perspectives



Producing a stratified cartilaginous construct