



HAL
open science

Palynological and sedimentological records since 8.5 ka BP on the southern Brittany platform (NW Europe): complex responses to sealevel, rapid climate and anthropogenic changes

Ophélie David, Aurélie Penaud, Muriel Vidal, Wiem Fersi, Clément Lambert, Evelyne Goubert, Maiwenn Herledan, Pierre Stéphan, Yvan Pailler, Jean-François Bourillet, et al.

► To cite this version:

Ophélie David, Aurélie Penaud, Muriel Vidal, Wiem Fersi, Clément Lambert, et al.. Palynological and sedimentological records since 8.5 ka BP on the southern Brittany platform (NW Europe): complex responses to sealevel, rapid climate and anthropogenic changes. EGU General Assembly 2022, May 2022, Vienne, Austria. hal-03908047

HAL Id: hal-03908047

<https://hal.science/hal-03908047>

Submitted on 20 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU22-1003

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Palynological and sedimentological records since 8.5 ka BP on the southern Brittany platform (NW Europe): complex responses to sea-level, rapid climate and anthropogenic changes

Ophélie David^{1,2}, Aurélie Penaud¹, Muriel Vidal¹, Wiem Fersi¹, Clément Lambert², Evelyne Goubert², Maiwenn Herledan³, Pierre Stéphan⁴, Yvan Pailler⁵, Jean-François Bourillet⁶, and Agnes Baltzer⁷

¹Laboratoire Géo-Océan, Université de Bretagne Occidentale, IUEM, Plouzané, France

²Laboratoire Géo-Océan, Université de Bretagne Sud, Vannes, France

³Laboratoire d'Océanologie et de Géosciences, Université de Lille, Villeneuve d'Ascq, France

⁴Littoral Environnement Télédétection Géomatique, IUEM, Plouzané, France

⁵Chaire Armerie, Inrap, Littoral Environnement Télédétection Géomatique, Université de Bretagne Occidentale, IUEM, Plouzané, France

⁶Laboratoire Géo-Océan, Ifremer, Plouzané, France

⁷Littoral Environnement Télédétection Géomatique, Université de Nantes, Nantes, France

New results acquired in the south-Brittany shelf (core MD08-3204 CQ: Bay of Quiberon and core VK03-58bis: south-Glénan) allow depicting Holocene paleoenvironmental changes from 8.5 ka BP to present through a multi-proxy dataset including sedimentological and palynological data.

First, grain-size analyses and AMS-¹⁴C dates depict a common sedimentary history for both study cores. After the post-glacial sea-level (RSL) rise and related high sedimentation rates, the parallel slowdown of the RSL rise and the drop of sedimentation rates occurred between 8.3 and 5.7 ka BP. This interval leads to the establishment of a shell-condensed level, identified in the VK03-58bis core by the "*Turritella layer*" and interpreted as a marker for the establishment of the maximum flooding surface. Palynological data (pollen grains and dinocyst assemblages) acquired in the core MD08-3204 CQ argue for an amplification of the fluvial influence since 5.9 ka BP; the establishment of the highstand system tract (i.e. estuarine-type sedimentation on the platform) then accompanying the slowdown of the RSL rise. On the shelf, the Anthropogenic Pollen Indicators (API) amplification, is detected since 4.2 ka BP, due to the fluvial influence becoming predominant in the context of the Late Holocene.

In addition, the comparison of fluvial palynological tracers, including API, over the last 7 kyrs, with coastal-marines sites subjected to northern vs. southern Loire catchment areas, allowed to discuss a major hydro-climatic effect on the reconstructed palynological signals. Strengthened subpolar gyre dynamics (SPG), combined with recurrent positive North Atlantic Oscillation (NAO) configurations, are well-known to favour increased winter precipitation and fluvial discharge in northern Europe, such as Brittany, and conversely during weakened SPG the winter fluvial discharge is intensified over southern Europe. Interestingly, we record, at an infra-orbital

timescale, major peaks of API during periods of strengthened (/weakened) SPG dynamics in sites whose catchment areas are located north (/south) of the Loire.