



**HAL**  
open science

## Past hydraulics influence microhabitat selection by invertebrates and fish in hydropeaking rivers

Clarisse Judes, Hervé Capra, Véronique Gouraud, Hervé Pella, Nicolas Lamouroux

► **To cite this version:**

Clarisse Judes, Hervé Capra, Véronique Gouraud, Hervé Pella, Nicolas Lamouroux. Past hydraulics influence microhabitat selection by invertebrates and fish in hydropeaking rivers. *River Research and Applications*, 2023, 39, pp.375-388. 10.1002/rra.3981 . hal-03910907

**HAL Id: hal-03910907**

**<https://hal.science/hal-03910907>**

Submitted on 12 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 Past hydraulics influence microhabitat 2 selection by invertebrates and fish in 3 hydropeaking rivers

4 Clarisse Judes<sup>1,2,3\*</sup>, Hervé Capra<sup>2</sup>, Véronique Gouraud<sup>1,3</sup>, Hervé Pella<sup>2</sup>, Nicolas Lamouroux<sup>2</sup>

5 1: EDF R&D LNHE - Laboratoire National d'Hydraulique et Environnement, Chatou, France.

6 2: INRAE RiverLy, Villeurbanne, France.

7 3: HYNES team (INRAE-EDF E&D), Chatou, France.

8 \*Corresponding author, [clarisse.judes@gmail.com](mailto:clarisse.judes@gmail.com).

9

## 10 Abstract

11 Hydropeaking hydropower plants are the major source of renewable energy meeting sub-daily  
12 peaks in electricity demand. They induce rapid artificial flow variations, highly variable  
13 velocities, drift and stranding risks for aquatic organisms. In hydropeaking reaches,  
14 microhabitat selection likely depends on both present and past hydraulics (flow velocity and  
15 water depth); this study aims to assess their relative impact. For this purpose, we used  
16 observations of fish abundance in 1,180 microhabitats (507 sampled by electrofishing, 673 by  
17 snorkeling) and of invertebrate abundance in 36 microhabitats (hyporheic and benthic) in a  
18 medium-sized hydropeaking river. We described past hydraulics of microhabitats over the 15  
19 days preceding sampling, using a 2D hydrodynamic model, by identifying microhabitats  
20 dewatering (drying during > 10h) or with high-velocity conditions (>1.3 m s<sup>-1</sup> during > 10h).  
21 Invertebrates guilds (defined based on their selection of present hydraulics in rivers without  
22 hydropeaking) responded significantly to past hydraulics, with abundances 3.5-15.3 times  
23 lower in dewatering habitats. Selection for present hydraulics by invertebrates was different  
24 from that observed in rivers without hydropeaking. For more mobile fish, responses were  
25 weaker and different, with a "bank" guild selecting dewatering microhabitats and, secondarily,  
26 a "midstream" guild avoiding them. Selection of present hydraulics by fish was similar to that  
27 observed in rivers without hydropeaking. Overall, past hydraulics influenced microhabitat

28 selection, with stronger effects on invertebrates and stronger effects of dewatering than of  
29 high past velocities. However, high past velocities force fish to move and invertebrates to  
30 experience a large range of velocity.

## 31 **Keywords**

32 Habitat preferences, Flow variations, 2D hydrodynamic model, Behavior, Dewatering  
33 responses, Hydropower

## 34 **Introduction**

35 Hydropeaking hydropower plants are the major source of renewable energy that meets  
36 sub-daily peaks in electricity demand. By producing electricity on demand, they create frequent  
37 rapid flow variations known as hydropeaking. Hydropeaking influences the habitats of aquatic  
38 organisms, with strong spatial and temporal variations in point flow velocity, water depth and  
39 shear stress (hydraulic conditions). In particular, near-shore areas may be subject to  
40 dewatering during base flow (i.e., become temporarily dry when turbines are shut down),  
41 which may result in stranding of individuals using shallow and slow-flowing habitats (Halleraker  
42 et al., 2003; Saltveit et al., 2001). Midstream habitats may be subject to high flow velocity  
43 during peak flow (when turbines are on), which may cause forced drift of individuals with low  
44 swimming ability (Bruno et al., 2013).

45 Hydropeaking often results in a decrease in abundance, biomass and species diversity in  
46 both fish and invertebrate communities (Kjaerstad et al., 2018; Schmutz et al., 2015; Vila-  
47 Martínez et al., 2019). However, these negative effects are not systematic and may depend on  
48 hydropeaking hydraulic characteristics. For example, Judes et al. (2021) showed, in rivers with  
49 moderate hydraulic variations, that hydropeaking could have weaker effects than floods on fish  
50 community dynamics and did not greatly modify the organization of fish communities along  
51 longitudinal gradients. To improve our mechanistic understanding of hydropeaking influences  
52 on biota, we need to better understand individual responses to changes in the spatial and  
53 temporal variations of point hydraulic conditions (i.e. microhabitat scale). The microhabitat  
54 scale (here corresponding to fixed patches of  $\sim 7\text{m}^2$  for fish,  $0.05\text{ m}^2$  for benthic invertebrates

55 and 6 L of hyporheic water for hyporheic invertebrates following Stubbington et al., 2016) is  
56 particularly suited for identifying the key hydraulic drivers of biological response to  
57 hydropeaking. Several studies quantified key hydraulic conditions in microhabitats for fish and  
58 invertebrates drifting and stranding processes (Auer et al., 2017; Halleraker et al., 2003; Saltveit  
59 et al., 2001). For example, Halleraker et al., (2003) showed that stranding of juvenile brown  
60 trout (*Salmo trutta*) occurs with dewatering vertical water level gradients higher than 0.10 cm  
61 h<sup>-1</sup>. By contrast, little is known concerning how microhabitat selection of fish and invertebrates  
62 is influenced by variations in point hydraulic conditions.

63 In rivers without hydropeaking and associated sub-daily variations, microhabitat  
64 selection by fish and invertebrates is usually predicted from present hydraulic conditions and  
65 substrate size (Forcellini et al., in press.; Lamouroux et al., 1999; Plichard et al., 2020). In  
66 hydropeaking rivers, past hydraulic variability during the weeks preceding sampling is expected  
67 to influence microhabitat selection by aquatic organisms. Indeed, experiments on fish  
68 individual behavior (Green, 1971; Roy & Bhat, 2018) indicated that fish can memorize  
69 environmental conditions over durations up to two weeks. Individuals perceive temporal  
70 variations in hydraulic conditions (Patton & Braithwaite, 2015) and may avoid habitats that are  
71 frequently dewatered or have major variations in flow velocity. For example, using telemetry in  
72 the Rhône River (France) to monitor 18 individual fish, Capra et al. (2017) suggested that fish  
73 memorize spatial and temporal environmental variations during the two weeks preceding  
74 sampling so as to use the “least constraining” microhabitats. Organisms may also have strong  
75 habitat fidelity even after their habitats become unsuitable (Kemp et al., 2003). Several studies  
76 showed that some fish species are reluctant to move across the river during sudden flow  
77 increases and consequently use habitats with higher flow velocity and/or deeper habitats  
78 (Kemp et al., 2003; Pert & Erman, 1994; Shirvell, 1994). This behavior may be explained by the  
79 energy cost associated with displacement and/or a higher risk of predation. Less mobile taxa,  
80 which include many invertebrates, may be unable to respond instantaneously to a rapid shift in  
81 habitat conditions (Blinn et al., 1995). Some invertebrate taxa may also be removed from  
82 hydropeaking reaches by passive or active drift during periods of low or high flow, altering the  
83 taxonomic and functional composition of benthic assemblages in hydropeaking reaches (Bruno  
84 et al., 2013). However, in adapting to naturally variable microhabitat conditions, some taxa  
85 have developed morphological and behavioral adaptations to maximize colonization of

86 available habitat and can withstand extreme habitat conditions. For example, due to behavior  
87 or/and morphological adaptation to high current velocity Simuliidae and Hydropsychidae are  
88 less likely to passive drift due to hydropeaking and can be better able to use microhabitat  
89 undergoing variable current velocity (Bruno et al. 2010, 2016; Mochizuki et al. 2006, Sidler et  
90 al. 2018). Other invertebrate taxa may also colonize subsurface hyporheic habitats to escape  
91 from variable benthic hydraulic conditions (Dole-Olivier et al., 1997; Vadher et al., 2017).

92         Studying the effects of past hydraulics on microhabitat selection by aquatic organisms  
93 from field data is challenging. A first challenge is to describe past hydraulic conditions in  
94 microhabitats. Spatially explicit hydrodynamic models, such as two-dimensional (2D) models  
95 that map vertically averaged flow velocities at various discharge rates, are useful for this  
96 purpose. However, calibrating and testing 2D models is time-consuming, expensive, and  
97 requires an appropriate amount of quality field measurements. A second challenge is to sample  
98 organisms at several discharge rates in order to have a sufficient number of microhabitat  
99 samples with various combinations of past and present hydraulic conditions. Addressing this  
100 challenge requires extensive fieldwork.

101         In this study, we combined extensive biological sampling in microhabitats at various  
102 discharge rate with results of a calibrated 2D hydrodynamic model in a hydropeaking reach to  
103 weight the relative effects of past and present hydraulic conditions on microhabitat selection.  
104 Specifically, we recorded fish taxa abundance in 1,187 microhabitats and the abundance of  
105 benthic and hyporheic invertebrate taxa in 36 microhabitats along a 6 km reach of a  
106 hydropeaking reach of a medium-sized French river (the Ain River). Due to a limited statistical  
107 power, we studied the responses of selected taxonomic guilds that had significant and  
108 contrasting microhabitat selection in rivers without hydropeaking, according to the literature.  
109 The 2D hydrodynamic model was used to estimate past hydraulics at each microhabitat during  
110 the 15 days preceding sampling (Capra et al., 2017).

111         For fish, we expected that fish using shallow and slow-flowing habitats along the banks  
112 would be forced to use dewatering habitats at high flows. We expected “midstream” fish taxa,  
113 using fast-flowing and deep habitats, to be less forced to use risky dewatering areas at high  
114 flows because they should sustain higher flow velocity variations.

115 For invertebrates, given their reduced mobility, we expected that past hydraulic  
116 conditions would have a stronger influence on their microhabitat than for fish. They should be  
117 disfavored by past dewatering and high flow velocity periods. Consequently, we also expected  
118 invertebrate selection for present hydraulics to be different in hydropeaking rivers than those  
119 without hydropeaking. Finally, we expected different responses in benthic and in hyporheic  
120 invertebrates, with a weaker effect of dewatering and high flow velocities on hyporheic  
121 invertebrates, the hyporheic area acting as shelter.

## 122 **Materials and methods**

### 123 **Study reach**

124 The Ain River in eastern France flows from the Jura mountains to the Rhône River, with a  
125 catchment area of 3,630 km<sup>2</sup>. The study reach is 6 km long and 110 m wide, with a mean  
126 discharge of 103 m<sup>3</sup> s<sup>-1</sup>, an altitude of 230 m and a Strahler order of 5. It includes a diversity of  
127 morphological units (runs, riffles and pools), secondary channels and tributaries. Substrate is  
128 dominated by cobbles and the reach contains a few woody debris but little macrophyte cover.  
129 The reach is located 20 km downstream from the Allement hydropower plant (46° 06' 44" N, 5°  
130 25' 20" E), the last of a series of five hydropeaking hydropower plants along the river between  
131 Vouglans and the Rhône confluence. According to data measured by DREAL Rhône-Alpes  
132 (between 1960 and 2019) at Pont-d'Ain (7.5 km upstream of the study reach), mean daily  
133 discharge is generally between 13.5 m<sup>3</sup> s<sup>-1</sup> (exceeded 95% of the time) and 278 m<sup>3</sup> s<sup>-1</sup> (exceeded  
134 5% of the time). The study reach is subject to frequent sub-daily discharge variations, with base  
135 flow generally between 14 m<sup>3</sup> s<sup>-1</sup> and (occasionally) 150 m<sup>3</sup> s<sup>-1</sup> and peak flow generally between  
136 40 m<sup>3</sup> s<sup>-1</sup> and 200 m<sup>3</sup> s<sup>-1</sup> (Figure 1). Discharge is more stable during weekends and low-flow  
137 periods (June-September).

### 138 **Reach hydraulics: the 2D model**

139 A 2D unsteady hydrodynamic model (mapping flow velocities averaged over the vertical; Rubar  
140 2D; Bazin et al. 2017) was built, based on a digital elevation model obtained from topographic  
141 and bathymetric LiDAR surveys carried out in 2015 and 2016. Model calibrations used water  
142 surface levels derived from the LiDAR surveys at 16 m<sup>3</sup> s<sup>-1</sup> and from field surveys at 90 m<sup>3</sup> s<sup>-1</sup>.

143 Calibrations were made so that the absolute difference between the water level simulated by  
144 the model and that measured in the field was less than 10 cm at any measuring point. We used  
145 the model to translate the hourly flow time-series given by the gauging station of Pont d'Ain,  
146 seven kilometers upstream of the study site (from <http://hydro.eaufrance.fr>), into maps of  
147 hourly hydraulic time-series (flow velocity, water level and water depth) over the 403,591  
148 nodes of the 2 m-edge rectangular mesh of the model.

## 149 **Sampling**

### 150 **Fish sampling**

151 Fish species abundance was estimated by electrofishing in 507 microhabitats and by snorkeling  
152 in 673 microhabitats. With both protocols, the sampled surface area of microhabitats was  
153 approximately 7 m<sup>2</sup>, fish being attracted (electrofishing) or observable (snorkeling) within a  
154 radius of about 1.5 m (although this distance can depend on fish size; Regis et al., 1981)

155         Electrofishing was conducted by a team of three or four operators from a motorboat in  
156 deep areas and by wading in shallow areas, approaching the microhabitat as discreetly as  
157 possible to minimize fish escape. An anode was immersed and held steady, and all fish around  
158 the anode were captured with a landing net, identified, measured and released. Microhabitat  
159 positions were recorded by GPS with a precision of 5 m. Electrofishing surveys covered  
160 different flow conditions (Figure 2) and microhabitat locations were chosen to cover diverse  
161 combinations of past and present hydraulic conditions. In practice, the 2D model was used to  
162 draw maps of simulated past hydraulic conditions corresponding to a typical hydropeaking  
163 week, to help finding these different combinations of past and present hydraulics in the field.  
164 Sampling at different flows also reduced the correlation between present and past hydraulic  
165 conditions (e.g., the expected negative correlation between dewatering frequency and water  
166 depth).

167         Snorkeling observations targeted only large individuals of species that are difficult to  
168 sample by electrofishing (estimated length > 25 cm) (Plichard et al., 2017). Observations were  
169 made along six longitudinal transects, regularly spaced laterally across the reach, in the  
170 direction of flow over the entire study reach, during base flow only to ensure fish identification  
171 (~16 m<sup>3</sup> s<sup>-1</sup>). Two persons drifted along the transect in a downstream direction: (1) an

172 experienced snorkeler detecting and identifying fish, followed by (2) an hydrospeeder  
173 recording the observations and GPS positions with a waterproof recorder. As reported by  
174 others (Chamberland et al., 2014), we did not frequently observed strong escape behavior  
175 while snorkeling and judged that most fish were observed by snorkelers. Underwater visibility,  
176 estimated as the distance at which snorkelers could see their bright swimming fins, was 5 m.  
177 Fish were observed and identified when in a 1.5 m radius around the snorkeler, and several fish  
178 individuals were associated to the same GPS position when observed simultaneously.  
179 Therefore, each GPS position in the dataset was considered as a microhabitat of  $\sim 7\text{m}^2$ ,  
180 containing one or several fish. When drifting, snorkelers observed large fish (estimated length  $>$   
181 25 cm) of eight species (*Barbus barbus*, *Squalius cephalus*, *Chondrostoma nasus*, *Thymallus*  
182 *thymallus*, *Perca fluviatilis*, *Esox lucius*, *Cyprinus carpio*, *Salmo trutta fario*) and smaller  
183 individuals of *Thymallus thymallus* only (estimated length  $<$  25 cm).

184 All snorkeling microhabitats were presence-only (microhabitats with fish presence),  
185 unlike the electrofishing and invertebrate microhabitats. To enable common statistical  
186 processing of all data, we created fictive snorkeling microhabitats with fish absence (“pseudo-  
187 absence microhabitats”; Manly et al. 2002). In practice, for each taxon  $\times$  microhabitat  
188 combination, we randomly picked  $n = 10$  pseudo-absence microhabitats among the nodes of  
189 the 2D model network. At this step, we considered only nodes with water depth  $> 30$  cm, as  
190 snorkeling was not conducted in shallower habitats. The choice of  $n = 10$  pseudo-absence  
191 microhabitats was intended to provide a global prevalence comparable with that of the  
192 electrofishing data. We tested the sensitivity of results to this choice by repeating the analysis  
193 for  $n=20$ .

#### 194 **Invertebrate sampling**

195 We sampled benthic and hyporheic invertebrates in 36 microhabitats, under different flow  
196 conditions (Figure 2). Similarly to the procedure used for fish, we used the 2D model to draw  
197 maps of simulated past hydraulic conditions corresponding to a typical hydropeaking week, to  
198 help finding different combinations of past and present hydraulics in the field. We chose  
199 microhabitats in six cross-sections with low and six with high velocity, distributed along the  
200 reach to have different present and past flow velocity. Within each cross-section, we selected



201 three points with low ( $0 \leq \text{depth} < 0.2$  m), medium ( $0.2 \leq \text{depth} < 0.5$  m) and high water depth  
202 ( $0.5 \leq \text{depth} < 0.8$  m) (and thus different dewatering frequencies).

203 We sampled benthic invertebrates with a Hess sampler (surface area  $0.05 \text{ m}^2$ ,  $250 \mu\text{m}$   
204 mesh size) and recorded their positions by GPS. Then, within a radius of one meter around the  
205 Hess sampler, hyporheic invertebrates were sampled with a Bou-Rouch pump (Bou & Rouch,  
206 1967; Dole-Olivier et al., 2014), which extracted 6L of water from 30 cm below the stream bed  
207 (using a planted steel pipe) to catch organisms living in the interstices of substrate particles.  
208 Benthic samples were filtered through a sieve with  $250 \mu\text{m}$  mesh size to be consistent with the  
209 benthic sampling. Invertebrates were preserved in 96% ethanol and were sorted, counted and  
210 identified to the lowest practical taxonomic level using a microscope in the laboratory (list of  
211 references used for invertebrate identification in Supporting Information, list S1).

## 212 **Classification in species guilds**

213 We grouped fish and invertebrate individuals into a few habitat guilds with comparable habitat  
214 selection according to Plichard et al. (2020) and Forcellini et al. (in press), who synthesized a  
215 large part of the available information on fish and invertebrate microhabitat selection in a wide  
216 range of mostly non-hydropeaking European rivers. This grouping by guilds was used to  
217 increase the statistical power of our analyses and the potential transferability of our results in  
218 rivers with other species composition.

219 Fish at electrofishing points were first grouped by size class, using the taxa definitions of  
220 Plichard et al. (2020). These taxa combined a specific code and one or several size classes (1:  $< 8$   
221 cm; 2: 8–18 cm, 3: 18–30 cm; 4:  $> 30$  cm; Table 1). For example, taxon Lel\_cl123 corresponds to  
222 *Leuciscus Leuciscus* (Lel) fish with sizes  $< 30$  cm. We defined two fish guilds for which we  
223 expected different influence of past hydraulics on microhabitat selection. The ‘bank’ guild  
224 grouped together taxa selecting shallow and slow-flowing habitats, thus incurring high  
225 dewatering risk. Conversely, the ‘midstream’ guild grouped together species selecting deep and  
226 fast-flowing habitats, typically found in the center channel, which should better sustain high  
227 flow velocity variations. Observed graylings (*Thymallus thymallus*) were also assigned to the  
228 midstream guild (Mallet et al. 2000). Plichard et al. (2020) summarized the microhabitat  
229 selection by fish taxa for microhabitat velocity  $V$  using the statistic  $AGV_V$ , which in short

230 corresponds to the average microhabitat velocity that would be used by the taxa if velocities  
231 between 0 and 0.94 m s<sup>-1</sup> were uniformly available in the river. Similarly, AVG\_D summarizes  
232 selection for depth D over the depth range 0-3 m. Our midstream guild comprised the nine taxa  
233 with the highest value for the product AVG\_V\*AVG\_D, and our bank guild the eight taxa with  
234 the lowest value. The number of taxa per guild was a compromise between guild abundance  
235 and magnitude of habitat selection.

236 For invertebrates, we defined two guilds according to selection for bottom shear stress  
237 reported in Forcellini et al. (in press), where shear stress was measured using hemispheres  
238 (FST, Statzner & Müller, 1989) numbered from 0 to 19 (low to high shear stress). As done by  
239 Plichard et al. (2020) for fish, Forcellini et al. (in press) summarized microhabitat selection by  
240 invertebrates, using AVG\_V and AVG\_FST, which is the equivalent of AVG\_V but over the FST  
241 range 0-19. The AVG\_FST given by Forcellini et al. (in press) were calculated by species or  
242 higher levels, without accounting for variations in habitat use with individual size. Therefore,  
243 we used the same guild definitions for both benthic and hyporheic taxa, although individuals in  
244 hyporheic samples were generally smaller. The 'limnophilic' guild grouped together taxa  
245 selecting low shear stress (AVG\_FST < 5), and thus expected to be negatively affected by high  
246 flow velocity variations. Conversely, the 'rheophilic' guild corresponded to taxa selecting high  
247 shear stress (AVG\_FST > 8).

## 248 **Present and past microhabitat hydraulics**

249 In each electrofishing and invertebrates sampling microhabitat (i.e. a couple  
250 benthic/hyporheic), we measured the present water depth (D in m), and present flow velocity  
251 ( $V_{40}$  in m s<sup>-1</sup>; measured at 40% of water depth from the bottom with an electromagnetic 30 Hz  
252 current meter and averaged over 10 seconds, using a Marsh McBirney FLO MATE 2000).

253 We described past hydraulic conditions using the 2D hydraulic model simulations for  
254 flows observed during the 15 days preceding sampling. The duration of 15 days was consistent  
255 with previous experiments on fish individual learning (Green, 1971; Roy & Bhat, 2018) and  
256 previous field studies on fish in hydropeaking reaches (Capra et al., 2017). This 15 days duration  
257 also allowed to account for typical discharge variations observed in the Ain River (Figure 2).

258 Nevertheless, we tested the sensitivity of results to our choice of a duration of 15 days, by  
259 repeating our analyses for durations of 5 and 10 days before sampling.

260 For estimating past hydraulics at a given microhabitat (electrofishing point or  
261 invertebrate sample), we associated the sampled microhabitat to a node of the 2D model that  
262 (1) had comparable water depth, and (2) was as close as possible from the sampled  
263 microhabitat. We judged the primary use of a “depth similarity criteria” as essential for an  
264 optimum estimation of the dewatering frequency of the microhabitat. Indeed, for a given  
265 measurement discharge, water depth and past dewatering frequency are expected to be  
266 strongly correlated. In addition, even if the morphology of the river had changed locally  
267 between the 2D model calibration and the biological survey, we expected that velocity  
268 variations for a given water depth was well represented by the 2D model. In the Ain River,  
269 mesohabitats (e.g., riffles, run, pools) have surface areas of several hundred m<sup>2</sup>, and the 2D  
270 model is expected to well reflect hydraulic behavior of these mesohabitats, event it can be  
271 locally imprecise (Guay et al., 2000). In practice, we selected the closest 2D node that had a  
272 modeled depth differing from the measured depth at the microhabitat by less than 10 cm. For  
273 snorkeling observations, depth measurement was not available and we assigned snorkeling  
274 microhabitats to the closest 2D node.

275 The past hydraulic conditions during the 15 days preceding sampling were described  
276 with two variables coded as boolean categories:

277 (1) Duration of dewatering: we considered the microhabitat as “dewatering” if the dewatering  
278 time (depth = 0 cm) over the 15-days period before sampling was >10 hours, and as “non-  
279 dewatering” if ≤10 hours. We tested the sensitivity of results to our choice of a duration of 10  
280 hours, by repeating our analyses for a duration of 1 hour.

281 (2) Duration of high flow velocity conditions (>1.3 m s<sup>-1</sup>): We considered that microhabitats had  
282 “high past velocities” if the total duration with velocity >1.3 m s<sup>-1</sup> was >10 hours, and “low past-  
283 velocities” if ≤10 hours. We chose a threshold of 1.3 m.s<sup>-1</sup> because this value is above the velocity  
284 used by most aquatic organisms (Plichard et al., 2020; Forcellini et al., in press), and is also  
285 frequently reached in many microhabitats of the Ain River. Here again, we tested the sensitivity

286 of results to our choice of a threshold of 1.3 m.s<sup>-1</sup>, by repeating our analyses for a threshold of  
287 0.7 m.s<sup>-1</sup>.

## 288 **Microhabitat selection models by guild**

289 For the two fish (electrofishing, snorkeling) datasets and the benthic invertebrates dataset, we  
290 related guild abundance to past and present microhabitat characteristics (measured when  
291 available, modeled otherwise). We derived similar microhabitat selection models based on the  
292 hyporheic dataset, for testing if the influence of past hydraulics observed on benthic  
293 invertebrates was still observed in hyporheic samples, or if the hyporheic area played a  
294 particular role for these guilds. For each dataset, following Plichard et al. (2020) and Forcellini  
295 et al. (in press), we used GLMs with B-spline transformations of the present hydraulic variable  
296 (enabling non-linear responses) and assuming a negative binomial distribution of abundance  
297 (accounting for abundance overdispersion). We used splines with two degrees of freedom, with  
298 a single knot positioned at the median value (Plichard et al., 2020). To avoid  
299 overparameterization and due to limited statistical power, we had to consider models involving  
300 combinations of a single present hydraulic variable (*PresHyd* =  $V_{40}$  or  $D$ ) and a single past  
301 hydraulic variable (*PastHyd*=  $Dew$  or  $HV$ ). Consistently, we looked at the inter-correlations  
302 between present and past hydraulic variables to interpret the results.

303 For each *PresHyd* × *PastHyd* combination, the abundance  $Y_i$  of a guild in microhabitat  $i$  was  
304 assumed to follow a negative binomial distribution of mean  $\mu_i$  and dispersion parameter  $\Theta$ :

$$305 Y_i \sim NB(\mu_i, \Theta)$$

306 Three GLMs of increasing complexity were compared to explain  $\mu_i$  as a function of hydraulics:

$$307 \mathbf{M0 (no microhabitat selection):} \log(\mu_i) = \beta_0$$

$$308 \mathbf{M1 (habitat selection with present hydraulic conditions only):} \log(\mu_i) = \beta_0 + f(\text{PresHyd}_i)$$

$$309 \mathbf{M2 (habitat selection with present and past hydraulic conditions):} \log(\mu_i) = \beta_0 + f(\text{PresHyd}_i) + \beta_1$$

310 *PastHyd* <sub>$i$</sub>

311 where  $\beta_0$  is the intercept,  $f()$  is a spline transformation of the present hydraulic variable  
312 *PresHyd* <sub>$i$</sub> , and  $\beta_1$  the coefficient of the past hydraulics effect. In M0, microhabitat variables have

313 no influence on abundance, in M1 present hydraulic conditions have an influence, and in M2  
314 there is an additive effect of past hydraulics.

315 We fitted all models using the R software (R Development Core Team 2018) and the  
316 glm.nb function of the 'MASS' package (Venables & Ripley, 2002), combined with spline  
317 functions of the 'splines' package. We used likelihood ratio tests comparing M0 vs. M1 to test  
318 the influence of present hydraulics on microhabitat selection, and tests comparing M1 vs. M2  
319 to assess for an additive effect of past hydraulics. Following Plichard et al. (2020), we used non-  
320 parametric Spearman Rho rank correlation (Spearman, 1904) to assess fits; this is an  
321 appropriate statistic for overdispersed data such as our abundance data, based on the  
322 correlation between the ranks of predicted and observed abundance.

### 323 **Influence of hydropeaking on taxa selection for present hydraulics**

324 For the fish electrofishing dataset and benthic invertebrates, we computed an average  
325 preferred velocity AGV\_V and depth AVG\_D as defined by Plichard et al. (2020), but  
326 corresponding to the fits of our M1 model by taxa. We then compared our values with those of  
327 Plichard (2020) for fish and Forcellini et al. (in press) for invertebrates, to estimate how  
328 hydropeaking modifies the selection of present microhabitat hydraulics generally observed in  
329 rivers without hydropeaking. We used only electrofishing data for the fish comparison, because  
330 fish size was not available in snorkeling data, preventing comparison with the size classes of  
331 Plichard et al. (2020). We used only the benthic invertebrate data, to be consistent with  
332 Forcellini et al. (in press).

## 333 **Results**

### 334 **Biological assemblages**

335 We sampled 3,642 fish of 23 species by electrofishing. The bank guild contained 2,344  
336 individuals and was composed mainly of *Phoxinus phoxinus* cl12 (N=1,785). The midstream  
337 guild contained 73 individuals, mainly *Barbus barbus* cl34 (N=27) (Table 1). We sampled 1,610  
338 fish of 12 species by snorkeling (mainly *Thymallus thymallus*, N=444; *Barbus barbus*, N=437 and  
339 *Chondrostoma nasus*, N=238). In the snorkeling dataset, large *Barbus barbus*, large

340 *Chondrostoma nasus* and all *Thymallus thymallus* belonged to the midstream guild, with many  
341 more individuals sampled than with electrofishing (N =356) (Table 1).

342 We sampled 48,275 invertebrates: 38,196 in benthic samples and 10,079 in hyporheic  
343 samples. The full list of taxa of both datasets are available as Supporting Information (Table S1).  
344 For the benthic dataset, the limnophilic guild contained 687 individuals (principally composed  
345 of *Chironomini*, N=612) and the rheophilic guild contained 2988 individuals (principally  
346 composed of *Esolus*, N=2,021) (Table 2). For the hyporheic dataset, the limnophilic guild  
347 contained 170 individuals (principally composed of *Chironomini*, N=140) and the rheophilic  
348 guild contained 961 individuals (principally composed of *Esolus*, N=754) (Table S2). Note that  
349 only 10% of all benthic individuals and 11% of all hyporheic individuals belong to a habitat guild  
350 (i.e. documented and significant habitat selection).

### 351 Present and past hydraulic variables and their intercorrelation (Figure 3)

352 Present depth and velocity typically ranged between 0-4 m and 0-2 m s<sup>-1</sup> in the fish  
353 datasets (Figure 3A). By contrast, depth was generally shallower than 0.8 m and velocity lower  
354 than 1 m s<sup>-1</sup> in the invertebrate dataset, due to sampling limits. Water depth and flow velocity  
355 were poorly correlated (and with different directions) for the fish data (electrofishing: r=0.13,  
356 P<0.05; snorkeling: r=-0.48, P<0.05) (Figure 3A). The correlation was greater in the invertebrate  
357 dataset (r=0.57; P<0.05).

358 When calculating past hydraulics using the 2D hydrodynamic model, 76% of the  
359 electrofishing microhabitats and 75% of the invertebrate microhabitats were associated with a  
360 model node within a radius of 5.5 m around the sampled microhabitat. In the electrofishing  
361 dataset, 102/507 microhabitats were dewatering habitats and 255/507 had high past velocities  
362 (Figure 3B). The snorkeling dataset contained no dewatering microhabitats and 342/673  
363 microhabitats with high past velocities. A little less than half of the invertebrate samples  
364 (32/72) were dewatering and a higher proportion (44/72) had past low velocities. For fish  
365 (electrofishing) and invertebrates, all combinations of dewatering × high past velocity habitats  
366 were sampled (Figure 3B). Nevertheless, dewatering habitats were generally those with low  
367 past velocity for both fish and invertebrates (Xhi2 test, P<0.05).

368 Correlations between present and past hydraulics were limited ( $r^2 < 0.30$  in all cases).  
369 The strongest correlation was between flow velocity and high past velocities (Figure 3C),  
370 especially for the electrofishing and the invertebrate datasets. Dewatering habitats also often  
371 had lower flow velocities and depths (Figure 3D).

## 372 **Microhabitat selection models by guilds**

373 As expected from the fish guild definitions, the midstream guild used higher present velocities  
374 and depths than the bank guild, according to both electrofishing and snorkeling data. The bank  
375 guild significantly avoided fast-flowing and deep microhabitats (Table 3, Figure 4). Differences  
376 in response to flow velocity between guilds, however, were less clear for benthic invertebrates  
377 than for fish (Table 3, Figure 4A). The rheophilic benthic invertebrate guild significantly avoided  
378 shallow microhabitats (Table 3, Figure 4A). Similar patterns were observed for hyporheic  
379 invertebrates (Table S3, Figure S1).

380 Several significant additive effects of past hydraulics were observed (indicated by  
381 asterisks in Figure 4). Benthic and hyporheic invertebrates responded more significantly to past  
382 hydraulics, all invertebrate guilds being 3.5-15.3 times less abundant in dewatering habitats  
383 (Figure 4A, Figure S1). Consistently, Spearman Rho values for benthic invertebrate M2 models  
384 involving dewatering (between 0.27 and 0.61, Table 3) were much higher than the  
385 corresponding values for M1 models (between 0.04 and 0.35, Table 3). Again, similar patterns  
386 were observed for hyporheic invertebrates (Table S3). Secondarily, benthic rheophilic  
387 invertebrates selected microhabitats with high past velocities (3.9 times more abundant in high  
388 past velocity microhabitats in average) and hyporheic limnophilic invertebrates tend to avoid  
389 them (0.4 times less abundant, only significant with the model including water depth as  
390 present hydraulic variable) (Figure 4A, Figure S1). The influence of high past velocities was not  
391 significant for hyporheic rheophilic invertebrates and benthic limnophilic invertebrates. For fish  
392 (electrofishing), responses were weaker and clearly different, with the bank guild selecting  
393 dewatering microhabitats (Figure 4A) and, less evidently, the midstream guild avoiding  
394 dewatering microhabitats. Midstream fish of the snorkeling dataset selected microhabitats  
395 with high past velocities.

396 The sensitivity tests concerning the choice of a 15 days duration for calculating past  
397 hydraulics and a 10 hours duration for defining dewatering microhabitats did not modify any of  
398 our results (significance tests in Table 3 and Figure 4).

399 By contrast, a threshold of 0.7 m.s<sup>-1</sup> instead of 1.3 m.s<sup>-1</sup> for defining microhabitats with  
400 high past velocities and, a number of 20 instead of 10 pseudo-absence when building model on  
401 the snorkeling dataset modified some of our (secondary) results on the effects of high past  
402 velocities. With a lower 0.7 m.s<sup>-1</sup> threshold, the bank fish guild and hyporheic rheophilic  
403 invertebrates significantly selected high past velocities (M2 models) and the midstream fish  
404 (electrofishing) guild avoided them (M2 model with velocity). In addition, the selection of high  
405 past velocities by midstream fish (snorkeling) was no longer significant. The selection of high  
406 past velocities by midstream fish (snorkeling) was also no longer significant with a number of  
407 20 instead of 10 pseudo-absence when building models.

#### 408 **Influence of hydropeaking on taxa selection for present hydraulics**

409 Microhabitat selection for present hydraulics could be compared between the present study  
410 and the literature for nine fish taxa and thirteen benthic invertebrate taxa. Present velocity and  
411 water depth selection by fish taxa corresponded to those reported in the literature (Figure 5A)  
412 and had comparable Spearman Rho values (Figure 5B). By contrast, present velocity and water  
413 depth selection by benthic invertebrate taxa did not match values reported in the literature  
414 (Figure 5C) and had lower Spearman Rho values here (Figure 5D).

#### 415 **Discussion**

416 By studying the combined influence of present and past hydraulics on microhabitat selection,  
417 we improved our mechanistic understanding of the impact of hydropeaking on fish and  
418 invertebrate guilds. We found that both benthic limnophilic and rheophilic invertebrates were  
419 less abundant in dewatering microhabitats. This confirms findings such as those of Blinn et al.  
420 (1995), who reported a four-fold lower invertebrate biomass in dewatering habitats of the  
421 Colorado River. Consistently with this result, we also showed that the average velocity and  
422 water depth preferences of some invertebrate taxa differed from those previously established  
423 in mainly non-hydropeaking rivers (Figure 5). This suggests that invertebrates are not able to



424 maintain habitat preferences when flow changes too rapidly or frequently: they likely stay at  
425 the same location, undergoing different hydraulic conditions according to flow conditions.

426 Similarly, to benthic invertebrates, we found that hyporheic invertebrates are also less  
427 abundant in dewatering microhabitat. Yet, we expected the opposite since we suggested that  
428 hyporheic area provide shelter under dewatering conditions in the Ain River (Williams & Hynes,  
429 1974). Others studies find some comparable results suggested that benthic macroinvertebrates  
430 may not always use the hyporheic area as shelter in response to a discharge reduction (del  
431 Rosario and Resh 2000; James et al. 2008). In our case, since some microhabitats could have  
432 been dewatered for several days before sampling, we cannot exclude that in some case the  
433 hyporheic area undergone dewatering or low oxygen levels resulting in death of some  
434 individuals. Furthermore, invertebrates may have migrate rapidly through the benthic area just  
435 after flow increases and before sampling (Bruno et al., 2020). Sampling during the dry period  
436 could allow to test if the hyporheic area plays the role of shelter.

437 In contrast to invertebrates, fish followed their preferred habitat preferences observed  
438 in other rivers and were less influenced by past hydraulics. Nevertheless, fish from the “bank”  
439 guild probably stayed along the bank and moved to dewatering microhabitats when flow  
440 increased, in order to remain within their strict range of preferred hydraulic conditions. This  
441 result contrasts with many studies concerning salmonid juveniles, also living near the banks,  
442 and indicating that only a fraction of individuals relocate when flow changes (Kemp et al., 2003;  
443 Korman & Campana, 2009; Shirvell, 1994). High velocities in the Ain River and the different  
444 species considered here (mostly cyprinids) may explain these differences. In the Ain River, fish  
445 from the bank guild are probably exposed to a high risk of stranding, particularly because  
446 cobble bars have very low slopes (Tuhtan et al., 2012). It would be interesting to quantify the  
447 influence of horizontal down-ramping rate (water surface retreat rate) on fish stranding near  
448 these cobble bars. The fish “midstream” guild also followed their preferred range of habitat,  
449 and tended to avoid dewatering and stressful microhabitats (Capra et al., 2017; Reeb, 1996).  
450 Although it concerned other species (large barbel, catfish and chub), the telemetry experiment  
451 by Capra et al. (2017) also reported that fish could avoid dewatering microhabitats. However,  
452 due to the small number (N=73) of fish involved in this present result, it is not possible to  
453 generalize.

454 Observed microhabitat selection by fish and invertebrates was not negatively affected  
455 by high past velocities, suggesting a weaker influence of high past velocities compared to  
456 dewatering. In addition, the statistical significance of the weak responses to high past  
457 velocities depended on the threshold chosen for defining high past velocities, indicating  
458 unstable results. However, considering that hydropeaking has occurred since the 1930s in the  
459 Ain River, environmental filtering may have selected only species that can cope with rapid and  
460 frequent variations in hydraulic conditions. Furthermore, our results suggest that high  
461 velocities force fish to move (sticking to their general hydraulic preference) and invertebrates  
462 to find flow-shelter or attach to the substrate (e.g. with silk draglines, tarsal claws, or a  
463 muscular foot; Holomuzki et al. 2000). Moving (fish) or staying in sub-optimal habitat  
464 conditions (invertebrates) may create energetic costs that influence community structure on  
465 the long term and explain reach-scale effects of hydropeaking (Judes et al., 2021). More  
466 detailed observations of behavior at high flows (e.g. by snorkeling or with cameras),  
467 experimental/physiologic experiments in flumes, or additional comparisons between reaches  
468 subjected to different degrees of hydropeaking (Schmutz et al., 2015; Judes et al., 2021) will be  
469 useful to confirm this interpretation.

470 We identify five major limits and ways of improvement in our study. First, we pooled  
471 microhabitat samples under low and high flow conditions due to a limited statistical power.  
472 However, habitat availability strongly changes according to flow conditions. For example, in the  
473 Ain, 77% of the wetted surface is less than 1 m deep at  $14 \text{ m}^3 \text{ s}^{-1}$  versus 33% at  $214 \text{ m}^3 \text{ s}^{-1}$ . Such  
474 differences in habitat availability could influence species microhabitat selection (Arthur et al.,  
475 1996; Schooley, 1994). Further sampling could target a more detailed analysis of the effects of  
476 habitat availability. Similarly, improved analyses should also consider the effects of woody  
477 debris, macrophyte cover, tributary proximity and/or thermal refugia. Secondly, here, guilds  
478 are composed only of taxa with significant habitat selection documented in the literature. In  
479 the future, it would be interesting to be more representative of the community in place. In  
480 particular, it would be interesting to study ubiquitous taxa, to account for size distributions  
481 within taxa, or to analyze responses of functional traits. Thirdly, discharge before and during  
482 our sampling period combined hydropeaks and artificial releases of intermediate flows (around  
483  $100 \text{ m}^3 \text{ s}^{-1}$ , Figure 2) to compensate for low discharge in the Rhône River. The duration of  
484 intermediate flow may influence observed fish and invertebrates microhabitat selection. Future

485 sampling will need to be implemented during different hydropeaking regimes, especially for  
486 invertebrates sampling, which was conducted at only three flow conditions here. Fourthly,  
487 because our results were little sensitive to a number of our methodological choices (duration  
488 considered for past hydraulics, threshold used for defining dewatering habitats and high past  
489 velocities), additional experiments/data would be useful to specify the key quantitative drivers  
490 of biological responses. Finally, although the 2D hydrodynamic model allowed us to estimate  
491 past hydraulics at each microhabitat, its accuracy and the methods used for associating field  
492 observations with model nodes would deserve further investigation (see Guay et al., 2000).

493 In conclusion, our microhabitat-scale study shows a stronger effect of dewatering than  
494 high past velocities on invertebrates, and secondarily on fish. Although this study failed to  
495 identify critical past velocity thresholds for hydropeaking management, the results suggests  
496 that fish are forced to move and invertebrate to hide due to these high past velocities.  
497 Comparing our results with data collected in other rivers and at a larger variety of flows,  
498 combined with experimental studies, should help further identifying the key hydropeaking  
499 characteristics to consider in mitigation policies.

## 500 **Acknowledgments**

501 We are grateful for the financial support of EDF- Hydro Alpes (Gérald Ramos) for the LiDAR and  
502 water line surveys. We thank EDF-CIH (Le Bourget du Lac, France) and Dimitri Lague  
503 (Géosciences Rennes, UMR 6118, France) for collaboration in developing the Digital Elevation  
504 Model derived from LiDAR surveys. We gratefully acknowledge all the people who contributed  
505 to the field work.

## 506 **References**

- 507 Arthur, S. M., Manly, B. F., McDonald, L. L., & Garner, G. W. (1996). Assessing habitat selection  
508 when availability changes. *Ecology*, 77(1), 215-227.  
509 <https://doi.org/https://doi.org/10.2307/2265671>
- 510 Auer, S., Zeiringer, B., Führer, S., Tonolla, D., & Schmutz, S. (2017). Effects of river bank  
511 heterogeneity and time of day on drift and stranding of juvenile European grayling

512 (Thymallus thymallus L.) caused by hydropeaking. *Science of the Total Environment*, 575,  
513 1515–1521. <https://doi.org/10.1016/j.scitotenv.2016.10.029>

514 Bazin, P., Mignot, E., Paquier, A., Bazin, P., Mignot, E., & Paquier, A. (2017). Computing flooding  
515 of crossroads with obstacles using a 2D numerical model. *Journal of Hydraulic Research*,  
516 55(5). <https://doi.org/https://doi.org/10.1080/00221686.2016.1217947>

517 Blinn, D. W., Shannon, J. P., Stevens, L. E., & Carder, J. P. (1995). Consequences of fluctuating  
518 discharge for lotic communities. *Journal of the North American Benthological Society*,  
519 14(2), 233–248. <https://doi.org/10.2307/1467776>

520 Bou, C., & Rouch, R. (1967). Un nouveau champ de recherches sur la faune aquatique  
521 souterraine. *C. R. Academie Hebdomadaire Des Sciences*, 265, 369–370.

522 Bruno, M. C., Siviglia, A., Carolli, M., & Maiolini, B. (2013). Multiple drift responses of benthic  
523 invertebrates to interacting hydropeaking and thermopeaking waves. *Ecohydrology*, 6(4),  
524 511–522. <https://doi.org/10.1002/eco.1275>

525 Capra, H., Plichard, L., Bergé, J., Pella, H., Ovidio, M., Mcneil, E., & Lamouroux, N. (2017). Fish  
526 habitat selection in a large hydropeaking river: strong individual and temporal variations  
527 revealed by telemetry. *Science of the Total Environment*, 578, 109–120.  
528 <https://doi.org/10.1016/j.scitotenv.2016.10.155>

529 Chamberland, J. M., Lanthier, G., & Boisclair, D. (2014). Comparison between electrofishing and  
530 snorkeling surveys to describe fish assemblages in Laurentian streams. *Environmental*  
531 *Monitoring and Assessment*, 186(3), 1837–1846. [https://doi.org/10.1007/s10661-013-](https://doi.org/10.1007/s10661-013-3497-4)  
532 3497-4

533 Dole-Olivier, M. J., Maazouzi, C., Cellot, B., Fiers, F., Galassi, D. M. P., Claret, C., Martin, D.,  
534 Méricoux, S., & Marmonier, P. (2014). Assessing invertebrate assemblages in the  
535 subsurface zone of stream sediments (0-15 cm deep) using a hyporheic sampler. *Water*  
536 *Resources Research*, 50(1), 453–465. <https://doi.org/10.1002/2012WR013207>

537 Dole-Olivier, M. J., Marmonier, P., & Befly, J. L. (1997). Response of invertebrates to lotic  
538 disturbance: Is the hyporheic zone a patchy refugium? *Freshwater Biology*, 37(2), 257–

- 539 276. <https://doi.org/10.1046/j.1365-2427.1997.00140.x>
- 540 Forcellini, M., Plichard, L., Dolédec, S., Méricoux, S., Olivier, J.-M., Cauvy-Fraunié, S., &  
541 Lamouroux, N. (in press). Microhabitat selection by macroinvertebrates: generality among  
542 rivers and functional interpretation. *Journal of Ecohydraulics*.  
543 <https://doi.org/10.1080/24705357.2020.1858724>
- 544 Green, J. (1971). High Tide Movements and Homing Behaviour of the Tidepool Sculpin  
545 *Oligocottus maculosus*. *Journal of the Fisheries Board of Canada*, 28(250), 3.  
546 <https://doi.org/10.1139/f71-051>
- 547 Guay, J. C., Boisclair, D., Rioux, D., Leclerc, M., Lapointe, M., & Legendre, P. (2000).  
548 Development and validation of numerical habitat models for juveniles of Atlantic salmon  
549 (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, 57(10), 2065–2075.  
550 <https://doi.org/10.1139/cjfas-57-10-2065>
- 551 Halleraker, J. H., Saltveit, S. J., Harby, A., Arnekleiv, J. V., Fjeldstad, H. P., & Kohler, B. (2003).  
552 Factors influencing stranding of wild juvenile brown trout (*Salmo trutta*) during rapid and  
553 frequent flow decreases in an artificial stream. *River Research and Applications*, 19(5–6),  
554 589–603. <https://doi.org/10.1002/rra.752>
- 555 Holomuzki, J., & Biggs, B. J. F. (2000). Taxon-specific responses to high-flow disturbance in  
556 streams. *Journal of the North American Benthological Society*, 19(4), 670–679.  
557 <https://doi.org/https://doi.org/10.2307/1468125>
- 558 Judes, C., Gouraud, V., Capra, H., Maire, A., Barillier, A., & Lamouroux, N. (2021). Consistent but  
559 secondary influence of hydropeaking on stream fish assemblages in space and time.  
560 *Journal of Ecohydraulics*. <https://doi.org/10.1080/24705357.2020.1790047>
- 561 Kemp, P. S., Gilvear, D. J., & Armstrong, J. D. (2003). Do juvenile Atlantic salmon parr track local  
562 changes in water velocity? *River Research and Applications*, 19(5–6), 569–575.  
563 <https://doi.org/10.1002/rra.727>
- 564 Kjaerstad, G., Arnekleiv, J. V., Speed, J. D. M., & Herland, A. K. (2018). Effects of hydropeaking  
565 on benthic invertebrate community composition in two central Norwegian rivers. *River*

566        *Research and Applications*, 34(3), 218–231. <https://doi.org/10.1002/rra.3241>

567 Korman, J., & Campana, S. E. (2009). Effects of Hydropeaking on Nearshore Habitat Use and  
568        Growth of Age-0 Rainbow Trout in a Large Regulated River. *Transactions of the American*  
569        *Fisheries Society*, 138(1), 78–87. <https://doi.org/10.1577/T08-026.1>

570 Lamouroux, N., Capra, H., Pouilly, M., & Souchon, Y. (1999). Fish habitat preferences in large  
571        streams of southern France. *Freshwater Biology*, 42(4), 673–687.  
572        <https://doi.org/10.1046/j.1365-2427.1999.00521.x>

573 Manly, B. F., McDonald, L., Thomas, D., McDonald, T. L., & Erickson, W. P. (2002). *Resource*  
574        *Selection by Animals: Statistical Design and Analysis for Field Studies*. Springer  
575        Netherlands.

576 Patton, B. W., & Braithwaite, V. A. (2015). Changing tides: Ecological and historical perspectives  
577        on fish cognition. *Wiley Interdisciplinary Reviews: Cognitive Science*, 6(2), 159–176.  
578        <https://doi.org/10.1002/wcs.1337>

579 Pert, E. J., & Erman, D. (1994). Habitat use by adult rainbow trout under moderate artificial  
580        fluctuations in flow. *Transactions of the American Fisheries Society*, 123(6), 913–923.  
581        [https://doi.org/10.1577/1548-8659\(1994\)123<0913](https://doi.org/10.1577/1548-8659(1994)123<0913)

582 Plichard, L., Capra, H., Mons, R., Pella, H., & Lamouroux, N. (2017). Comparing electrofishing  
583        and snorkelling for characterizing fish assemblages over time and space. *Canadian Journal*  
584        *of Fisheries and Aquatic Sciences*, 74(1), 75–86. <https://doi.org/10.1139/cjfas-2015-0578>

585 Plichard, L., Forcellini, M., Le Coarer, Y., Capra, H., Carrel, G., Ecochard, R., & Lamouroux, N.  
586        (2020). Predictive models of fish microhabitat selection in multiple sites accounting for  
587        abundance overdispersion. *River Research and Applications*, 36(7), 1056–1075.  
588        <https://doi.org/10.1002/rra.3631>

589 Reeb, S. G. (1996). Time-place learning in golden shiners (Pisces: Cyprinidae). *Behavioural*  
590        *Processes*, 36(3), 253–262. [https://doi.org/10.1016/0376-6357\(96\)88023-5](https://doi.org/10.1016/0376-6357(96)88023-5)

591 Regis, J., Pattee, E., & Lebreton, J. (1981). A new method for evaluating the efficiency of electric  
592        fishing. *Archiv Für Hydrobiologie*, 93, 68–82. <https://doi.org/10.1016/j.fishres.2011.11.011>

- 593 Roy, T., & Bhat, A. (2018). Divergences in learning and memory among wild zebrafish : Do sex  
594 and body size play a role ? *Learn Behav*, 46(2), 124–133. [https://doi.org/10.3758/s13420-](https://doi.org/10.3758/s13420-017-0296-8)  
595 017-0296-8
- 596 Saltveit, S. J., Halleraker, J. H., Arnekleiv, J. V., & Harby, A. (2001). Field experiments on  
597 stranding in juvenile Atlantic Salmon (*Salmo salar*) and Brown Trout (*Salmo trutta*) during  
598 rapid flow decreases caused by hydropeaking. *Regulated Rivers: Research & Management*,  
599 17(4–5), 609–622. <https://doi.org/10.1002/rrr.652>
- 600 Schmutz, S., Bakken, T. H., Friedrich, T., Greimel, F., Harby, A., Jungwirth, M., Melcher, A.,  
601 Unfer, G., & Zeiringer, B. (2015). Response of fish communities to hydrological and  
602 morphological alterations in hydropeaking rivers of Austria. *River Research and*  
603 *Applications*, 31, 919–930. <https://doi.org/10.1002/rra>
- 604 Schooley. (1994). Annual Variation in Habitat Selection : Patterns Concealed by Pooled Data.  
605 *The Journal of Wildlife Management*, 58(2), 367–374.  
606 <https://doi.org/https://doi.org/10.2307/3809404>
- 607 Shirvell, C. S. (1994). Effect of changes in streamflow on the microhabitat use and movements  
608 of sympatric juvenile coho salmon (*Oncorhynchus kisutch*) and chinook salmon  
609 (*O.tshawytscha*) in a natural stream. *Canadian Journal of Fisheries and Aquatic Sciences*,  
610 51(7), 1644–1652. <https://doi.org/10.1139/f94-165>
- 611 Spearman, C. (1904). The proof and measurement of association between two things.  
612 *International Journal of Epidemiology*, 15(1), 72–101. <https://doi.org/10.1093/ije/dyq191>
- 613 Stubbington, R., Dole-Olivier, M. J., Galassi, D. M. P., Hogan, J.-P., & Wood, P. J. (2016).  
614 Characterization of macroinvertebrate communities in the hyporheic zone of river  
615 ecosystems reflects the pump-sampling technique used. *PLOS One*, 1–27.  
616 <https://doi.org/10.1371/journal.pone.0164372>
- 617 Tachet, P., Richoux, P., Bournaud, M., & Usseglio-Polatera, P. (2002). Invertébrés d'eau douce:  
618 systématique, biologique et écologie. *CNRS Editions*.
- 619 Tuhtan, J. A., Noack, M., & Wieprecht, S. (2012). Estimating stranding risk due to hydropeaking

- 620 for juvenile European grayling considering river morphology. *KSCE Journal of Civil*  
 621 *Engineering*, 16(2), 197–206. <https://doi.org/10.1007/s12205-012-0002-5>
- 622 Vadher, A. N., Leigh, C., Millett, J., Stubbington, R., & Wood, P. J. (2017). Vertical movements  
 623 through subsurface stream sediments by benthic macroinvertebrates during experimental  
 624 drying are influenced by sediment characteristics and species traits. *Freshwater Biology*,  
 625 62(10), 1730–1740. <https://doi.org/10.1111/fwb.12983>
- 626 Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S*. Springer, fourth  
 627 edition. <http://www.stats.ox.ac.uk/pub/MASS4>
- 628 Vila-Martínez, N., Caiola, N., Ibáñez, C., Benejam, L., & Brucet, S. (2019). Normalized abundance  
 629 spectra of fish community reflect hydro-peaking on a Mediterranean large river. *Ecological*  
 630 *Indicators*, 97, 280–289. <https://doi.org/10.1016/j.ecolind.2018.10.014>

631

## 632 Figure legends

633 Figure 1. Location of the Ain River study reach and maps of its flow velocity and water depth for the  
 634 maximum and the minimum flow discharge encountered over the study period.

635 Figure 2. Hourly discharge at Pont d'Ain (7.5 km upstream the study reach) during the 15 days preceding  
 636 sampling and during sampling (2018). The vertical lines represent indicate sampling days at noon.

637 Figure 3. Relationship (A) among present hydraulic variables, (B) among past hydraulic variables and  
 638 (C,D) between present and past hydraulic variables for invertebrates ( $\Rightarrow$ ), one value for a  
 639 benthic/hyporheic pair), fish sampled by electrofishing ( $\bullet$ ), and fish sampled by snorkeling ( $\circ$ ). Points  
 640 represent microhabitats.

641 Figure 4. Fits of M2 models relating guild abundance to present hydraulics and including an additive  
 642 effect of dewatering (A) or past velocity (B). Red and green curves reflect the effect of past hydraulic  
 643 variables, when significant. Graphs are shown for the electrofishing ( $\bullet$ ) snorkeling ( $\circ$ ) and benthic  
 644 invertebrate ( $\Rightarrow$ ) datasets. Stars show the significance of the past hydraulics effect (\*\*)  $P < 0.01$ ; (\*)  
 645  $P < 0.05$ .

646 Figure 5. Comparison of the average preferred velocity (AVG\_V, corresponding to model M3) obtained  
 647 in this study with data from Plichard et al. (2020) for fish and from Forcellini et al. (in press) for benthic  
 648 invertebrates. Axis regressions (full lines) are shown as well as the  $y = x$  lines (dotted lines). For Plichard  
 649 et al. (2020), AVG\_V:  $y = -0.76x - 0.09$ ,  $R^2 = 0.76$ ,  $P < 0.01$ ; Spearman Rho:  $-0.48x + 0.33$ ,  $R^2 = 0.13$ ,  $P = 0.34$ .  
 650 AVG\_D:  $y = 0.54x + 0.08$ ,  $R^2 = 0.21$ ,  $P = 0.22$ ; Spearman Rho:  $-0.13x + 0.26$ ,  $R^2 = 0.03$ ,  $P = 0.66$ . For Forcellini et  
 651 al. (in press), AVG\_V:  $y = 0.02x + 0.68$ ,  $R^2 = 0.01$ ,  $p\text{-value} = 0.92$ ; Spearman Rho:  $0.56x + 0.091$ ,  $R^2 = 0.03$ ,  
 652  $P = 0.63$ , AVG\_D:  $y = -0.09x + 0.52$ ,  $R^2 = 0.01$ ,  $P = 0.84$ ; Spearman Rho:  $0.05x + 0.20$ ,  $R^2 = 0.00$ ,  $P = 0.79$ . GPUL:



653 *Gammarus pulex*; HYDROP: *Hydropsyche contubernalis*; TANYT: *Tanytarsini* (tribe); LIMN: *Limnomysis*  
654 *benedeni*; OVIL: *Orectochilus villosus*; VCRI: *Valvata cristata*; PISI: *Pisidium* (genus); TIPUL: *Tipula*  
655 (genus). See table 1 and 2 for other taxa names.

656

657

658




Figure 1.

659

660






Figure 2.


(A) Relationship among **present hydraulic variables**



(B) Relationship among **past hydraulic variables**



(C) Relationship **between past velocity and present hydraulic variables**



(D) Relationship **between past dewatering and present hydraulic variables**





Figure 3.

(A) M2 models relating guild abundance to present hydraulics and including an additive effect of dewatering



(B) M2 models relating guild abundance to present hydraulics and including an additive effect of past velocity






Figure 4.

## Electrofishing data



663

664

Figure 5

Table 1. Fish guilds, species, taxa codes, with total abundance and occurrence in microhabitats.

Dataset	Guild	Family	Scientific name	Common name	Taxa code	Abundance	Occurrence
Electrofishing	Bank N=2344	Cyprinidae	<i>Phoxinus phoxinus</i>	Minnow	PhP_cl12	1785	134
		Cyprinidae	<i>Squalius cephalus</i>	Chub	SqC_cl1	217	47
		Cyprinidae	<i>Rutilus rutilus</i>	Roach	RuR_cl1	122	15
		Balitoridae	<i>Barbatula barbatula</i>	Stone loach	BaBa_cl12	116	57
		Cyprinidae	<i>Telestes soufia</i>	Blageon	TeS_cl1	57	17
		Cyprinidae	<i>Gobio gobio</i>	Gudgeon	GoG_cl1	30	16
		Percidae	<i>Perca fluviatilis</i>	Perch	PER_cl12	11	11
	Centrarchidae	<i>Lepomis gibbosus</i>	Pumpkinseed	LeG_cl123	6	5	
	Midstream N=73	Cyprinidae	<i>Barbus barbus</i>	Barbel	BaBu_cl34	27	17
		Salmonidae	<i>Thymallus thymallus</i>	Grayling	ThT_cl2	15	10
		Cyprinidae	<i>Chondrostoma nasus</i>	Nase	ChN_cl34	12	7
		Cyprinidae	<i>Barbus barbus</i>	Barbel	BaBu_cl2	8	8
		Cyprinidae	<i>Alburnus alburnus</i>	Bleak	AlA_cl23	4	5
Salmonidae		<i>Thymallus thymallus</i>	Grayling	ThT_cl4	3	2	
Salmonidae		<i>Thymallus thymallus</i>	Grayling	ThT_cl3	2	3	
Snorkeling	Midstream N= 1119	Salmonidae	<i>Thymallus thymallus</i>	Grayling	ThT_cl234	444	347
		Cyprinidae	<i>Barbus barbus</i>	Barbel	BaBu_cl34	437	164
		Cyprinidae	<i>Chondrostoma nasus</i>	Common nase	ChN_cl34	238	43

665

666

Table 2. Benthic macroinvertebrate guilds, family, genus, taxa codes, with total abundance and occurrence in microhabitats.

Guild	Family	Tribe	Genus	Taxa	Code taxa	Abundance	Occurrence
Limnophilic c N=687	Chironomidae	Chironomini			CHIR	612	23
	Sphaeriidae		<i>Pisidium spp.</i>		PISI	70	15
	Tabanidae				TABA	3	3
	Baetidae (non Baetis)		<i>Procloeon spp.</i>	<i>Procloeon bifidum</i>	PBIF	2	1
Rheophilic N=2988	Elmidae		<i>Esolus spp.</i>		ESOL	2021	34
	Hydropsychidae		<i>Cheumatopsyche spp.</i>	<i>Cheumatopsyche lepida</i>	CLEP	393	14
	Psychomyiidae		<i>Psychomyia spp.</i>	<i>Psychomyia pusilla</i>	PPUS	214	14
	Elmidae		<i>Elmis spp.</i>		ELMI	192	17
	Baetidae		<i>Baetis spp.</i>	<i>Baetis fuscatus</i>	BFUS	102	14
	Hydropsychidae		<i>Hydropsyche spp.</i>	<i>Hydropsyche incognita</i>	HINC	48	10
	Psychodidae				PSYC	7	4
	Heptageniidae		<i>Heptagenia spp.</i>	<i>Heptagenia sulphurea</i>	HSUL	6	3
	Baetidae		<i>Baetis spp.</i>	<i>Baetis vardarensis</i>	BVAR	3	1
	Hydropsychidae		<i>Hydropsyche spp.</i>	<i>Hydropsyche exocellata</i>	HEXO	1	1
	Baetidae		<i>Baetis spp.</i>	<i>Baetis rhodani</i>	BRHO	1	1

667

668



Table 3. Fitting statistics for models M0 (no microhabitat selection), M1 (effects of present hydraulics) and M2 (additional effect of past hydraulics), including AIC and Spearman Rho values. M2 models correspond to those shown in Figure 4. Asterisks indicate significance of likelihood ratio tests comparing M1 vs. M0 or M2 vs. M1, with (\*) for  $P < 0.05$  and (\*\*) for  $P < 0.01$ . In bold: abundance is significantly higher in past high velocity or dewatering habitat; Underlined: abundance is lower.

Model type	Hydraulic variables involved	Dataset									
		Electrofishing data				Snorkeling data		Benthic macroinvertebrate data			
		Bank guild		Midstream guild		Midstream guild		Limnophilic guild		Rheophilic guild	
		AIC	Spea. Rho	AIC	Spea. Rho	AIC	Spea. Rho	AIC	Spea. Rho	AIC	Spea. Rho
M0		1780		392		4108		256		273	
M1	<b>Flow velocity</b>	<b>1670**</b>	0.41	395	0.11	<b>4847*</b>	0.16	258	0.09	<b>370*</b>	0.13
M2	<b>Flow velocity + Dewatering</b>	<b>1650**</b>	0.44	<u>392*</u>	0.14			<u>256*</u>	0.27	<u>353*</u>	0.61
	<b>Flow velocity + Past velocity</b>	1672	0.40	395	0.11	<b>4843*</b>	0.15	259	0.09	<b>369*</b>	0.31
M1	<b>Water depth</b>	<b>1734**</b>	0.43	<b>385**</b>	0.10	<u>4933</u>	0.04	259	0.04	<b>370*</b>	0.35
M2	<b>Water depth + Dewatering</b>	<b>1725**</b>	0.43	386	0.13			<u>258*</u>	0.44	<u>358*</u>	0.53
	<b>Water depth + Past velocity</b>	1734	0.44	384	0.12	<b>4931*</b>	0.03	259	0.15	<b>366*</b>	0.46