HAL
open science

NO-REGRET ALGORITHMS IN ON-LINE LEARNING, GAMES AND CONVEX OPTIMIZATION

Sylvain Sorin

To cite this version:

Sylvain Sorin. NO-REGRET ALGORITHMS IN ON-LINE LEARNING, GAMES AND CONVEX OPTIMIZATION. 2022. hal-03911132

HAL Id: hal-03911132

https://hal.science/hal-03911132v1

Preprint submitted on 22 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NO-REGRET ALGORITHMS IN ON-LINE LEARNING, GAMES AND CONVEX OPTIMIZATION

SYLVAIN SORIN

ABSTRACT. The purpose of this article is to underline the links between some no-regret algorithms used in on-line learning, games and convex optimization and to compare the continuous and discrete time versions.

CONTENTS

1. Introduction 2
1.1. The model: no-regret condition in continuous and discrete time 2
1.2. The analysis 3
1.3. No-regret in on-line learning, games and convex optimization 3
1.4. Summary 6
2. Basic properties of the closed form 6
2.1. Definition and notations 6
2.2. Results 8
3. Continuous time 10
3.1. Level functions and their properties 10
3.2. Positive correlation 11
3.3. Projected dynamics 12
3.4. Mirror descent : differential/incremental approach 16
3.5. Dual averaging: integral/cumulative approach 18
3.6. Comments on the continuous time dynamics framework 19
4. Discrete time: general case 20
4.1. Projected dynamics 21
4.2. Mirror descent 22
4.3. Dual averaging 24

Date: October 2022.

Part of this research has been supported by a PGMO grant (2015): "COGLED - Convergence of gradient-like and evolutionary dynamics", with Jerôme Bolte.

Partial successive versions have been presented at the following events: PGMO Days, October 2015; Tutorial "Topics on strategic learning", IMS, Singapore, November 2015; JFCO, Toulouse, July 2017; GAMENET, Krakow, September 2018; Variational Day, Ecole Polytechnique, December 2018; Operator Splitting Methods in Data Analysis, Flatiron Institute, NY, March 2019; GAMENET, Cluj, April 2019; MAPLE 2019, Politecnico Milano, September 2019; SPOT, Toulouse, January 2020; SPO, Paris, October 2020; One World Optimization Seminar, March 2021.

I would like to thank J. Bolte, P. Combettes, J. Kwon, R. Laraki, V. Perchet, G. Vigeral for nice discussions and interesting comments.

4.4. Comments on the discrete dynamics framework 26
5. Discrete time: Regularity 26
5.1. Projected dynamics 26
5.2. Mirror descent 29
5.3. Dual averaging 30
5.4. Comments on the regular case 31
5.5. Mirror prox 32
6. Concluding remarks 35
References 36

1. Introduction

General learning algorithms associate to an observation a decision and evaluate the result through a specific criteria. On-line procedures involve time hence a process of observations but require that successive decisions are taken at each instant. We will study here an example of such instance where the criteria is a "regret function".
The specificity of the current presentation lies mainly in the two following aspects:
1) We will establish properties for the general observation process and then study the so-called closed model where the observation is a function of the actual decision. A typical example will be the joint process of players choosing actions and observing each their induced private payoff. Another basic application is a first order method in convex optimization.
While these three frameworks seem very different : no hypothesis on the observation in the first case, collection of unilateral procedures in parallel in the second, gradient based dynamics in the third, we will see that general optimal properties are shared and similar main concepts and tools apply.
2) We will consider both the continuous and discrete time versions and compare the basic notions, tools, ideas of proofs and results in both cases. Here also some common properties will emerge.

1.1. **The model: no-regret condition in continuous and discrete time.**

The general framework is as follows:
V is a normed vector space, finite dimensional, with dual V^* and duality map $\langle V^*|V\rangle$, X is a non-empty convex compact subset of V (while this last hypothesis is not needed in convex optimization - but $argminf$ non empty is often required - it is almost necessary in the analysis for games).

The aim is to study properties of algorithms that associate to a continuous time process $\{u_t \in V^*, t \geq 0\}$, a similar procedure $\{x_t \in X, t \geq 0\}$, where x_t is function of the past $\{(x_s, u_s), 0 \leq s < t\}$.
The process corresponds to the observation, the procedure to the induced trajectory of choices.

The adequation of $\{x_t\}$ to $\{u_t\}$ is measured by a *regret function* defined on X by:

$$R_t(x) = \int_0^t \langle u_s | x - x_s \rangle ds \tag{1}$$

which compares the integral of the instantaneous score $\langle u_s | x_s \rangle$ to the one induced by a fixed action $\langle u_s | x \rangle$. (Notice that the regret can be positive or negative for some x.)
One will study procedures satisfying the *no-regret condition*:

$$\sup_{x \in X} R_t(x) \leq o(t) \tag{2}$$

which means that the (positive part of the) time average regret $\frac{R_t(x)}{t}$ vanishes.

Similarly in discrete time, $m \in I\!N$, given $\{u_m\}$ and $\{x_m\}$, with $\{x_m\}$ depending on $\{x_1, u_1, ..., x_{m-1}, u_{m-1}\}$, one defines:

$$R_n(x) = \sum_{m=1}^{n} \langle u_m | x - x_m \rangle$$

and requires:

$$\sup_{x \in X} R_n(x) \leq o(n). \tag{3}$$

1.2. **The analysis.**

A) We compare the performance of the algorithms in terms of evaluation of the regret under three (increasing) assumptions:
(I) *general case*: the information $\{u_t\}$ is a bounded measurable process in V^*,
(II) *closed form*: the information is determined by the choice, thus $u_t = g(x_t)$ for a continuous vector field $g : X \to V^*$,
(III) *convex gradient*: $u_t = -\nabla f(x_t)$, where f is a $\mathcal{C}^1$ convex function : $X \to \mathbb{R}$,
(with similar properties in discrete time).

B) We consider three different procedures:
- Projected dynamics (PD),
- Mirror descent (MD),
- Dual averaging (DA)

that will satisfy the no-regret property and we will discuss their performances.

C) We analyze the relations between the continuous and discrete time processes (in particular in terms of speed of convergence to 0 of the average regret).
Recall that in continuous time u_t is not observed at time t and the information process is only measurable; moreover u_t may depend on x_t.

D) In classes (II) and (III) it makes sense to consider the trajectories of $\{x_t\}$ or $\{x_n\}$ and we will study their convergence.

1.3. **No-regret in on-line learning, games and convex optimization.**

1.3.1. Framework (I) corresponds to the usual model of on-line learning where at each time t the agent observes the process $\{u_s, s < t\}$ and chooses the action x_t.
Note that, for K a finite set with cardinal $|K|$, if $V = \mathbb{R}^{|K|}$ and the agent selects a component $k_t \in K$ at random, then X is the simplex on K, denoted $\Delta(K)$, x_t stands for the law of k_t and the regret is expressed in terms of conditional expectation.
Recall that since no hypothesis is made on the process u_t, no prediction on its future values makes sense, but to satisfy the no-regret criteria expresses a desirable a-posteriori property in the spirit of dynamic best-reply.

The litterature on this topic is very large and almost impossible to cover. Let us mention few aspects.

The notion of regret appears in Hannan (1957) [40], Blackwell (1956) [15] in a game theoretical set-up. Algorithms and properties are studied in this spirit in Foster and Vohra (1993) [31], Fudenberg and Levine (1995) [36], Foster and Vohra (1999) [34], Hart and Mas-Colell (2000) [42], Lehrer (2003) [60], Benaim, Hofbauer and Sorin (2005)[14], Cesa-Bianchi and Lugosi (2006) [25], Viossat and Zapechelnyuk, 2013 [102], ... among others.

Similar tools and properties occur in statistics and in the learning community: Vvok (1990) [103], Cover (1991) [28], Littlestone and Warmuth (1994) [63], Freund and Shapire (1999) [35], Auer P., Cesa-Bianchi N., Freund Y. and R.E. Shapire (2002) [7], Cesa-Bianchi and Lugosi (2003) [24], Stoltz and Lugosi (2005) [96], Kalai and Vempala, 2005 [53], Blum and Mansour, 2007 [16], ...

This topic is analyzed in the following books:
FUDENBERG D. AND D. LEVINE (1998) [37] *The Theory of Learning in Games*, MIT Press,
CESA-BIANCHI N. AND G. LUGOSI (2006) [25] *Prediction, Learning and Games*, Cambridge University Press,
HART S. AND A. MAS-COLELL (2013) [45] *Simple Adaptive Strategies: From Regret-Matching to Uncoupled Dynamics*, World Scientific Publishing,
and the connection with related notions of approachability and consistency is well presented in the survey by Perchet (2014) [79], see also Abernethy, Bartlett and Hazan (2011) [1] .

The next two paragraphs describe more specific cases where the observation u_t is a function of the action x_t.

1.3.2. In particular, framework (II) *closed form* is relevant for game dynamics and variational inequalities as explained now.
Consider a strategic game with a finite set of players I, where $X^i \subset V^i$ is the strategy set of player i and $X = \prod_i X^i$. Assume that the equilibrium set E is obtained as the set of solutions $x = (x^i) \in X$ of the following variational inequalities:

$$\langle g^i(x)|x^i - y^i\rangle \geq 0, \qquad \forall y^i \in X^i, \forall i \in I.$$

where $g^i : X \to V^{i*}$ is the "evaluation" function of player i.
Examples include (mixed extension of) finite games where g^i is the "vector payoff" function of player i, games with smooth concave payoffs (the payoff function $F^i : X \to \mathbb{R}$ of player i is concave w.r.t. x^i and $g^i = \nabla_{x_i} F^i$), but also nonatomic population

games and Wardrop equilibria (I is then the set of populations), see Sorin and Wan (2016) [95].
Notice that an equivalent formulation of the equilibrium condition is through the single variational inequality, with $g = \{g^i\}$:

$$\langle g(x)|x-y\rangle = \sum_i \langle g^i(x)|x^i - y^i\rangle \geq 0, \qquad \forall y \in X. \tag{4}$$

We will denote by $\Gamma(g)$ a game for which equilibria are solution of (4).
For each player i, the reference/observation process is specific, namely $u_t^i = g^i(x_t)$ which, as a function of x_t, is determined by the behavior of all players. In particular, in a concave smooth game, the observation of player i is the gradient w.r.t. x_i : $u_t^i = \nabla_{x_i} F^i(x_t)$.
Note that the overall global dynamics of the profile of actions $\{x_t\}$ is generated by a family of unilateral procedures since for each i, x_t^i depends on (u^i, x^i) only. In particular for each player i, the knowledge of $g^j, j \neq i$ is not assumed. Thus for each participant individually the situation is similar to (I) *general case* since the observation is not controlled, while these private observations of the participants are linked via x_t.

We will analyze the consequences on the process $\{x_t\}$ assuming only that each player uses a procedure satisfying the no-regret condition (2) or (3). Obviously the (global) algorithm associated to $g = \{g^i\}$ as a vector field on a product space associated to (4) will also share the no-regret property since:

$$\int_0^t \langle g^i(x_s)|x^i - x_s^i\rangle ds \leq o(t), \qquad \forall x^i \in X^i$$

implies:

$$\int_0^t \langle g(x_s)|x - x_s\rangle ds \leq o(t), \qquad \forall x \in X$$

but in addition it is "decentralized" in the sense that x^i depends upon g^i only.

1.3.3. Framework (III) covers the case of convex optimization where the observation, given the choice x_t, is the gradient of the (unknown) convex function, explicitly one has $u_t = -\nabla f(x_t)$.
The recent research in this area is very wide and links basic optimisation algorithms Polyak (1987) [81], Nemirovski and Yudin (1983) [73], Nesterov (2004) [75], to on-line procedures, see Zinkevich (2003) [106].
Let us mention the recent books:
BUBECK S. (2015) [22] Convex optimization: Algorithms and complexity, *Fondations and Trends in Machine Learning*, **8**, 231-357.
HAZAN E. (2011) [46] The convex optimization approach to regret minimization, *Optimization for machine learning*, S. Sra, S. Nowozin, S. Wright eds, MIT Press, 287-303.
HAZAN E. (2015) [47] Introduction to Online Convex Optimization, *Fondations and Trends in Optimization*, **2**, 157-325.
HAZAN E. (2019) [48] Optimization for Machine Learning , https://arxiv.org/pdf/1909.03550.pdf.

SHALEV-SHWARTZ S. (2012) [90] Online Learning and Online Convex Optimization, *Foundations and Trends in Machine Learning*, **4** , 107-194.

1.3.4. One should add that related algorithms have been developped in Operations Research (transportation, networks), see e.g. Harker and Pang (1990) [39], Dupuis and Nagurney (1993) [29], Nagurney and Zhang (1997) [71].
Note that each community (learning, game theory, optimization) has its own terminology and point of view. As a consequence specific properties may inherit different names and will be rediscovered at several occasions. One of the aim of the current work is to clarify the relations between these approaches and results and unify the analysis.
In particular we will show that few basic principles are in use and we will underline the analogy between continuous and discrete time.
However there are subtle differences between the different algorithms and their properties that we will discuss (see 3.6, 4.4 and 5.4).

1.4. **Summary.**

Section 2 is devoted to the *closed form* framework (II) and explores the links between no-regret, solutions of variational inequalities and convex optimization.
Section 3 deals with continuous time dynamics. After introducing "level functions" and "positive correlation" we describe the three algorithms (PD, MD, DA), prove that they satisfy the no-regret property and compare their performances.
Section 4 is the discrete time analog of Section 3.
Section 5 considers basically framework (III) under an additional regularity hypothesis on the convex function f. Subsection 5.4 on "Mirror prox" recalls related results using similar tools.
Concluding comments are in Section 6.

2. BASIC PROPERTIES OF THE CLOSED FORM

We describe first some relations with variational inequalities, when the observation process has a *closed form* : $u = g(x)$. Then we exhibit general properties of no-regret procedures.

2.1. **Definition and notations.**

2.1.1. *iS: set of internal solutions.*

Notation: iS is the set of solutions $x \in X$, (internal since it involves the value of g at $x \in X$), of the variational inequality:

$$\langle g(x)|y-x\rangle \leq 0, \qquad \forall y \in X. \tag{5}$$

Note that if g is associated to a game $\Gamma(g)$, see 1.3.2, (4) shows that iS corresponds to the set of equilibria.

Recall also that in an Hilbertian framework property (5) is equivalent to:

$$\Pi_X(x+\lambda g(x)) = x, \qquad \lambda > 0 \tag{6}$$

where Π_C denotes the (Hilbertian) projection operator on a closed convex set C, and (5) defines also the solutions $x \in X$ of:

$$\Pi_{TX(x)}(g(x)) = 0 \tag{7}$$

where $TX(x)$ is the tangent cône to X at x, see e.g. Kinderlehrer and Stampacchia (1980) [54], Facchinei and Pang (2007) [30].

The minimization of a $\mathcal{C}^1$ convex function f on X corresponds to the variational inequality (5) with $g = -\nabla f$.
This case presents two specific properties: i) g is a gradient, ii) g is dissipative.
We will consider now the analogous general definitions for vector fields.

2.1.2. *Fields with potential.*

Definition: $g : X \to V^*$ is a vector field with *potential* G, if $G : X' \to \mathbb{R}$, where X' is an open neighborhood of X, satisfies:

$$\langle \nabla G(x) - g(x) | y - x \rangle = 0, \forall x, y \in X$$

or more generally if there exist a strictly positive function μ on X, such that:

$$\left\langle \nabla G(x) - \mu(x) g(x), y - x \right\rangle = 0, \quad \forall x \in X, \forall y \in X, \tag{8}$$

see Sorin and Wan (2016) [95].
Then local maxima of G on X belong to iS and if G is concave its maxima on X coincide with iS.
If g is defined through a game $\Gamma(g)$, see 1.3.2, the game is a *potential game*, Monderer and Shapley (1996) [69], Sandholm (2001) [85].

2.1.3. *Dissipative fields.*

Definition: g is *dissipative* ($-g$ is *monotone*) if it satisfies:

$$\langle g(x) - g(y) | x - y \rangle \leq 0, \qquad \forall x, y \in X. \tag{9}$$

A fundamental example is the following: let F, defined on $X = X_1 \times X_2$, be a $\mathcal{C}^1$ concave/convex function, (like the payoff function of a two-person 0-sum game, $F = F^1 = -F^2$), then $g = \{\nabla_{x_1} F, -\nabla_{x_2} F\}$ is dissipative, Rockafellar (1970), [83].

If g is dissipative and defined through a game $\Gamma(g)$, see 1.3.2, the game is dissipative, as introduced by Rosen (1965) [84]. The terminology is "stable" in Hofbauer and Sandholm (2009) [50], "contractive" in Sandholm (2015) [87] and "dissipative" in Sorin and Wan (2016) [95].

It will be usefull to consider the following set.

2.1.4. *eS: set of external solutions.*

Notation: eS is the set of solutions $x \in X$ (external since it involves the values of g at alternative points $y \in X$), of the variational inequality:

$$\langle g(y) | y - x \rangle \leq 0, \qquad \forall y \in X. \tag{10}$$

Observe that eS is either convex or empty, independently on the properties of g.
If g is dissipative, $eS \neq \emptyset$ follows from the - finite version of the - minmax theorem, Minty, 1967 [68].

2.1.5. *Internal vs external solutions.*

Let us recall the relations between internal and external solutions:
If g is dissipative, the following (weak) inclusion holds :

$$iS \subset eS$$

and if g is continuous (or simply $t \mapsto g(x + t(y - x))$ continuous in $t, \forall x, y \in X$), the reverse (weak) inclusion is satisfied:

$$eS \subset iS$$

see Kinderlehrer and Stampacchia (1980) [54], Facchinei and Pang (2007) [30].
Thus, if g defined through a game $\Gamma(g)$, see 1.3.2, is dissipative and continuous then $iS = eS = E$; in particular for a smooth zero-sum game this corresponds to the set of optimal strategies.
If $iS = eS$ we will also use the notation S for this set.

2.2. **Results.**

Assume in this subsection that the procedure x_t or x_n satisfies the no-regret property (2) or (3).

2.2.1. *iS and convergence.*
A first property deals with convergent trajectories $\{x_t\}$.

Proposition 1.
If g is continuous and x_s converges to x, then $x \in iS$.

Proof:
Since $R_t(y) = \int_0^t \langle g(x_s)|y - x_s\rangle ds$:

$$\frac{R_t(y)}{t} \to \langle g(x)|y - x\rangle, \qquad \forall y \in X. \tag{11}$$

and $R_t(y) \leq o(t)$ implies $x \in iS$. ∎

In particular, if x is a *stationary point* for the discrete or continuous time procedure, i.e. $x_t = x$ if $x_0 = x$, then $x \in iS$.

Concerning the set eS, clearly one has:

Lemma 1.
If $x^* \in eS$, then $R_t(x^*) \geq 0$ for all $t \geq 0$.

2.2.2. *Time average.*
Define the time average trajectories as follows:

$$\bar{x}_t = \frac{1}{t}\int_0^t x_s ds \qquad \text{and} \qquad \bar{x}_n = \frac{1}{n}\sum_1^n x_m.$$

Proposition 2.
If g is dissipative, the accumulation points of $\{\bar{x}_t\}$ or $\{\bar{x}_n\}$ are in eS.

Proof:

$$\frac{R_t(y)}{t} = \frac{1}{t}\int_0^t \langle g(x_s)|y-x_s\rangle \geq \frac{1}{t}\int_0^t \langle g(y)|y-x_s\rangle = \langle g(y)|y-\bar{x}_t\rangle.$$

Hence under (2) or (3) any accumulation point $\hat{x}$ of $\{\bar{x}_t\}$ will satisfy $\langle g(y)|y-\hat{x}\rangle \leq 0$. ∎

Note that this result implies the non-emptiness of eS for dissipative g.
In particular the minmax theorem (in the $\mathcal{C}^1$ case) follows from the existence of no-regret procedures. This is in the spirit of the previous proofs based on dynamics: e.g. Brown and von Neumann (1950) [20], Hofbauer and Sorin (2006) [52], Lehrer and Sorin (2007) [61], Hofbauer (2018) [49].

2.2.3. *Class (III).*
Since $u_t = g(x_t) = -\nabla f(x_t)$ with f being $\mathcal{C}^1$ and convex, this corresponds to a specific case of dissipative and continuous vector field g, hence $eS = iS = S = \operatorname{argmin}_X f$.
Use the basic convexity inequality:

$$\langle \nabla f(x_t)|y-x_t\rangle \leq f(y) - f(x_t)$$

to deduce with $u_t = -\nabla f(x_t)$ in (1):

$$\int_0^t [f(x_s) - f(y)]\, ds \leq \int_0^t \langle -\nabla f(x_s)|y-x_s\rangle ds = R_t(y)$$

which implies by Jensen's inequality:

$$f(\bar{x}_t) - f(y) \leq \frac{1}{t}\int_0^t [f(x_s) - f(y)]\, ds \leq \frac{R_t(y)}{t}. \tag{12}$$

Similarly in discrete time with $u_m = -\nabla f(x_m)$:

$$n[f(\bar{x}_n) - f(y)] \leq \sum_{m=1}^n f(x_m) - f(y) \leq \sum_{m=1}^n \langle \nabla f(x_m)|x_m - y\rangle = R_n(y).$$

In particular one obtains:

Proposition 3.
i) The accumulation points of $\{\bar{x}_t\}$ or $\{\bar{x}_n\}$ belong to S.
ii) If $t \mapsto f(x_t)$ (resp. $n \mapsto f(x_n)$) is decreasing, the accumulation points of $\{x_t\}$ or $\{x_n\}$ belong to S.

Note that i) is a particular instance of Proposition 2.

One can also deal with the case $u_m = -\lambda_m \nabla f(x_m)$ with $\lambda_m \geq 0, \sigma_m = \sum_{k\leq m} \lambda_k$ and $\sigma_n \hat{x}_n = \sum_{m\leq n} \lambda_m x_m$ to obtain:

$$\sigma_n[f(\hat{x}_n) - f(y)] \leq \sum_{m\leq n} \lambda_m [f(x_m) - f(y)] \leq \sum_{m\leq n} \langle \lambda_m \nabla f(x_m)|x_m - y\rangle \leq R_n(y).$$

This allows to compare the regret for discrete and continuous time trajectories, see Kwon and Mertikopoulos (2017) [58], Section 6.2.

3. Continuous time

We describe in this section three procedures in continuous time that satisfy the no-regret property. Their discrete time counterparts will be analyzed in the next section. As usual, discrete time dynamics are easier to describe but their mathematical properties are more difficult to establish. This explains why we choose to start with the continuous time versions.
In addition a very useful tool in the form of a "level function" is available in this set-up and we start by analyzing it.

3.1. **Level functions and their properties.**

Definition: $P : \mathbb{R}^+ \times X \to \mathbb{R}^+$ is a *level function* (for $\{u_t, x_t\}$) if $P(t; y)$ is bounded and satisfies:

$$\langle u_t, x_t - y\rangle \geq \frac{d}{dt}P(t; y), \quad \forall t \in \mathbb{R}^+, \forall y \in X. \tag{13}$$

Note that P is not defined on the trajectory $\{x_t\}$ alone, but is a function of the joint processes $\{u_t, x_t\}$.

Proposition 4.
If there exists a level function, R_t is upper bounded; hence $R_t(y)/t \leq O(1/t)$.

Proof:

$$R_t(y) = \int_0^t \langle u_s | y - x_s\rangle ds \leq P(0; y) - P(t; y) \leq P(0; y).$$

■

Hence the existence of a level function implies the no-regret property.

Proposition 5.
Consider class (II).
Assume $y^* \in eS$, then $t \mapsto P(t; y^*)$ is decreasing.

Proof:

$$\frac{d}{dt}P(t; y^*) \leq \langle g(x_t) | x_t - y^*\rangle \leq 0.$$

■

Thus a level function evaluated at an external solution in eS is a weak Lyapounov function.

Lemma 2.
Consider class (III).
If $\{x_t\}$ is a descent procedure (meaning that $\frac{d}{dt}f(x_t) \leq 0$), then:

$$E(t; y) = t(f(x_t) - f(y)) + P(t; y)$$

is decreasing, for all $y \in X$.

Proof:

$$\begin{aligned}\frac{d}{dt}E(t;y) &= f(x_t) - f(y) + t\frac{d}{dt}f(x_t) + \frac{d}{dt}P(t;y)\\ &\leq f(x_t) - f(y) + \langle \nabla f(x_t)|y - x_t\rangle \leq 0\end{aligned}$$

■

We recover the fact that the accumulation points of $\{x_t\}$ are in $S = \mathrm{argmin}_X\ f$ and that the speed of convergence of $f(x_t)$ to $\min f$ is $O(1/t)$.

3.2. **Positive correlation.**

Given a first order dynamics of the form $\dot{x}_t = A(x_t)$, f decreases on trajectories if:

$$\frac{d}{dt}\, f(x_t) = \langle \nabla f(x_t)|\dot{x}_t\rangle \leq 0.$$

The analogous property for a vector field g is:

$$\langle g(x_t)|\dot{x}_t\rangle \geq 0. \tag{14}$$

In the framework of games, a similar condition was described in discrete time as Myopic Adjustment Dynamics, Swinkels (1993) [99] : if $x^i_{n+1} \neq x^i_n$ then $G^i(x^i_{n+1}, x^{-i}_n) > G^i(x^i_n, x^{-i}_n), i \in I$.

The corresponding property in continuous time corresponds to *positive correlation*, (between the dynamics and the vector field), Sandholm (2010) [86]:

$$\dot{x}^i_t \neq 0 \Longrightarrow \langle g^i(x_t)|\dot{x}^i_t\rangle > 0, i \in I. \tag{15}$$

The use of this notion for vector fields with potential is a follows:

Proposition 6.
Consider a vector field g with potential G.
If the dynamics satisfies positive correlation, then G is a strict (increasing) Lyapunov function.
All ω-limit points are rest points.

Proof:
Let $V_t = G(x_t)$ for $t \geq 0$. Then:

$$\dot{V}_t = \langle \nabla G(x_t)|\dot{x}_t\rangle = \sum_{i\in I}\langle \nabla_{x_i} G(x_t)|\dot{x}^i_t\rangle = \sum_{i\in I}\mu^i(x)\langle g^i(x_t)|\dot{x}^i_t\rangle \geq 0.$$

Moreover $\langle g^i(x_t)|\dot{x}^i_t\rangle = 0, \forall i \in I$, holds if and only if $\dot{x}_t = 0$.

One concludes by using Lyapunov's theorem (e.g. Theorem 2.6.1 in [51]). ■

This result is proved by Sandholm (2001) [85] for his version of potential population game, see extensions in Benaim, Hofbauer and Sorin (2005) [14].

A similar property for fictitious play in discrete time is established in Monderer and Shapley (1996) [69].

We will show that this property (15) holds for the three dynamics defined below.

We now introduce and study three dynamics:
- Projected dynamics (PD),
- Mirror descent (MD),
- Dual averaging (DA).
In each case we first define the dynamics, then control the evaluation of the regret by exhibiting a level function and finally study the trajectories for class (II) and (III).

3.3. **Projected dynamics.**

We assume in this subsection that $V = V^*$ is an Euclidean space with scalar product denoted by $\langle , \rangle$.

3.3.1. *Dynamics.*
The continuous time analog of the generalization of the *Projected Gradient Descent*, Levitin and Polyak (1966) [62], Polyak (1987) [81], see also Section 4.1, is defined by $x_t \in X$ satisfying:

$$\boxed{\langle u_t - \dot{x}_t, y - x_t \rangle \leq 0, \forall y \in X. \qquad (16)}$$

which is:

$$\dot{x}_t = \Pi_{T_X(x_t)}(u_t) \qquad (17)$$

since $T_X(x_t)$ is a cône.

3.3.2. *Values.*
Let:

$$V(t; y) = \frac{1}{2}\|x_t - y\|^2, \quad y \in X. \qquad (18)$$

Proposition 7.
V is a level function.

Proof:
One has:

$$\frac{d}{dt}V(t; y) = \langle \dot{x}_t, x_t - y \rangle \leq \langle u_t, x_t - y \rangle$$

by (16). ∎

Thus the properties of section 3.1 hold.

3.3.3. *Trajectories.*
• Consider class (II) : $u_t = g(x_t)$.
One has the following convergence result:

Proposition 8.
Assume g dissipative.
Then $\{\bar{x}_t\}$ converges to a point in eS.

Proof:
- The limit points of $\{\bar{x}_t\}$ are in eS by Proposition 2.
- $\|x_t - y^*\|$ converges when $y^* \in eS$ by Propositions 5 and 7.
Hence using Opial's Lemma (1967) [78], which states:

(19) *In an Hilbert space, if for any weak accumulation point $\hat{x}$ of $\{x_t\}$ (resp. $\{\bar{x}_t\}$), $\|x_t - \hat{x}\|$ has a limit as $t \to \infty$, then $\{x_t\}$ (resp. $\{\bar{x}_t\}$) weakly converges.*

it follows that $\bar{x}_t$ converges to a point in eS. ∎

Proposition 9.
Positive correlation holds.

Proof:

$$\langle g(x_t), \dot{x}_t \rangle = \|\dot{x}_t\|^2$$

since $u_t = g(x_t)$ and $\langle u_t - \dot{x}_t, \dot{x}_t \rangle = 0$, by Moreau's decomposition, Moreau (1965) [70].
∎

• Consider class (III) : $u_t = -\nabla f(x_t)$.

Proposition 10.
i) $\{x_t\}$ converges to a point in S.
ii) $f(x_t)$ decreases to $\min f$ with speed $O(1/t)$.

Proof:
i) Both Proposition 3 or Lemma 2, and Proposition 9 imply that the accumulation points of $\{x_t\}$ are in S. Then using Proposition 5, Opial's Lemma (19) applies.
ii) Follows from Lemma 2. ∎

3.3.4. *Hilbert case.*

We assume in this subsection that V is an Hilbert space with scalar product $\langle , \rangle$ and that $X \subset V$ is non-empty, convex and closed.
The results of Section 2 and 3.1 will extend, while considering weak accumulation points and assuming eS non empty.

• Consider class (II).

Lemma 3.
Assume $eS \neq \emptyset$. Then the trajectory $\{x_t\}$ is bounded.

Proof:
By Proposition 5, for $y^* \in eS$, $V(t; y^*)$ is decreasing. ∎

In particular this implies the following convergence result:

Proposition 11.
Assume $eS \neq \emptyset$ and g dissipative.
Then $\{\bar{x}_t\}$ converges weakly to a point in eS.

Proof:
- $\{\bar{x}_t\}$ is bounded by Lemma 3 hence has weak accumulation points.
Then follow the proof of Proposition 8. ■

Recall that if g is dissipative and X is bounded, $eS \neq \emptyset$ by Lemma 2.

• Consider class (III).
With a proof as above, Proposition 12 extends to:

Proposition 12.
Assume $S \neq \emptyset$.
i) $\{x_t\}$ weakly converges to a point in S.
ii) $f(x_t)$ decreases to $\min f$ with speed $O(1/t)$.

• Maximal monotone operators.

The preceeding results show strong links with properties of maximal monotone operators, see e.g. Brezis (1973) [19].
Recall that A is a monotone operator on V if $Ax \subset V$ and

$$\langle u - v, x - y \rangle \geq 0, \quad \forall x \in V, u \in Ax, y \in V, v \in Ay.$$

A is a maximal monotone operator if in addition its graph $G_A = \{(x, u); x \in V, u \in Ax\}$ is maximal for the inclusion among monotone operators.
If A is a maximal monotone operator with domain $\{x \in V; Ax \neq \emptyset\} = \text{dom } A$, the set of external solutions associated to $X = \text{dom } A$ and $-A$, that will play the role of eS, is the set T with:

$$T = A^{-1}(0) = \{x \in V; 0 \in Ax\} = \{x \in V; \langle Ay, y - x \rangle \geq 0, \forall y \in \text{dom } A\} \tag{20}$$

where as usual $\langle Ay, z \rangle \geq 0$ means $\langle u, z \rangle \geq 0, \ \forall u \in Ay$.
Let $w_t \in \text{dom } A$ be an absolutely continuous trajectory that satisfies a.e. (for the existence, see Brezis (1973) [19]) :

$$\dot{w}_t \in -Aw_t, t \in \mathbb{R}^+; w_0 \in \text{dom } A. \tag{21}$$

The following result is due to Baillon and Brezis (1976) [10] and is the exact analog of Proposition 11 in terms of statements and proof.

Proposition 13.
Assume $T \neq \emptyset$ and A maximal monotone. Then $\{\bar{w}_t\}$ converges weakly to a point in T.

Proof:
Let $z \in \text{dom } A$. Then :

$$\frac{d}{dt}\|w_t - z\|^2 = \langle \dot{w}_t, w_t - z \rangle \in \langle -Aw_t, w_t - z \rangle \tag{22}$$

Hence if $z^* \in T$:

$$\frac{d}{dt}\|w_t - z^*\|^2 \leq 0 \tag{23}$$

and $\|w_t - z^*\|^2$ is decreasing, thus $\{w_t\}$ is bounded hence has weak accumulation points. From monotonicity:

$$\frac{d}{dt}\|w_t - z\|^2 \in \langle -Aw_t, w_t - z\rangle \leq \langle Az, z - w_t\rangle \tag{24}$$

one deduces:

$$\langle -Az, z - \bar{w}_t\rangle = \frac{1}{t}\int_0^t \langle -Az, z - w_t\rangle \leq \frac{1}{t}\|w_0 - z\|^2 \tag{25}$$

so that any weak accumulation point $\hat{w}$ of $\{\bar{w}_t\}$ satisfies :

$$\langle -Az, z - \hat{w}\rangle \leq 0, \quad \forall z \in \text{dom } A$$

hence belongs to T.
Thus by Opial's Lemma (19), $\bar{w}_t$ converges weakly to a point in T. ∎

Similarly the following result, due to Bruck (1975) [21], and its proof, corresponds to Proposition 12. Let $f : V \to \mathbb{R} \cup \{+\infty\}$ be convex, l.s.c. and proper. A is the subdifferential ∂f and we consider the dynamics (21).

Proposition 14.
i) $f(w_t)$ decreases to $\inf_V f$ with speed $O(1/t)$.
ii) If $T = \text{argmin}_V f \neq \emptyset$, $\{w_t\}$ weakly converges to a point in T.

Proof:
The subdifferential inequality writes :

$$f(w_s) - f(w_t) \geq \langle -\dot{w}_t, w_s - w_t\rangle$$

so that :

$$\limsup_{s \to t^-} \frac{f(w_t) - f(w_s)}{t - s} \leq -\|\dot{w}_t\|^2$$

hence $f(w_t)$ is decreasing.
It follows that for any $y \in V$ and $s \in [0, t]$:

$$f(y) \geq f(w_s) + \langle \dot{w}_s, w_s - y\rangle \geq f(w_t) + \frac{1}{2}\frac{d}{ds}\|w_s - y\|^2$$

hence by integration :

$$tf(y) \geq tf(w_t) + \frac{1}{2}\|w_t - y\|^2 - \frac{1}{2}\|w_0 - y\|^2$$

so that :

$$f(w_t) \leq f(y) + \frac{1}{2t}\|w_0 - y\|^2, \quad \forall y \in V.$$

This gives i).
The proof of the previous Proposition 13 and i) implies that the weak accumulation points of $\{w_t\}$ are in T.
Using Opial's Lemma 19 then gives ii). ∎

To compare with Proposition 12 take $f_X = f + \mathbf{1}_X$ with $int(\text{dom} f \cap X) \neq \emptyset$, where $\mathbf{1}_X$ is the indicator function of X.

3.4. **Mirror descent : differential/incremental approach.**

We study here the continuous version of the extension of the *mirror descent algorithm* studied in convex optimization, Nemirovski and Yudin (1983) [73], Beck and Teboulle (2003) [13], see also section 4.2.

3.4.1. *Dynamics.*

The assumptions are:
H is a strictly convex, $\mathcal{C}^1$ function from V to $\mathbb{R} \cup \{+\infty\}$.
$X \subset V$ is nonempty, compact, convex and $X \subset \text{dom} H$.
The continuous time procedure satisfies $x_t \in X$ with:

$$\langle u_t - \frac{d}{dt}\nabla H(x_t) | y - x_t \rangle \leq 0, \forall y \in X. \tag{26}$$

Recall that the *Bregman distance* associated to H is:

$$D_H(y, x) = H(y) - H(x) - \langle \nabla H(x) | y - x \rangle \, (\geq 0). \tag{27}$$

3.4.2. *Values.*

The use of the Bregman distance is the following:

Proposition 15.
$P(t; y) = D_H(y, x_t)$ is a level function.

Proof:
Note the relation:

$$\frac{d}{dt} D_H(y, x_t) = -\langle \nabla H(x_t) | \dot{x}_t \rangle - \frac{d}{dt} \langle \nabla H(x_t) | y - x_t \rangle = \langle \frac{d}{dt} \nabla H(x_t) | x_t - y \rangle \tag{28}$$

so that (26) implies

$$\frac{d}{dt} D_H(y, x_t) \leq \langle u_t | x_t - y \rangle. \tag{29}$$

■

Hence the properties of section 3.1 apply.

The previous analysis of Section 3.3 corresponds to the Euclidean case with the regularization function:

$$H(x) = \frac{1}{2} \|x\|^2$$

for the dynamics and the level function.

3.4.3. *Interior trajectory.*

The use of a specfic function H adapted to X, with $domH = X$, H $\mathcal{C}^2$ on $intX$ and $\|\nabla H(x)\| \to +\infty$ as $x \to \partial X$ allows to produce a trajectory that remains in $intX$.
In this case (26) leads to an equality:

$$\frac{d}{dt}\nabla H(x_t) = u_t \tag{30}$$

thus:

$$\nabla H(x_t) = \int_0^t u_s ds \tag{31}$$

or, with H^* being the Fenchel conjugate of H:

$$x_t = \nabla H^*(\int_0^t u_s ds) \tag{32}$$

and then:

$$\dot{x}_t = \nabla^2 H(x_t)^{-1} u_t. \tag{33}$$

$\nabla^2 H(x)$ induces a Riemannian metric, see Alvarez, Bolte and Brahic (2004) [2], Mertikopoulos and Sandholm (2018) [66].

In this framework one has a monotonic algorithm for a vector field g with potential G, since (recall 3.2.):

Proposition 16.
Positive correlation holds.

Proof:

$$\langle g(x_t)|\dot{x}_t\rangle = \langle g(x_t)|\nabla^2 H(x_t)^{-1} g(x_t)\rangle \geq 0.$$

since H is convex and $\dot{x}_t = \nabla^2 H(x_t)^{-1} g(x_t) \neq 0$ implies $\langle g(x_t)|\nabla^2 H(x_t)^{-1} g(x_t)\rangle > 0$.

■

Consider now class (III).
By Proposition 3, the accumulation points of $\{x_t\}$ are in S.
To prove convergence one introduces the following :
Hypothesis [H1]: if $z^k \to y^* \in S$ then $D_H(y^*, z^k) \to 0$.
For example H L-smooth (see e.g. Nesterov (2004) [75] Section 1.2.2.) and then:

$$0 \leq D_H(x, y) \leq \frac{L}{2}\|x - y\|^2.$$

Hypothesis [H2]: if $D_H(y^*, z^k) \to 0, y^* \in S$ then $z^k \to y^*$.
For example H β-strongly convex (see e.g. Nesterov (2004) [75] Section 2.1.3.) and then:

$$D_H(x, y) \geq \frac{\beta}{2}\|x - y\|^2.$$

Proposition 17.
Consider class (III). If H is smooth and strongly convex or more generally if [H1] and [H2] hold, $\{x_t\}$ converges to some $x^* \in S$.

Proof:
Let x^* be an accumulation point of $\{x_t\}$. Then $x^* \in S$ by Proposition 3 and thus $D_H(x^*, x_t)$ is decreasing by Proposition 5 and Proposition 15. Since this sequence is decreasing to 0 on a subsequence $x_{t_k} \to x^*$ by [H1], it is decreasing to 0, hence by [H2] $x_t \to x^*$. ∎

3.5. **Dual averaging: integral/cumulative approach.**

We consider here the continuous version of the extension of *dual averaging*, Nesterov (2009) [76], see also section 4.3.
We follow the analysis and results in Kwon and Mertikopoulos (2017) [58].

3.5.1. *Dynamics.*
The assumption is:
the regularization function h is a bounded strictly convex l.s.c. function from V to $\mathbb{R} \cup \{+\infty\}$ with $\operatorname{dom} h = X$.
Let $h^*(w) = \sup_{x \in V} \langle w|x\rangle - h(x)$ be the Fenchel conjugate of h. h^* is differentiable since h is strictly convex.
Introduce the integral:

$$U_t = \int_0^t u_s ds$$

and let us define x_t by:

$$x_t = argmax\{\langle U_t|x\rangle - h(x); x \in V\} = argmax\{\langle U_t|x\rangle - h(x); x \in X\}.$$

The dynamics can be written as:

$$\boxed{x_t = \nabla h^*(U_t) \in X. \qquad (34)}$$

3.5.2. *Values.*
Define, for $y \in X$:

$$W(t; y) = h^*(U_t) - \langle U_t|y\rangle + h(y) \qquad (35)$$

which corresponds to the Fenchel coupling between the cumulative input $U_t \in V^*$ and a reference point $y \in X \subset V$.

Proposition 18.
W is a level function.

Proof:
$W(t; y) \geq 0$ by Fenchel inequality.
Use that:

$$\frac{d}{dt} h^*(U_t) = \langle u_t|\nabla h^*(U_t)\rangle = \langle u_t|x_t\rangle \qquad (36)$$

by (34), thus:

$$\frac{d}{dt} W(t; y) = \langle u_t|x_t - y\rangle.$$

■

In particular one has:

$$R_t(y) = \int_0^t \langle u_s | y - x_s \rangle ds = W(0;y) - W(t;y) \leq [-\inf_X h + h(y)] \leq r_X(h) \tag{37}$$

with $r_X(h) = \sup_X h(x) - \inf_X h(x)$.

Note that due to the integral formulation of the dynamics (34) (x_t as a function of U_t) compared to the differential formulation (26) ($\dot{x}_t$ as a function of u_t) the level function is expressed through the dual space, however properties of section 3.1 applies as well.

3.5.3. *Trajectories.*

Proposition 19.
Positive correlation holds.

Proof:

$$\frac{d}{dt} G(x_t) = \langle \nabla G(x_t) | \dot{x}_t \rangle = \langle g(x_t) | \nabla^2 h^*(U_t)(u_t) \rangle$$

using (34) with $u_t = g(x_t)$. ■

Hence in class (III), using Proposition 3 the accumulation points of $\{x_t\}$ are in S.

3.5.4. *Remarks.*
In the interior smooth case both dynamics and level functions of sections 3.4 and 3.5 are the same, since one has:

$$x_t = \nabla h^*(U_t), \qquad \nabla h(x_t) = U_t, \qquad h^*(U_t) + h(x_t) = \langle U_t | x_t \rangle$$

and

$$\begin{aligned} D_h(y, x_t) &= h(y) - h(x_t) - \langle \nabla h(x_t) | y - x_t \rangle \\ &= h(y) + h^*(U_t) - \langle U_t | x_t \rangle - \langle \nabla h(x_t) | y - x_t \rangle \\ &= h(y) + h^*(U_t) - \langle U_t | y \rangle. \end{aligned}$$

3.6. **Comments on the continuous time dynamics framework.**

1) One obtains the existence of a level function and same speed of convergence of the no-regret values in classes (I), (II) or (III) : $O(\frac{1}{t})$.
2) Hence by Section 2 the accumulation points of the average $\{\bar{x}_t\}$ in class (II) with g dissipative are in eS.
3) In addition one has convergence of the average $\{\bar{x}_t\}$ in class (II) with g dissipative, with (PD), via Opial's Lemma.
The linear aspect of the derivative of the level function seems crucial to obtain this property.

Note that the Hilbertian structure is used first in the definition of the dynamics, then for the level function and in the reference to Opial's Lemma.

4) Similarly convergence of $\{x_t\}$ in case (III) holds for (PD), and (MD) with an adapted penalization function H.

5) Positive correlation holds for the three dynamics (under conditions on H for (MD)).

6) For vector fields g with potential G, $G(x_t)$ is strictly decreasing in (PD) and (DA), and under conditions on H for (MD).

7) In the framework of games the entropy function:

$$h(x) = \sum_{p \in S} x^p Log x^p$$

defined on the simplex $X = \Delta(S)$ leads (via (MD) or (DA)) to the *replicator dynamics* on $int\ X$, Taylor and Jonker (1978) [100], Hofbauer and Sigmund (1998) [51], Sorin (2009) [92], (2020) [94].
(MD) gives the differential version:

$$\dot{x}_t^p = x_t^p(u_t^p - \langle x_t, u_t \rangle), \quad p \in S$$

while (DA) corresponds to the integral representation:

$$x_t^p = \frac{e^{U_t^p}}{\sum_{s \in S} e^{U_t^s}}, \quad p \in S.$$

The corresponding Riemannian metric $[a, b]_x = \sum_{s \in S} \frac{1}{x^s} a^s b^s$, $x \in int\ X$, is introduced in Shahshahani, 1979 [89].
Recall also that the replicator dynamics is the continuous time version of the *multiplicative weight algorithm*, Littlestone and Warmuth, 1994 [63], Vovk, 1990 [103], Sorin, 2009 [92], 2020 [94].

On the other hand, $h(x) = \frac{1}{2}\|x\|^2$ leads to the *local/direct projection dynamics*, for a comparison, see Lahkar and Sandholm, 2008 [59], Sandholm, Dokumaci and Lahkar, 2008 [88].

8) There is an important literature on continuous time dynamics enjoying similar features, see e.g. :
- in convex optimization: Attouch and Teboulle, 2004 [5], Attouch, Bolte, Redont and Teboulle, 2004 [3], Auslender and Teboulle, 2006 [8], 2009 [9], Bolte and Teboulle (2003) [17], Teboulle (2018) [101] ...
- in game theory: Hofbauer and Sandholm, 2009 [50], Coucheney, Gaujal and Mertikopoulos, 2015 [27], Mertikopoulos and Sandholm, 2016 [65], Mertikopoulos and Sandholm (2018) [66], Mertikopoulos and Zhou (2019) [67] ...

4. Discrete time: general case

We consider now discrete time algorithms.

Remark that the dynamics depends on an additional parameter, the *step size*.

4.1. **Projected dynamics.**

Recall that V is Euclidean and let $m(X)$ denote the diameter of X.
Assumption: $\|u_m\| \leq M, \forall m \in \mathbb{N}$.

4.1.1. *Dynamics.*
The standard discrete dynamics (*Gradient projection method* in class (III), Levitin and Polyak (1966) [62], Polyak (1987) [81]) is given by:

$$\boxed{(38) \qquad x_{m+1} = argmax_X\{\langle u_m, x\rangle - \frac{1}{2\eta_m}\|x - x_m\|^2\},}$$

with $\eta_m > 0$ decreasing, which corresponds to:

$$x_{m+1} = \Pi_X[x_m + \eta_m u_m], \tag{39}$$

or with variational characterization, $x_{m+1} \in X$ and :

$$\langle x_m + \eta_m u_m - x_{m+1}, y - x_{m+1}\rangle \leq 0, \forall y \in X, \tag{40}$$

which is :

$$\langle u_m - \frac{x_{m+1} - x_m}{\eta_m}, y - x_{m+1}\rangle \leq 0, \forall y \in X,$$

and leads to (16) as $\eta_m \to 0$.

4.1.2. *Values.*
Recall that :

$$R_n(x) = \sum_{m=1}^{n} \langle u_m | x - x_m\rangle.$$

Proposition 20.

$$R_n(x) \leq \frac{1}{2\eta_n} m(X)^2 + \frac{M^2}{2}\sum_{m=1}^{n} \eta_m.$$

Hence with $\eta_n = 1/\sqrt{n}$:

$$R_n(x) \leq O(\sqrt{n}).$$

Proof:
Let $x_{m+1} = \Pi_X(y_{m+1})$ with $y_{m+1} = x_m + \eta_m u_m$.
So that:

$$2\eta_m\langle u_m, x - x_m\rangle = 2\langle y_{m+1} - x_m, x - x_m\rangle = \|y_{m+1} - x_m\|^2 + \|x - x_m\|^2 - \|x - y_{m+1}\|^2.$$

Note that for $x \in X$, (40) implies $\|y_{m+1} - x\|^2 \geq \|x_{m+1} - x\|^2$ hence:

$$2\eta_m\langle u_m, x - x_m\rangle \leq \eta_m^2\|u_m\|^2 + \|x - x_m\|^2 - \|x - x_{m+1}\|^2 \tag{41}$$

which is the discrete analog of the level function property in Proposition 7.
This gives:

$$\begin{aligned} R_n(x) &= \sum_{m=1}^{n} \langle u_m, x - x_m \rangle \\ &\leq \frac{1}{2\eta_1}\|x - x_1\|^2 - \frac{1}{2\eta_n}\|x - x_{n+1}\|^2 + \sum_{m=2}^{n}[\frac{1}{2\eta_m} - \frac{1}{2\eta_{m-1}}]\|x - x_m\|^2 + \frac{M^2}{2}\sum_{m=1}^{n}\eta_m. \end{aligned}$$

Thus, with $m(X)$ being the diameter of X:

$$R_n(x) \leq \frac{1}{2\eta_n} m(X)^2 + \frac{M^2}{2}\sum_{m=1}^{n}\eta_m$$

and the choice of $\eta_m = \frac{1}{\sqrt{m}}$ gives:

$$.R_n(x) \leq O(\sqrt{n}).$$

■

4.1.3. *Trajectories.*
Consider class (II).

Lemma 4.
For $x^* \in eS$, $\|x_m - x^*\|$ converges if $\{\eta_n\} \in \ell^2$.

Proof:
If $x^* \in eS$ then:

$$\|x^* - x_{m+1}\|^2 \leq \|x^* - (x_m + \eta_m g(x_m))\|^2 \leq \eta_m^2 \|g(x_m)\|^2 + \|x^* - x_m\|^2$$

so that $\|x_m - x^*\|$ converges if $\{\eta_n\} \in \ell^2$. ■
This corresponds to Proposition 5 in this framework.

Lemma 5.
If g is dissipative and $\{\eta_n\} \in \ell^2$, $\{\bar{x}_n\}$ converges to a point in eS.

Proof:
The limit points of $\{\bar{x}_n\}$ are in eS by Lemma 2, hence by Lemma 4 and Opial's Lemma (19), $\{\bar{x}_n\}$ converges to a point in eS. ■

This property is the counterpart of Proposition 8.

4.2. **Mirror descent.**

Assumptions:
a) H is a $\mathcal{C}^1$ function from V to $\mathbb{R} \cup \{+\infty\}$, L-strongly convex for some norm $\|.\|$ on $V = \mathbb{R}^n$ and $X \subset$ dom H,
b) $\|u_n\|_* \leq M, \forall n \in \mathbb{N}$.

4.2.1. *Dynamics.*

The classical *mirror descent algorithm*, introduced for class (III) in Nemirovski and Yudin (1983) [73], see also Beck and Teboulle (2003) [13], is given by (recall (27)):

$$\boxed{x_{m+1} = argmax_X\{\langle u_m|x\rangle - \frac{1}{\eta_m}D_H(x,x_m)\}. \qquad (42)}$$

The variational expression takes the form:

$$\langle \nabla H(x_m) + \eta_m u_m - \nabla H(x_{m+1})|x - x_{m+1}\rangle \leq 0, \forall x \in X. \tag{43}$$

which leads to the continuous formulation (26).
Note that $D_H(x,y)$ plays the rôle of $\frac{1}{2}\|x-y\|^2$ in (38).

4.2.2. *Values.*

We will use the identity:

$$D_H(x,z) - D_H(x,y) - D_H(y,z) = \langle \nabla H(y) - \nabla H(z)|x-y\rangle. \tag{44}$$

which is a direct consequence of the definition of D_H.

Proposition 21.
Let the step size $\eta_n = \frac{1}{\sqrt{n}}$, then:

$$R_n(x) \leq O(\sqrt{n}).$$

Proof:

$$\begin{aligned} \langle \eta_n u_n|x-x_n\rangle &= \langle \eta_n u_n|x-x_{n+1}\rangle + \langle \eta_n u_n|x_{n+1}-x_n\rangle \\ &\leq \langle \nabla H(x_{n+1}) - \nabla H(x_n)|x-x_{n+1}\rangle + \langle \eta_n u_n|x_{n+1}-x_n\rangle \\ &= D_H(x,x_n) - D_H(x,x_{n+1}) - D_H(x_{n+1},x_n) + \langle \eta_n u_n|x_{n+1}-x_n\rangle \end{aligned} \tag{45}$$

by using (44).
Now, H is L strongly convex, hence:

$$D_H(x_{n+1},x_n) \geq \frac{L}{2}\|x_{n+1}-x_n\|^2 \tag{46}$$

and moreover :

$$\langle \eta_n u_n|x_{n+1}-x_n\rangle - \frac{L}{2}\|x_n-x_{n+1}\|^2 \leq M\eta_n\|x_n-x_{n+1}\| - \frac{L}{2}\|x_n-x_{n+1}\|^2 \leq \frac{(\eta_n M)^2}{2L}$$

so that one obtains the analogous of the level function property in Proposition 15:

$$\langle \eta_n u_n|x-x_n\rangle \leq D_H(x,x_n) - D_H(x,x_{n+1}) + \frac{(\eta_n M)^2}{2L}.$$

Summing leads to :

$$R_n(x) \leq \sum_m [D_H(x,x_m)(\frac{1}{\eta_{m+1}} - \frac{1}{\eta_m}) + \eta_m\frac{M^2}{2L}]. \tag{47}$$

Hence the bound like in Proposition 20. ∎

4.2.3. *Trajectories.*
Consider class (II).

Lemma 6.
For $x^* \in eS$, $D_H(x^*, x_n)$ converges if $\{\eta_n\} \in \ell^2$.

Proof:
Start with (45) for $x^* \in eS$ and use $D_H \geq 0$ to get:

$$D_H(x^*, x_{m+1}) \leq D_H(x^*, x_m) - \langle \eta_m u_m | x_m - x_{m+1} \rangle \tag{48}$$

thus it remains to control $\langle \eta_m u_m | x_m - x_{m+1} \rangle$.
H being L strongly convex implies:

$$\langle \nabla H(x_m) - \nabla H(x_{m+1}) | x_m - x_{m+1} \rangle \geq L \| x_{m+1} - x_m \|^2$$

but one has by (43):

$$\langle \nabla H(x_m) - \nabla H(x_{m+1}) | x_m - x_{m+1} \rangle \leq \langle -\eta_m u_m | x_m - x_{m+1} \rangle \leq \| \eta_m u_m \|_* \| x_m - x_{m+1} \|.$$

It follows that:

$$\| x_m - x_{m+1} \| \leq \frac{1}{L} \| \eta_m u_m \|_*$$

hence:

$$\langle -\eta_m u_m | x_m - x_{m+1} \rangle \leq \frac{1}{L} \| \eta_m u_m \|_*^2$$

Altogether this implies from (48) that $D_H(x^*, x_m)$ converges if $\{\eta_n\} \in \ell^2$. ∎

This is the counterpart of Proposition 5.

4.3. **Dual averaging.**

Assumptions :
a) h is a l.s.c. function from V to $\mathbb{R} \cup \{+\infty\}$, L-strongly convex for some norm $\|.\|$ on $V = \mathbb{R}^n$, with dom $h = X$.
b) $\|u_m\|_* \leq M, \forall n \in \mathbb{N}$.

4.3.1. *Dynamics.*
We extend the formulation in Nesterov (2009) [76]. The starting point is again a maximization property:

$$x_{m+1} = argmax_X \{ \langle U_m | x \rangle - \frac{1}{\eta_m} h(x) \}, \tag{49}$$

with $U_m = \sum_{k=1}^m u_k$ and where $\{\eta_m\}$ is decreasing.
Note that there is an explicit form without using a variational formulation. Hence the *dual averaging* algorithm is given by:

$$\boxed{x_{m+1} = \nabla h^*(\eta_m U_m). \tag{50}}$$

This cumulative representation corresponds to the integral dynamics (34) while the incremental algorithm (43) is associated to the differential dynamics (26).

4.3.2. *Values.*

A direct proof, see Xiao (2010) [105] or a discrete approximation of (34), see Kwon and Mertikopoulos (2017) [58], allows to obtain:

Proposition 22.

$$R_n(x) = \sum_{m=1}^{n} \langle u_m | x - x_m \rangle \leq \frac{r_X(h)}{\eta_n} + \frac{\sum_{m=1}^{n} \eta_{m-1} \|u_m\|_*^2}{2L} \tag{51}$$

Proof:
Fenchel inequality:

$$\langle \eta_n U_n | x \rangle \leq h^*(\eta_n U_n) + h(x)$$

implies:

$$\langle U_n | x \rangle \leq \frac{h^*(0)}{\eta_0} + \sum_{m=1}^{n} \left(\frac{h^*(\eta_m U_m)}{\eta_m} - \frac{h^*(\eta_{m-1} U_{m-1})}{\eta_{m-1}} \right) + \frac{\max_X h}{\eta_n}. \tag{52}$$

Now:

$$\begin{aligned} \frac{h^*(\eta_m U_m)}{\eta_m} &= \sup_X [\langle U_m | x \rangle - \frac{h(x)}{\eta_m}] \\ &\leq \sup_X [\langle U_m | x \rangle - \frac{h(x)}{\eta_{m-1}}] + \sup_X [-\frac{h(x)}{\eta_m} + \frac{h(x)}{\eta_{m-1}}] \\ &= \frac{h^*(\eta_{m-1} U_m)}{\eta_{m-1}} + (\frac{1}{\eta_{m-1}} - \frac{1}{\eta_m}) \min_X h, \end{aligned}$$

so that replacing in (52) gives:

$$\langle U_n | x \rangle \leq \frac{h^*(0)}{\eta_0} + \sum_{m=1}^{n} \frac{1}{\eta_{m-1}} [h^*(\eta_{m-1} U_m) - h^*(\eta_{m-1} U_{m-1})] + \min_X h (\frac{1}{\eta_0} - \frac{1}{\eta_n}) + \frac{\max_X h}{\eta_n}. \tag{53}$$

h is L strongly convex for $\|.\|$, so that h^* is $1/L$ smooth for $\|.\|_*$ hence:

$$\begin{aligned} & h^*(\eta_{m-1} U_m) - h^*(\eta_{m-1} U_{m-1}) - \langle \eta_{m-1} U_m - \eta_{m-1} U_{m-1} | \nabla h^*(\eta_{m-1} U_{m-1}) \rangle \\ = \; & h^*(\eta_{m-1} U_m) - h^*(\eta_{m-1} U_{m-1}) - \langle \eta_{m-1} u_m | x_m \rangle \\ \leq \; & \frac{\eta_{m-1}^2}{2L} \|u_m\|_*^2. \end{aligned} \tag{54}$$

This leads to a property similar to the level function property in Proposition 18 since:

$$\begin{aligned} \langle u_m | x - x_m \rangle &\leq \langle u_m | x \rangle + \frac{1}{\eta_{m-1}} [h^*(\eta_{m-1} U_{m-1}) - h^*(\eta_{m-1} U_m)] + \frac{\eta_{m-1}}{2L} \|u_m\|_*^2 \\ &\leq \frac{1}{\eta_{m-1}} [h^*(\eta_{m-1} U_{m-1}) + h(x) - \langle \eta_{m-1} U_{m-1}, x \rangle] \\ &\quad - \frac{1}{\eta_m} [h^*(\eta_m U_m) + h(x) - \langle \eta_m U_m, x \rangle] \\ &\quad + \frac{\eta_{m-1}}{2L} \|u_m\|_*^2 + (\frac{1}{\eta_{m-1}} - \frac{1}{\eta_m})(\min_X h - h(x)). \end{aligned}$$

Now inserting (54) in (53) gives:

$$\langle U_n|x\rangle - \sum_{m=1}^{n}\langle u_m|x_m\rangle \leq \frac{h^*(0)}{\eta_0} + \sum_{m=1}^{n}\frac{\eta_{m-1}}{2L}\|u_m\|_*^2 + (\frac{1}{\eta_0} - \frac{1}{\eta_n})\min_X h + \frac{\max_X h}{\eta_n}$$

and recall that $h^*(0) = -\min_X h$. ■

Hence the convergence rate $O(\sqrt{n})$ with time varying parameters $\eta_n = 1/\sqrt{n}$.

4.4. **Comments on the discrete dynamics framework.** .

1) The three algorithms achieve the same bound $O(1/\sqrt{n})$ for the speed of convergence of the average regret, which is optimal already in class (III), Nesterov (2004), using time varying step sizes $\eta_n = 1/\sqrt{n}$.

2) The no-regret property holds for the three algorithms hence the results of Section 2.2 apply for the closed form.

3) More precise properties concerning the trajectories are available only in the (PD) set-up. The results are similar to the ones in the continuous case, Section 3.2 if $\eta_n \in \ell^2$, for class (II). (Compare to the analysis in Peypouquet and Sorin [80] for dynamics induced by maximal monotone operators in discrete and continuous time, see Section 3.3.4.).

4) There is no discrete counterpart of "positive correlation".

5) For vector fields g with potential one does not have the crucial property $G(x_n)$ decreasing.

5. Discrete time: Regularity

This section deals mainly with class (III) *convex gradient*, where in addition f satisfies some regularity properties.
Recall that f is β smooth if:

$$|f(y) - f(x) - \langle \nabla f(x)|y - x\rangle| \leq \frac{\beta}{2}\|x - y\|^2. \tag{55}$$

Alternatively ∇f is β-Lipschitz.

A last part is devoted to the so-called mirror-prox procedure, class (II) with g β-Lipschitz.

5.1. **Projected dynamics.**

Assumption: f is β smooth.

The algorithm is like (40) with constant step size $\eta_m = 1/\beta$.

$$x_{m+1} = argmax_X\{\langle -\nabla f(x_m), x\rangle - \frac{1}{2\beta}\|x - x_m\|^2\}, \tag{56}$$

which gives:

$$\boxed{x_{m+1} = \Pi_X[x_m - \frac{1}{\beta}\nabla f(x_m)]. \qquad (57)}$$

The analysis in this section is standard, see e.g. Nesterov (2004) [75].

5.1.1. *Preliminaries.*

Define $Tx = \Pi_X[x - \frac{1}{\beta}\nabla f(x)]$ and $v(x) = \beta(x - Tx)$ (which plays the role of $\nabla f(x)$, corresponding to $X = V$).
The projection property gives:

$$\langle x - \frac{1}{\beta}\nabla f(x) - Tx, y - Tx\rangle \leq 0, \quad \forall y \in X \tag{58}$$

so that:

$$\langle \nabla f(x), Tx - y\rangle \leq \langle v(x), Tx - y\rangle, \quad \forall y \in X. \tag{59}$$

Now one has:

$$\begin{aligned} f(Tx) - f(y) &= f(Tx) - f(x) + f(x) - f(y) \\ &\leq \langle \nabla f(x), Tx - x\rangle + \frac{\beta}{2}\|Tx - x\|^2 + \langle \nabla f(x), x - y\rangle \qquad f\ \beta \text{ smooth, convex} \\ &= \langle \nabla f(x), Tx - y\rangle + \frac{1}{2\beta}\|v(x)\|^2 \\ &\leq \langle v(x), Tx - y\rangle + \frac{1}{2\beta}\|v(x)\|^2, \quad \forall y \in X \qquad \text{by } (59) \end{aligned}$$

hence:

$$f(Tx) - f(y) \leq \langle v(x), x - y\rangle - \frac{1}{2\beta}\|v(x)\|^2, \quad \forall y \in X. \tag{60}$$

The following decreasing property is crucial and shows the difference with the general non-smooth case.

Lemma 7. *Descent lemma*

$$f(x_{m+1}) - f(x_m) \leq -\frac{1}{2\beta}\|v(x_m)\|^2 = -\frac{\beta}{2}\|x_{m+1} - x_m\|^2 \tag{61}$$

Proof:
By the previous inequality (60) with $x_m = x = y$. ∎
Note that in the previous lemma the convexity of f is actually not usued.

A consequence of (61) is:

$$\frac{1}{2\beta}\sum_{m=1}^{n}\|v(x_m)\|^2 \leq f(x_1) - f(x_{n+1}) \leq f(x_1) - f^*$$

hence $\{\|v(x_n)\|\} \in \ell^2$.

5.1.2. *Values.*

Proposition 23.

$$f(x_n) - f^* \leq O(\frac{1}{n}).$$

Proof:
Consider the algorithm defined by $\{z_n\}$ and the process $\{v_n\}$ with $z_1 = x_1$, $v_n = -v(z_n)$ and $z_{n+1} = z_n + \eta v_n$ with $\eta = 1/\beta$. Clearly $z_n = x_n$.
From section 3.1, Proposition 20 with η_m constant, one obtains:

$$R_n^v(y) = \sum_{m=1}^{n} \langle v_m, y - z_m \rangle \leq \frac{1}{2\eta}\|y - z_1\|^2 + \frac{\eta}{2}\sum_{m=1}^{n}\|v_m\|^2 \tag{62}$$

which is bounded since $\{\|v(x_n)\|\} \in \ell^2$. This implies, using f decreasing and (60):

$$n[f(x_{n+1}) - f(y)] \leq R_n^v(y) - \frac{1}{2\beta}\|\sum_{m=1}^{n}\|v_m\|^2 = \frac{\beta}{2}\|y - x_1\|^2.$$

■

5.1.3. *Trajectories.*

Lemma 8.
For $y^* \in S$, $\|x_n - y^*\|$ is decreasing.

Proof:
From (60) one has:

$$0 \leq f(x_{n+1}) - f(y^*) \leq \langle v(x_n), x_n - y^* \rangle - \frac{1}{2\beta}\|v(x_n)\|^2$$

hence :

$$\langle v(x_n), x_n - y^* \rangle \geq \frac{1}{2\beta}\|v(x_n)\|^2.$$

So that:

$$\begin{aligned} \|x_{n+1} - y^*\|^2 &= \|x_{n+1} - x_n\|^2 + \|x_n - y^*\|^2 + 2\langle x_{n+1} - x_n, x_n - y^* \rangle \\ &= \frac{1}{\beta^2}\|v(x_n)\|^2 + \|x_n - y^*\|^2 + 2\langle -\frac{1}{\beta}v(x_n), x_n - y^* \rangle \\ &\leq \|x_n - y^*\|^2. \end{aligned}$$

■

Proposition 24.
$\{x_n\}$ converges to a point in S.

Proof: Since $f(x_n)$ decreases, the accumulation points of $\{x_n\}$ are in S and by the previous Lemma 8, Opial's Lemma (19) applies. ■

5.2. **Mirror descent.**

We first assume only that H and f are $\mathcal{C}^1$.
The next analysis follows Bauschke, Bolte and Teboulle (2017) [12].

The main assumption in this section is the existence of a constant $L > 0$ such that (recall that D_H is the Bregman distance (27)):

$$(A) \qquad L\, D_H - D_f \geq 0$$

which is equivalent to : $LH - f$ convex.
Note that if H is strongly convex and f is smooth (not assumed convex), there exists L such that (A) holds. However f is not required to be smooth.
(A similar pre-oder on convex functions appears in Nguyen (2017) [77]).
Recall the procedure (43) with constant step size λ:

$$\boxed{(63) \quad \langle \lambda \nabla f(x_n) + \nabla H(x_{n+1}) - \nabla H(x_n) | x - x_{n+1} \rangle \geq 0, \forall x \in X}$$

and the identity:

$$(64) \qquad D_H(x,z) - D_H(x,y) - D_H(y,z) = \langle \nabla H(y) - \nabla H(z) | x - y \rangle.$$

Let us consider a step size satisfying $2\lambda L = 1$.

5.2.1. *Values.*

Lemma 9.

$$(65) \quad \lambda[f(x_{n+1}) - f(y)] \leq D_H(y, x_n) - D_H(y, x_{n+1}) - \frac{1}{2} D_H(x_{n+1}, x_n) - \lambda D_f(y, x_n), \forall y \in X.$$

Proof :
Since:

$$(66) \qquad D_f(x,z) = D_f(y,z) + f(x) - f(y) - \langle \nabla f(z) | x - y \rangle,$$

one has, by (A):

$$f(x) \leq f(y) + \langle \nabla f(z) | x - y \rangle + L D_H(x,z) - D_f(y,z)$$

Let $x = x_{n+1}, z = x_n$

$$f(x_{n+1}) - f(y) \leq \langle \nabla f(x_n) | x_{n+1} - y \rangle + L D_H(x_{n+1}, x_n) - D_f(y, x_n)$$

so that by (63):

$$\lambda[f(x_{n+1}) - f(y)] \leq \langle \nabla H(x_{n+1}) - \nabla H(x_n) | y - x_{n+1} \rangle + \lambda L D_H(x_{n+1}, x_n) - \lambda D_f(y, x_n).$$

Uses then (64):

$$\lambda[f(x_{n+1}) - f(y)] \leq D_H(y, x_n) - D_H(y, x_{n+1}) - D_H(x_{n+1}, x_n) + \lambda L D_H(x_{n+1}, x_n) - \lambda D_f(y, x_n)$$

Hence the result since $2\lambda L = 1$. ∎

Proposition 25.
Assume H convex . Then:
1) $f(x_n)$ is decreasing.
2) $\sum D_H(x_{n+1}, x_n) < +\infty$.
Assume f convex. Then :
3)

$$f(x_n) - f(y) \leq \frac{2L}{n} D_H(y, x_1), \quad \forall y \in X.$$

Proof:
1) and 2) Take $y = x_n$ in (65).
3) Use $f(x_n)$ decreasing, $D_f \geq 0$ and the telescoping sum in (65). ∎

5.2.2. *Trajectories.*

Proposition 26.
Assume f convex.
1) $y \in S$ implies: $D_H(y, x_n)$ decreases.
2) Assume:
$[H1] : x^k \to x^* \in S \Rightarrow D_H(x^*, x^k) \to 0$,
$[H2] : x^* \in S, D_H(x^*, x^k) \to 0 \Rightarrow x^k \to x^*$,
then $\{x_n\}$ converges to a point in S.

Proof:
1) follows from (65).
2) By the previous Proposition 25, the accumulation points of $\{x_n\}$ are in S. Let thus $x_{n_k} \to x^* \in S$.
By $H1$, $D_H(x^*, x_{n_k}) \to 0$ then by 1) $D_H(x^*, x_n) \to 0$. Now use $H2$. ∎
Compare with the proof of Proposition 17.

Note that the result is more precise than in the continuous case, Section 3.3 where there was no decreasing property (for general H).

Let us finally mention the very recent result due to Bui and Combettes (2020) [23] Theorem 3.9., where the use of variable metrics H_n allows to reach $f(x_n) - f^* = o(1/n)$.

5.3. **Dual averaging.**

We follow the analysis in Lu, Freund and Nesterov (2018) [64].
Recall that we consider class (III) : f convex and $\mathcal{C}^1$ and as in the previous Section 4.2. the main hypothesis is the existence of $L > 0$ with:

$$(A) \qquad L\, D_h - D_f \geq 0.$$

but we will simply assume here $L\, h - f$ convex,
where $h : V \to \mathbb{R} \cup \{+\infty\}$ is l.s.c. with dom $h = X$. Let $x_0 = argmin_X h(x)$ and assume $h(x_0) = 0$.
Define :

$$G_k(x) = \sum_{i=0}^{k-1} [\langle \nabla f(x_i), x - x_i \rangle + f(x_i)] + Lh(x) \tag{67}$$

and as in (49):

$$\boxed{(68) \quad x_k = argmin_X G_k(x) = argmax_X \{\langle U_{k-1}, x\rangle - Lh(x)\}}$$

with as usual $u_k = -\nabla f(x_k)$ and $U_m = \sum_{k=1}^m u_k$.

Proposition 27.

$$f(\bar{x}_k) - f(x) \leq \frac{L}{k} h(x), \tag{69}$$

$$\min_{i=0,\ldots,k} f(x_i) - f(x) \leq \frac{L}{k} h(x). \tag{70}$$

Proof:
By definition:

$$G_{k+1}(x_{k+1}) = G_k(x_{k+1}) + \langle \nabla f(x_k), x_{k+1} - x_k\rangle + f(x_k). \tag{71}$$

Note that (A) implies that each $G_k - f$ is convex hence:

$$G_k(x_{k+1}) - f(x_{k+1}) \geq G_k(x_k) - f(x_k) + \langle \partial G_k(x_k) - \nabla f(x_k), x_{k+1} - x_k\rangle. \tag{72}$$

Thus one has:

$$G_{k+1}(x_{k+1}) \geq f(x_{k+1}) + G_k(x_k) + \langle \partial G_k(x_k), x_{k+1} - x_k\rangle \tag{73}$$

but there exists $u_k \in \partial G_k(x_k)$ with :

$$\langle u_k, x - x_k\rangle \geq 0, \forall x \in X$$

by the choice of x_k.
Finally, with $g_{k+1} = G_{k+1}(x_{k+1})$ one obtains:

$$g_{k+1} \geq f(x_{k+1}) + g_k. \tag{74}$$

Using f convex, thus $G_k(x) \leq kf(x) + Lh(x)$ by (67), this implies that:

$$\sum_{i=0}^{k+1} f(x_i) \leq g_{k+1} \leq (k+1)f(x) + Lh(x) \tag{75}$$

hence the result. ∎

5.4. **Comments on the regular case.**

1) In the three cases (PD), (MD) and (DA) the speed of convergence of the values is $O(1/n)$ and the algorithms use a constant step parameter.
2) Using (PD) with f smooth implies $f(x_n)$ decreasing and the convergence of $\{x_n\}$.
3) The approach in Section 5.2 shows that similar results can be obtained using (MD) without assuming f with Lipschitz gradient if the regularization function H is adapted to f: condition (A).
4) Analogous results for the values are much simpler to obtain in the (DA) framework.

However the properties concern the value at the average $f(\bar{x}_n)$ and no result is available on the trajectories.

5.5. **Mirror prox.**

We follow in this section the model introduced by Korpelevich (1976) [55], and further studied by Nemirovski (2004) [72].
We present it for two main reasons: first it gives faster convergence than in Section 4 for a Lipschitz vector field g; second the tools used in the proofs are very similar to the one used in the previous sections.

5.5.1. *Dynamics.*
The algorithm corresponds to a two-stage procedure.
The first step produces y_{n+1} from x_n via usual (MD) (43) applied to $u_n = g(x_n)$ with constant step size λ:

$$\langle -\lambda g(x_n) + \nabla H(y_{n+1}) - \nabla H(x_n) | x - y_{n+1} \rangle \geq 0, \forall x \in X. \tag{76}$$

The second step produces x_{n+1} from x_n via (MD) (43) applied to $v_n = g(y_{n+1})$ with constant step size λ:

$$\langle -\lambda g(y_{n+1}) + \nabla H(x_{n+1}) - \nabla H(x_n) | x - x_{n+1} \rangle \geq 0, \forall x \in X. \tag{77}$$

Altogether, the decision after x_n uses a more refined information $g(y_{n+1})$ rather than the usual (in (MD)) $g(x_n)$.

5.5.2. *Values.*
The result concerns the process $\{y_n\}$.

Proposition 28. *Nemirovski (2004)* [72]
Assume H α strongly convex and g β-Lipschitz with $\alpha \geq \lambda\beta$, then:

$$R_n(x) = \sum_{m=1}^{n} \langle g(y_m) | x - y_m \rangle \leq \frac{1}{\lambda}[D_H(x, x_1) - D_H(x, x_n)].$$

Proof:
Decompose:

$$\begin{aligned} \langle g(y_{n+1}) | x - y_{n+1} \rangle &= \langle g(y_{n+1}) | x - x_{n+1} \rangle \\ &\quad + \langle g(x_n) | x_{n+1} - y_{n+1} \rangle \\ &\quad + \langle g(y_{n+1}) - g(x_n) | x_{n+1} - y_{n+1} \rangle. \end{aligned}$$

The first term gives via (77):

$$\begin{aligned} \lambda \langle g(y_{n+1}) | x - x_{n+1} \rangle &\leq \langle \nabla H(x_{n+1}) - \nabla H(x_n) | x - x_{n+1} \rangle \\ &= D_H(x, x_n) - D_H(x, x_{n+1}) - D_H(x_{n+1}, x_n) \end{aligned}$$

using (44).
Invoking (76), the second term leads to:

$$\lambda \langle g(x_n) | x_{n+1} - y_{n+1} \rangle \leq \langle \nabla H(x_n) - \nabla H(y_{n+1}) | y_{n+1} - x_{n+1} \rangle$$

$$= \quad D_H(x_{n+1}, x_n) - D_H(x_{n+1}, y_{n+1}) - D_H(y_{n+1}, x_n)$$

and for the last one, using g β-Lipschitz :

$$\begin{aligned}\langle g(y_{n+1}) - g(x_n) | x_{n+1} - y_{n+1}\rangle &\leq \|g(y_{n+1}) - g(x_n)\|_* \|x_{n+1} - y_{n+1}\| \\ &\leq \beta \|y_{n+1} - x_n\| \|y_{n+1} - x_{n+1}\| \\ &\leq \frac{\beta}{2}[\|y_{n+1} - x_n\|^2 + \|y_{n+1} - x_{n+1}\|^2].\end{aligned}$$

So that:

$$\begin{aligned}\lambda\langle g(y_{n+1}) | x - y_{n+1}\rangle &\leq D_H(x, x_n) - D_H(x, x_{n+1}) - D_H(x_{n+1}, y_{n+1}) \\ &\quad - D_H(y_{n+1}, x_n) + \frac{\lambda\beta}{2}[\|y_{n+1} - x_n\|^2 + \|y_{n+1} - x_{n+1}\|^2]. \qquad (78)\end{aligned}$$

Hence if H is α strongly convex and $\alpha \geq \lambda\beta$ one has a telescopic majorant leading to:

$$\lambda R_n(x) = \lambda \sum_{m=1}^{n} \langle g(y_m) | x - y_m\rangle \leq D_H(x, x_1) - D_H(x, x_n).$$

■

Comments :
If $eS \neq \emptyset$, take $x \in eS$ in (78) and $\alpha > \lambda\beta$ to get:
- $\sum_n \|y_{n+1} - x_n\|^2 + \|y_{n+1} - x_{n+1}\|^2 < \infty$
- $D_H(x, x_n)$ decreasing.

Notice that g was not assumed dissipative.
Adding g dissipative gives:

$$\langle g(x) | x - \bar{y}_n\rangle \leq R_n(x)$$

hence the accumulation points of $\{\bar{y}_n\}$ are in S.

5.5.3. *Trajectories.*
The result involves the trajectory $\{x_n\}$.

Proposition 29. *Korpelevich (1976)* [55]
Assume g β-Lipschitz and dissipative.
(i) (PD) case:
for $\lambda < 1/\beta$, x_n converges to a point in S.
(ii) (MD) case:
Assume H α strongly convex with $\alpha > \lambda\beta$. $y \in S$ implies: $D_H(y, x_n)$ decreases.
Assume:
$[H1] : x^k \to x^* \in S \Rightarrow D_H(x^*, x^k) \to 0$,
$[H2] : D_H(z^k, x^k) \to 0 \Rightarrow \|z^k - x^k\| \to 0$,
then $\{x_n\}$ converges to a point in S.

Proof:
(i) The property of the projection gives:

$$\|z - y\|^2 \geq \|z - \Pi_X(z)\|^2 + \|\Pi_X(z) - y\|^2, \qquad \forall y \in X$$

hence with $z = x_n + \lambda g(y_{n+1})$, $\Pi_X(z) = x_{n+1}$ by (77) and one deduces:

$$\|x_{n+1} - y\|^2 \quad \leq \quad \|x_n + \lambda g(y_{n+1}) - y\|^2 - \|x_n + \lambda g(y_{n+1}) - x_{n+1}\|^2$$

$$= \|x_n - y\|^2 - \|x_n - x_{n+1}\|^2 + 2\lambda\langle g(y_{n+1}), x_{n+1} - y\rangle. \tag{79}$$

Taking $y = x^* \in eS = S$ and adding $2\lambda\langle g(y_{n+1}), x^* - y_{n+1}\rangle \geq 0$, we obtain:

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \|x_n - x^*\|^2 - \|x_n - x_{n+1}\|^2 + 2\lambda\langle g(y_{n+1}), x_{n+1} - y_{n+1}\rangle \\ &= \|x_n - x^*\|^2 - \|x_n - y_{n+1}\|^2 - \|y_{n+1} - x_{n+1}\|^2 \\ &\quad -2\langle x_n - y_{n+1}, y_{n+1} - x_{n+1}\rangle + 2\lambda\langle g(y_{n+1}), x_{n+1} - y_{n+1}\rangle. \end{aligned} \tag{80}$$

Now by (76) one has:

$$\langle x_n + \lambda g(x_n) - y_{n+1}, x_{n+1} - y_{n+1}\rangle \leq 0 \tag{81}$$

replacing in (80) and using:

$$\begin{aligned} \langle g(y_{n+1}) - g(x_n), x_{n+1} - y_{n+1}\rangle &\leq \|g(y_{n+1}) - g(x_n)\|\|x_{n+1} - y_{n+1}\| \\ &\leq \beta\|y_{n+1} - x_n\|\|x_{n+1} - y_{n+1}\| \end{aligned} \tag{82}$$

one finally achieves:

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \|x_n - x^*\|^2 - \|x_n - y_{n+1}\|^2 - \|y_{n+1} - x_{n+1}\|^2 \\ &\quad +2\lambda\beta\|y_{n+1} - x_n\|\|x_{n+1} - y_{n+1}\| \\ &\leq \|x_n - x^*\|^2 - (1 - \lambda^2\beta^2)\|y_{n+1} - x_n\|^2. \end{aligned} \tag{83}$$

This implies :

$$\|y_{n+1} - x_n\|^2 \to 0. \tag{84}$$

By (83) the sequence $\{x_n\}$ is bounded. Let $\hat{x}$ be an accumulation point. Properties (76) and (84) imply that $\hat{x}$ satisfies:

$$\langle g(\hat{x}), \hat{x} - y\rangle \geq 0, \qquad \forall y \in X$$

hence $\hat{x} \in iS = S$, since g is dissipative. Taking now $x^* = \hat{x}$ in (83) implies that $\|x_n - \hat{x}\|$ is decreasing, but going to 0 on a subsequence, it converges to 0.

(II) Let us start with $x^* \in S$ and the identity:

$$D_H(x^*, x_{n+1}) = D_H(x^*, x_n) - D_H(x_{n+1}, x_n) - \langle \nabla H(x_{n+1}) - \nabla H(x_n)|x^* - x_{n+1}\rangle. \tag{85}$$

Denote $A = -\langle \nabla H(x_{n+1}) - \nabla H(x_n)|x^* - x_{n+1}\rangle$ and from (77) with $x = x^*$ one deduces:

$$A \leq \langle -\lambda g(y_{n+1})|x^* - x_{n+1}\rangle$$

But $x^* \in S$ hence:

$$\langle -\lambda g(y_{n+1})|x^* - y_{n+1}\rangle \leq 0$$

so that:

$$A \leq \langle -\lambda g(y_{n+1})|y_{n+1} - x_{n+1}\rangle.$$

Adding (76) with $x = x_{n+1}$ gives:

$$A \leq \langle \lambda[g(x_n) - g(y_{n+1})] - \nabla H(y_{n+1}) + \nabla H(x_n)|y_{n+1} - x_{n+1}\rangle.$$

Use that:

$$\langle -\nabla H(y_{n+1}) + \nabla H(x_n)|y_{n+1} - x_{n+1}\rangle = D_H(x_{n+1}, , x_n) - D_H(x_{n+1}, y_{n+1}) - D_H(y_{n+1}, x_n)$$

and coming back to (85) we obtain:

(86)

$$D_H(x^*, x_{n+1}) = D_H(x^*, x_n) - D_H(x_{n+1}, y_{n+1}) - D_H(y_{n+1}, x_n) + \lambda\langle g(x_n) - g(y_{n+1})|y_{n+1} - x_{n+1}\rangle.$$

Recall that H is α strongly convex so that:

$$D_H(u,v) \geq \frac{\alpha}{2}]\|u-v\|^2$$

and use that g is β- Lipschitz to bound:

$$\begin{aligned}\langle g(x_n)-g(y_{n+1})|y_{n+1}-x_{n+1}\rangle &\leq \|g(x_n)-g(y_{n+1})\|_*\|y_{n+1}-x_{n+1}\| \\ &\leq \beta\|x_n-y_{n+1}\|\|y_{n+1}-x_{n+1}\| \\ &\leq \frac{\beta}{2}[\rho\|x_n-y_{n+1}\|^2+\frac{1}{\rho}\|y_{n+1}-x_{n+1}\|^2].\end{aligned}$$

Replacing in (86) with $\rho=\frac{\lambda\beta}{\alpha}$ gives:

$$D_H(x^*,x_{n+1}) \leq D_H(x^*,x_n)-(1-(\frac{\lambda\beta}{\alpha})^2)D_H(y_{n+1},x_n).$$

Taking $\lambda<\frac{\alpha}{\beta}$ implies $D_H(y_{n+1},x_n)\to 0$ and $D_H(x^*,x_n)$ decreasing.
Finally let $\tilde{x}$ be an accumulation point of $\{x_n\}$ and $x_{n_k}\to\tilde{x}$. By [H2], $y_{n_k}\to\tilde{x}$ as well.
Using (76) this implies $\tilde{x}\in iS=S$ since g is dissipatif. But then $D_H(\tilde{x},x_n)$ decreases hence goes to 0, since it is the case on $\{x_{n_k}\}$ by [H1].
Using again [H2] $x_n\to\tilde{x}$ hence the result. ∎

The result has the same flavour than Proposition 26.

Notice that the algorithm is well defined for any vector field g but the interpretation in the decentralized framework of games is difficult. The main point is that x^i_{n+1} is determined via $g^i(y_{n+1})$ where all the components of y_{n+1} appear, hence some coordination is required.

6. Concluding remarks

For the three dynamics PG, MD and DA 1), 2) and 3) holds:

1) In continuous time the speed of convergence of the average regret to 0, of the order $O(1/t)$, is not better in the general gradient convex case than in the on-line learning case. This means that the specificity of the observation process cannot be exploited.

2) Similarly in discrete time the speed of convergence of the average regret to 0, of the order $O(1/\sqrt{n})$ is not better in the general gradient convex case than in the on-line learning case.

3) Adding a smoothness hypothesis on the convex function does not change the speed in the continuous time framework but allow a better convergence rate in discrete time from $O(1/\sqrt{n})$ to $O(1/n)$, and with a simpler algorithm : constant step size. Basically an additional property is required to get a descent lemma.

4) A similar phenomena appears with the so-called acceleration procedures following Nesterov (1983) [74].
In the continuous time case a second order ODE leads to a speed of convergence $f(x_t)-f(x^*)\leq O(\frac{1}{t^2})$ with no further hypothesis on f, see Su, Boyd and Candes (2014) [97] (2016) [98], Krichene, Bayen and Bartlett (2015) [56], (2016) [57], Attouch, Chbani, Peypouquet and Redont (2018) [4].
To obtain a similar property in discrete time, namely $f(x_n)-f(x^*)\leq O(\frac{1}{n^2})$ one has to

assume f smooth.
The same remark apply to the (weak) convergence of the trajectory, where f smooth is needed in discrete time and not in continuous time, Chambolle and Dossal (2015) [26], Attouch, Chbani, Peypouquet, Redont (2018) [4].

5) Concerning the link between discrete and continuous time dynamics, there are no direct results of the form: no-regret in continuous time imply no-regret in discrete time but analogy of the tools used and ad-hoc choice of the step parameter, see Sorin (2009) [92], Kwon and Mertikopoulos (2017) [58]. Similar properties hold in the closed form for Lyapounov functions in Krichene, Bayen and Bartlett (2015) [56], (2016) [57], Wibisono, Wilson and Jordan (2016) [104], see also Bansal and Gupta (2019) [11].

6) The Hilbert framework for (PD) allows to obtain convergence results on the trajectories. The two other algorithms ((MD and (DA)) are more flexible and can achieve better explicit speed of convergence of the values by choosing an adequate norm, see the discussion in Bauschke, Bolte and Teboulle (2017), [12]. For (MD) specific regularization functions H can also lead to convergence of the trajectories. (DA) is much simpler to implement due to its integral formulation. However no convergence properties of the trajectories are available.

7) In the framework of games, general positive results are obtained in the class of dissipative games. Accumulation points of the average trajectory are equilibria.

8) For potential games or more generally vector fields with potential the three continuous time algorithms imply the inclusion of the set of accumulation point in the set of rest points.
A further study in the framework of tame functions and invoving quasi-gradients, in the spirit of Bégout, Bolte and Jendoubi (2015) [18], would be very interesting.

9) Obviously potential games with a concave potential P will share the properties of class (III) since basically one can replace for each i, $\langle g^i(x), y^i - x^i\rangle$ by $\langle \nabla_i P(x), y^i - x^i\rangle$.

References

[1] Abernethy J., P. L. Bartlett and E. Hazan (2011) Blackwell approachability and no-regret learning are equivalent, *Proceedings of Machine Learning Research*, **19**, 27-46.

[2] Alvarez F., J. Bolte and O. Brahic (2004) Hessian Riemannian gradient flows in convex programming, *SIAM Journal on Control and Optimization*, **43**, 477-501.

[3] Attouch H., Bolte J., Redont P. and M. Teboulle (2004) Singular Riemannian barrier methods and gradient-projection dynamical systems for constrained optimization, *Optimization*, **53**, 435-454.

[4] Attouch H., Chbani Z., Peypouquet J., Redont P. (2018) Fast convergence of inertial dynamics and algorithms with asymptotic vanishing damping, *Mathematical Programming* 168 , no. 1-2, 123-175.

[5] Attouch H. and M. Teboulle (2004) Regularized Lotka-Volterra dynamical system as continuous proximal-like method in optimization, *JOTA*, **121**, 541-570.

[6] Aubin J.-P. and A. Cellina (1984) *Differential inclusions*, Springer.

[7] AUER P., CESA-BIANCHI N., FREUND Y. AND R.E. SHAPIRE (2002) The nonstochastic multiarmed bandit problem, *SIAM J. Comput.*, **32**, 48-77.
[8] AUSLENDER A. AND M. TEBOULLE (2006) Interior gradient and proximal methods for convex and conic optimization, *SIAM J Optimization*, **16**, 697-725.
[9] AUSLENDER A. AND M. TEBOULLE (2009) Projected subgradient methods with non Euclidean distances for non- differentiable convex minimization and variational inequalities, *Mathematical Programming*, **120**, 27-48.
[10] BAILLON J. B. AND H. BRÉZIS (1976) Une remarque sur le comportement asymptotique des semigroupes non linéaires, *Houston J. Math.*, **2**, 5-7.
[11] BANSAL N. AND A. GUPTA (2019) Potential-function proofs for gradient methods, *Theory of Computing*, **15**, 1-32.
[12] BAUSCHKE H., BOLTE J. AND M. TEBOULLE (2017) A descent Lemma beyond Lipschitz gradient continuity: first-order methods revisited and applications, *Mathematics of Operations Research*, **42**, 330-348.
[13] BECK A. AND M. TEBOULLE (2003) Mirror descent and nonlinear projected subgradient methods for convex optimization, *Oper. Res. Lett.*, **31**, 167-175.
[14] BENAIM M., J. HOFBAUER AND S. SORIN (2005) Stochastic approximations and differential inclusions, *SIAM Journal on Control and Optimization*, **44**, 328-348.
[15] BLACKWELL D. (1956) An analog of the minimax theorem for vector payoffs, *Pacific Journal of Mathematics*, **6**, 1-8.
[16] BLUM A. AND Y. MANSOUR (2007) From external to internal regret, *Journal of Machine Learning Reserach*, **8**, 1307-1324.
[17] BOLTE J. AND M. TEBOULLE (2003) Barrier operators and associated gradient-like dynamical systems for constrained minimization problems, *SIAM Journal on Control and Optimization*, **42**, 1266-1292.
[18] BÉGOUT P., J. BOLTE AND M.A. JENDOUBI (2015) On damped second-order gradient systems, *J. Differential Equations*, **259**, 3115-3143.
[19] BRÉZIS H. (1973) "Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert". North Holland Publishing Company, Amsterdam.
[20] BROWN G.W. AND J. VON NEUMANN (1950) Solutions of games by differential equations, in *Contibutions to the Theory of Games, I* (ed. H.W. Kuhn and A.W. Tucker), *Ann. Math. Studies*, **24**, 73–79.
[21] BRUCK R. E. (1975) Asymptotic convergence of nonlinear contraction semigroups in Hilbert space, *J. Funct. Anal.*, **18**, 15-26.
[22] BUBECK S. (2015) Convex optimization: Algorithms and complexity, *Fondations and Trends in Machine Learning*, **8**, 231-357.
[23] BUI M.N. AND P.L. COMBETTES (2020) Bregman forward-backward operator splitting, *Set-Valued and Variational Analysis*, https://doi.org/10.1007/s11228-020-00563-z.
[24] CESA-BIANCHI N. AND G. LUGOSI (2003) Potential-based algorithms in on-line prediction and game theory, *Machine Learning*, **51**, 239-261.
[25] CESA-BIANCHI N. AND G. LUGOSI (2006) *Prediction, Learning and Games*, Cambridge University Press.
[26] CHAMBOLLE A. AND C. DOSSAL (2015) On the convergence of the iterates of FISTA, *Journal of Optimization Theory and Applications*, **166**, 968-982.
[27] COUCHENEY P., B. GAUJAL AND P. MERTIKOPOULOS (2015) Penalty-regulated dynamics and robust learning procedures in games, *Mathematics of Operations Research*, **40**, 611-633.
[28] COVER T. (1991) Universal Portfolios, *Math. Finance*, **1**, 1-29.
[29] DUPUIS P. AND A. NAGURNEY (1993) Dynamical systems and variational inequalities, *Ann. Oper. Res.*, **44**, 9–42.
[30] FACCHINEI F. AND J. PANG (2007) *Finite-Dimensional Variational Inequalities and Complementarity Problems*, Springer.
[31] FOSTER D. AND R. VOHRA, (1993) A randomization rule for selecting forecasts, *Operations Research*, **41**, 704-707.

[32] FOSTER D. AND R. VOHRA (1997) Calibrated leaning and correlated equilibria, *Games and Economic Behavior*, **21**, 40-55.
[33] FOSTER D. AND R. VOHRA (1998) Asymptotic calibration, *Biometrika*, **85**, 379-390.
[34] FOSTER D. AND R. VOHRA (1999) Regret in the on-line decision problem, *Games and Economic Behavior*, **29**, 7-35.
[35] FREUND Y. AND R.E. SCHAPIRE (1999) Adaptive game playing using multiplicative weights, *Games and Economic Behavior*, **29**, 79-103.
[36] FUDENBERG D. AND D. K. LEVINE (1995) Consistency and cautious fictitious play, *Journal of Economic Dynamics and Control*, **19**, 1065-1089.
[37] FUDENBERG D. AND D. K. LEVINE (1998) *The Theory of Learning in Games*, MIT Press.
[38] FUDENBERG D. AND D. K. LEVINE (1999) Conditional universal consistency, *Games and Economic Behavior*, **29**, 104-130.
[39] HARKER P.T. AND J.S. PANG (1990) Finite-dimensional variational inequality and nonlinear complementarity problems: A survey of theory, algorithms, and applications, *Mathematical Programming*, **48**, 161-220.
[40] HANNAN J. (1957) Approximation to Bayes risk in repeated plays, *Contributions to the Theory of Games, III*, Drescher M., A.W. Tucker and P. Wolfe eds., Princeton University Press, 97-139.
[41] HART S. (2005) Adaptive heuristics, *Econometrica*, **73**, 1401-1430.
[42] HART S. AND A. MAS-COLELL (2000) A simple adaptive procedure leading to correlated equilibria, *Econometrica*, **68**, 1127-1150.
[43] HART S. AND A. MAS-COLELL (2001) A general class of adaptive strategies, *Journal of Economic Theory*, **98**, 26-54.
[44] HART S. AND A. MAS-COLELL (2003) Regret-based continuous time dynamics, *Games and Economic Behavior*, **45**, 375-394.
[45] HART S. AND A. MAS COLELL (2013) *Simple Adaptive Strategies: From Regret-Matching to Uncoupled Dynamics*, World Scientific Publishing.
[46] HAZAN E. (2011) The convex optimization approach to regret minimization, *Optimization for machine learning*, S. Sra, S. Nowozin, S Wright eds, MIT Press, 287-303.
[47] HAZAN E. (2015) Introduction to Online Convex Optimization, *Fondations and Trends in Optimization*, **2**, 157-325.
[48] HAZAN E. (2019) Optimization for Machine Learning , https://arxiv.org/pdf/1909.03550.pdf.
[49] HOFBAUER J. (2018) Minmax via replicator dynamics, *Dynamic Games and Applications*, **8**, 637-640.
[50] HOFBAUER J. AND W. H. SANDHOLM (2009) Stable games and their dynamics, *Journal of Economic Theory*, **144**, 1710-1725.
[51] HOFBAUER J. AND K. SIGMUND (1998) *Evolutionary Games and Population Dynamics*, Cambridge U.P.
[52] HOFBAUER J. AND S. SORIN (2006) Best response dynamics for continuous zero-sum games, *Discrete and Continuous Dynamical Systems-series B*, **6**, 215-224.
[53] KALAI A. AND S. VEMPALA (2005) Efficient algorithms for online decision problems, *Journal of Computer and System Sciences*, **71**, 291-307.
[54] KINDERLEHRER D. AND G. STAMPACCHIA (1980) *An Introduction to Variational Inequalities and Their Applications*, Academic Press.
[55] KORPELEVICH G. (1976) The extragradient method for finding saddle points and other problems, *Ekonomika i Matematicheskie Metody*, **12**, 747-756 (English version, (1977), *Matekon*, **13**, 35-49).
[56] KRICHENE W., A. BAYEN, AND P. BARTLETT (2015) Accelerated mirror descent in continuous and discrete time. In NIPS, 2015.
[57] KRICHENE W., A. BAYEN, AND P. BARTLETT (2016) Adaptive Averaging in Accelerated Descent Dynamics, In NIPS, 2016.
[58] KWON J. AND P. MERTIKOPOULOS (2017) A continuous time approach to on-line optimization, *Journal of Dynamics and Games*, **4**, 125-148.
[59] LAHKAR R. AND W. H. SANDHOLM (2008) The projection dynamic and the geometry of population games, *Games and Economic Behavior*, **64**, 565-590.

[60] LEHRER E. (2003) A wide range no-regret theorem, *Games Economic Behavior*, **42**, 101- 115.
[61] LEHRER E. AND S. SORIN (2007) Minmax via differential inclusion, *Journal of Convex Analysis*, **14**, 271-274.
[62] LEVITIN E.S. AND B.T. POLYAK (1966) Constrained minimization methods, *USSR Computational Mathematics and Mathematical Physics*, **6**, 1-50.
[63] LITTLESTONE N. AND M.K. WARMUTH (1994) The weighted majority algorithm, *Information and Computation*, **108**, 212-261.
[64] LU H., FREUND AND Y. NESTEROV (2018) Relatively smooth convex optimization by first-order methods, and applications, *Siam J Opt*, **28**, 333-354.
[65] MERTIKOPOULOS P. AND W. H. SANDHOLM (2016) Learning in games via reinforcement and regularization, *Mathematics of Operations Research*, **41**, 1297-1324.
[66] MERTIKOPOULOS P. AND W. H. SANDHOLM (2018) Riemannian game dynamics, *Journal of Economic Theory*, **177**, 315-364.
[67] MERTIKOPOULOS P. AND Z. ZHOU (2019) Learning in games with continuous action sets and unknown payoff functions, *Mathematical Programming*, **173**, 465-507.
[68] MINTY G. J. (1967) On the generalization of a direct method of the calculus of variations, *Bulletin A.M.S.*, **73**, 315–321.
[69] MONDERER D. AND L. S. SHAPLEY (1996) Potential games, *Games Econom. Behav.*, **14**, 124–143.
[70] MOREAU J.J. (1965) Proximité et dualité dans un espace Hilbertien, *Bull. Soc. Math. France*, **93**, 273–299.
[71] NAGURNEY A.AND D. ZHANG (1997) Projected dynamical systems in the formulation, stability analysis, and computation of fixed demand traffic network equilibria, *Transportation Sci.*, **31**, 147–158.
[72] NEMIROVSKI, A.(2004) Prox-method with rate of convergence O(1/t) for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems, *SIAM J Opt*, **15**, 229-251.
[73] NEMIROVSKI A. AND D. YUDIN (1983) *Problem Complexity and Method Efficiency in Optimization*, Wiley.
[74] NESTEROV Y. (1983) A method of solving a convex programming problem with convergence rate $O(1/k^2)$. *Soviet Mathematics Doklady*, **27**, 372-376.
[75] NESTEROV Y. (2004) *Introductory Lectures on Convex Optimization*, Kluwer.
[76] NESTEROV Y. (2009) Primal-dual subgradient methods for convex problems, *Mathematical Programming*, **120**, 221-259.
[77] NGUYEN Q.V. (2017) Forward-Backward splitting with Bregman distances, *Vietnam J. Math.*, **45**, 519-539.
[78] OPIAL Z. (1967) Weak Convergence of the sequence of successive approximations for nonexpansive mappings, *Bull. Amer. Math. Soc.*, **73**, 591-597.
[79] PERCHET V. (2014) Approachability, regret and calibration: implications and equivalences, *Journal of Dynamics and Games*, **1**, 181-254.
[80] PEYPOUQUET J. AND S. SORIN (2010) Evolution equations for maximal monotone operators: asymptotic analysis in continuous and discrete time, *Journal of Convex Analysis*, **17**, 1113-1163.
[81] POLYAK B. (1987) *Introduction to optimization*, Optimization software.
[82] ROCKAFELLAR R.T. (1970) *Convex Analysis*, Princeton University Press.
[83] ROCKAFELLAR R.T. (1970) Monotone operators associated with saddle-functions and minmax problems, *Nonlinear Functional Analysis, F. Browder, ed.*, Proceedings of Symposia in Pure Math, **18**, AMS , 241-250.
[84] ROSEN J.B. (1965) Existence and uniqueness of equilibrium points for concave N-person games. *Econometrica*, **33**, 520-534.
[85] SANDHOLM W. H. (2001) Potential games with continuous player sets, *Journal of Economic Theory*, **97**, 81-108.
[86] SANDHOLM W. H. (2010) *Population Games and Evolutionary Dynamics*, MIT Press.

[87] SANDHOLM W. H. (2015) Population games and deterministic evolutionary dynamics, H. P. Young, S. Zamir, eds *Handbook of Game Theory IV*, Elsevier, 703-778.
[88] SANDHOLM W. H., E. DOKUMACI AND R. LAHKAR (2008) The projection dynamic and the replicator dynamic, *Games and Economic Behavior*, **64**, 666-683.
[89] SHAHSHAHANI S. (1979) A new mathematical framework for the study of linkage and selection, *Memoirs of the American Mathematical Society*, 211.
[90] SHALEV-SHWARTZ S. (2012) Online Learning and Online Convex Optimization, *Foundations and Trends in Machine Learning*, **4** , 107-194.
[91] SMITH M.J. (1984) A descent algorithm for solving monotone variational inequalities and monotone complementarity problems, *J. Optim. Theory Appl.*, **44**, 485–496.
[92] SORIN S. (2009) Exponential weight algorithm in continuous time, *Mathematical Programming*, Ser. B, **116**, 513-528.
[93] SORIN S. (2011) On some global and unilateral adaptive dynamics, in *Evolutionary Game Dynamics*, K. Sigmund ed., Proceedings of Symposia in Applied Mathematics, **69**, A.M.S., 81-109.
[94] SORIN S. (2020) Replicator dynamics: Old and new, *Journal of Dynamics and Games*, **7**, 365-385.
[95] SORIN S. AND C. WAN (2016) Finite composite games: equilibria and dynamics, *Journal of Dynamics and Games*, **3**, 101-120.
[96] STOLTZ G. AND G. LUGOSI (2005) Internal Regret in On-Line Portfolio Selection, *Machine Learning*, **59**, 125-159.
[97] W. SU, S. BOYD, AND E. CANDES (2014) A differential equation for modeling Nesterov's accelerated gradient method: Theory and insights. In NIPS, 2014.
[98] SU W. , S. BOYD, AND E. CANDES (2016) A differential equation for modeling Nesterov's accelerated gradient method: Theory and insights, *Journal of Machine Learning Research*, **17**, 1-43.
[99] SWINKELS J.M. (1983) Adjustment dynamics and rational play in games, *Games and Economic Behavior*, **5**, 455–484.
[100] TAYLOR P. AND L. JONKER (1978) Evolutionary stable strategies and game dynamics, *Mathematical Biosciences*, **40**, 145-156.
[101] TEBOULLE M. (2018) A simplified view of first order methods for optimization, *Math Programming*, **170**, 67-96.
[102] VIOSSAT Y. AND A. ZAPECHELNYUK (2013) No-regret dynamics and fictitious play, *Journal of Economic Theory*, **148**, 825-842.
[103] VOVK V. (1990) *Aggregating strategies*, Proceedings of the 3rd Annual Conference on Computational Learning Theory, 371-383.
[104] WIBISONO A., A. C. WILSON, AND M. I. JORDAN (2016) A variational perspective on accelerated methods in optimization, *PNAS* , **113**, 7351-7358.
[105] XIAO L. (2010) Dual averaging methods for regularized stochastic learning and online optimization, *Journal of Machine Learning Research*, **11**, 2543-2596.
[106] ZINKEVICH M. (2003) Online convex programming and generalized infinitesimal gradient ascent, *Proceedings of the 20th International Conference on Machine Learning*, 928-936.

INSTITUT DE MATHÉMATIQUES JUSSIEU-PRG, SORBONNE UNIVERSITÉ, CAMPUS PIERRE ET MARIE CURIE, CNRS UMR 7586
E-mail address: `sylvain.sorin@imj-prg.fr`