HAL
open science

DNA MELTING STRUCTURES IN THE GENERALIZED POLAND-SCHERAGA MODEL

Quentin Berger, Maha Khatib

To cite this version:

Quentin Berger, Maha Khatib. DNA MELTING STRUCTURES IN THE GENERALIZED POLAND-SCHERAGA MODEL. ALEA : Latin American Journal of Probability and Mathematical Statistics, 2018, 15 (2), pp.993-1025. 10.30757/alea.v15-37 . hal-03911997

HAL Id: hal-03911997

https://hal.science/hal-03911997v1

Submitted on 23 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DNA MELTING STRUCTURES IN THE GENERALIZED POLAND-SCHERAGA MODEL

QUENTIN BERGER, GIAMBATTISTA GIACOMIN, AND MAHA KHATIB

Abstract. The Poland-Scheraga model for DNA denaturation, besides playing a central role in applications, has been widely studied in the physical and mathematical literature over the past decades. More recently a natural generalization has been introduced in the biophysics literature (see in particular [10]) to overcome the limits of the original model, namely to allow an *excess of bases* – i.e. a different length of the two single stranded DNA chains – and to allow slippages in the chain pairing. The increased complexity of the model is reflected in the appearance of configurational transitions when the DNA is in double stranded form. In [12] the generalized Poland-Scheraga model has been analyzed thanks to a representation in terms of a bivariate renewal process. In this work we exploit this representation further and fully characterize the path properties of the system, making therefore explicit the geometric structures – and the configurational transitions – that are observed when the polymer is in the double stranded form. What we prove is that when the excess of bases is not absorbed in a homogeneous fashion along the double stranded chain – a case treated in [12] – then it either condensates in a single *macroscopic* loop or it accumulates into an unbound single strand free end.

2010 *Mathematics Subject Classification: 60K35, 82D60, 92C05, 60K05, 60F10*

Keywords: DNA Denaturation, Polymer Pinning Model, Two-dimensional Renewal Processes, Sharp Deviation Estimates, non-Cramér regime, Condensation phenomena

1. Introduction

The Poland-Scheraga (PS) model [18, 4, 7] played and still plays a central role in the analysis of DNA denaturation (or melting): double stranded DNA *melts* into two single stranded DNA polymer chains at high temperature. The success of the model is partly due to the fact that it is exactly solvable when the heterogeneous character of the DNA is neglected. Moreover, solvability has an interest on its own, from a more theoretical standpoint: phase transition and critical phenomena in the PS model are completely understood [8, 11]. However, the PS model is an oversimplification in many respects: it deals with two strands of equal length and it does not allow slippages of the two chains. These simplifications make the model one dimensional, and solvability becomes less surprising. What is instead surprising is that a natural generalization [9, 10, 17] – called generalized Poland-Scheraga (gPS) model – fully overcomes these limitations, retaining the solvable character in spite of the substantially richer variety of structures that it displays. In [12] a mathematical approach to the gPS model is developed and it is pointed out that it can be represented in terms of a two dimensional renewal process, much like the PS model can be represented in terms of a one dimensional renewal. The solvable character of both models is then directly related to their renewal structure. The growth in complexity from PS to gPS models is nevertheless considerable: the key feature of PS and gPS is the presence of a localization transition, corresponding to the passage from separated to bound strands, and for the gPS there are three, not only one, types

of localized trajectories (or configurations). This has been first pointed out, at least in part, in [17], where one can find theoretical arguments (based also on a Bose-Einstein condensation analogy) and numerical evidence that *"suggest that a temperature-driven conformational transition occurs before the melting transition"* [17, p.3].

In this work we fully characterize the possible localized configurations. The transitions between different types of configurations have been already studied at the level of the free energy in [12] where these phenomena have been mathematically identified and interpreted in a Large Deviations framework in terms of *Cramér* and *non-Cramér* strategies. This will be explained in detail below. Here we content ourselves with pointing out that a full analysis of the Cramér regimes is given in [12]. However, the non-Cramér regime, where the condensation phenomena happen, requires a a substantially finer analysis – moderate deviations and local limit estimates – at the level of the bivariate renewals. These estimates, to which much attention has been devoted in the literature in the one dimensional set-up (see [1, 5] and references therein), are lacking to the best of our knowledge for higher dimensional renewals and they are not straightforward generalizations. They represent the technical core of this paper.

1.1. **The Model and some basic results.** We introduce the model in detail only from the renewal representation. The link with the original representation of the model is summed up in Fig. 1 and its caption, and we refer to [12] for more details.

We consider a persistent bivariate renewal process $\tau = \{(\tau_n^{(1)}, \tau_n^{(2)})\}_{n\geq 0}$, that is a sequence of random variables such that $\tau_0 = (0,0)$, $\{\tau_n - \tau_{n-1}\}_{n=0,2,\dots}$ is IID and such that the inter-arrival law – i.e. the law of τ_1 –, takes values in $\mathbb{N}^2 := \{1, 2, \cdots\}^2$.

We set $\mathbf{P}(\tau_1 = (n,m)) = K(n+m)$ with

$$K(n) := \frac{L(n)}{n^{2+\alpha}}\,, \tag{1.1}$$

for some $\alpha > 0$ and some slowly varying function $L(\cdot)$. Moreover $\sum_{n,m} K(n+m) = 1$ since we assumed the process to be persistent.

We consider two versions of the model: constrained and free. The partition function of the constrained model, or *constrained partition function*, can be written as

$$Z^c_{N,M,h} := \sum_{n=1}^{N\wedge M} \sum_{\substack{\underline{l}\in\mathbb{N}^n \\ |\underline{l}|=N}} \sum_{\substack{\underline{t}\in\mathbb{N}^n \\ |\underline{t}|=M}} \prod_{i=1}^{n} \exp(h) K(l_i + t_i)\,, \tag{1.2}$$

where $h \in \mathbb{R}$ is the binding energy, or pinning parameter.

The partition function of the free model, or *free partition function*, is defined by

$$Z^f_{N,M,h} := \sum_{i=0}^{N} \sum_{j=0}^{M} K_f(i) K_f(j) Z^c_{N-i,M-j,h}\,, \tag{1.3}$$

where $K_f(n) := \overline{L}(n) n^{-\overline{\alpha}}$ for some $\overline{\alpha} \in \mathbb{R}$ and slowly varying function $\overline{L}(\cdot)$. We assume that $K_f(0) = 1$ just to prevent this constant from popping up in various formulas: this choice has the side effect of making clear that $K_f(\cdot)$ is not a probability.

In [12] it is shown that for every h and every $\gamma > 0$

$$\mathrm{F}_\gamma(h) := \lim_{\substack{N,M\to\infty \\ M/N\to\gamma}} \frac{1}{N} \log Z^c_{N,M,h} = \lim_{\substack{N,M\to\infty \\ M/N\to\gamma}} \frac{1}{N} \log Z^f_{N,M,h} < \infty\,, \tag{1.4}$$



FIGURE 1. A configuration of the free gPS model, with one strand containing 22 bases and the other 14, is represented in two fashions: the *natural* (or polymer) one and the renewal one. In particular we see that $(1,1)$ renewal increments (or inter-arrivals) correspond to bound base pairs and all other increments (i,j) correspond to unbound regions in the bulk, that we call *loops* (of total length $i+j$, with length i in the first strand and j in the second strand). The term *unbound* is rather reserved to the terminal portion of the polymer: we refer to the free ends as unbound strands. Throughout this work, a polymer trajectory is always given in the renewal representation: it is therefore just a point process in the plane.

which says that the free energy (density) of free and constrained models, with binding energy h and strand length asymptotic ratio equal to γ, coincide. A number of basic properties of $h \mapsto \mathrm{F}_\gamma(h)$ are easily established, notably that it is a convex non decreasing function, equal to zero for $h \le 0$ and positive for $h > 0$. This already establishes that $h = 0$ is a critical point, in the sense that $\mathrm{F}_\gamma(\cdot)$ is not analytic at the origin.

But [12] is not limited to results on the free energy: associated to $Z^c_{N,M,h}$ and $Z^f_{N,M,h}$ there are two probability measures, that we denote respectively by $\mathbf{P}^c_{N,M,h}$ and $\mathbf{P}^f_{N,M,h}$. They are point measures, like the renewal processes on which they are built. It is standard to see that $\partial_h \mathrm{F}_\gamma(h)$ (which exists except possibly for countably many values of h) yields the $N \to \infty$ limit of the expected density of points (under $\mathbf{P}^c_{N,M,h}$ or $\mathbf{P}^f_{N,M,h}$). Hence for $h < 0$ the density is zero, while for $h > 0$ the density is positive. This tells us that we are stepping from a regime in which the two strands are essentially fully unbound to a regime in which they are tightly bound. In [12] results go well beyond this: it is in particular proven that for $h < 0$ the number of renewal points is $O(1)$ and these points are all close to $(0,0)$ or (N,M) (see Fig. 2). In the polymer representation, this means that the two DNA strands are completely unbound, except for a few contacts between the bases just close to the extremities. More precisely, it was found in [12] that in the free case, if $\overline{\alpha} < 1 + \alpha/2$ the two strands are free except for $O(1)$ contacts close to the origin, and if $\overline{\alpha} > 1 + \alpha/2$ the two free ends are of length $O(1)$ and a large loop appears in the system, see Fig 2.



FIGURE 2. A schematic image of the two types of observable trajectories of the free gPS model in the delocalized (denaturated, melted) regime, according to whether the exponent $\overline{\alpha}$ is smaller (left picture) or larger (right picture) than $1+\alpha/2$. In the constrained case only the trajectory on the right is observed, and the small free tails are reduced to zero. This case is treated in [12].

On the other hand for $h>0$ the situation is radically different. This has been analyzed in [12] but only in the Cramér regime. We are now going to discuss this in details.

1.2. **Binding strategies.** A way to get a grip on what is going on for $h>0$ is to observe that we can make the elementary manipulation: for every non negative λ_1 and λ_2

$$Z^c_{N,M,h} = e^{N\lambda_1+M\lambda_2} \sum_{n=1}^{N\wedge M} \sum_{\substack{\underline{l}\in\mathbb{N}^n \\ |\underline{l}|=N}} \sum_{\substack{\underline{t}\in\mathbb{N}^n \\ |\underline{t}|=M}} \prod_{i=1}^{N} \exp\left(h-\lambda_1 l_i-\lambda_2 t_i\right) K(l_i+t_i)\,. \tag{1.5}$$

Since $h>0$ we identify a family, in fact a curve in $[0,\infty)^2$, of values of (λ_1,λ_2) such that

$$\sum_{l,t=1}^{\infty} \exp\left(h-\lambda_1 l-\lambda_2 t\right) K(l+t) \;=\; 1\,, \tag{1.6}$$

and (1.6) clearly defines a probability distribution that is an inter-arrival distribution for a new bivariate renewal process. At this point is not too difficult to get convinced that $Z^c_{N,M,h}$ is equal to $e^{N\lambda_1+M\lambda_2}$ times the probability that this new renewal hits (N,M) (we call this probability *target probability*). If we are able to choose (λ_1,λ_2) so that the logarithm of the target probability is $o(N)$, then of course $\mathrm{F}_\gamma(h)=\lambda_1+\gamma\lambda_2$. This can actually be done: it amounts to solving a variational problem and the uniqueness of the optimal (λ_1,λ_2) follows by convexity arguments. However the solution may be qualitatively different for different values of h:

(1) the optimal (λ_1,λ_2) belong to $(0,\infty)^2$, so both components of the inter-arrival law of the arising renewal have distributions that decay exponentially. We call this *Cramér regime* because the tilt of the measure (in both components) is efficient in targeting the point (N,M) to which we are aiming at;
(2) either λ_1 or λ_2 is zero, so only one component of the arising inter-arrival law is exponentially tight. For the sake of conciseness we call this for now *non-Cramér regime* because the tilt of the measure (in only one of the component) is only partially successful in making the renewal τ reach and visit the point (N,M). To be precise there is in reality a boundary region between the two regimes, and the

notion of non-Cramér regime will be made more precise just below – this regime is the main issue of this work – so we will not dwell further on this right now.

A full treatment of the Cramér regime is given in [12], and the results can be resumed as follows: all loops are small, in fact the largest is $O(\log N)$, and the unbound strands (the free ends) are of length $O(1)$ – see the leftmost case in Fig 3. In the present work, we focus on the non-Cramér regime and the reader who wants to have an anticipation on the results can have a look Fig 3.

1.3. **The non-Cramér regime.** In order to make as explicit as possible for which values of $h > 0$ the system is in the non-Cramér regime, let us define $\mathrm{N}(h) > 0$ as the unique solution of

$$\sum_{n,m=1}^{\infty} K(n+m)\exp\left(-n\mathrm{N}(h)\right) \,=\, \exp(-h)\,. \tag{1.7}$$

This computation amounts to solving the variational problem we were after, in the case in which the problem is not solvable in $(0,\infty)^2$ and the optimal tilt of the measure involves only one of the two components. From (1.7) one can extract a number of properties of $\mathrm{N}(\cdot)$: it is a real analytic, positive, convex and increasing function [12]. We insist on the fact that, in spite of being defined for every $h > 0$, $\mathrm{N}(h)$ is not always equal to the free energy. More precisely in [12] it is shown that $\mathrm{F}_\gamma(h) = \mathrm{N}(h)$ if and only if $\gamma \notin (1/\gamma_c(h), \gamma_c(h))$, where

$$\gamma_c(h) \,:=\, \frac{\sum_{n,m} mK(n+m)\exp(-n\mathrm{N}(h))}{\sum_{n,m} nK(n+m)\exp(-n\mathrm{N}(h))}\,. \tag{1.8}$$

We refer to [12] for more details on the form of the function $\gamma_c(\cdot)$ and the switching phenomena between the Cramér and the non-Cramér regime. In this work, and without loss of generality (by symmetry), we will consider only the case $\gamma > \gamma_c(h)$. To be precise we will rather consider the case $\gamma \geq \gamma_c(h)$ because the phenomenology observed for $\gamma > \gamma_c(h)$, that is for $M - \gamma_c(h)N \geq cN$ for some $c > 0$ persists also in a part of the window $M - \gamma_c(h)N = o(N)$ and we will analyze the model also in this window. In different terms: the analysis in the Cramér regime is a Large Deviations analysis, but the whole non-Cramér regime is equivalent from the Large Deviations viewpoint (the issues there are about sharp deviations). So there isn't much conceptual difference between $M - \gamma_c(h)N \geq cN$ and $M - \gamma_c(h)N = o(N)$, up to when $M - \gamma_c(h)N$ grows too slowly, as we shall see.

Crucial for us is the probability distribution $\hat{\mathrm{K}}_h(\cdot,\cdot)$ defined by

$$\hat{\mathrm{K}}_h(n,m) = K(n+m)e^{h-n\mathrm{N}(h)}\,, \tag{1.9}$$

which, as announced informally just above, allows to write the partition function as

$$Z^c_{N,M,h} = \exp(N\mathrm{N}(h))\mathbf{P}\left((N,M) \in \hat{\tau}_h\right)\,, \tag{1.10}$$

where $\hat{\tau}_h$ is the bivariate renewal process with inter-arrival distribution $\hat{\mathrm{K}}_h(\cdot,\cdot)$, *i.e.*

$$\mathbf{P}\big((\hat{\tau}_h)_1 = (n,m)\big) = \hat{\mathrm{K}}_h(n,m) = \frac{L(n+m)}{(n+m)^{1+\alpha}}e^{h-n\mathrm{N}(h)}\,. \tag{1.11}$$

Next, we are going to have a closer look at this renewal process.

1.4. **On the bivariate renewal $\hat\tau_h$.** Let us write for conciseness $\mathrm{N}(h) = \mathrm{N}_h$ (a practice that we will pick up again in the proofs), and drop the dependence on h in $\hat\tau_h$: $\hat\tau = (\hat\tau^{(1)}, \hat\tau^{(2)})$. In view of (1.11), it is clear that the distribution of this process is not symmetric, we have the marginals

$$\begin{aligned} \mathbf{P}\big(\hat\tau_1^{(1)} = n\big) &= \sum_{m\geq 1} K(n+m)\exp(h-\mathrm{N}_h n) \overset{n\to\infty}{\sim} \frac{\exp(h)}{(1+\alpha)}\,\frac{L(n)}{n^{1+\alpha}}\,e^{-\mathrm{N}_h n}\,, \\ \mathbf{P}\big(\hat\tau_1^{(2)} = m\big) &= \sum_{n\geq 1} K(n+m)\exp(h-\mathrm{N}_h n) \overset{m\to\infty}{\sim} \frac{\exp(h)}{\exp(\mathrm{N}_h)-1}\,\frac{L(m)}{m^{2+\alpha}}\,. \end{aligned} \tag{1.12}$$

Let us also denote (the dependence in h is implicit)

$$\hat\mu_1 := \mathbf{E}[\hat\tau_1^{(1)}] < +\infty, \qquad \hat\mu_2 := \mathbf{E}[\hat\tau_1^{(2)}] < +\infty\,, \tag{1.13}$$

so that $\gamma_c(h) = \hat\mu_2/\hat\mu_1$, cf. (1.8).

We notice that the process $\hat\tau^{(1)}$ has moments of all orders, and so $\{\hat\tau_n^{(1)}\}_{n=0,1,\dots}$ is in the domain of attraction of a normal law: we denote $a_n^{(1)} := \sqrt{n}$ the scaling sequence for $\hat\tau_n^{(1)}$. On the other hand, the process $\{\hat\tau_n^{(2)}\}_{n=0,1,\dots}$ is in the domain of attraction of an α_2-stable law, with $\alpha_2 := (1+\alpha)\wedge 2 > 1$: its scaling sequence $a_n^{(2)}$ verifies

$$L(a_n^{(2)})(a_n^{(2)})^{-\alpha_2} \sim \frac{1}{n} \ \text{ if } \alpha_2 < 2 \qquad \text{and} \qquad \sigma(a_n^{(2)})(a_n^{(2)})^{-2} \sim \frac{1}{n} \ \text{ if } \alpha_2 = 2 \tag{1.14}$$

where

$$\sigma(n) := \mathbf{E}\big[\big(\hat\tau_1^{(2)}\big)^2 \mathbf{1}_{\{\hat\tau_1^{(2)} \leqslant n\}}\big]\,, \tag{1.15}$$

and diverges as a slowly varying function if $\mathbf{E}[(\hat\tau_1^{(2)})^2] = +\infty$ (with $\sigma(n)/L(n) \to +\infty$ [3]). Note that $\sigma(n)$ converges if $\alpha > 1$. In particular, $a_n^{(2)}$ is regularly varying with exponent $1/\alpha_2 = (1+\alpha)^{-1} \vee (1/2)$.

As an additional relevant definition, we select a sequence $\{m_n^{(2)}\}_{n=1,2,\dots}$ satisfying

$$\mathbf{P}\big(\hat\tau_1^{(2)} > m_n^{(2)}\big) \overset{n\to\infty}{\sim} \frac{1}{n}, \tag{1.16}$$

so that $m_n^{(2)}$ gives the order of $\max_{1\leqslant j\leqslant n}\{\hat\tau_j^{(2)} - \hat\tau_{j-1}^{(2)}\}$. We stress that $m_n^{(2)}$ is regularly varying with exponent $(1+\alpha)^{-1}$, and that $m_n^{(2)}/a_n^{(2)} \in [1/c, c]$ for some $c \geq 1$ if $\alpha < 1$, but $m_n^{(2)}/a_n^{(2)} \to 0$ when $\alpha \geqslant 1$: in any case, there is a constant $c > 0$ such that $m_n^{(2)} \leqslant c a_n^{(2)}$ for every n, i.e. $m_n^{(2)} = O(a_n^{(2)})$.

1.5. **Non-Cramér regime and big-jump domain.** We drop the dependence of $\gamma_c(h)$ on h, and we set

$$t_N := M - \gamma_c N\,. \tag{1.17}$$

Of course, having $\gamma > \gamma_c$ means that $t_N/N \geq c$ for some $c > 0$. But it is natural and essentially not harder to tackle the problem assuming only

$$t_N/a_N^{(2)} \to +\infty \ \text{ as } N \to \infty, \tag{1.18}$$

with additionally, in the case $\alpha \geqslant 1$ (recall the definition of $\sigma(n)$ after (1.14))

$$\Big(\frac{t_N}{a_N^{(2)}}\Big)^2 \frac{\sigma(a_N)}{\sigma(t_N)} \overset{N\to\infty}{\sim} \frac{t_N^2}{N\sigma(t_N)} \geqslant C_0 \log N \ \text{ for a suitable choice of } C_0 > 0\,. \tag{1.19}$$

If $\alpha > 1$, as well as if $\alpha = 1$ and $\sigma(n) = O(1)$ (i.e. if $\mathbf{E}[(\hat{\tau}_1^{(2)})^2] < \infty$), (1.19) simply means that $t_N \geqslant C_0' \sqrt{N \log N}$ with C_0' easily related to C_0. Note also that (1.18) implies $t_N \gg \sqrt{N \log N}$ if $\mathbf{E}[(\hat{\tau}_1^{(2)})^2] = \infty$.

We stress that the constants C_0 depends only on $K(\cdot)$ and, for the interested reader, it can be made explicit by tracking the constants in (3.34) and (**??**) where the value of C_0 is used. This assumption is made to be sure that we lie in the so called *big-jump domain*, as studied for example in the one-dimensional setting in [5]: in our model it simply means that deviations – and the event we focus on is $(N, M) \in \hat{\tau}$ – are realized by an atypical deviation on just one of the increment variables $\hat{\tau}_{i+1} - \hat{\tau}_i$. As we shall, this happens just under the assumption (1.18) for $\alpha < 1$ and this condition is optimal (see Appendix B.1). For the case $\alpha \geq 1$ the extra condition (1.19) is not far from being optimal, but it is not: we discuss this point in Appendix B.2, but we do not treat it in full generality because it is a technically demanding issue that leads far from our main purposes.

1.6. **Mais results I: polymer trajectories.** Let us now introduce two fundamental events in an informal, albeit precise, fashion. The two events will be rephrased in a more formal way in (2.9), once further notations will have been introduced. Choose sequences of positive numbers $\{u_N\}_{N=1,2,\dots}$, $\{m_N^+\}_{N=1,2,\dots}$, $\{a_N^+\}_{N=1,2,\dots}$ and $\{\widetilde{a}_N^+\}_{N=1,2,\dots}$ such that

$$u_N \gg 1\,, \quad t_N \gg m_N^+ \gg m_N^{(2)}\,, \quad t_N \gg a_N^+ \gg a_N^{(2)} \quad \text{and} \quad t_N \gg \widetilde{a}_N^+ \gg a_N^{(2)}\,. \tag{1.20}$$

In practice, and to optimize the result that follows, u_N, $m_N^+/m_N^{(2)}$, $a_N^+/a_N^{(2)}$ and $\widetilde{a}_N^+/a_N^{(2)}$ should be chosen tending to ∞ in an arbitrarily slow fashion.

We then define the *Big Loop* event $E_{\mathtt{BL}}^{(N)}$ to be the set of trajectories such that

(1) there is one loop of size larger than $t_N - a_N^+$ and smaller than $t_N + a_N^+$, so that, to leading order, it is of size t_N;
(2) all other loops are smaller than m_N^+ (hence there is only one largest loop);
(3) the length of neither of the two unbound strands is larger than u_N.

The (*large* or *macroscopic*) *Unbound Strand* event $E_{\mathtt{US}}^{(N)}$ is instead the set of trajectories such that

(1) all loops are smaller than m_N^+;
(2) the length of the unbound portion of the shorter strand does not exceed u_N;
(3) the length of the unbound portion of the longer strand is larger than $t_N - \widetilde{a}_N^+$ and smaller than $t_N + \widetilde{a}_N^+$, so that, to leading order, it is of size t_N.

Note that $E_{\mathtt{BL}}^{(N)} \cap E_{\mathtt{US}}^{(N)} = \emptyset$ except, possibly, for finitely many N: the two conditions (1) are incompatible. We refer to Fig. 3 for a schematic image of these two events.

Theorem 1.1. *Under assumptions* (1.18), (1.19) *and* (1.20) *we have that*

$$\lim_{N\to\infty} \mathbf{P}^f_{N,M,h}\big(E_{\mathtt{BL}}^{(N)} \cup E_{\mathtt{US}}^{(N)}\big) \,=\, 1\,. \tag{1.21}$$

(1) *Moreover, if* $\overline{\alpha} < 1$ *(and hence* $\sum_j K_f(j) = \infty$*) then*

$$\lim_{N\to\infty} \mathbf{P}^f_{N,M,h}\big(E_{\mathtt{US}}^{(N)}\big) \,=\, 1\,. \tag{1.22}$$

(2) *If* $\sum_j K_f(j) < \infty$ *(and hence* $\overline{\alpha} \geq 1$*) then*

$$\mathbf{P}^f_{N,M,h}\big(E_{\mathtt{US}}^{(N)}\big) \overset{N\to\infty}{\sim} \frac{1}{1+Q_N} \quad \textit{and} \quad \mathbf{P}^f_{N,M,h}\big(E_{\mathtt{BL}}^{(N)}\big) \overset{N\to\infty}{\sim} \frac{Q_N}{1+Q_N}\,, \tag{1.23}$$

with

$$Q_N := c_h N\,(t_N)^{-(2+\alpha)+\overline{\alpha}}\,\frac{L(t_N)}{\overline{L}(t_N)} \quad \textit{and } c_h := \frac{e^h \sum_{j=0}^{\infty} K_f(j)}{\hat{\mu}_1(e^{\mathrm{N}(h)}-1)}\,. \tag{1.24}$$

For conciseness the case $\overline{\alpha}=1$ with $\sum_j K_f(j)=+\infty$ is not included in Theorem 1.1, but its is treated in full in Appendix A. It is a marginal case in which an anomalous behavior appears: a big loop and a large unbound strand may coexist.



FIGURE 3. Schematic image of the observable trajectories of the free gPS model in the Cramér regime (left), and in the non-Cramér regime (cf. Theorem 1.1): the Large Unbound Strand event (center, occuring when $\overline{\alpha}<\alpha+1$) and the Big Loop event (right, occuring when $\overline{\alpha}>\alpha+1$). In the constrained case the Unbound Strand event is not observed, and the free tails are of course absent. What cannot be appreciated in this schematic view is the fact that the small loop distribution has exponential tail in the Cramér regime (hence the largest is $O(\log N)$) and that it has power law tail in the non-Cramér regime (hence the largest is $O(N^a)$ for some $a\in(0,1)$: $O(m_N)$ to be precise).

It is worth pointing out that, *in most of the cases*, the expressions in (1.23) have a limit – at least if $\{t_N\}_{N=1,2,\dots}$ is not too wild (regularly varying is largely sufficient) – and it is either one or zero. In particular when $t_N\sim cN$ for some $c>0$ we have

$$\lim_{N\to\infty}\mathbf{P}^f_{N,M,h}\left(E^{(N)}_{\mathtt{US}}\right) = \begin{cases} 1 & \text{if } \overline{\alpha}<\alpha+1 \text{ or if } \overline{\alpha}=\alpha+1 \text{ and } \overline{L}(t) \overset{t\to\infty}{\gg} L(t),\\ 0 & \text{if } \overline{\alpha}>\alpha+1 \text{ or if } \overline{\alpha}=\alpha+1 \text{ and } \overline{L}(t) \overset{t\to\infty}{\ll} L(t). \end{cases} \tag{1.25}$$

(This is true also in the case $\overline{\alpha}=1$ with $\sum_j K_f(j)=+\infty$, see (A.6)-(A.7)). Note that in the case in which $\overline{\alpha}=\alpha+1$ and the ratio of the two slowly varying function has a limit which is neither 0 nor ∞, the limit of the probability of the unbound strand event exists and it is an explicit value in $(0,1)$.

Let us also give the following complement to Theorem 1.1 for the constrained case.

Theorem 1.2. *Under assumptions* (1.18) *(and additionally* (1.19) *if* $\alpha\geq 1$*) and* (1.20) *we have that*

$$\lim_{N\to\infty}\mathbf{P}^c_{N,M,h}\left(E^{(N)}_{\mathtt{BL},0}\right) = 1\,, \tag{1.26}$$

where $E^{(N)}_{\mathtt{BL},0}$ *is the event* $E^{(N)}_{\mathtt{BL}}$ *with the more stringent condition that the the length of both the unbound strands is* 0.

2. Main results II: sharp estimates on the partition functions

In this section, we give the asymptotic behavior of $e^{-N\mathrm{N}_h} Z^c_{N,M,h} = \mathbf{P}((N,M) \in \hat\tau)$ in the big-jump domain. Then we present the asymptotic behavior of $Z^f_{N,M,h}$. Both in the constrained and free case we also give more technical estimates that identify some events to which we can restrict the partition functions without modifying them in a relevant way. Theorem 1.1 turns out to be a corollary of these technical estimates, as we explain in the final part of the section.

In this section and in the rest of the paper we deal with order statistics and we introduce here the relative definitions. Consider the (non-increasing) order statistics $\{\mathcal{M}_{1,k}, \mathcal{M}_{2,k}, \dots, \mathcal{M}_{k,k}\}$ of the IID family $\{\hat\tau^{(2)}_j - \hat\tau^{(2)}_{j-1}\}_{j=1,\dots,k}$. In particular $\mathcal{M}_{1,k}$ is a maximum of this finite sequence. We will consider the order statistics also for k random, notably for $k = \kappa_N$, with

$$\kappa_N := \max\{i \,:\, \hat\tau^{(1)}_i \in [0,N]\}. \tag{2.1}$$

2.1. **On the constrained partition function.** We start with an important estimate for the constrained partition function (more precisely for $\mathbf{P}((N,M) \in \hat\tau)$, the the renewal mass function), that is essential for the study of the free partition function, as one can imagine from its definition (1.3). It is worth insisting on the link between $\mathbf{P}$ and the measure we are interested in for the constrained case:

$$\mathbf{P}^c_{N,M,h}(\cdot) \,=\, \mathbf{P}(\,\cdot\,|(N,M) \in \hat\tau)\,. \tag{2.2}$$

Theorem 2.1. *Assume that $\alpha > 0$ and that* (1.18) *holds. Moreover if $\alpha \geqslant 1$ assume also* (1.19)*. Recall that $\hat\tau$ has inter-arrival distribution* (1.11)*. Then (recall that $M = \gamma_c N + t_N$) we have that*

$$\mathbf{P}\big((N,M) \in \hat\tau\big) = \mathbf{P}\Big(\hat\tau^{(1)}_{\kappa_N} = N,\, \hat\tau^{(2)}_{\kappa_N} = M\Big) \overset{N\to\infty}{\sim} \frac{N}{\hat\mu_1^2}\mathbf{P}\Big(\hat\tau^{(2)}_1 = \lceil t_N \rceil\Big). \tag{2.3}$$

Moreover, for every $\eta \in (0,1)$ there exist $\varepsilon_0 > 0$ such that for every $\varepsilon \in (0,\varepsilon_0)$ and N sufficiently large (how large may depend on ε), we have

$$\mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} - t_N \in \Big[-\frac{a^{(2)}_N}{\varepsilon}, \frac{a^{(2)}_N}{\varepsilon}\Big],\, \mathcal{M}_{2,\kappa_N} \le \frac{m^{(2)}_N}{\varepsilon},\, \hat\tau^{(1)}_{\kappa_N} = N,\, \hat\tau^{(2)}_{\kappa_N} = M\Big) \ge$$
$$\ge (1-\eta)\frac{N}{\hat\mu_1^2}\mathbf{P}\Big(\hat\tau^{(2)}_1 = \lceil t_N \rceil\Big)\,. \tag{2.4}$$

Let us stress that the bivariate renewal process $\hat\tau_h$ falls in the domain of attraction of an $(\alpha_1 = 2, \alpha_2)$ stable distribution (see e.g. [19] or [13]). We have, as $n \to \infty$, that $\Big\{\Big(\frac{\hat\tau^{(1)}_n - \hat\mu_1 n}{a^{(1)}_n}, \frac{\hat\tau^{(2)}_n - \hat\mu_2 n}{a^{(2)}_n}\Big)\Big\}_{n=1,2,\dots}$ converges in distribution to a non-degenerate $(2,\alpha_2)$-bivariate stable law.

We mention that a bivariate local limit theorem is given in [6] and multivariate (d-dimensional) renewals are further studied in [2]: local large deviation estimates are given, as well as strong renewal theorems, i.e. asymptotics of $\mathbf{P}((n,m) \in \tau)$ as $(n,m) \to \infty$, when (n,m) is *close to* the *favorite direction* – i.e. the line $t \mapsto t\mathbf{E}[\tau_1]$, *close to* meaning at distance of the order of the fluctuations around that direction. We refer to [2] for further details (rough estimates when (n,m) is *away from* the favorite direction are also given). We stress that the multivariate case we consider is very special, and that here we need and obtain estimates that are (much) sharper than what one finds in the literature we have cited.

2.2. **On the free partition function.** We now give the behavior of the free partition function and identify trajectories contributing the most to it. Let us introduce some notations:

$$V_1^{(N)} := N - \hat{\tau}^{(1)}_{\kappa_N}, \qquad V_2^{(N)} := M - \hat{\tau}^{(2)}_{\kappa_N}, \tag{2.5}$$

the lengths of the free parts, see Fig 1.

For a set A of allowed trajectories, we define $Z^f_{N,M,h}(A)$ the partition function restricted to trajectories in A (by restricting the summation over subsets of $\{1,\dots N\}\times\{1,\dots M\}$ to those in A). For example, $Z^f_{N,M,h}((N,M)\in\hat{\tau}) = Z^c_{N,M,h}$.

We set $\overline{K} := \sum_{j=1}^{\infty} K_f(j)$ if the sum is finite, and we set $\overline{K}=0$ if $\sum_j K_f(j) = +\infty$.

Theorem 2.2. *Assume that $\alpha>0$ and (1.18). Moreover if $\alpha \geqslant 1$ assume also (1.19) and if $\overline{\alpha}=1$ assume that $\sum_j K_f(j)<\infty$. Then for $N\to\infty$*

$$\begin{aligned} e^{-N_{\mathrm{N}}(h)} Z^f_{N,M,h} = (1+o(1))\overline{K}\,\frac{N}{\hat{\mu}_1^2}\Big(\sum_{i\geq 0} K_f(i) e^{-i_{\mathrm{N}}(h)}\Big)\mathbf{P}\Big(\hat{\tau}_1^{(2)} = \lceil t_N\rceil\Big) \\ + (1+o(1))\frac{1}{\hat{\mu}_1}\Big(\sum_{i\,\geqslant\, 0} K_f(i) e^{-i_{\mathrm{N}_h}}\Big) K_f\left(t_N\right) . \end{aligned} \tag{2.6}$$

Moreover, for every $\eta\in(0,1)$ there exist $\varepsilon_0>0$ such that for every $\varepsilon\in(0,\varepsilon_0)$ and N sufficiently large (how large may depend on ε), such that

$$\begin{aligned} e^{-N_{\mathrm{N}_h}} Z^f_{N,M,h}\Big(V_1^{(N)}, V_2^{(N)} \leqslant \frac{1}{\varepsilon}, \mathcal{M}_{1,\kappa_N}\in\Big[t_N - \frac{a_N^{(2)}}{\varepsilon}, t_N + \frac{a_N^{(2)}}{\varepsilon}\Big], \mathcal{M}_{2,\kappa_N}\leq \frac{m_N^{(2)}}{\varepsilon}\Big) \\ \geqslant (1-\eta)\overline{K}\,\frac{N}{\hat{\mu}_1^2}\Big(\sum_{i\,\geqslant\,0} K_f(i) e^{-i_{\mathrm{N}_h}}\Big)\mathbf{P}\left(\hat{\tau}_1^{(2)} = t_N\right) ; \end{aligned} \tag{2.7}$$

$$\begin{aligned} e^{-N_{\mathrm{N}_h}} Z^f_{N,M,h}\Big(V_1^{(N)} \leqslant \frac{1}{\varepsilon}, V_2^{(N)}\in\Big[t_N - \frac{a_N^{(2)}}{\varepsilon}, t_N + \frac{a_N^{(2)}}{\varepsilon}\Big], \mathcal{M}_{1,\kappa_N}\leqslant \frac{m_N^{(2)}}{\varepsilon}\Big) \\ \geqslant (1-\eta)\frac{1}{\hat{\mu}_1}\Big(\sum_{i\,\geqslant\,0} K_f(i) e^{-i_{\mathrm{N}_h}}\Big) K_f\left(t_N\right) . \end{aligned} \tag{2.8}$$

We remark that when $\overline{\alpha}<1$ (so $\overline{K}=0$), the right-hand side of (2.6) reduces to one term and (2.7) becomes trivial. The case $\overline{\alpha}=1$ with $\sum_j K_f(j)=+\infty$ is treated in Theorem A.1.

2.3. **Back to the Big Loop and Unbound Strand events.** The notations we have introduced allow a compact formulation of the two key events of Theorem 1.1:

$$\begin{aligned} E^{(N)}_{\mathtt{BL}} &= \Big\{\mathcal{M}_{1,\kappa_N}\in[t_N - a_N^+, t_N + a_N^+],\, \mathcal{M}_{2,\kappa_N} < m_N^+,\, \max\Big(V_1^{(N)}, V_2^{(N)}\Big)\leq u_N\Big\} , \\ E^{(N)}_{\mathtt{US}} &= \Big\{\mathcal{M}_{1,\kappa_N} < m_N^+,\, V_1^{(N)}\leq u_N,\, V_2^{(N)}\in[t_N - \widetilde{a}_N^+, t_N + \widetilde{a}_N^+]\Big\} . \end{aligned} \tag{2.9}$$

Proof of Theorems 1.1 and 1.2. This is just a book-keeping exercise using the three estimates in Theorem 2.2 (or Theorem 2.1 for Theorem 1.2), together with the definition of $K_f(t_N)$ and the estimate of $\mathbf{P}(\hat{\tau}_1^{(2)} = \lceil t_N\rceil)$ in (1.12). □

2.4. **A word about the arguments of proof and organization of the remaining sections.** As we pointed out at the beginning of the introduction, *condensation phenomena* are widely studied in the mathematical literature (see [5, 1] and references therein), but not in the multivariate context. The full multivariate context is the object of [2], where renewal estimates, i.e. estimates on $\mathbf{P}((n_1,\dots,n_d)\in\hat\tau)$, are given: the big-jump domain in treated in much greater generality, but not with the precision we require. Here we deal only with the very special bivariate case motivated by the application – which has the simplifying feature of being highly asymmetric so in the end we can reduce to a one dimensional renewal – and we need and find the exact local asymptotic behavior of the renewal function. One of the main difficulties we face is that, on the event $(N,M)\in\hat\tau$, the number κ_N of renewal points is random and highly constrained by this event. We show that in the big-jump domain considered in Section 1.5, the main contribution to the probability $\mathbf{P}((N,M)\in\hat\tau)$ comes from trajectories with a number of renewal points that is approximately $\kappa_N=N/\hat\mu_1+O(\sqrt{N})$. For this number κ_N, $\hat\tau^{(1)}_{\kappa_N}$ does not have to deviate from its typical behavior to be equal to N, but $\hat\tau^{(2)}_{\kappa_N}$ has to deviate from its typical behavior to reach M and it does so by making one single big jump, of order $t_N+O(a^{(2)}_N)$. In this sense, if we accept that κ_N is forced to be $N/\hat\mu_1+O(\sqrt{N})$ by the condition $N\in\hat\tau^{(1)}$, we can focus on $M\in\hat\tau^{(2)}$ and the problem becomes *almost one dimensional*. This turns out to be a lower bound strategy: for a corresponding upper bound we have to show that all other trajectories bring a negligible contribution to $\mathbf{P}((N,M)\in\hat\tau)$.

In the rest of the paper, we estimate separately the constrained and free partition functions. We deal with the constrained partition function in Section 3: the main term (2.4) in Section 3.1 and the remaining negligible contributions in Section 3.2. The free partition function is dealt with in Section 4: the main terms (2.7) and (2.8) in Section 4.1 and the remaining negligible contributions in Section 4.2. In Appendix A we complete the analysis of the case $\overline{\alpha}=1$. In Appendix B we discuss the transition from the big-jump regime (a single big jump, with a big deviation of just one of the two components) to the Cramér deviation strategy (no big jump).

To keep things simpler in the rest of the paper, and with some abuse of notation, we will systematically omit the integer part in the formulas.

3. The constrained partition function: proof of Theorem 2.1

3.1. **Proof of the lower bound** (2.4)**.** We start by decomposing the event of interest according to $\kappa_N=k$. The probability of such an event, restricted to $\{\kappa_N=k\}$, becomes (recall that $M=\gamma_cN+t_N$)

$$\mathbf{P}\Big(|\mathcal{M}_{1,k}-t_N|\le\frac{a^{(2)}_N}{\varepsilon},\,\mathcal{M}_{2,k}\le\frac{m^{(2)}_N}{\varepsilon},\,\hat\tau^{(1)}_k=N,\,\hat\tau^{(2)}_k=M\Big)=$$
$$\mathbf{P}\Big(\bigcup_{j=1}^{k}\Big\{\hat\tau^{(2)}_j-\hat\tau^{(2)}_{j-1}\in t_N+I_N,\,\max_{i\neq j}\Big(\hat\tau^{(2)}_i-\hat\tau^{(2)}_{i-1}\Big)\le\frac{m^{(2)}_N}{\varepsilon},\,\hat\tau^{(1)}_k=N,\,\hat\tau^{(2)}_k=M\Big\}\Big),\quad(3.1)$$

where we defined $I_N:=\{-a^{(2)}_N/\varepsilon,\dots,a^{(2)}_N/\varepsilon\}$. Since $t_N-a^{(2)}_N/\varepsilon$ is larger than $m^{(2)}_N/\varepsilon$ (recall that $m^{(2)}_n\leqslant ca^{(2)}_n$ for every n, and (1.18)) for N sufficiently large, the union in the right-hand side of (3.1) is a union of disjoint events that have all the same probability.

This term is equal to

$$k\,\mathbf{P}\Big(\hat\tau_1^{(2)}\in t_N+I_N,\ \max_{i=2,\dots,k}\Big(\hat\tau_i^{(2)}-\hat\tau_{i-1}^{(2)}\Big)\le\frac{m_N^{(2)}}{\varepsilon},\ \hat\tau_k^{(1)}=N,\ \hat\tau_k^{(2)}=\gamma_cN+t_N\Big)\ =$$
$$k\sum_{y\in I_N}\sum_{x\in\mathbb{N}}\mathbf{P}\Big(\hat\tau_1^{(2)}=t_N+y,\ \hat\tau_1^{(1)}=x\Big)\mathbf{P}\Big(\mathcal{M}_{1,k-1}\leqslant\frac{m_N^{(2)}}{\varepsilon},\hat\tau_{k-1}^{(1)}=N-x,\ \hat\tau_{k-1}^{(2)}=\gamma_cN-y\Big)\,, \tag{3.2}$$

where we have used that $\{(\hat\tau_j^{(1)}-\hat\tau_1^{(1)},\hat\tau_j^{(2)}-\hat\tau_1^{(2)})\}_{j=2,\dots,k}$ and $\{(\hat\tau_j^{(1)},\hat\tau_j^{(2)})\}_{j=1,\dots,k-1}$ have the same law.

Since we are after a lower bound we may and do restrict the sum over x between 1 and $1/\varepsilon$ and $y\in I_N:=\{-a_N^{(2)}/\varepsilon,\dots,a_N^{(2)}/\varepsilon\}$. And using that $\mathbf{P}(\hat\tau_1^{(2)}=n)$ is regularly varying, we have that uniformly for such x and $y\in I_N$

$$\mathbf{P}\Big(\hat\tau_1^{(2)}=t_N+y,\ \hat\tau_1^{(1)}=x\Big)\ \ge(1-\delta_N)\,\mathbf{P}\Big(\hat\tau_1^{(2)}=t_N\Big)\mathbf{P}\Big(\hat\tau_1^{(1)}=x\,\Big|\,\hat\tau_1^{(2)}=t_N+y\Big)\,, \tag{3.3}$$

where $\delta_N=\delta_N(\varepsilon)\ge0$ is such that $\lim_{N\to\infty}\delta_N=0$. If now we set $p_h(x):=(e^{\mathrm{N}_h}-1)e^{-x\mathrm{N}_h}$, by using (1.12) we have that

$$\begin{aligned}\mathbf{P}\Big(\hat\tau_1^{(1)}=x\,\big|\,\hat\tau_1^{(2)}=&t_N+y\Big)\ =\ \frac{\mathbf{P}\big(\hat\tau_1^{(1)}=x,\ \hat\tau_1^{(2)}=t_N+y\big)}{\mathbf{P}\big(\hat\tau_1^{(2)}=t_N+y\big)}\\ &\ge\ (1-\delta_N)\,\frac{p_h(x)K\left(t_N+x+y\right)}{K\left(t_N+y\right)}\ \ge\ (1-\delta_N)^2\,p_h(x)\,,\end{aligned} \tag{3.4}$$

possibly for a different choice of $\delta_N=\delta_N(\varepsilon)$. Therefore, going back to (3.1) we see that (again, by redefining δ_N)

$$\mathbf{P}\Big(\mathcal{M}_{1,k}\in t_N+I_N,\ \mathcal{M}_{2,k}\le\frac{m_N^{(2)}}{\varepsilon},\ \hat\tau_k^{(1)}=N,\ \hat\tau_k^{(2)}=M\Big)\ \ge$$
$$(1-\delta_N)k\,\mathbf{P}\big(\hat\tau_1^{(2)}=t_N\big)\sum_{x=1}^{1/\varepsilon}p_h(x)\mathbf{P}\Big(\mathcal{M}_{1,k-1}\leqslant\frac{m_N^{(2)}}{\varepsilon},\hat\tau_{k-1}^{(1)}=N-x,\ \hat\tau_{k-1}^{(2)}-\gamma_cN\in I_N\Big)\,. \tag{3.5}$$

We now sum over the values of k and we restrict to $k\in[(N/\hat\mu_1)-\sqrt{N/\varepsilon},(N/\hat\mu_1)+\sqrt{N/\varepsilon}]\cap\mathbb{Z}:=J_N$. Hence, redefining δ_N, the left-hand side of (2.4) is bounded from below by

$$(1-\delta_N)\mathbf{P}\Big(\hat\tau_1^{(2)}=t_N\Big)\frac{N}{\hat\mu_1}\sum_{x=1}^{1/\varepsilon}p_h(x)P_\varepsilon(x)\,, \tag{3.6}$$

where we defined, with $n_N^+:=\max J_N$,

$$P_\varepsilon(x):=\sum_{k\in J_N}\mathbf{P}\Big(\max_{i=1,\dots,n_N^+}\Big(\hat\tau_i^{(2)}-\hat\tau_{i-1}^{(2)}\Big)\leqslant\frac{m_N^{(2)}}{\varepsilon},\hat\tau_{k-1}^{(1)}=N-x,\ \hat\tau_{k-1}^{(2)}-\gamma_cN\in I_N\Big) \tag{3.7}$$

For $P_\varepsilon(x)$, we observe right away that by introducing also $n_N^- := \min J_N$ – note that $n_N^\pm$ are equal to $(N/\hat\mu_1) \pm \sqrt{N/\varepsilon}$ – we have

$$\begin{aligned} P_\varepsilon(x) \geq & \sum_{k\in J_N} \mathbf{P}\Big(\mathcal{M}_{1,n_N^+} \leqslant \frac{m_N^{(2)}}{\varepsilon}, \hat\tau_{k-1}^{(1)} = N-x\Big) \\ & - \sum_{k\in J_N} \mathbf{P}\Big(\mathcal{M}_{1,n_N^+} \leqslant \frac{m_N^{(2)}}{\varepsilon}, \hat\tau_{k-1}^{(1)} = N-x,\, \hat\tau_{n_N^+}^{(2)} - \gamma_c N > \frac{a_N^{(2)}}{\varepsilon}\Big) \\ & - \sum_{k\in J_N} \mathbf{P}\Big(\mathcal{M}_{1,n_N^+} \leqslant \frac{m_N^{(2)}}{\varepsilon}, \hat\tau_{k-1}^{(1)} = N-x,\, \hat\tau_{n_N^-}^{(2)} - \gamma_c N < \frac{a_N^{(2)}}{\varepsilon}\Big) \\ & \geqslant \mathbf{P}\left(E_1 \cap E_2(x)\right) - \mathbf{P}\left(E_3^+\right) - \mathbf{P}\left(E_3^-\right), \end{aligned} \tag{3.8}$$

where

$$E_1 := \Big\{\mathcal{M}_{1,n_N^+} \leqslant \frac{m_N^{(2)}}{\varepsilon}\Big\}, \qquad E_2(x) := \Big\{\exists k \in J_N \text{ such that } \hat\tau_{k-1}^{(1)} = N-x\Big\}, \tag{3.9}$$

and

$$E_3^+ := \Big\{\hat\tau_{n_N^+}^{(2)} - \gamma_c N > \frac{a_N^{(2)}}{\varepsilon}\Big\}, \qquad E_3^- := \Big\{\hat\tau_{n_N^-}^{(2)} - \gamma_c N < \frac{a_N^{(2)}}{\varepsilon}\Big\}. \tag{3.10}$$

We now estimate separately the probability of these events.

E_1 has probability close to one. For this, we use that $\mathbf{P}(\hat\tau_1^{(2)} > n)$ is regularly varying with index $(1+\alpha)^{-1}$ together with the definition (1.16) of $m_n^{(2)}$ to obtain that for N larger than some constant $N_0 = N_0(\varepsilon)$ we have $\mathbf{P}\left(\hat\tau_1^{(2)} > \frac{1}{\varepsilon} m_N^{(2)}\right) \leqslant 2\varepsilon^{1+\alpha} N^{-1}$. Therefore, we have for $N \geqslant N_0$

$$\mathbf{P}\left(E_1\right) \geqslant \left(1 - 2\varepsilon^{1+\alpha} N^{-1}\right)^{n_N^+} \geqslant e^{-3\varepsilon^{1+\alpha}}. \tag{3.11}$$

where we used that $n_N^+ \leqslant N$ and ε small.

$E_2(x)$ has probability close to $1/\hat\mu_1$. The probability of $E_2(x)$ is estimated by writing

$$\begin{aligned} \mathbf{P}\left(E_2(x)\right) &= \mathbf{P}(N-x \in \hat\tau^{(1)}) - \sum_{k<n_N^-} \mathbf{P}\left(\hat\tau_k^{(1)} = N-x\right) - \sum_{k>n_N^+} \mathbf{P}\left(\hat\tau_k^{(1)} = N-x\right) \\ &\geqslant \mathbf{P}(N-x \in \hat\tau^{(1)}) - \mathbf{P}\Big(\hat\tau_{n_N^-}^{(1)} \geqslant N - \tfrac{1}{\varepsilon}\Big) - \mathbf{P}\Big(\hat\tau_{n_N^+}^{(1)} \leqslant N\Big), \end{aligned} \tag{3.12}$$

where for the second term we used that $\mathbf{P}\big(\exists\, k < n_N^- \text{ s.t. } \hat\tau_k^{(1)} = N-x\big) \leqslant \mathbf{P}\big(\hat\tau_{n_N^-}^{(1)} \geqslant N-x\big)$ together with the fact that $x \leqslant 1/\varepsilon$ (and similarly for the last term).

First, because the inter-arrivals of $\hat\tau^{(1)}$ are exponentially integrable, $|\mathbf{P}(N \in \hat\tau^{(1)}) - 1/\hat\mu_1| \leq \exp(-cN)$ for $N \geq N_0$ with $c > 0$ and N_0 that depend on the inter-arrival law [15]. Therefore, uniformly in $x = 1, \dots, 1/\varepsilon$, we have that for N sufficiently large $\mathbf{P}(N-x \in \hat\tau^{(1)}) \geqslant 1/\hat\mu_1 - e^{-cN/2}$.

For the remaining terms in (3.12) it is just a matter of using the Central Limit Theorem. In fact, recalling that $n_N^- = N/\hat\mu_1 - \sqrt{N/\varepsilon}$, we have

$$\mathbf{P}\Big(\hat\tau_{n_N^-}^{(1)} \geqslant N - \tfrac{1}{\varepsilon}\Big) = \mathbf{P}\Big(\hat\tau_{n_N^-}^{(1)} - \hat\mu_1 n_N^- \geqslant \hat\mu_1 \varepsilon^{-1/2}\sqrt{N} - \tfrac{1}{\varepsilon}\Big) \leqslant e^{-c\varepsilon^{-1}}, \tag{3.13}$$

for N larger than some $N_0 = N_0(\varepsilon)$. On the other hand, we also have that

$$\mathbf{P}\Big(\hat\tau^{(1)}_{n_N^+} \leqslant N\Big) = \mathbf{P}\Big(\hat\tau^{(1)}_{n_N^+} - \hat\mu_1 n_N^+ \leqslant -\hat\mu_1 \varepsilon^{-1/2}\sqrt{N}\Big) \leqslant e^{-c'\varepsilon^{-1}}, \tag{3.14}$$

provided again that N is large enough.

Therefore we have proven that for every $\eta \in (0,1)$ there exists ε_0 and $N_0 : (0,1) \to \mathbb{N}$ such that for $\varepsilon \in (0,\varepsilon_0)$ and $N \geq N_0(\varepsilon)$ we have

$$\min_{x=1,\dots,1/\varepsilon} \mathbf{P}\left(E_2(x)\right) \geq \frac{1-\eta}{\hat\mu_1}. \tag{3.15}$$

$E_3^\pm$ have a small probability. This is a consequence of the convergence to stable limit law. In fact, using that $\gamma_c = \hat\mu_2/\hat\mu_1$ so that $\gamma_c N = \hat\mu_2 n_N^+ - \hat\mu_2\sqrt{N/\varepsilon}$, we get

$$E_3^+ = \left\{\hat\tau^{(2)}_{n_N^+} - \hat\mu_2 n_N^+ > \frac{a_N^{(2)}}{\varepsilon} - \hat\mu_2\sqrt{N/\varepsilon}\right\} \subset \left\{\hat\tau^{(2)}_{n_N^+} - \hat\mu_2 n_N^+ > \frac{a_N^{(2)}}{2\varepsilon}\right\}, \tag{3.16}$$

where the last inclusion holds provided that ε is sufficiently small, since there is a constant c such that $a_N^{(2)} \geqslant c\sqrt{N}$ for all N (we actually simply need N to be large if $a_N^{(2)}/\sqrt{N} \to +\infty$ for $N \to \infty$, which is the case when $\mathbf{E}[(\hat\tau_1^{(2)})^2] = +\infty$). Very much in the same way we get also to

$$E_3^- \subset \left\{\hat\tau^{(2)}_{n_N^-} - \hat\mu_2 n_N^- < -\frac{a_N^{(2)}}{2\varepsilon}\right\}. \tag{3.17}$$

Since $(\hat\tau^{(2)}_{n_N^\pm} - \hat\mu_2 n_N^\pm)/a^{(2)}_{n_N^\pm}$ converges in law for $N \to \infty$ to a stable limit variable Y, and using that $a_N^{(2)}/a^{(2)}_{n_N^\pm} \to \hat\mu_1^{1/\alpha_2}$ (since $n_N^\pm \sim N/\mu_1$ and $a_N^{(2)}$ is regularly varying with exponent α_2^{-1}, recall $\alpha_2 := \min(1+\alpha, 2)$), it is straightforward to see that

$$\limsup_{N\to\infty} \mathbf{P}(E_3^+) \leq \mathbf{P}\left(Y \geq \frac{\hat\mu_1^{1/\alpha_2}}{2\varepsilon}\right) \quad\text{and}\quad \limsup_{N\to\infty} \mathbf{P}(E_3^-) \leq \mathbf{P}\left(Y \leq -\frac{\hat\mu_1^{1/\alpha_2}}{2\varepsilon}\right), \tag{3.18}$$

which are both vanishing as $\varepsilon \searrow 0$.

We therefore see that (3.11), (3.15) and (3.18) yield that, provided that ε_0 is small enough, for every $\varepsilon < \varepsilon_0$ and N large enough (how large depends on ε), $P_\varepsilon(x) \geqslant (1-\eta)/\hat\mu_1$ uniformly for $x \in \{1,\dots,1/\varepsilon\}$. If we now go back to (3.6) and (3.8), and using that $\sum_{x\geqslant 1} p_h(x) = 1$, we obtain(2.4). □

3.2. **Proof of** (2.3)**.** In view of (2.4), we simply need to give an upper bound on the probability $\mathbf{P}\left((N, \gamma_c N + t_N) \in \hat\tau\right)$. Fix some $\varepsilon > 0$.

First step. We control

$$\begin{aligned}&\mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} > (1-\varepsilon)t_N, \hat\tau^{(1)}_{\kappa_N} = N, \hat\tau^{(2)}_{\kappa_N} = \gamma_c N + t_N\Big)\\ &\quad\leq \sum_{k=1}^N k \sum_{y>-\varepsilon t_N}\sum_{x\in\mathbb{N}} \mathbf{P}\left(\hat\tau_1^{(2)} = t_N + y,\, \hat\tau_1^{(1)} = x\right)\mathbf{P}\left(\hat\tau_{k-1}^{(1)} = N - x, \hat\tau_{k-1}^{(2)} = \gamma_c N - y\right).\end{aligned} \tag{3.19}$$

Recalling (1.11) and (1.12), we have that there is some $N_0 = N_0(\varepsilon)$ and some $\eta = \eta_\varepsilon$ (with $\eta_\varepsilon \to 0$ as $\varepsilon \downarrow 0$), such that for all $N \geqslant N_0$, $x \leqslant 1/\varepsilon$ and $y > -\varepsilon t_N$ we have

$$\mathbf{P}\Big(\hat\tau_1^{(2)} = t_N + y,\, \hat\tau_1^{(1)} = x\Big) = \frac{L\left(t_N + y + x\right)}{(t_N + y + x)^{2+\alpha}} e^{h - x_{\mathrm{N}_h}} \leqslant (1+\eta)\mathbf{P}\Big(\hat\tau_1^{(2)} = t_N\Big) p_h(x), \tag{3.20}$$

where we recall that $p_h(x) := (e^{\mathrm{N}_h} - 1)e^{-x\mathrm{N}_h}$. Note that we also have that there is a constant c such that uniformly for $x \in \mathbb{N}$

$$\mathbf{P}\Big(\hat{\tau}_1^{(2)} = z,\, \hat{\tau}_1^{(1)} = x\Big) \leqslant cL(z)z^{-(2+\alpha)}p_h(x)\,. \tag{3.21}$$

We can use that to get that uniformly for $y \geqslant -t_N/2$ (so that $t_N + y \geqslant t_N/2$) we have that for any $x \geqslant 1$

$$\mathbf{P}\Big(\hat{\tau}_1^{(2)} = t_N + y,\, \hat{\tau}_1^{(1)} = x\Big) \leqslant c' p_h(x)\mathbf{P}\Big(\hat{\tau}_1^{(2)} = t_N\Big). \tag{3.22}$$

Then, dividing (3.19) according to whether $x \leq 1/\varepsilon$ or $x > 1/\varepsilon$ (and summing over $y > \varepsilon t_N$), we obtain the following upper bound

$$\begin{aligned}(1+\eta)\sum_{x=1}^{1/\varepsilon} p_h(x)\mathbf{P}\Big(\hat{\tau}_1^{(2)} = t_N\Big)\sum_{k=1}^{N} k\mathbf{P}\Big(\hat{\tau}_{k-1}^{(1)} = N - x\Big)\\ + c\sum_{x=1/\varepsilon}^{N} p_h(x)\mathbf{P}\Big(\hat{\tau}_1^{(2)} = t_N\Big)\sum_{k=1}^{N} k\mathbf{P}\Big(\hat{\tau}_{k-1}^{(1)} = N - x\Big)\,.\end{aligned} \tag{3.23}$$

The second term is bounded from above by (using also $k \leqslant n$)

$$cN\mathbf{P}\Big(\hat{\tau}_1^{(2)} = t_N\Big)\sum_{x>1/\varepsilon} p_h(x) = ce^{-\mathrm{N}_h/\varepsilon} \times N\mathbf{P}\Big(\hat{\tau}_1^{(2)} = t_N\Big)\,. \tag{3.24}$$

In the first term (3.23), we split the sum according to whether k is smaller or greater than $(1+\varepsilon)N/\hat{\mu}_1$: we get that

$$\begin{aligned}&\sum_{k=1}^{N} k\mathbf{P}\Big(\hat{\tau}_{k-1}^{(1)} = N - x\Big)\\ &\qquad \leq (1+\varepsilon)\frac{N}{\hat{\mu}_1}\mathbf{P}\Big(N - x \in \hat{\tau}^{(1)}\Big) + N\mathbf{P}\Big(\exists k > (1+\varepsilon)N/\hat{\mu}_1 \ \text{ s.t. } \hat{\tau}_{k-1}^{(1)} = N - x\Big)\\ &\qquad \leq (1+\varepsilon)^2\frac{N}{\hat{\mu}_1^2} + N\mathbf{P}\Big(\hat{\tau}^{(1)}_{(1+\varepsilon)N/\hat{\mu}_1} \leqslant N - x\Big)\,,\end{aligned} \tag{3.25}$$

where we used that in the first part $k \leqslant (1+\varepsilon)N/\hat{\mu}_1$, and the renewal theorem to get that $\mathbf{P}\left(N - x \in \hat{\tau}^{(1)}\right) \leqslant (1+\varepsilon)N/\hat{\mu}_1$ uniformly for $x \leq 1/\varepsilon$ and N large enough (how large depends on ε). The second term is exponentially small since it is a large deviation for $\hat{\tau}^{(1)}$ (x here is bounded by $1/\varepsilon$). Recalling that $\sum p_h(x) = 1$, the first term (3.23) is therefore bounded from above by

$$(1+\eta)\left((1+\varepsilon)^2 + e^{-c_\varepsilon N}\right)\frac{N}{\hat{\mu}_1^2}\mathbf{P}\left(\hat{\tau}_1^{(2)} = t_N\right)\,, \tag{3.26}$$

In the end, the left-hand side of (3.19) is bounded by

$$\left(1+\eta'_\varepsilon\right)\frac{N}{\hat{\mu}_1^2}\mathbf{P}\left(\hat{\tau}_1^{(2)} = t_N\right) \qquad \text{with } \eta'_\varepsilon \overset{\varepsilon\to 0}{\to} 0\,. \tag{3.27}$$

Second step. It remains to control

$$\begin{aligned}\mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} \leqslant (1-\varepsilon)t_N, \hat\tau^{(1)}_{\kappa_N} = N, \hat\tau^{(2)}_{\kappa_N} = \gamma_c N + t_N\Big)& \\ = \mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} \in (\varepsilon t_N, (1-\varepsilon)t_N)\,, \hat\tau^{(1)}_{\kappa_N} = N, \hat\tau^{(2)}_{\kappa_N} = \gamma_c N + t_N\Big)& \quad (3.28)\\ + \mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} \le \varepsilon t_N, \hat\tau^{(1)}_{\kappa_N} = N, \hat\tau^{(2)}_{\kappa_N} = \gamma_c N + t_N\Big)\,.& \quad (3.29)\end{aligned}$$

The first term in the right-hand side, that is (3.28), is smaller than

$$\begin{aligned}&\sum_{k=1}^N k \sum_{z=\varepsilon t_N}^{(1-\varepsilon)t_N} \sum_{x\in\mathbb{N}} \mathbf{P}\Big(\hat\tau^{(2)}_1 = z, \hat\tau^{(1)}_1 = x\Big)\mathbf{P}\Big(\hat\tau^{(2)}_{k-1} = \gamma_c N + t_N - z, \hat\tau^{(1)}_{k-1} = N - x\Big)\\ &\leqslant c\varepsilon^{-(2+\alpha)} N \mathbf{P}\Big(\hat\tau^{(2)}_1 = t_N\Big) \sum_{x\in\mathbb{N}} p_h(x) \sum_{k=1}^N \mathbf{P}\Big(\hat\tau^{(2)}_{k-1} \geqslant \gamma_c N + \varepsilon t_N, \hat\tau^{(1)}_{k-1} = N - x\Big)\end{aligned} \quad (3.30)$$

where we used (3.21) uniformly for $z \geqslant \varepsilon t_N$ and then summed over z to get the first inequality. Then, we split the last sum into two parts. For $k \leqslant k_N^{(\varepsilon)} := N/\hat\mu_1 + \varepsilon^2 t_N$, we have

$$\sum_{k=1}^{k_N^{(\varepsilon)}} \mathbf{P}\left(\hat\tau^{(2)}_{k-1} \geqslant \gamma_c N + \varepsilon t_N, \hat\tau^{(1)}_{k-1} = N - x\right) \leqslant \mathbf{P}\Big(\hat\tau^{(2)}_{k_N^{(\varepsilon)}} \geqslant \gamma_c N + \varepsilon t_N\Big)\,, \quad (3.31)$$

since the events on the left side are disjoints. Then, provided that ε has been fixed small enough so that $\gamma_c N + \varepsilon t_N \geqslant \hat\mu_2 k_N^{(\varepsilon)} + \frac12 \varepsilon t_N$, and since $t_N/a_N^{(2)} \to +\infty$ (and $a^{(2)}_{k_N^{(\varepsilon)}} \leqslant a_N^{(2)}$), we have

$$\limsup_{N\to\infty} \mathbf{P}\Big(\hat\tau^{(2)}_{k_N^{(\varepsilon)}} \geqslant \gamma_c N + \varepsilon t_N\Big) = 0\,. \quad (3.32)$$

On the other hand, for $k \geqslant k_N^{(\varepsilon)}$, we have

$$\sum_{k=k_N^{(\varepsilon)}+1}^N \mathbf{P}\Big(\hat\tau^{(1)}_{k-1} = N - x\Big) \leqslant \mathbf{P}\Big(\hat\tau^{(1)}_{k_N^{(\varepsilon)}} \leqslant N - x\Big) \leqslant \mathbf{P}\Big(\hat\tau^{(1)}_{k_N^{(\varepsilon)}} \leqslant \hat\mu_1 k_N^{(\varepsilon)} - \hat\mu_1 \varepsilon^2 t_N\Big), \quad (3.33)$$

and since $t_N/\sqrt{N} \to +\infty$, also this terms goes to 0 as $N \to \infty$. In the end, we get that the term (3.28) is negligible compared to $N\mathbf{P}\big(\hat\tau^{(2)}_1 = t_N\big)$.

Then, it remains to bound (3.29), and a first observation is that we can restrict it to having $\kappa_N \leqslant k_N^+ := N/\hat\mu_1 + t_N/(4\hat\mu_2)$. Indeed, we have that

$$\begin{aligned}&\mathbf{P}\big(\mathcal{M}_{1,\kappa_N} \leqslant \varepsilon t_N, \hat\tau^{(1)}_{\kappa_N} = N,\, \hat\tau^{(2)}_{\kappa_N} = \gamma_c N + t_N, \kappa_N \geqslant k_N^+\big)\\ &\qquad \le \mathbf{P}\big(\hat\tau^{(1)}_{\kappa_N} = N, \kappa_N \geqslant k_N^+\big) \leqslant \mathbf{P}\big(\hat\tau^{(1)}_{k_N^+} \leqslant N = \hat\mu_1 k_N^+ - (N - \hat\mu_1 k_N^+)\big)\\ &\qquad \le \exp\big(-c\,(N - \hat\mu_1 k_N^+)^2/k_N^+\big) \le \exp\big(-c'\,(t_N)^2/N\big).\end{aligned} \quad (3.34)$$

We mention that the bound for $\mathbf{P}\big(\hat\tau^{(1)}_{k_N^+} \leqslant \hat\mu_1 k_N^+ - j\big)$ comes from the fact that $-\hat\tau^{(1)}_1$ is bounded (so it has exponential moments and one can use a Chernov bound, leading to this Gaussian bound).

Now, the last term decays faster than $N\mathbf{P}\big(\hat\tau^{(2)}_1 = t_N\big)$ because of assumption (1.19), provided that C_0 had been chosen large enough.

It remains to control

$$\sum_{k=1}^{k_N^+} \mathbf{P}\Big(\mathcal{M}_{1,k} \leqslant \varepsilon t_N, \hat{\tau}_k^{(1)} = N,\, \hat{\tau}_k^{(2)} = \gamma_c N + t_N\Big)\,. \tag{3.35}$$

We write that each term in the sum is

$$\sum_{j=\log_2(1/\varepsilon)}^{q_n} \mathbf{P}\left(\mathcal{M}_{1,k} \in (2^{-(j+1)}t_N, 2^{-j}t_N], \hat{\tau}_k^{(1)} = N,\, \hat{\tau}_k^{(2)} = \gamma_c N + t_N\right)\,, \tag{3.36}$$

where q_N is the smallest integer such that $2^{-(q_N+1)}t_N < 1$, so $q_N = O(\log_2 N)$. Then, using (3.21), each term in the sum (i.e. for every k and j) is bounded by a constant (not depending on j and k) times

$$\begin{aligned} k \sum_{z=2^{-(j+1)}t_N}^{2^{-j}t_N} \sum_{x\in\mathbb{N}} \frac{L(2^{-j}t_N)}{(2^{-j}t_N)^{(2+\alpha)}} p_h(x) \times \\ \mathbf{P}\left(\hat{\tau}_{k-1}^{(1)} = N - x, \hat{\tau}_{k-1}^{(2)} = \gamma_c N + t_N - z, \mathcal{M}_{1,k-1} \leqslant 2^{-j}t_N\right) \\ \leqslant \frac{NL(t_N)}{t_N^{2+\alpha}} \sum_{x\in\mathbb{N}} p_h(x) 2^{j(3+\alpha)} \mathbf{P}\left(\hat{\tau}_{k-1}^{(1)} = N - x, \hat{\tau}_{k-1}^{(2)} \geqslant \gamma_c N + \frac{t_N}{2}, \mathcal{M}_{1,k-1} \leqslant 2^{-j}t_N\right), \end{aligned} \tag{3.37}$$

where we used that provided that t_N is large enough, $L(2^{-j}t_N) \leqslant 2^j L(t_N)$ (this is a direct consequence of Potter's bound for slowly varying functions [3, Th. 1.5.6]) and summed over z. Recovering the sum over k and j, we therefore need to show that

$$\sum_{k=1}^{k_N^+} \sum_{j=\log_2(1/\varepsilon)}^{q_n} \sum_{x\in\mathbb{N}} p_h(x) 2^{j(3+\alpha)} \mathbf{P}\Big(\hat{\tau}_{k-1}^{(1)} = N - x, \hat{\tau}_{k-1}^{(2)} \geqslant \gamma_c N + t_N/2, \mathcal{M}_{1,k-1} \leqslant 2^{-j}t_N\Big) \tag{3.38}$$

is small for N large.

Then, for every j, we define $\{\bar{\tau}_k\}_{k=0,1,\dots}$ (with distribution $\bar{\mathbf{P}}^{(j)}$, carrying the dependence on j) as an i.i.d. sum of k variables with distribution $(\hat{\tau}_1^{(1)}, \hat{\tau}_1^{(2)} \mathbf{1}_{\{\hat{\tau}_1^{(2)} \leqslant 2^{-j}t_N\}})$: we therefore obtain that for $k \leqslant k_N^+$

$$\begin{aligned} &\mathbf{P}\Big(\hat{\tau}_{k-1}^{(1)} = N - x, \hat{\tau}_{k-1}^{(2)} \geqslant \gamma_c N + t_N/2, \mathcal{M}_{1,k-1} \leqslant 2^{-j}t_N\Big) \\ &\quad \leqslant \bar{\mathbf{P}}^{(j)}\Big(\bar{\tau}_{k-1}^{(1)} = N - x, \bar{\tau}_{k-1}^{(2)} \geqslant \gamma_c N + t_N/2\Big) \leqslant \bar{\mathbf{P}}^{(j)}\Big(\bar{\tau}_{k-1}^{(1)} = N - x, \bar{\tau}_{k_N^+}^{(2)} \geqslant \gamma_c N + t_N/2\Big)\,. \end{aligned} \tag{3.39}$$

Using this inequality and summing it over k in (3.38), (and then using that $\sum_x p_h(x) = 1$), we obtain that (3.38) is smaller than

$$\sum_{j=\log_2(1/\varepsilon)}^{q_n} 2^{j(3+\alpha)} \bar{\mathbf{P}}^{(j)}\Big(\bar{\tau}_{k_N^+}^{(2)} \geqslant \hat{\mu}_2 k_N^+ + t_N/4\Big)\,, \tag{3.40}$$

where we used that $\gamma_c N \geq \hat{\mu}_2 k_N^+ - \frac{1}{4}t_N$. Then, we may use a Fuk-Nagaev inequality, see for example in [16], to control the last probability – we regroup the inequalities we need under the following lemma.

Lemma 3.1. *Let $\{X_i\}_{i=1,2,\dots}$ be a sequence of i.i.d. non negative r.v. with $\mathbf{P}(X_1 > x) \sim \varphi(x)x^{-\rho}$ with $\rho > 1$ and $\varphi(\cdot)$ a slowly varying function. Denote $\mu := \mathbf{E}[X]$ and $\sigma(y) = \mathbf{E}[X_1^2 \mathbf{1}_{\{X_1 \leqslant y\}}]$. We have that there exist constants $c, c' > 0$ such that for any $y \leqslant x$*

$$\mathbf{P}\Big(\sum_{i=1}^n X_i \mathbf{1}_{\{X_i \leqslant y\}} - \mu n \geqslant x\Big) \leqslant \Big(1 + c' \frac{x}{n\varphi(y)y^{1-\rho}}\Big)^{-cx/y} + e^{-cx^2/n\sigma(y)} \mathbf{1}_{\{\rho \geqslant 2\}}. \quad (3.41)$$

Moreover, for any $u > 0$ there is a constant c_u such that, if $x/(n\varphi(y)y^{1-\rho}) \leqslant u$, then

$$\mathbf{P}\Big(\sum_{i=1}^n X_i \mathbf{1}_{\{X_i \leqslant y\}} - \mu n \geqslant x\Big) \leqslant \exp\Big(-c_u \frac{x^2}{n\sigma(y)}\Big). \quad (3.42)$$

The first term on the right hand side of (3.41) roughly corresponds to the probability to make x/y anomalous jumps of order y, and the second term to a *collective* Cramér strategy, with no exceptional jump.

Lemma 3.1 can be extracted from [16]: (3.41) is taken from [16, Theorem 1.2] ($\rho \in (1,2)$) and [16, Corollary 1.6] ($\rho \geqslant 2$); (3.42) comes from Theorem 1.3 in [16]—we use that $\sigma(y) = \mathbf{E}[X_1^2 \mathbf{1}_{\{X_1 \leqslant y\}}] \geqslant c'' y^{2-\rho}\varphi(y)$, so that $\log(1 + c'x/(n\varphi(y)y^{1-\rho})) \geqslant \delta_u xy/n\sigma(y)$ if $c'x/(n\varphi(y)y^{1-\rho}) \leqslant u$ (this is condition (1.4) in [16, Theorem 1.3]).

Applying (3.41) to $X_1 = \hat\tau_1^{(2)}$ (*i.e.* $\rho = 1 + \alpha$ and $\varphi(\cdot)$ a constant times $L(\cdot)$), and $x = \varepsilon t_N$, $y = 2^{-j} t_N$ (so $y \le x$ in the summation range for j, cf. (3.40)), we get that (using that $\sigma(y) \le \sigma(x)$ and $k_N^+ \leqslant N$ for the term $\alpha \geqslant 1$)

$$\bar{\mathbf{P}}^{(j)}\Big(\bar\tau_{k_N^+}^{(2)} \geqslant \mu_2 k_N^+ + t_N/4\Big) \leqslant \left(1 + c' \frac{\varepsilon 2^j}{N\mathbf{P}(\tau_1^{(2)} > 2^{-j} t_N)}\right)^{-c\varepsilon 2^j} + e^{-c''\varepsilon^2 \frac{t_N^2}{N\sigma(t_N)}} \mathbf{1}_{\{\alpha \geqslant 1\}}. \quad (3.43)$$

The second term poses no issue: summing it for $j \in [\log_2(1/\varepsilon), q_n]$, we get that

$$\sum_{j=\log_2(1/\varepsilon)}^{q_n} 2^{j(3+\alpha)} e^{-c''t_N^2/N\sigma(t_N)} \mathbf{1}_{\{\alpha \geqslant 1\}} \leqslant N^{3+\alpha} e^{-c''t_N^2/N\sigma(t_N)} \mathbf{1}_{\{\alpha \geqslant 1\}}, \quad (3.44)$$

and this is small when $N \to +\infty$ thanks to assumption (1.19), provided that the constant C_0 has been fixed large enough in the case $\alpha \geqslant 1$.

For the first term in (3.43), let $j_\varepsilon = j_\varepsilon(t_N)$ be the first j such that $N\mathbf{P}\big(\tau_1^{(2)} > 2^{-j} t_N\big) \geqslant c'\varepsilon^A 2^j$ for some constant A large enough (the precise choice is made below): note that $2^{-j_\varepsilon} t_N$ is larger than $a_N^{(2)}$ (since $N\mathbf{P}(\tau_1^{(2)} > a_N^{(2)}) \to 1$), so that $j_\varepsilon \geqslant \log_2(t_n/a_N^{(2)}) \to +\infty$. Summing over $j \in [\log_2(1/\varepsilon), j_\varepsilon)$ the first term in (3.43), we get with the choice $c(A-1) \geqslant 4 + \alpha$ and ε sufficiently small

$$\begin{aligned}
\sum_{j=\log_2(1/\varepsilon)}^{j_\varepsilon - 1} 2^{j(3+\alpha)} \big(1 + \varepsilon^{1-A}\big)^{-c\varepsilon 2^j} &\leqslant \sum_{j=\log_2(1/\varepsilon)}^{j_\varepsilon - 1} 2^{j(3+\alpha)} \varepsilon^{c(A-1)\varepsilon 2^j} \\
&\leqslant \sum_{j=0}^{\infty} 2^{(3+\alpha)(j+\log_2(1/\varepsilon)) - (4+\alpha)(2^j-1)\log_2(1/\varepsilon) - (4+\alpha)\log_2(1/\varepsilon)} \\
&= \varepsilon \sum_{j=0}^{\infty} 2^{(3+\alpha)j - (4+\alpha)(2^j-1)\log_2(1/\varepsilon)} \leqslant 2\varepsilon. \quad (3.45)
\end{aligned}$$

For $j \geqslant j_\varepsilon$, we use (3.42). In the case $\alpha \geqslant 1$, we therefore have the same reasoning as in (3.44): the sum for $j \in [j_\varepsilon, q_n]$ is small. If $\alpha < 1$, then $\sigma(y) = \mathbf{E}[X_1^2 \mathbf{1}_{\{X_1 \leqslant y\}}] \leqslant \widetilde{c} y^2 \varphi(y) y^{-\rho}$, and (3.43) becomes, for $j \geqslant j_\varepsilon$

$$\bar{\mathbf{P}}^{(j)}\Big(\bar{\tau}^{(2)}_{k_N^+} \geqslant \mu_2 k_N^+ + t_N/4\Big) \leqslant \exp\big(-c_\varepsilon 2^{2j}(N\mathbf{P}(\tau_1^{(2)} > 2^{-j}t_N))^{-1}\big) \leqslant \exp\big(-c'_\varepsilon 2^{2j}\big),$$

where we used in the second inequality that $(N\mathbf{P}(\tau_1^{(2)} > 2^{-j}t_N))^{-1}$ is bounded above by a constant if $j \geq j_\varepsilon$. Then, summing over $j \in [j_\varepsilon, q_N]$, we get

$$\sum_{j=j_\varepsilon(t_N)}^{q_N} 2^{j(3+\alpha)} e^{-c'_\varepsilon 2^{2j}} \to 0 \quad \text{as } N \to +\infty. \tag{3.46}$$

All together, we have shown that (3.40) is small. □

4. The free partition function: proof of Theorem 2.2

We will first prove (2.6) for the case $\sum_j K_f(j) < \infty$. Many estimates are in common with the case $\sum_j K_f(j) = \infty$ that we treat right after, and we will stress along the proof when the estimates are dependent or not on the fact that $\sum_j K_f(j) < \infty$. Also, the proof of the lower bounds (2.7) and (2.8) are contained in the proof of (2.6) as we explain along the way.

Proof of Theorem 2.2. As announced, we start with the proof of (2.6) and assume $\sum_j K_f(j) < \infty$. Let us fix $\eta > 0$, and $\varepsilon > 0$ small, how small depends on η as will be stressed in the proof.

We decompose the free partition function into several parts:

$$\begin{aligned} Z^f_{N,M,h} &= \mathrm{I} + \mathrm{II} + \mathrm{III} + \mathrm{IV} + \mathrm{V} \\ \text{with} \qquad \mathrm{I} &= Z^f_{N,M,h}\big(V_1^{(N)} \geqslant 1/\varepsilon\big) \\ \mathrm{II} &= Z^f_{N,M,h}\big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} \leqslant 1/\varepsilon\big) \\ \mathrm{III} &= Z^f_{N,M,h}\big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} \in (1/\varepsilon, t_N - \tfrac{1}{\varepsilon}a_N^{(2)})\big) \\ \mathrm{IV} &= Z^f_{N,M,h}\big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} \in [t_N - \tfrac{1}{\varepsilon}a_N^{(2)}, t_N + \tfrac{1}{\varepsilon}a_N^{(2)}]\big) \\ \mathrm{V} &= Z^f_{N,M,h}\big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} > t_N + \tfrac{1}{\varepsilon}a_N^{(2)}\big). \end{aligned} \tag{4.1}$$

The main contribution comes from the terms II and IV. We first estimate these terms, before showing that all the other ones are negligible compared to $\max(\mathrm{II}, \mathrm{IV})$.

4.1. Main terms, and proof of (2.7) and (2.8).

Analysis of II and proof of (2.7). Recall (1.3) and (1.10). The term II can be written as

$$\mathrm{II} := \sum_{i<1/\varepsilon} \sum_{j \leqslant 1/\varepsilon} K_f(i) K_f(j) e^{(N-i)\mathrm{N}_h} \mathbf{P}((N-i, M-j) \in \hat{\tau}), \tag{4.2}$$

and it is just a matter of estimating $\mathbf{P}((N-i, M-j) \in \hat{\tau})$ uniformly for $0 \leqslant i, j \leqslant 1/\varepsilon$. We have from Theorem 2.1, uniformly for $i, j \leqslant 1/\varepsilon$,

$$\mathbf{P}((N-i, M-j) \in \hat{\tau}) \overset{N\to\infty}{\sim} \frac{N-i}{\hat{\mu}_1^2} \mathbf{P}(\hat{\tau}_1^{(2)} = M - j - \gamma_c(N-i)) \sim \frac{N}{\hat{\mu}_1^2} \mathbf{P}(\hat{\tau}_1^{(2)} = t_N). \tag{4.3}$$

(Recall $t_n := M - \gamma_c N$ and (1.12).) Hence,

$$\mathrm{II} \sim \frac{N}{\hat{\mu}_1^2} \mathbf{P}(\hat{\tau}_1^{(2)} = t_N) e^{N \mathrm{N}_h} \sum_{i \leqslant 1/\varepsilon} e^{-i \mathrm{N}_h} K_f(i) \sum_{j \leqslant 1/\varepsilon} K_f(j) \,. \tag{4.4}$$

Hence, if $\overline{K} = \sum_{j \in \mathbb{N}} K_f(j) < +\infty$, we get that provided that ε has been fixed small enough (depending on η), for all N sufficiently large,

$$\begin{aligned} \mathrm{II} &\geq (1-\eta)\overline{K} \, \frac{N}{\hat{\mu}_1^2} e^{N \mathrm{N}_h} \Big(\sum_{i \geq 0} e^{-i \mathrm{N}_h} K_f(i) \Big) \mathbf{P}(\hat{\tau}_1^{(2)} = t_N) \,, \\ \mathrm{II} &\leq (1+\eta)\overline{K} \, \frac{N}{\hat{\mu}_1^2} e^{N \mathrm{N}_h} \Big(\sum_{i \geq 0} e^{-i \mathrm{N}_h} K_f(i) \Big) \mathbf{P}(\hat{\tau}_1^{(2)} = t_N) \,. \end{aligned} \tag{4.5}$$

The lower bound (2.7) is obtained simply by using the estimate (2.4) instead of (2.3) in (4.2): the straightforward details are left to the reader.

Analysis of IV and proof of (2.8). It can be written as

$$\mathrm{IV} = \sum_{i \leqslant 1/\varepsilon} \sum_{j = t_N - \frac{1}{\varepsilon} a_N^{(2)}}^{t_N + \frac{1}{\varepsilon} a_N^{(2)}} K_f(i) K_f(j) e^{(N-i)\mathrm{N}_h} \mathbf{P}((N-i, M-j) \in \hat{\tau}). \tag{4.6}$$

We have that uniformly for $j \in [t_N - \frac{1}{\varepsilon} a_N^{(2)}, t_N + \frac{1}{\varepsilon} a_N^{(2)}]$, $K_f(j) \sim K_f(t_N)$. We can therefore focus on estimating, uniformly for $i \leqslant 1/\varepsilon$

$$\begin{aligned} &\sum_{j = t_N - \frac{1}{\varepsilon} a_N^{(2)}}^{t_N + \frac{1}{\varepsilon} a_N^{(2)}} \mathbf{P}\big((N-i, M-j) \in \hat{\tau}\big) \\ &\qquad = \mathbf{P}\Big(\text{for some } k,\ \hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} \in [\gamma_c N - \tfrac{1}{\varepsilon} a_N^{(2)}, \gamma_c N + \tfrac{1}{\varepsilon} a_N^{(2)}]\Big) \,, \end{aligned}$$

and now we prove that this term is close to $1/\hat{\mu}_1$. In fact we have

$$\begin{aligned} &\mathbf{P}\Big(\text{for some } k,\ \hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} \in [\gamma_c N - \tfrac{1}{\varepsilon} a_N^{(2)}, \gamma_c N + \tfrac{1}{\varepsilon} a_N^{(2)}]\Big) \\ &= \mathbf{P}(N-i \in \hat{\tau}^{(1)}) - \mathbf{P}\Big(\text{for some } k,\ \hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} \notin [\gamma_c N - \tfrac{1}{\varepsilon} a_N^{(2)}, \gamma_c N + \tfrac{1}{\varepsilon} a_N^{(2)}]\Big), \end{aligned}$$

and we now show that, provided that ε had been fixed small enough, uniformly for $i \leqslant 1/\varepsilon$ and N large enough:

$$\mathbf{P}\Big(\hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} < \gamma_c N - \tfrac{1}{\varepsilon} a_N^{(2)} \text{ for some } k\Big) \leq \eta \,, \tag{4.7}$$

$$\mathbf{P}\Big(\hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} > \gamma_c N + \tfrac{1}{\varepsilon} a_N^{(2)} \text{ for some } k\Big) \leq \eta \,. \tag{4.8}$$

This will be enough, since by the Renewal Theorem we have that $\mathbf{P}(N-i \in \hat{\tau}^{(1)}) \to \hat{\mu}_1^{-1}$ uniformly for $i \leqslant 1/\varepsilon$.

To treat (4.7), define $k_N := \frac{1}{\hat\mu_1}N - \frac{1}{2\hat\mu_2\varepsilon}a_N^{(2)}$: we have uniformly for $i \le 1/\varepsilon$

$$\begin{aligned}
&\mathbf{P}\Big(\text{for some } k,\ \hat\tau_k^{(1)} = N-i, \hat\tau_k^{(2)} < \gamma_c N - \tfrac{1}{\varepsilon}a_N^{(2)}\Big)\\
&\qquad = \mathbf{P}\Big(\text{for some } k \leqslant k_N,\ \hat\tau_k^{(1)} = N-i, \hat\tau_k^{(2)} < \gamma_c N - \tfrac{1}{\varepsilon}a_N^{(2)}\Big)\\
&\qquad\qquad + \mathbf{P}\Big(\text{for some } k \geqslant k_N,\ \hat\tau_k^{(1)} = N-i, \hat\tau_k^{(2)} < \gamma_c N - \tfrac{1}{\varepsilon}a_N^{(2)}\Big)\\
&\qquad \leqslant \mathbf{P}\left(\hat\tau_{k_N}^{(1)} \geqslant N - 1/\varepsilon\right) + \mathbf{P}\left(\hat\tau_{k_N}^{(2)} < \gamma_c N - \tfrac{1}{\varepsilon}a_N^{(2)}\right).
\end{aligned} \tag{4.9}$$

Now, it is easy to see that the two terms in the last line are small: we indeed have that for arbitrary $\eta' > 0$, one can choose ε small enough so that for all N large enough,

$$\begin{aligned}
&\mathbf{P}\left(\hat\tau_{k_N}^{(1)} \geqslant N - 1/\varepsilon\right) = \mathbf{P}\left(\hat\tau_{k_N}^{(1)} \geqslant \hat\mu_1 k_N + \frac{\hat\mu_1}{2\hat\mu_2\varepsilon}a_N^{(2)} - 1/\varepsilon\right) \le \eta',\\
\text{and}\quad &\mathbf{P}\left(\hat\tau_{k_N}^{(2)} < \gamma_c N - \frac{1}{\varepsilon}a_N^{(2)}\right) = \mathbf{P}\left(\hat\tau_{k_N}^{(2)} < \hat\mu_2 k_N - \frac{1}{2\varepsilon}a_N^{(2)}\right) \le \eta'.
\end{aligned} \tag{4.10}$$

For the first line, we used that $\frac{\hat\mu_1}{2\hat\mu_2\varepsilon}a_N^{(2)} - 1/\varepsilon \ge \varepsilon^{-1/2}\sqrt{N} \ge \varepsilon^{-1/2}\sqrt{k_N}$ provided that N is large enough (and ε small), and then simply Chebichev's inequality. For the second line, we used that $\gamma_c = \hat\mu_2/\hat\mu_1$ to get that $\gamma_c N = \hat\mu_2 k_N + \frac{1}{2\varepsilon}a_N^{(2)}$, and then the approximation of $(a_{k_N}^{(2)})^{-1}(\hat\tau_{k_N} - k_N\hat\mu_2)$ by an α_2-stable distribution, as done in (3.18).

For (4.8), we define $k'_N = \frac{1}{\hat\mu_1}N + \frac{1}{2\hat\mu_2\varepsilon}a_N^{(2)}$, and similarly to what is done above, we have

$$\begin{aligned}
&\mathbf{P}\Big(\text{for some } k,\ \hat\tau_k^{(1)} = N-i, \hat\tau_k^{(2)} > \gamma_c N + \tfrac{1}{\varepsilon}a_N^{(2)}\Big)\\
&\qquad\qquad \leqslant \mathbf{P}\left(\hat\tau_{k'_N}^{(1)} \leqslant N-i\right) + \mathbf{P}\left(\hat\tau_{k'_N}^{(2)} > \gamma_c N + \tfrac{1}{\varepsilon}a_N^{(2)}\right),
\end{aligned}$$

and both terms are smaller than η' provided that ε had been fixed small enough and N is large, for the same reasons as in (4.10).

In the end, we get that provided that ε had been fixed small enough, for all sufficiently large N

$$\begin{aligned}
\mathrm{IV} &\ge (1-\eta)\frac{1}{\hat\mu_1}K_f(t_N)e^{N\mathrm{N}_h}\sum_{i\le 1/\varepsilon} e^{-i\mathrm{N}_h}K_f(i)\,,\\
\mathrm{IV} &\le (1+\eta)\frac{1}{\hat\mu_1}K_f(t_N)e^{N\mathrm{N}_h}\sum_{i\le 1/\varepsilon} e^{-i\mathrm{N}_h}K_f(i)\,.
\end{aligned} \tag{4.11}$$

Obviously, since the last sum converges, we can replace it with the infinite sum, and simply replace η by 2η provided that ε is small enough. This competes the analysis of IV.

For what concerns (2.8) we simply need to show that

$$\begin{aligned}
\mathrm{IVb} &:= Z^f_{N,M,h}\Big(V_1^{(N)} \le \frac{1}{\varepsilon}, V_2^{(N)} \in \big[t_N - \tfrac{1}{\varepsilon}a_N^{(2)}, t_N + \tfrac{1}{\varepsilon}a_N^{(2)}\big], \mathcal{M}_{1,\kappa_N} > \tfrac{1}{\varepsilon}m_N^{(2)}\Big)\\
&= \sum_{i\,\leqslant\,1/\varepsilon}\ \sum_{j=t_N-\frac{1}{\varepsilon}a_N^{(2)}}^{t_N+\frac{1}{\varepsilon}a_N^{(2)}} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}\mathbf{P}\Big((N-i, M-j)\in\hat\tau, \mathcal{M}_{1,\kappa_N} > \tfrac{1}{\varepsilon}m_N^{(2)}\Big).
\end{aligned}$$

is negligible compared to (4.11). But again, uniformly for the range of j considered, we have $K_f(j) \le 2K_f(t_N)$ (provided that N is large enough). Then, dropping the event $N-i \in \hat\tau^{(1)}$, and summing over j, we get that

$$\mathrm{IVb} \le 2K_f(t_N)e^{N\mathrm{N}_h}\Big(\sum_{i\le 1/\varepsilon} K_f(i)e^{-i\mathrm{N}_h}\Big)\mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} > \tfrac{1}{\varepsilon}m_N^{(2)}\Big).$$

Then, using that $\kappa_N \le N$, we get that

$$\mathbf{P}\Big(\mathcal{M}_{1,\kappa_N} > \tfrac{1}{\varepsilon}m_N^{(2)}\Big) \le \mathbf{P}\Big(\max_{1\le k\le N}(\hat\tau_k^{(2)} - \hat\tau_{k-1}^{(2)}) > \frac{1}{\varepsilon}m_N^{(2)}\Big) \le N\mathbf{P}\big(\hat\tau_1^{(2)} > \tfrac{1}{\varepsilon}m_N^{(2)}\big),$$

which can be made arbitrarily small by choosing ε small (uniformly in N), thanks to the definition (1.16) of $m_N^{(2)}$. Hence IVb is negligible compared to IV. We also stress here that to estimate IV – in particular to obtain (4.11) –, we did not make use of the assumption $\sum K_f(i) < +\infty$.

4.2. **Remaining terms.** It remains to estimate the terms I, III and V in (4.1), and show that they are negligible compared to (4.4) or (4.11). We start by parts III and V.

Analysis of III. Assume that N is large enough, so that $\frac{1}{\varepsilon}a_N^{(2)} \leqslant \frac{1}{2}t_N$ we write

$$\begin{aligned}\mathrm{III} \leqslant\ & Z^f_{N,M,h}\Big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} \in (1/\varepsilon, \tfrac{1}{2}t_N)\Big) \quad \text{(denoted IIIa)}\\ &+ Z^f_{N,M,h}\Big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} \in (\tfrac{1}{2}t_N, t_N - \tfrac{1}{\varepsilon}a_N^{(2)})\Big) \quad \text{(denoted IIIb)}\end{aligned} \tag{4.12}$$

The first term is

$$\mathrm{IIIa} = \sum_{i<1/\varepsilon}\sum_{j=1/\varepsilon}^{t_N/2} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}\mathbf{P}\big((N-i, M-j)\in\hat\tau\big). \tag{4.13}$$

Now we can bound, uniformly for $i < 1/\varepsilon$ and $j \leqslant t_N/2$ (so that $(M-j)-\gamma_c(N-i) \geqslant t_N/4$ for N sufficiently large, and we can apply Theorem 2.1)

$$\mathbf{P}\big((N-i, M-j)\in\hat\tau\big) \leqslant cN \sup_{m\,\geqslant\, t_N/4} \mathbf{P}(\hat\tau_1^{(2)} = m) \leqslant c'N\mathbf{P}(\hat\tau_1^{(2)} = t_N). \tag{4.14}$$

Hence we get

$$\mathrm{IIIa} \le c'N\mathbf{P}\left(\hat\tau_1^{(2)} = t_N\right)e^{N\mathrm{N}_h}\sum_i e^{-i\mathrm{N}_h}K_f(i)\sum_{j=1/\varepsilon}^{t_N/2}K_f(j), \tag{4.15}$$

and in the case when $\sum K_f(j) < +\infty$, the last sum can be made arbitrarily small by choosing ε small. Hence, recalling (4.4), we get that $\mathrm{IIIa} \le \eta \times \mathrm{II}$ for all N sufficiently large, provided that ε is small enough.

For the term IIIb, we use that $K_f(j) \leqslant cK_f(t_N)$ for j between $t_N/2$ and $t_N - \varepsilon^{-1}a_N^{(2)} \leqslant 2t_N$ ($K_f(\cdot)$ is regularly varying), to get that

$$\begin{aligned}\mathrm{IIIb} &\leq \sum_{i \leqslant 1/\varepsilon} \sum_{j=t_N/2}^{t_N - \frac{1}{\varepsilon}a_N^{(2)}} K_f(i) c K_f(t_N) e^{(N-i)\mathrm{N}_h} \mathbf{P}\left((N-i, M-j) \in \hat{\tau}\right) \\ &\leq c\, K_f(t_N) e^{N\mathrm{N}_h} \sum_{i \leqslant 1/\varepsilon} e^{-i\mathrm{N}_h} K_f(i) \mathbf{P}\big(\text{for some } k,\ \hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} > \gamma_c N + \tfrac{1}{\varepsilon}a_N^{(2)}\big).\end{aligned} \tag{4.16}$$

Since we have seen in (4.8) that the last probability is smaller than some arbitrary η' for all N large enough (provided that $\varepsilon > 0$ is small enough), uniformly for all $i \leq 1/\varepsilon$ we have that IIIb $\leq \eta \times$ IV (recall (4.11)). We stress that, here again, we do not make use of the assumption $\sum K_f(i) < +\infty$.

In the end, we obtain that III $\leq \eta \times$ (II + IV) (provided that N is large enough).

Analysis of V. Let us split V into two parts:

$$\begin{aligned}\mathrm{Va} &= Z_{N,M,h}^f\big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} \in (t_N + \tfrac{1}{\varepsilon}a_N^{(2)}, 2t_N)\big), \\ \mathrm{Vb} &= Z_{N,M,h}^f\big(V_1^{(N)} < 1/\varepsilon; V_2^{(N)} > 2t_N\big).\end{aligned}$$

Let us start with Va. We proceed analogously as above: using that $K_f(j) \leqslant cK_f(t_N)$ uniformly for $j \in (t_N, 2t_N)$, we get that

$$\begin{aligned}\mathrm{Va} &\leqslant \sum_{i \leqslant 1/\varepsilon} \sum_{j \geqslant t_N + \frac{1}{\varepsilon}a_N^{(2)}} K_f(i) c K_f(t_N) e^{(N-i)\mathrm{N}_h} \mathbf{P}\big((N-i, M-j) \in \hat{\tau}\big) \\ &\leqslant cK_f(t_N) e^{N\mathrm{N}_h} \sum_{i \leqslant 1/\varepsilon} e^{-i\mathrm{N}_h} K_f(i) \mathbf{P}\big(\text{for some } k,\ \hat{\tau}_k^{(1)} = N-i, \hat{\tau}_k^{(2)} < \gamma_c N - \tfrac{1}{\varepsilon}a_N^{(2)}\big).\end{aligned}$$

Now we again recall (4.7), which tells that the last probability is smaller than some arbitrary η' for all N large enough (provided that $\varepsilon > 0$ is small enough, uniformly for all $i \leq 1/\varepsilon$). In the end, in view of (4.11), we get that Va $\leq \eta \times$ IV, and here again we did not make use of the assumption $\sum K_f(i) < +\infty$.

For the term Vb, we write

$$\mathrm{Vb} = \sum_{i \leqslant 1/\varepsilon} \sum_{j \geqslant 2t_N} K_f(i) K_f(j) e^{(N-i)\mathrm{N}_h} \mathbf{P}\big((N-i, M-j) \in \hat{\tau}\big). \tag{4.17}$$

Then, we estimate the last probability: by setting $k_N := \frac{1}{\hat{\mu}_1}N - \frac{1}{2\hat{\mu}_2}j$ (for some small c_0), we obtain as in (4.9)-(4.10) that

$$\begin{aligned}\mathbf{P}\big((N-i, M-j) \in \hat{\tau}\big) &\leqslant \mathbf{P}\Big(\hat{\tau}_{k_N}^{(1)} \geqslant N - 1/\varepsilon\Big) + \mathbf{P}\Big(\hat{\tau}_{k_N}^{(1)} \leqslant \gamma_c N - j\Big) \\ &\leqslant \mathbf{P}\Big(\hat{\tau}_{k_N}^{(1)} \geqslant \hat{\mu}_1 k_N + \frac{\hat{\mu}_1}{2\hat{\mu}_2} j - 1/\varepsilon\Big) + \mathbf{P}\Big(\hat{\tau}_{k_N}^{(2)} \leqslant \hat{\mu}_2 k_N - j/2\Big).\end{aligned} \tag{4.18}$$

Then, the two terms are bounded by e^{-cj^2/k_N}: recall that $\hat{\tau}^{(1)}$ has an exponential tail, and that $-\hat{\tau}^{(2)}$ is bounded above, so that they have some exponential moments (and we

may use standard Chernov bounds, as in (3.34)). All together, we get that

$$\mathrm{Vb} \leqslant e^{N\mathrm{N}_h}\Big(\sum_{i\leqslant 1/\varepsilon} K_f(i)e^{-i\mathrm{N}_h}\Big)\Big(\sum_{j=2t_N}^{M} K_f(j)e^{-cj^2/N}\Big) \leqslant CN^{c+1}\exp\big(-ct_N^2/N\big)\,, \quad (4.19)$$

where we used that $K_f(j)$ is bounded by N^c for all $j \leqslant M$, for some $c > 0$. Then, if $\alpha < 1$ then there is a $\delta > 0$ such that $t_N \geqslant a_N^{(2)} \geqslant N^{1/2+\delta}$, so that $\exp(-c(t_N)^2/N)$ decays faster than any power of N. In the case $\alpha \geqslant 1$, we get thanks to (1.19) that $\exp(-c(t_N)^2/N)$ decays faster than $N^{-c'C_0}$: hence choosing C_0 arbitrarily large, we get that (4.19) is negligible compared to IV.

Analysis of I. We separate it into two parts: $V_1^{(N)} \geqslant (\log N)^2$, and $V_1^{(N)} \in (1/\varepsilon, (\log N)^2)$. We have

$$\begin{aligned} Z^f_{N,M,h}\Big(V_1^{(N)} \geqslant (\log N)^2\Big) &= \sum_{i\geqslant(\log N)^2}\sum_{j=0}^{M} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}\mathbf{P}\big((N-i,M-j)\in\hat\tau\big) \\ &\leqslant \Big(\sum_{j=0}^{M}K_f(j)\Big)e^{N\mathrm{N}_h}\sum_{i=(\log N)^2}^{N} e^{-i\mathrm{N}_h}K_f(i) \leqslant cN^{2c+2}e^{N\mathrm{N}_h}e^{-(\log N)^2\mathrm{N}_h}, \end{aligned} \quad (4.20)$$

where we first simply bounded the probability by 1, and also that there is some constant $c > 0$ such that $K_f(i) \le cN^c$ for $i \le N, M$. Clearly, in view of (4.11) (or (4.4)), we get that $Z^f_{N,M,h}(V_1^{(N)} \geqslant (\log N)^2) = o(\mathrm{IV})$, since $1/K_f(t_N)$ and $1/\mathbf{P}(\hat\tau_1^{(2)} = t_N)$ are $O(N^{c'})$ for some $c' > 0$. Again, we did not use that $\sum K_f(i) < +\infty$, even if it would have simplified the upper bound.

We now turn to the case when $V_1^{(N)} \le (\log N)^2$. We write

$$\begin{aligned} Z^f_{N,M,h}\Big(V_1^{(N)} \in [1/\varepsilon, (\log N)^2)\Big) \;=\; & Z^f_{N,M,h}\Big(V_1^{(N)} \in [1/\varepsilon, (\log N)^2), V_2^{(N)} \leqslant t_N/2\Big) \\ & + Z^f_{N,M,h}\Big(V_1^{(N)} \in [1/\varepsilon, (\log N)^2), V_2^{(N)} > t_N/2\Big)\,. \end{aligned} \quad (4.21)$$

For the first term, and using that $\mathbf{P}\big((N-i,M-j)\in\hat\tau\big) \leqslant cN\mathbf{P}(\hat\tau_1^{(2)} = t_N)$ uniformly for $i \leqslant (\log N)^2$ and $j \leqslant t_N/2$ (since then we have $M - j - N - i \ge t_N/4$ for N large enough, similarly to (4.14)), we have

$$\begin{aligned} & Z^f_{N,M,h}\Big(V_1^{(N)} \in [1/\varepsilon, (\log N)^2), V_2^{(N)} \leqslant t_N/2\Big) \\ & \qquad \leqslant \sum_{i\geqslant 1/\varepsilon}\sum_{j=0}^{t_N/2} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}cN\mathbf{P}(\hat\tau_1^{(2)} = t_N) \\ & \qquad\qquad \leqslant c\Big(\sum_{j=0}^{t_N/2}K_f(j)\Big)N\mathbf{P}(\hat\tau_1^{(2)} = t_N)e^{N\mathrm{N}_h}\sum_{i\ge 1/\varepsilon}K_f(i)e^{-i\mathrm{N}_h}\,. \end{aligned} \quad (4.22)$$

When $\sum_j K_f(j) < +\infty$, then recalling (4.4), this term is smaller than $\eta \times \mathrm{II}$ provided that ε is small enough.

For the second term, similarly to term V, we split it according to whether $V_2^{(N)} \in (t_N/2, 2t_N)$ or $V_2^{(N)} \geqslant 2t_N$. The first contribution is

$$\begin{aligned} Z^f_{N,M,h}\Big(V_1^{(N)} \in (1/\varepsilon, (\log N)^2), V_2^{(N)} \in (t_N/2, 2t_N)\Big) \\ &= \sum_{i \geqslant 1/\varepsilon} \sum_{j=t_N/2+1}^{2t_N-1} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}\mathbf{P}\big((N-i, M-j) \in \hat\tau\big) \\ &\leqslant cK_f(t_N)e^{N\mathrm{N}_h} \sum_{i \geq 1/\varepsilon} K_f(i)e^{-i\mathrm{N}_h}\,, \end{aligned} \tag{4.23}$$

where we used that $K_f(j) \leqslant cK_f(t_N)$ for $j \in (t_N/2, 2t_N)$, and that the sum over j of $\mathbf{P}\big((N-i, M-j) \in \hat\tau\big)$ is bounded by 1. Then, the last sum can be made arbitrarily small by choosing ε small, so that in view of (4.11), this term can be bounded by $\eta \times$ IV.

The second contribution is

$$\begin{aligned} Z^f_{N,M,h}\Big(V_1^{(N)} \in (1/\varepsilon, (\log N)^2), V_2^{(N)} > 2t_N\Big) \\ &= \sum_{i \geqslant 1/\varepsilon} \sum_{j=2t_N}^{M} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}\mathbf{P}\big((N-i, M-j) \in \hat\tau\big) \\ &\leqslant Ce^{N\mathrm{N}_h} \sum_{j=2t_N}^{M} K_f(j)e^{-cj^2/k_N} \leqslant Ce^{N\mathrm{N}_h}N^{c+1}e^{-ct_N^2/N}\,, \end{aligned} \tag{4.24}$$

where we used (4.18), and (4.19). Then, this term decays faster than IV, provided that C_0 has been fixed large in (1.19), as for the term Vb.

Conclusion in the case of $\sum_j K_f(j) < \infty$. We have therefore proven that for any $\eta > 0$, we can choose $\varepsilon > 0$ small such that, for all N large enough (how large depend on ε),

$$\mathrm{II} + \mathrm{IV} \leq Z^f_{N,M,h} \leq (1+2\eta)\mathrm{II} + (1+4\eta)\mathrm{IV} \tag{4.25}$$

and the two terms behave asymptotically respectively as (4.4) and (4.11): this proves (2.6) for $\sum_j K_f(j) < \infty$.

The case of $\sum_j K_f(j) = \infty$*, with* $\overline{\alpha} < 1$. This time we have to show that IV dominates. We go through the various terms, but as pointed out during the proof, we have not used $\sum_j K_f(j) < \infty$ in estimating IV, so (4.11) still holds. We retain, for local use, that IV behaves (and, in particular, is bounded from below by) a constant times $K_f(t_N)\exp(N\mathrm{N}_h)$.

The estimate (4.4) for II is still valid. This term can be dealt directly without troubles, but it is more practical to observe that II (and the right side of (4.15), which bounds IIIa) is bounded (for N large) by a constant times

$$N\mathbf{P}\Big(\hat\tau_1^{(2)} = t_N\Big)\exp(N\mathrm{N}_h)\sum_{j \leq t_N/2} K_f(j)\,. \tag{4.26}$$

Therefore, in view of the behavior of IV that we have just recalled, this term is negligible if

$$Nt_N\mathbf{P}\Big(\hat\tau_1^{(2)} = t_N\Big) \ll 1\,, \tag{4.27}$$

since $\sum_{j \leq t_N/2} K_f(j) \leq cst.t_N K_f(t_N)$ if $\overline{\alpha} < 1$.

But the left-hand side is equivalent to $NL(t_N)/t_N^{1+\alpha}$ and hence (4.27) directly follows by recalling the definition (1.14) of $a_N^{(2)}$ and that $t_N \gg a_N^{(2)}$. This shows that both II and of the right side of (4.15) (which bounds IIIa) are negligible compared to IV.

The estimates for IIIb and V, as already pointed out, are valid without assuming that $\sum_j K_f(j) < \infty$, so we are left with controlling I. Recall that we split the contribution of I into three parts: (4.20), (4.22) and (4.23). As noticed above, the fact that $\sum_j K_f(j) < \infty$ was not used in estimating (4.20) and (4.23). Moreover (4.22) we can be bounded like (4.15) (in fact, it is much smaller), that was found above to be negligible compared to IV. We therefore conclude that I is also negligible compared to IV, and this completes the analysis of the case $\sum_j K_f(j) = \infty$, and of the proof of Theorem 2.2. □

Acknowledgements

G.G. acknowledges the support of grant ANR-15-CE40-0020.

Appendix A. The case $\overline{\alpha} = 1$ and $\sum_j K_f(j) = +\infty$

Recall the definition (2.9) of the Big Loop and Unbound strand, and when $\overline{\alpha} = 1$ with $\sum_j K_f(j) = +\infty$, define the new event $E_{mixed}^{(N)}$

$$E_{mixed}^{(N)} = \Big\{ \frac{\mathcal{M}_{1,\kappa_N}}{t_N} \in \big[1-\varepsilon_N, 1+\varepsilon_N\big], \mathcal{M}_{2,\kappa_N} < m_N^+, V_1^{(N)} \leqslant u_N, V_2^{(N)} \in \big[v_N, \varepsilon_N t_N\big] \Big\} \tag{A.1}$$

where $v_N \gg 1$ and $\varepsilon_N \ll 1$ are chosen such that

$$\overline{K}(v_N) \ll \overline{K}(\varepsilon_N t_N)\,, \qquad \overline{K}(\varepsilon_N t_N) \overset{N\to\infty}{\sim} \overline{K}(t_N)\,. \tag{A.2}$$

The event $E_{mixed}^{(N)}$ is therefore a set of trajectories with both a big loop (of order t_N), and a large unbound strand (of large order, but much smaller than t_N) – to optimize the interval for the length of the unbound strand, one can take $v_N \to +\infty$ and $\varepsilon_N \to 0$ as fast as possible, with the limitation given by (A.2).

Theorem A.1. *Suppose that $\alpha > 0$ and (1.18), and if $\alpha \geqslant 1$ assume additionally (1.19). We assume that $\overline{\alpha} = 1$ and that $\sum_j K_f(j) = +\infty$, and we denote $\overline{K}(x) := \sum_{j=1}^x K_f(j)$. Then, as $N \to \infty$,*

$$Z_{N,M,h}^f = (1+o(1)) Z_{N,M,h}^f\Big(E_{mixed}^{(N)}\Big) + (1+o(1)) Z_{N,M,h}^f\Big(E_{US}^{(N)}\Big)\,, \tag{A.3}$$

with

$$e^{-N\mathrm{N}_h} Z_{N,M,h}^f\Big(E_{mixed}^{(N)}\Big) \overset{N\to\infty}{\sim} \frac{N}{\hat{\mu}_1^2} \mathbf{P}(\hat{\tau}_1^{(2)} = t_N) \overline{K}(t_N) \Big(\sum_{i\geq 0} K_f(i) e^{-i\mathrm{N}_h} \Big)\,, \tag{A.4}$$

$$e^{-N\mathrm{N}_h} Z_{N,M,h}^f\Big(E_{US}^{(N)}\Big) \overset{N\to\infty}{\sim} \frac{1}{\hat{\mu}_1} \Big(\sum_{i\,\geqslant\, 0} K_f(i) e^{-i\mathrm{N}_h} \Big) K_f\left(t_N\right)\,. \tag{A.5}$$

Obviously, this theorem can easily be translated in term of path properties. Indeed, since $\mathbf{P}(\hat{\tau}_1^{(2)} = t_N) \sim cst.\, t_N^{-1} \mathbf{P}(\hat{\tau}_1^{(2)} > t_N)$ and $K_f(t_N) = \overline{L}(t_N) t_N^{-1}$, we have the asymptotic of the ratio

$$\widetilde{Q}_N = \widetilde{Q}_N(t_N) := \frac{Z_{N,M,h}^f\big(E_{mixed}^{(N)}\big)}{Z_{N,M,h}^f\big(E_{US}^{(N)}\big)} \overset{N\to\infty}{\sim} cst.\, N \mathbf{P}(\hat{\tau}_1^{(2)} > t_N) \frac{\overline{K}(t_N)}{\overline{L}(t_N)}, \tag{A.6}$$

with $\overline{K}(x)/\overline{L}(x) \to +\infty$ as a slowly varying function. Therefore, we obtain

$$\mathbf{P}^f_{N,M,h}(E^{(N)}_{US}) \overset{N\to\infty}{\sim} \frac{1}{1+\widetilde{Q}_N} \quad \text{and} \quad \mathbf{P}^f_{N,M,h}(E^{(N)}_{mixed}) \overset{N\to\infty}{\sim} \frac{\widetilde{Q}_N}{1+\widetilde{Q}_N}\,. \tag{A.7}$$

We stress that, when $\alpha > 1$, the ratio $\widetilde{Q}_N$ always goes to 0 as $N \to \infty$: indeed, in that case $N\mathbf{P}(\hat{\tau}^{(2)}_1 > t_N)$ decays faster than any slowly varying function. However, in the case $\alpha \in (0,1]$, the ratio $\widetilde{Q}_N$ diverges when $t_N \to +\infty$ slowly enough, showing that there is a regime under which the mixed trajectories described in the event $E^{(N)}_{mixed}$ occur, in the sense that $\mathbf{P}^f_{N,M,h}(E^{(N)}_{mixed}) \to 1$ as $N \to \infty$.

Proof. We treat this case in a concise way because most of the technical work has already been done above. To summarize – recall the different contributions in (4.1) – the term IV is well estimated in (4.11) and the terms I, IIIb and V were found to be negligible compared to it – this was valid even when $\sum_j K_f(j) = +\infty$. When $\sum_j K_f(j) = +\infty$, then the term II is found to be negligible compared to the right side of (4.15) (which bounds IIIa), and we therefore focus on this last term.

We can again decompose IIIa into two contributions:

$$\text{IIIa} = Z^f_{N,M,h}\Big(V^{(N)}_1 < 1/\varepsilon; V^{(N)}_2 \in (1/\varepsilon, \varepsilon t_N)\Big) + Z^f_{N,M,h}\Big(V^{(N)}_1 < 1/\varepsilon; V^{(N)}_2 \in (\varepsilon t_N, t_N/2)\Big)\,.$$

The second one, exactly in the same manner as for IIIb, can be shown to be negligible compared to IV as $N \to \infty$. Then, the first term is equal to

$$\text{IIIa}' := \sum_{i<1/\varepsilon} \sum_{j=1/\varepsilon}^{\varepsilon t_N} K_f(i)K_f(j)e^{(N-i)\mathrm{N}_h}\mathbf{P}\big((N-i, M-j) \in \hat{\tau}\big).$$

Then, thanks to Theorem 2.1, for every $\eta > 0$ we can choose $\varepsilon > 0$ small enough and N_ε large enough so that uniformly for the range of i and j considered, and $N \geq N_\varepsilon$

$$\mathbf{P}\big((N-i, M-j) \in \hat{\tau}\big) \begin{cases} \geq (1-\eta)\frac{N}{\hat{\mu}_1^2}\mathbf{P}(\hat{\tau}^{(2)}_1 = t_N)\,, \\ \leq (1+\eta)\frac{N}{\hat{\mu}_1^2}\mathbf{P}(\hat{\tau}^{(2)}_1 = t_N)\,, \end{cases}$$

and we stress that the main contribution to this probability comes from a big loop event, of length larger than $(1-\varepsilon)t_N$. We therefore get that, for N large enough, and denoting $\overline{K}(x) := \sum_{j=1}^x K_f(j)$ which is a slowly varying function,

$$\begin{aligned} \text{IIIa}' &\geq (1-\eta)\frac{N}{\hat{\mu}_1^2}\mathbf{P}(\hat{\tau}^{(2)}_1 = t_N)e^{N\mathrm{N}_h}\Big(\sum_{j=1/\varepsilon}^{\varepsilon t_N} K_f(j)\Big)\sum_{i<1/\varepsilon} K_f(i)e^{-i\mathrm{N}_h} \\ &\geq (1-\eta)\frac{N}{\hat{\mu}_1^2}\mathbf{P}(\hat{\tau}^{(2)}_1 = t_N)\overline{K}(t_N)e^{N\mathrm{N}_h}\Big(\sum_{i\geq 0} K_f(i)e^{-i\mathrm{N}_h}\Big)\,, \end{aligned}$$

and similarly for an upper bound with $1-\eta$ replaced by $1+\eta$.

We are actually able to narrow the condition $V^{(N)}_2 \in (1/\varepsilon, \varepsilon t_N)$ in IIIa$'$ to a smaller interval $(v_N, \varepsilon_N t_N)$ without changing the estimates, provided that $v_N \to +\infty$ and $\varepsilon_N \to 0$ slowly enough, so that condition (A.2) holds. □

Appendix B. About the transition between Cramér and non-Cramér regimes

In this Appendix, we discuss the condition (1.18)-(1.19) ensuring that one lies in the big-jump regime described by Theorem 2.1. We focus on the constrained partition function – or rather the probability $\mathbf{P}((N,M)\in\hat\tau)$ – to study the transition between the condensation phenomenon that we highlighted and the Cramér regime, but all the observations made here could also apply to the other results. Like in Section 4 we omit integer parts, so $\gamma_c N$ stands for the (upper or lower, as one wishes) integer part of $\gamma_c N$.

B.1. **Between Cramér and non-Cramér regimes I.** If one sets $M=\gamma_c N$, or in other words if $t_N=0$, then [2] proves that

$$\mathbf{P}\big((N,\gamma_c N)\in\hat\tau\big) \overset{N\to\infty}{\sim} \frac{c_0}{a_N^{(2)}}\,, \tag{B.1}$$

where the constant $c_0>0$ is explicit. The heuristics of this result can be easily understood: the typical number of renewal is $k_N=N/\hat\mu_1+O(\sqrt{N})$ and, for each k in that range, Doney's Local Limit Theorem [6] gives that $\mathbf{P}(\hat\tau_k=(N,M))$ is equivalent up to a multiplicative constant to $(a_N^{(2)}\sqrt{N})^{-1}$. Hence, neither $\hat\tau^{(1)}$ nor $\hat\tau^{(2)}$ have to make an atypical deviation, and the term $(a_N^{(2)})^{-1}$ simply comes from a local limit theorem: there is no condensation phenomenon, i.e. the typical trajectories contributing to the event $(N,\gamma_c N)\in\hat\tau$ do not exhibit a big jump. However, we are not in the Cramér regime – one component of the inter-arrivals does not have exponential tails, so there are jumps that are luch larger than $\log N$ – and we can see this critical situation as a *moderate* Cramér regime, because (moderate) deviations are carried by both components, like in the Cramér regime the (large) deviations are carried by both components.

A behavior like (B.1) also holds when $t_N/a_N^{(2)}\to t\in\mathbb{R}$: the constant c_0 is simply replaced by a constant c_t depending on t. When $\alpha\in(0,1)$, the fact that one lies in the big-jump regime (and Theorem 2.1 holds) as soon as $t_N/a_N^{(2)}\to+\infty$ is optimal, in the sense that when $\sup_N t_N/a_N^{(2)}<+\infty$, then the typical trajectories do not exhibit a condensation phenomenon.

B.2. **Between Cramér and non-Cramér regimes II.** When $\alpha\ge 1$, the situation is more involved because the condition $t_N/a_N^{(2)}\to+\infty$ alone is not enough to ensure that the model is in the big-jump domain.

We conjecture that when $\alpha>1$, there is some $a_c=a_c(\alpha)$ – that we give explicitly below – such that the big-jump regime holds when $t_N>a\sqrt{N\log N}$ with $a>a_c$ (i.e. theorem 2.1 holds), and a moderate Cramér regime holds when $t_N<a\sqrt{N\log N}$ with $a<a_c$ (we give an explicit conjectured analogue of Theorem 2.1, see (B.5) below). Note that having $t_N>a\sqrt{N\log N}$ corresponds to our condition (1.19), except that our constant C_0 is not optimal. Finding the correct threshold when $\alpha=1$ is even more involved and we prefer to leave it aside.

So let us now focus on the case $\alpha>1$, and develop some heuristic arguments to conjecture the asymptotic behavior of $\mathbf{P}\big((N,M)\in\hat\tau\big)$, and the typical behavior of trajectories contributing to this event. We take $a_N^{(2)}=\sqrt{N}$, and we are considering the case $t_N/\sqrt{N}\to\infty$ (the case $t_N/\sqrt{N}\to t\in\mathbb{R}$ being given in Appendix B.1), with

$t_N \le C_0\sqrt{N\log N}$ (otherwise we already know we are in the big-jump domain). Writing

$$\mathbf{P}\big((N,\gamma_c N+t_N)\in\hat\tau\big)=\sum_{k=1}^{N}\mathbf{P}\Big(\hat\tau_k^{(1)}=N,\hat\tau_k^{(2)}=\gamma_c N+t_N\Big), \tag{B.2}$$

then, the k's bringing the main contribution to the sum are either $k=N/\hat\mu_1+O(\sqrt{N})$, in which case the deviation is entirely carried by $\hat\tau^{(2)}$; $k=N/\hat\mu_1+t_N/\hat\mu_2+O(\sqrt{N})$, in which case the cost is brought by $\hat\tau^{(1)}$; or more generally $k=N/\hat\mu_1+\theta t_N/\hat\mu_2+O(\sqrt{N})$ with some $\theta\in\mathbb{R}$ (it is natural to expect $\theta\in[0,1]$, but $\theta\notin[0,1]$ should not be excluded), in which case the cost is shared *jointly* by the two coordinates $\hat\tau^{(1)},\hat\tau^{(2)}$.

Then, having a look at Nagaev's Theorem 1.9 in [16] suggests that for any k, only two possible behavior can contribute to $\mathbf{P}(\hat\tau_k^{(1)}=N,\hat\tau_k^{(2)}=\gamma_c N+t_N)$: having one large jump (in which case, and since $\hat\tau^{(2)}$ has a heavier tail, the probability is maximal when $k=N/\hat\mu_1+O(\sqrt{N})$ so that only $\hat\tau^{(2)}$ has to make a large jump), or using a collective joint strategy with no big jump (i.e. a moderate Cramér regime). The first possible behavior is therefore the big-jump strategy that we already identified, and we would therefore have that

$$\begin{aligned}\mathbf{P}\Big((N,\gamma_c N+t_N)\in\hat\tau\Big)&=(1+o(1))\frac{N}{\hat\mu_1^2}\mathbf{P}\big(\hat\tau_1^{(2)}=t_N\big)\\&\quad+(1+o(1))\sum_{k=1}^{N}\mathbf{P}\big(\hat\tau_k^{(1)}=N,\hat\tau_k^{(2)}=\gamma_c N+t_N,\text{“with no big jump”}\big)\,,\end{aligned} \tag{B.3}$$

where by “no big jump” we mean that all jumps are $O(m_N)$.

Using a local moderate deviation theorem for the probability when no big jump occurs (such a local moderate deviation theorem should hold because t_N is not too large, $t_N\le C_0\sqrt{N\log N}$), we would have that, for $k=N/\hat\mu_1+\theta t_N/\hat\mu_2$

$$\begin{aligned}\mathbf{P}\Big(\hat\tau_k^{(1)}=N,\hat\tau_k^{(2)}=\gamma_c N+t_N,\text{“no big jump”}\Big)&=\frac{1+o(1)}{k}g\left(\frac{N-\hat\mu_1 k}{\sqrt{k}},\frac{\gamma_c N+t_N-\hat\mu_2 k}{\sqrt{k}}\right)\\&=\frac{(1+o(1))\hat\mu_1}{2\pi N\sqrt{(1-\rho^2)\sigma_1^2\sigma_2^2}}\exp\left(-\frac{\hat\mu_1 t_N^2}{2(1-\rho^2)N}\Big\{\frac{\theta^2}{\gamma_c^2\sigma_1^2}-2\rho\frac{\theta(1-\theta)}{\gamma_c\sigma_1\sigma_2}+\frac{(1-\theta)^2}{\sigma_2^2}\Big\}\right),\end{aligned} \tag{B.4}$$

where $g(\cdot,\cdot)$ is the bivariate normal density of the limit $\frac{1}{\sqrt{k}}\big(\hat\tau_k-(\hat\mu_1,\hat\mu_2)k\big)$, which is centered with normalized covariance $\rho=(\sigma_1\sigma_2)^{-1}\mathrm{Cov}(\hat\tau_1^{(1)},\hat\tau_1^{(2)})\in(-1,1)$ — and σ_1^2,σ_2^2 are the respective variances of $\hat\tau_1^{(1)},\hat\tau_1^{(2)}$. For the second equality, we used that $N-\hat\mu_1 k=\gamma_c^{-1}\theta t_N$, $\gamma_c N+t_N-\hat\mu_2 k=(1-\theta)t_N$ and $k=(1+o(1))N/\hat\mu_1$.

Hence, in the sum over k in (B.3), the main contribution should be for $k=\frac{N}{\hat\mu_1}+\frac{\theta_0}{\hat\mu_2}t_N+O(\sqrt{N})$, with θ_0 minimizing $Q(\theta):=\frac{\theta^2}{\gamma_c^2\sigma_1^2}-2\rho\frac{\theta(1-\theta)}{\gamma_c\sigma_1\sigma_2}+\frac{(1-\theta)^2}{\sigma_2^2}$ — after some calculation we find that $\min Q(\theta)=(1-\rho^2)(\gamma_c^2\sigma_1^2+2\rho\gamma_c\sigma_1\sigma_2+\sigma_2^2)^{-1}$. We end up with the following conjecture in the case $\alpha>1$, when $t_N/\sqrt{N}\to+\infty$

$$\mathbf{P}\Big((N,\gamma_c N+t_N)\in\hat\tau\Big)=(1+o(1))\frac{N}{\hat\mu_1^2}\mathbf{P}\Big(\hat\tau_1^{(2)}=t_N\Big)+(1+o(1))\frac{c_1}{\sqrt{N}}\exp\Big(-\mathbf{c}\,\frac{t_N^2}{N}\Big), \tag{B.5}$$

with $\mathbf{c}:=\frac{\hat\mu_1}{2}(\gamma_c^2\sigma_1^2+2\rho\gamma_c\sigma_1\sigma_2+\sigma_2^2)^{-1}$, and the constant c_1 could in principle be made explicit.

Plugging $t_N = a\sqrt{N \log N}$ in (B.5), we find that the first term is regularly varying with index $-\alpha/2$ and that the second term has index $-1/2 - \mathbf{c}a^2$. Hence, depending on a we can identify the dominant term in (B.5):

$$1^{\text{st}} \text{ term is dominant if } a > \sqrt{\frac{\alpha-1}{2\mathbf{c}}} \quad \text{and} \quad 2^{\text{nd}} \text{ term is dominant if } a < \sqrt{\frac{\alpha-1}{2\mathbf{c}}}\,.$$

We therefore interpret this as $a_c = \sqrt{(\alpha-1)/2\mathbf{c}}$, where a_c is the critical value mentioned at the beginning of Appendix B.2, separating a big-jump domain (when $t_N > a\sqrt{N \log N}$ with $a > a_c$) from a moderate Cramér regime (when $t_N < a\sqrt{N \log N}$ with $a < a_c$).



FIGURE 4. A schematic sum-up of the correspondence of the the values of $t_N = (\gamma - \gamma_c)N$ with the different regimes. We treat the big-jump domain and it is the one to the right of the right-most dashed line. We believe that to the right of the moderate Cramér regime there is the big-jump domain – put otherwise, the non-Cramér regime coincides with the big-jump domain – but this is proven only for $\alpha < 1$.

Notice that, when $t_N/\sqrt{N} \to -\infty$, one could develop an identical argument (except that the big-jump term disappears), provided that a local moderate deviation theorem as (B.4) holds – i.e. provided that $|t_N|/\sqrt{N}$ is not too large, how large depend mostly on the tail exponent $1+\alpha > 2$ of $\hat{\tau}^{(2)}$. In the end, the sharp asymptotics of $\mathbf{P}\big((N, \gamma_c N + t_N) \in \hat{\tau}\big)$ should also be given by the second term in (B.5) – as already seen in the case $t_N/\sqrt{N} \to t \in \mathbb{R}$ in Appendix B.1.

References

[1] I. Armendáriz and M. Loulakis, *Conditional distribution of heavy tailed random variables on large deviations of their sum*, Stoch. Proc. Appl. **121** (2011), 1138-1147.

[2] Q. Berger, *Strong renewal theorems and local large deviations for multivariate random walks and renewals*, arXiv:1807.03575

[3] N. H. Bingham, C. M. Goldie and J. L. Teugels, *Regular variation*, Cambridge University Press, Cambridge, 1987.

[4] R. D. Blake and S. G. Delcourt, *Thermal stability of DNA*, Nucleic Acids Research **26** (1998), 3323-3332.

[5] D. Denisov, A. B. Dieker and V. Shneer, *Large deviations for random walks under subexponentiality: the big-jump domain*, Ann. Probab. **36** (5), pp. 1946-199, 2008.

[6] R. A. Doney, *A bivariate Local Limit Theorem*, Jour. Multivariate Anal., **36**. pp. 95–102, 1991.

[7] T. R. Einert, H. Orland and R. R. Netz, *Secondary structure formation of homopolymeric single-stranded nucleic acids including force and loop entropy: implications for DNA hybridization*, Eur. Phys. J. E **34** (2011), 55 (15 pages).

[8] M. E. Fisher, *Walks, walls, wetting, and melting*, J. Statist. Phys. **34** (1984), 667-729.

[9] T. Garel and H. Orland, *On the role of mismatches in DNA denaturation*, arXiv:cond-mat/0304080, 2003

[10] T. Garel and H. Orland, *Generalized Poland-Scheraga model for DNA hybridization*, Biopolymers **75** (2004), 453-467.

[11] G. Giacomin, *Random polymer models*, Imperial College Press, World Scientific, 2007.

[12] G. Giacomin and M. Khatib, *Generalized Poland-Sheraga denaturation model and two dimensional renewal processes*, Stoch. Proc. Appl. **127** (2017), 526-573.

[13] L. de Haan, E. Omey and S. Resnick, *Domains of attraction and regular variation in* $\mathbb{R}^d$, J. Multivariate Anal. **14** (1984), 17-33.

[14] A. Kabakçıoğlu, A. Bar and D. Mukamel, *Macroscopic loop formation in circular DNA denaturation*, Phys. Rev. E **85** (2012), 051919.

[15] D. G. Kendall, *Unitary dilations of Markov transition operators and the corresponding integral representation of transition probability matrices*, In Probability and Statistics (U. Grenander, ed.) 138?161. Almqvist and Wiksell, Stockholm (1959).

[16] A. V. Nagaev, *Large deviations of sums of independent random variables*, Ann. Probab. **7** (1979) , 745–789.

[17] R. A. Neher and U. Gerland, *Intermediate phase in DNA melting*, Phys. Rev. E **73** (2006), 030902R.

[18] D. Poland and H. A. Scheraga, *Theory of helix-coil transitions in biopolymers;: Statistical mechanical theory of order-disorder transitions in biological macromolecules*, Academic Press, 1970.

[19] S. Resnick, P. E. Greenwood, *A bivariate stable characterization and domains of attraction*, J. Multivariate Anal. **9** (1979), 206-221.

Sorbonne Université, Laboratoire de Probabilités Statistique et Modélisation, UMR 8001, F- 75205 Paris, France

Université Paris Diderot, Sorbonne Paris Cité, Laboratoire de Probabilités Statistique et Modélisation, UMR 8001, F- 75205 Paris, France

Université Libanaise, Laboratoire de Mathématiques-EDST, Beyrouth, Liban