



HAL
open science

A statistical study of dipolarization fronts observed by MMS

Soboh Alqeeq, Olivier Le Contel, Patrick Canu, Alessandro Retinò, Hugo Breuillard, Thomas Chust, Alexandra Alexandrova, Laurent Mirioni, Yuri Khotyaintsev, Rumi Nakamura, et al.

► **To cite this version:**

Soboh Alqeeq, Olivier Le Contel, Patrick Canu, Alessandro Retinò, Hugo Breuillard, et al.. A statistical study of dipolarization fronts observed by MMS. EGU22, the 24th EGU General Assembly, May 2022, Vienne, Austria. 10.5194/egusphere-egu22-9532 . hal-03916798

HAL Id: hal-03916798

<https://hal.science/hal-03916798>

Submitted on 20 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PREVIEW

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



A statistical study of dipolarization fronts observed by MMS

Soboh ALQEEQ¹, olivier Le Contel¹, Patrick Canu¹, Alessandro Retinò¹, Hugo Breuillard¹, Thomas Chust¹, Alexandra Alexandrova¹, Laurent Mirioni, Yuri Khotyaintsev², Rumi Nakamura³, Frederick Wilder⁴, Hanying Wei⁵, David Fischer³, Daniel Gershman⁷, James Burch⁸, Roy Torbert⁶, Barbara Giles⁷, Stephen Fuselier⁸, Robert Ergun⁴, Per arne Lindqvist⁹, and the C.T Russell (5), R. J. Strangeway (5), D. L. Turner (10), I. J. Cohen (11)*

¹Laboratoire de Physique des Plasmas, Paris, France

²Swedish Institute of Space Physics, Uppsala, Sweden

³Space Research Institute, Austrian Academy of Sciences, Graz, Austria

⁴Laboratory of Atmospheric and Space Physics, Colorado, USA

⁵Institute of Geophysics and Planetary Physics, Los Angeles, USA

⁶Space Science Center and Department of Physics, University of New Hampshire, Durham, USA

⁷NASA Goddard Space Flight Center, Greenbelt, MD, USA

⁸Southwest Research Institute, San Antonio, Texas, USA

⁹Space and Plasma Group, Royal Institute of Technology, Stockholm, Sweden

*A full list of authors appears at the end of the abstract

In the present work, we consider 49 dipolarization fronts (DF) detected by the Magnetospheric Multiscale (MMS) mission on 2017, near the Earth's magnetotail equator ($B_x < 5\text{nT}$). Criteria for selecting DF using an AIDapy routine are based on difference of maximum and minimum values computed with a 306 s sliding window. They request a B_z increase, an ion velocity increase and a density decrease. This first automatic selection is then adjusted manually with the following criteria : B_z increase larger than 5 nT, ion velocity larger than 150 km/s, density decrease and both ion and electron temperature increases. All these events belong to the most common category (A) defined by Schmid et al., 2015 in term of density decrease and temperature increase at the DF. However, based on a superposed epoch analysis of DF basic properties (magnetic field, density, velocity, ...) we distinguish two subcategories of events depending on the shape of the DF. The first subcategory (55.1%) corresponds to a slow decrease of the magnetic field after the DF and is associated with smaller ion velocity and hotter plasma. The second subcategory (44.9%) has the same time scale for the rising and the falling of the magnetic field (a bump) associated with a decrease of ion and electron pressures and faster velocity as shown in Alqeeq et al. 2021. For both categories we found that ions are mostly decoupled from the magnetic field by the Hall fields. The electron pressure gradient term is also contributing to the ion decoupling and likely responsible for an electron decoupling at DF. We also analyzed the energy conversion process. For the first subcategory we found that the energy in the spacecraft frame is transferred from the electromagnetic field to the plasma ($\mathbf{J} \cdot \mathbf{E} > 0$) ahead or at the DF. For the second subcategory, we found the same behavior ahead or at the DF whereas it is the opposite ($\mathbf{J} \cdot \mathbf{E} < 0$) behind the front. In the fluid frame, we found that the energy is mostly transferred from the plasma to the

electromagnetic field ($\mathbf{J} \cdot \mathbf{E}' < 0$) ahead or at the DF for both subcategories but energy dissipation ($\mathbf{J} \cdot \mathbf{E}' > 0$) only occurs behind the front for the second subcategory. The possible origin of these two subcategories is discussed.

C.T Russell (5), R. J. Strangeway (5), D. L. Turner (10), I. J. Cohen (11): MMS Team