



HAL
open science

Significant SAR11 removal by a hard-bottom community

Ayelet Dadon-Pilosof, Keats Conley, Marcelino T Suzuki

► **To cite this version:**

Ayelet Dadon-Pilosof, Keats Conley, Marcelino T Suzuki. Significant SAR11 removal by a hard-bottom community. 2023. hal-03917304

HAL Id: hal-03917304

<https://hal.science/hal-03917304v1>

Preprint submitted on 1 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1

2

Significant SAR11 removal by a hard-bottom community

3

4

Ayelet Dadon-Pilosof^{1,2} and Keats Conley³ Marcelino T. Suzuki⁴

5

6

7

8

¹The Faculty of Marine Science, Ruppin Academic Center, Michmoret, Israel

9

²Department of Ecology, Evolution & Behavior, The Hebrew University of Jerusalem,

10

Israel. <https://orcid.org/0000-0002-4108-4111>

11

³University of Oregon, Oregon Institute of Marine Biology - 1585 E 13th Ave, Eugene,

12

OR, USA 97403 <https://orcid.org/0000-0002-2966-1844>

13

⁴Sorbonne Université, CNRS, Laboratoire de Biodiversité et Biotechnologies

14

Microbiennes, LBBM, Observatoire Océanologique, F-66650, Banyuls-sur-mer, France.

15

<https://orcid.org/0000-0003-3868-6362>

16

17

Corresponding author: Ayelet Dadon-Pilosof, ayelet@ruppin.ac.il

18

19 **Abstract**
20

21 Microorganisms are a key component of the marine food webs through the microbial
22 loop. In previous work, we have shown that some bacteria, including *Candidatus*
23 *Pelagibacter* spp (SAR11)—the most abundant bacterium in the ocean—can evade filtration
24 by benthic and planktonic tunicates. Here we tested whether differential removal of microbial
25 taxa by benthic filter-feeders can be observed in the distribution and abundance of microbial
26 taxa from hard-bottom subtidal communities, a common coastal habitat in the Eastern
27 Mediterranean Sea towards the open sea. The abundance of microbial groups along cross-
28 shore transects was measured based on combined flow cytometry and SSU rRNA gene
29 metabarcoding. Our results show that most groups were depleted (up to 50%) over the hard-
30 bottom compared to the open sea, but unexpectedly we did not observe a clear differential
31 removal of different taxa, SAR 11 notably. This study indicates a strong top-down control of
32 the abundance of pelagic microorganisms over shallow hard-bottom where suspension
33 feeders are common.

34

35 Introduction

36

37 Marine microbial communities form the basis of the ocean food web and mediate
38 most of the energy and material fluxes in the ocean (Glöckner et al. 2012). Microorganisms
39 constitute a large fraction of the living biomass in the sea (Pomeroy et al. 2007), and the
40 structure and function of their populations are shaped by a delicate balance between growth
41 and mortality (Pernthaler 2005). Grazing and virus-driven lysis constitute the main sources of
42 mortality (Sánchez et al. 2020), countered by the capabilities of microorganisms to avoid
43 grazing (Matz and Kjelleberg 2005), survive digestion or resist viral lysis. Grazing or
44 predation on microorganisms by either planktonic or benthic organisms is an important
45 mortality factor in many habitats (e.g., Verity 1991; Gili and Coma 1998; Gorsky et al. 1999;
46 Riisgard and Larsen 2001; Pernthaler 2005; Patten et al. 2011a). In the pelagic realm, protists
47 commonly dominate the guild of grazers of bacteria (Calbet and Landry 2004; Matz and
48 Kjelleberg 2005). However, in some habitats and seasons, grazing by metazoan and lysis by
49 phages may dominate mortality (Hahn and Höfle 2001).

50 Bacteria form some 65-86% of the biomass of microorganisms in the upper ocean
51 (Morris et al. 2002). However, only a small number of bacterial groups dominate that guild
52 (Teeling et al. 2012). In the oligotrophic eastern Mediterranean, pico-cyanobacteria
53 (*Synechococcus* and *Prochlorococcus*) and a few members of *Candidatus Pelagibacter*
54 *ubique* (SAR11) clade dominate microbial communities, accounting for >70% of the total
55 bacterial biomass (Partensky et al. 1999, Dadon-Pilosof et al. 2017). SAR11 is a clade of
56 heterotrophic bacteria, which constitutes 15–60% of total bacteria in the upper ocean (Morris
57 et al. 2002,2012; Rappé et al. 2002; Eiler et al. 2009; Giovannoni 2017). It is one of the
58 smallest free-living bacteria in the sea and is thought to be the most abundant group in the
59 world ocean. Within that group, some ecotypes thrive in oligotrophic environments while
60 others in more productive waters (Morris et al. 2002, Salter et al 2015). Members of the

61 SAR11 clade also have the lowest nucleic acid content (LNA) among all non-photosynthetic
62 bacteria (Mary et al. 2006). Dadon-Pilosof et al. (2017) reported that some bacteria,
63 especially members of the SAR11 clade, can effectively evade grazing from both pelagic and
64 benthic tunicates and this lack of grazing pressure on SAR11 could partially explain its
65 abundance and ubiquity.

66 A diverse guild of benthic invertebrate suspension feeders, including sponges,
67 bivalves, cirripedians, cnidarians, bryozoans, and tunicates, often dominates subtidal hard
68 substrates (Topçu et al. 2010). Their diet ranges from consumption of dissolved organic
69 matter (DOM) through grazing on microorganisms such as phytoplankton, virioplankton,
70 archaea and bacteria, as well as feeding on zooplankton and detritus (Gili and Coma 1998;
71 Topçu et al. 2010). Hard-bottom subtidal communities along the Mediterranean Sea are
72 diverse and are undergoing dramatic changes in the recent decades due to the combined effect
73 of global warming, overfishing, and the introduction of invasive species (Rilov et al. 2019).
74 Within these communities, retention of small particles is an adaptive advantage since
75 picoplankton often dominate the planktonic community biomass (Topçu et al. 2010).

76 Grazing by benthic suspension feeders on microorganisms is also an important
77 component of the benthic-pelagic coupling in coral reefs (Yahel et al. 1998; Genin et al.
78 2002, 2009; Patten et al. 2011). This grazing pressure on the microbial community is not
79 necessarily uniform as it depends on the spatial heterogeneity of the distribution of different
80 suspension feeders and their respective diets (Yahel et al. 2006, 2009; Hanson et al. 2009;
81 Dadon-Pilosof et al. 2017). Differential capture of particles from the ambient water based on
82 their size, concentration or morphological features is therefore expected to be reflected in the
83 prey distribution (Gili and Coma 1998).

84 The goal of the current study was to indirectly evaluate the effect of the whole benthic
85 community removal on the distribution of microorganisms across a shallow, subtropical

86 rocky coast. Following the study of Yahel et al. (1998) across a coral reef, a study that
87 preceded the demonstration of the role of phytoplankton grazing in the trophic dynamics of
88 coral reefs (Genin et al. 2009; Monismith et al. 2010), we sought to test the hypothesis that
89 differential removal on microbes is also reflected in the cross-shore distribution of bacteria
90 and other picoplankton.

91 Methods

92 Study site

93 Sampling was conducted off Michmoret, Israel (32° 24'N, 34° 52'E) in the Eastern
94 Mediterranean Sea. The oligotrophic Israeli shoreline is characterized by extremely low
95 nutrient levels (Nitrate 0.5-1 μ M, Phosphate 0.05-0.1 μ M, Krom and Suari 2015), high
96 salinity (38.3 to 40.0 PSU) and relatively warm waters (16.5 to 30.8°C) (Suari et al. 2019)
97 that is dominated in terms of numbers and biomass by pico- and nanoplankton sized
98 organisms (Herut et al. 2000, Raveh et al. 2015). Water stratification appears usually in
99 spring following a deep winter mixing that is enhanced by the high salinity of surface waters.

100 Eight transects were performed between December 2015 and April 2017 (Table S1).
101 Each transect encompassed eight stations spanning from 1.5 m depth at the hard-bottom
102 subtidal towards the “open-sea” (~ 0.9 km off-shore, water depth >12 m, Fig. 1). During the
103 sampling periods the average temperature (\pm SD) was 19.7 \pm 0.7 °C, salinity was high
104 (39.25 \pm 0.2 PSU), and the water column was fully oxygenated (dissolved oxygen
105 concentration 216 \pm 6 μ M). Due to the need to work very close to the bottom at the rocky
106 shoreline, sampling dates were dictated by sea conditions and were limited to days of calm
107 sea (wave height < 30 cm) and weak winds (<6m s⁻¹). To get a larger scale context within the
108 eastern Mediterranean shelf offshore of the high-resolution cross-shore transects, six cross-
109 shelf transects were sampled during March 2018, spanning from one km offshore, equivalent

110 to the farthest station in the small scale transects, to 42 km west to the shelf edge at 750 m
111 bottom depth (Fig S1).




Figure 1: The location of the transect line in the Eastern Mediterranean (a) transects location along the Israeli shore (b) transect showing the bottom (yellow), sampling stations (white numbered lines), dotted white line (1.5 m depth) representing sampling depth.

112
113 The shallowest sampling station in each transect was located ~150 m offshore at 1.6-
114 1.8 m depth where Niskin bottle was placed just above the rocky bottom (c.a. 10 cm) to
115 collect a sample. The 1.5 m sampling depth was retained along the transect as the water
116 deepened down to 12 m at the furthest offshore station, ~0.9 km offshore. The four
117 shallowest stations (hereafter Stations 1-4) were located above hard bottom (rocky area), the
118 5th station was located at the boundary between rocks and sand, while stations 6 to 8 were
119 above sandy bottom.

120 Sampling methods

121 Prior to sampling, each station was marked with a moored surface buoy and profiles
122 of temperature, salinity, fluorescence, and oxygen were measured using a Seabird

123 SBE19plusV2 CTD equipped with an *in vivo* chlorophyll fluorimeter (Cyclop7, Turner
124 Designs) and a dissolved oxygen sensor (SBE43, Seabird). The CTD was lowered in a
125 horizontal position at a slow pace (about a meter per minute) so that the water column was
126 adequately profiled even in the shallow-most stations. Due to the need to work close to the
127 bottom at the rocky shoreline, sampling was conducted from either a kayak or a small
128 inflatable skiff.

129 Seawater was collected simultaneously with CTD profiles, using a 5 L Niskin bottle.
130 At each station, samples were collected directly from the Niskin bottle for all the required
131 analyses: flow cytometry, DNA extraction and Chl-a measurements. Water samples were
132 kept on ice in the dark for further processing in the laboratory within 2-3 hours. Samples for
133 DNA were collected only during three of the eight sampling times.

134 Sample analysis

135 Flow cytometry

136 Flow cytometry was used to quantify the concentrations and the cell characteristics of
137 non-photosynthetic microorganisms (hereafter referred to as non-photosynthetic bacteria),
138 and the following four dominant autotrophic groups: *Prochlorococcus* (Pro), *Synechococcus*
139 (Syn), pico-eukaryotic algae (PicoEuk), and nano-eukaryotic algae (NanoEuk). We used an
140 Attune® Acoustic Focusing Flow Cytometer (Applied Biosystems) equipped with a syringe-
141 based fluidic system that allows a precise adjustment of the injected sample volume and
142 hence high precision of the measurements of cell concentrations ($\pm 5\%$). The instrument's
143 optical unit contained violet and blue lasers (405 and 488 nm, respectively) and was further
144 adapted for the analysis of marine ultra-plankton samples as described below.

145 Aliquots of 1.8 mL were collected from each water sample and transferred into 2 mL
146 cryotubes (Corning cat No. 430659). Samples were first incubated for 15 min at room
147 temperature with Glutaraldehyde 50% (electron microscopy grade, Sigma-Aldrich, cat No.

148 340855) at 0.1% (final concentration). Samples were frozen in liquid nitrogen (at least 60
149 min) and then stored at -80 °C until analysis (within a few days).

150 Each sample was analyzed twice. First, 600 µl of the sample water was analyzed at a
151 high flow rate (100 µL min⁻¹) for the determination of ultra-phytoplankton with a dual
152 threshold (trigger) on the red fluorescence channels of the violet and blue lasers. A second
153 run was used to analyze cells with no autofluorescence, i.e., non-photosynthetic microbes. To
154 visualize these cells, a 300 µL aliquot of the sampled water was incubated with the nucleic
155 acid stain SYBR Green I (20-120 min dark incubation at room temperature, 1:10⁴ of the
156 SYBR Green commercial stock). For this run, we used a low flow rate of 25 µL min⁻¹ and the
157 instrument was set to high sensitivity mode. Seventy-five µL of the sample water was
158 analyzed with a dual threshold (trigger) on green fluorescence channels of the violet and blue
159 lasers. The taxonomic identification was based on orange fluorescence (B12, 574±13 nm) of
160 phycoerythrin and red fluorescence (B13, 690±20 nm and VL3, 685±20 nm) of chlorophylls;
161 side-scatter (SSC), provided a proxy of cell surface complexity and cell volume (Marie et al.
162 1999), and forward-scatter (FSC) was a proxy of cell size (Cunningham and Buonaccorsi
163 1992; Simon et al. 1994).

164 Where possible, the non-photosynthetic bacteria were further divided based on their
165 green fluorescence (proxy for nucleic acid content) and forward scatter (proxy for size) into
166 three groups: LNA, low nucleic acid non-photosynthetic bacteria; HNA-Ls, high nucleic acid
167 low-scatter non-photosynthetic bacteria; HNA-Hs, high nucleic acid high-scatter non-
168 photosynthetic bacteria (Zubkov et al. 2004). Similarly, the eukaryotic algae were separated
169 to pico- and nano-phytoplankton (Simon et al. 1994). The size of *Synechococcus* is still
170 somewhat controversial, indicating a range of 0.3 to 1.2 µm (e.g., Uysal 2001; Garcia et al.
171 2016). For pico-eukaryotic algae, we followed Worden and Not (2008), who suggested a size
172 range of up to 3.0 µm. Larger cells were termed nano-eukaryotic algae (2.0-20 µm). As a

173 rough proxy of cell size, we used the ratio of the median forward scatter of each cell
174 population to that of the median forward scatter of reference beads (Polysciences™, cat#
175 23517, Flow Check High-Intensity Green Alignment 1.0 µm) that were used as an internal
176 standard in each sample. See Dadon-Pilosof et al. (2019) for a further discussion of the
177 accuracy of size estimates.

178 Chlorophyll measurements

179 Water samples (~300 mL) were collected directly from the Niskin bottles at each
180 station into dark volumetric BOD glass bottles (Wheaton 227667) and maintained on ice in
181 dark cool box. In the lab, samples were prefiltered through a 100 µm mesh (to remove large
182 zooplankton and/or aggregates and suspended pieces of benthic algae) and filtered using low
183 vacuum onto a 25 mm Whatman GF/F filter. Filters were kept frozen at -20°C in 20 mL
184 scintillation vials until further processing. To ensure complete chlorophyll *a* extraction of
185 coastal phytoplankton, we used a hot dimethyl sulfoxide (DMSO) extraction method
186 (Burnison 1980). Briefly, 2 mL of DMSO were added into each vial containing the frozen
187 filter. Vials were then incubated for 20 min at 65°C, then cooled in a dark box to room
188 temperature (approximately 1 hr). Four mL of buffered Acetone (90% Acetone, 10%
189 saturated MgCO₃) were added to the vial and thoroughly mixed. Vials were then left to settle
190 for few minutes and 3 mL sample was drawn from the vial to a fluorometer cuvette.
191 Fluorescence was measured with a calibrated Trilogy fluorometer (Turner Designs) using
192 the non-acidification method (Welschmeyer and Naughton 1994).

193 DNA extraction

194 The relative abundance of prominent microbial taxa (phylotypes) in the seawater was
195 estimated using next-generation sequencing (NGS) of SSU rRNA genes to evaluate any
196 differential removal by suspension feeders benthic community. Ten mL of seawater collected
197 from each station and filtered on a 25 mm, 0.2 µm polycarbonate membrane (GE Healthcare

198 Biosciences, cat. No. 110606) under low vacuum and frozen in 1.5 mL micro-tubes at -20°C
199 until analysis. DNA from each filter was extracted using the DNeasy ‘blood & tissue kit’
200 (QIAGEN, Cat. No. 69504) with the following modifications to the manufacturer’s protocol:
201 ATL buffer (180 µL) and 20 µL of proteinase K were added and samples were incubated at
202 56°C for 1 hr. Then 200 µL of AL Buffer and 200 µL of 95-100% ethanol was added to the
203 sample and the mixture was pipetted into spin columns and placed in a 2 mL collection tube.
204 Tubes were centrifuged at 6000 RCF for 1 min. The flow-through was discarded and 500 µL
205 of AW1 buffer was added to the column, centrifuged at 6000 RCF for 1 min, and the flow-
206 through again discarded. This step was repeated for the third time, with 500 µL Buffer AW2
207 and a spin of 18,000 RCF for 1 min to dry the membrane before elution. For the elution step,
208 the spin column was placed on a new collection tube. Two hundred µL of buffer AE
209 preheated to 56°C was pipetted at three steps (50 µL, 50 µL, and 100 µL) into the column and
210 each step was followed by 6000 RCF centrifugation for 1 min. The sample was then
211 incubated at room temperature for at least a minute and stored at -20°C.

212 [Next-generation sequencing](#)

213 Samples were sequenced by Research and Testing Laboratories (Lubbock TX). The
214 SSU rRNA genes were amplified for sequencing using a forward and reverse fusion primers
215 (515F-Y - 926R (Parada et al. 2016). The forward primer was constructed with (5’-3’) the
216 Illumina i5 adapter (AATGATACGGCGACCACCGAGATCTACAC), an 8-10bp barcode, a
217 primer pad, and the 5’- GAGTTTGATCNTGGCTCAG -3’ primer. The reverse fusion
218 primer was constructed with (5’-3’) the Illumina i7 adapter
219 (CAAGCAGAAGACGGCATAACGAGAT), an 8-10bp barcode, a primer pad, and the 5’-
220 GTNTTACNGCGGCKGCTG -3’ primer. Primer pads were designed to ensure the primer
221 pad/primer combination had a melting temperature of 63°C-66°C, according to methods
222 developed by Patrick Schloss’ laboratory ([9](http://www.mothur.org/w/images/0/0c/Wet-</p></div><div data-bbox=)

223 lab_MiSeq_SOP.pdf). Amplifications were performed in 25 μ L reactions with Qiagen
224 HotStar Taq master mix (Qiagen Inc, Valencia, California), 1 μ L of each 5 μ M primers, and 1
225 μ L of the template. Reactions were performed on ABI Veriti thermocyclers (Applied
226 Biosystems, Carlsbad, California) under the following cycle conditions: 95°C for 5 min, then
227 35 cycles of 94°C for 30 sec, 54°C for 40 sec, 72°C for 1 min, followed by one cycle of 72°C
228 for 10 min and a final 4°C hold.

229 Amplification products were visualized with eGels (Life Technologies, Grand Island,
230 New York). Products were then pooled equimolarly and each pool was size-selected in two
231 rounds using Agencourt AMPure XP (BeckmanCoulter, Indianapolis, Indiana) in a 0.7 ratio
232 for both rounds. Size-selected pools were then quantified using a Qubit 2.0 fluorometer (Life
233 Technologies) and loaded on an Illumina MiSeq (Illumina, Inc. San Diego, California) 2x300
234 flow cell at 10 pM.

235 Sequence data analysis

236 SSU rRNA sequences were treated using a qiime2 (v. 2018-8) and biom-format (v.
237 2.1.6) deployed through a bash scripts (Suppl. File). Briefly demultiplexed forward and
238 reverse reads were imported into qiime2 artifacts and ASV tables were generated using *qiime*
239 *dada2* using options "--p-trim-left-f 19 --p-trim-left-r 20 --p-trunc-len-f 300 --p-trunc-len-r
240 250" The resulting ASVs were identified using *qiime feature-classifier* using an in house
241 version of the Silva132 (arb-silva.de) including only the region flanked by the primers, and a
242 taxonomy files extracted from this database (see data availability below for details).

243 Since the primers we used amplify both 16S rRNA from prokaryotes and chloroplasts
244 as well as 18S rRNA from eukaryotes, the ASV table and sequence files were filtered using
245 *qiime taxa* to generate three sets of table/sequences, one with 16S rRNA, one with
246 chloroplasts and one with eukaryotic sequences. ASV tables were modified to include

247 taxonomy using *sed* and *gawk* and *biom-format* and exported in ".tsv" format for further
248 analyses. Here we describe only results concerning 2388 prokaryotic ASVs.

249 Data analysis

250 Due to temporal changes in microbial communities, and assuming the entire transect
251 represented a single water mass the concentrations along each transect were normalized to the
252 seaward-most station and presented as “% of open sea”. The significance level of cross-shore
253 trends was tested using the “Page test” for ordered alternatives (Page 1963). This non-
254 parametric test is a modified version of the Kruskal-Wallis one-way ANOVA for ranked data.
255 Nearshore depletion of microbial taxa was tested (each taxon per season) with H_1 as an
256 ordered decrease in concentrations from the “open water” toward the shore. Due to missing
257 sampling points, only 3 complete transects toward the hard-bottom were used for the test in
258 each season.

259 Data collected with the CTD was converted, aligned, and binned (at 0.1 m) with the
260 SBE DataProcessing software (Version 7.2). Since salinity differences along the transects
261 were negligible (<0.01 PSU), density differences were driven solely by temperature. The
262 vertical and horizontal change in temperature and *in-vivo* chlorophyll fluorescence along the
263 cross-shore transects (hereafter ‘anomaly’) was calculated within each transect as anomalies
264 (the difference of each data point from the average temperature or fluorescence along the
265 transect). All seasonal anomalies were plotted together using Ocean Data View (Version 5).
266 Interpolation was made with the Weighted-average gridding function of Ocean Data View
267 using a seeking distance of 0.25 m along the vertical axis and 100 m along the horizontal
268 axis. Due to the low (N=4), data are presented as a seasonal average \pm standard error (SE)
269 unless otherwise indicated.

270 Removal of specific ASVs toward the shore was calculated by multiplying the relative
271 abundance of each ASV and the total bacterial cell counts obtained by flow cytometry and

272 then calculation as a percentage of the open sea station. Normalized removal was calculated
273 by normalizing measured removal to the ASV with the highest removal within the same
274 transect. Implementation of this approach provides a powerful tool to indirectly evaluate the
275 effect of benthic removal of microbial prey at the ASV level. Hereafter, the use of the terms
276 “selectivity” and “preference” are limited to their technical definition (Chesson 1978, 1983),
277 i.e., the removal of a prey type in higher proportion than its proportional presence in the
278 environment, relative to other food types present.

279 **Data availability**

280 Raw sequencing reads are available through the NCBI SRA under accession number
281 PRJNA912166. The analysis pipeline and associated files including scripts, mapping files,
282 taxonomic identification databases and ASV tables are available through a github repository
283 (github.com/suzumar/transect_ms).

284 **Results**

285 During winter, a temperature gradient was found along the transects with colder water
286 at the shallow stations above the hard-bottom (stations 1-5, $19.4 \pm 0.03^\circ\text{C}$, Fig 2a) in
287 comparison to the "open sea" (stations 6-8, $20.1 \pm 0.03^\circ\text{C}$, Fig 2a). During spring, no such
288 temperature gradient was observed along the transects but the beginning of stratification was
289 noticeable, with a half-degree Celsius warmer surface layer (0-3 m, $20.1^\circ\text{C} \pm 0.07$, Fig 2b)
290 than the deeper water (3-12m, $19.6^\circ\text{C} \pm 0.02$, Fig 2b). The warmer, nearshore surface layer
291 (10-25 m depth, Fig S1) showed typical coastal enrichment, with higher chlorophyll
292 concentrations (Fig S1) and cell counts (Fig S2) than the open sea. Salinity was similar along
293 the transect (1.5m depth) and with depth (0-12m) during winter and spring (39.26-39.31 PSU
294 and 39.35-39.28 PSU, respectively, during winter and, 39.29-39.23 PSU and 39.06-39.21
295 PSU, respectively, during spring).




Figure 2: Temperature anomaly along the cross-shore transects. (a) Winters of 2015-2016, n=4. (b) Springs of 2016-2017, n=4. Sampling stations are shown as vertical black lines. Grey shading approximates the bottom depths based on the bottom depth of each cast. In most cases, the CTD was lower all the way to the bottom (~0.2 mab).

296 In the cross-shore transects, the concentration of all microbial populations decreased
297 toward shore in a highly significant gradient (Page test, $p < 0.01$, Fig. 3a-d). In most cases, and
298 for most of the microbial populations, cell concentrations were similar along the ~700 m
299 sandy section of the transects, between the open sea and the outer boundary of the rocky area
300 (Stations 6 to 8). Sometimes even an increase was observed along that section (Table S1, S2).
301 Most of the depletion occurred above the rocky habitat (Stations 1 to 5). The depletion of the
302 microbial populations over the rocky section range between 25-50% of their percentage of
303 the open sea (Fig 3a-d). *In vivo* chlorophyll fluorescence (Fig S3) and the concentration of
304 extracted chlorophyll a also showed a strong depletion above the hard-bottom area compared
305 to the open sea (Fig 3e).




Figure 3: Cross-shore transects showing the percent change in the concentrations of several microbial cell populations (counted by flow cytometry) from their “open sea” concentration ~900 m offshore. (a) Bact, non-photosynthetic bacteria, (b) Euk, eukaryotic algae, (c) Syn, *Synechococcus sp.* (d) PLP, *Prochlorococcus*-like particles, (e) Extracted chlorophyll *a* concentration along 3 transects. Blue, n=4, transects performed during winter. Orange, n=4, transects performed during spring. The significance of the Page test for a gradient of decreasing concentrations toward the shore is denoted as **<0.01, ***<0.001

341

342 Differential changes in relative abundances between ASVs and even within the same

343 ASV in different transects were observed (Fig 4a-c). For example, ASVs belonging to

344 cyanobacteria group were on average 8%, 35% and 3% of their percentage at the open sea
 345 (Fig 4a-c accordingly).




Figure 4: Cross-shore transects showing the percent change in the concentrations of the 20 most abundant prokaryotic ASVs in the water relative to their “open sea” concentration ~900 m offshore: (a-c) transects above rocky hard bottom (January 2016, March 2016 and, April 2016). The vertical dashed line represents the concentration in the open sea (100%). Pink indicates members of SAR11 clade, green indicates autotrophs, and blue indicates other non-photosynthetic bacteria.

347 Within the SAR11 clade, the percentage of four ASVs decreased along the transects,
348 in similar trends (Fig 5). Transects showed $83\% \pm 7\%$ (Mean \pm SD, Fig 6a-c) normalized
349 removal for the 20 most abundant ASVs. Abundant ASVs including *Synechococcus* CC9901
350 (27% relative abundance, transect during January 2016) and SAR11-Ia (25%, transect during
351 April 2016) were removed at similar percentages (95% and 84% accordingly) compared to
352 less abundant ASVs (e.g. SAR11-Ib 1.3% relative abundance and 87% removal, SAR86
353 2.3% relative abundance and 86% removal) (Fig 6).
354




Figure 5: Cross-shore transects showing the percent change in the concentrations of four ASVs belonging to SAR11 clade in the water relative to their “open sea” concentration ~900 m offshore: (a-c) transects above rocky hard bottom (January 2016, March 2016 and, April 2016). The vertical dashed line represents the concentration in the open sea (100%).

355

356

357

358

359

360

361

362

363

364

365

366

367




Figure 6: Normalized removal of the 20 most abundant ASVs from the open sea toward the shore. (a-c) transects above rocky hard bottom (January 2016, March 2016 and, April 2016). Pink indicates members of SAR11 clade, green indicates autotrophs, and blue indicates other non-photosynthetic bacteria. Dashed vertical line represents the expected retention assuming equal retention probability for all cells. Size of circles represents relative abundance in the open water (Station 8) during sampling., with the two white circles plotted for scaling.

Discussion

368
369 This study was designed to evaluate whether the effect of differential removal of
370 microorganisms by benthic suspension feeders can be detected at the community level. Most
371 of the microbial populations were depleted above the hard-bottom area compared to the open
372 sea. Differential removal was observed at low extent (between and within species). A similar
373 trend was also observed for chlorophyll *a* concentration, suggesting the formation of a
374 depleted boundary layer over the hard-bottom. The hard-bottom community at the study site
375 included a diverse assemblage of suspension feeders, including sponges, bivalves, ascidians,
376 polychaetes, hydrozoans, and bryozoans (Rilov et al. 2018). Suspensions feeders on the hard-
377 bottom occur in different densities, and hence the competition for available prey is between
378 species, within species and even between different taxa. Niche speciation is expected in such
379 diverse and dense community where different taxa utilize different filtration mechanisms and
380 presumably, different organic carbon sources in their diet. If the depletion of microbial cells
381 is the outcome of grazing by benthic suspension feeders, near-bottom depletion should
382 generate a shore-wise depletion above the hard-bottom section of the transects compared to
383 the sandy section of the transect (e.g., Genin et al. 2009; Jones et al. 2009). In case other
384 processes determine the cross-shore trend (e.g., runoff, eutrophication, and coastal pollution),
385 the depleted zone is expected to extend throughout the water column and no difference is
386 expected along the transects. Differential grazing of phytoplankton by dense populations of
387 benthic suspension feeders was also reported in San-Francisco bay where it was attributed to
388 different sinking rates of the microalgae (Lucas et al. 2016). Such a phenomenon is very
389 unlikely in the east Mediterranean where the planktonic community is dominated by very
390 small cells (<10 μm) with negligible sinking rates (Siokou-Frangou et al. 2010). This body of
391 prior work supports the assumption that the depletion reported here is due to benthic
392 suspension grazing although it was not measured directly.

393 Physical and biological processes are the major factors controlling changes in particle
394 concentrations throughout the ocean, and advective processes could be responsible for the
395 pico and nanoplankton depletion we observed. Monismith et al. (2006) showed that shallow
396 regions nearshore experience larger temperature changes than deeper regions offshore. When
397 the water warms during the daytime (e.g., in the spring), the shallow near-shore water body
398 tends to warm faster than the nearby open sea. The warmer water expands and flows
399 offshore, causing deeper and cooler water to flow onshore at depth to replace it. This ‘thermal
400 flow’ (Fig 7) leads to an upwelling of deeper water and material to the nearshore region (Fig
401 7). The opposite process occurs during winter or cold nights, when a faster cooling of the
402 shallow, near-shore waterbody initiates near-shore downwelling of cold surface water that
403 reverses the direction of the “thermal flow” cycle and induces onshore flow of surface water
404 (Fig 7). In the Eastern Mediterranean Sea, the water column is usually stably stratified, and
405 the numbers and biomass of surface water plankton is lower than subsurface layers (Suari et
406 al 2019). Onshore transport of surface water from the offshore during the cold phase of the
407 thermal flow cycle and the nearshore downwelling of "plankton poor" seawater is expected to
408 yield a benthic zone of depleted of plankton close to shore (Labiosa et al. 2003). However,
409 this physical mechanism would not be expected to change the relative abundance of different
410 microbial groups along the transects, nor should it differentially affect rocky and sandy
411 sections of the transects. Moreover, during the warm phase of the thermal flow cycle (in
412 spring for example), local upwelling is expected to bring plankton-rich water onshore,
413 countering removal, and the benthic zone should be as rich if not richer nearshore. Our results
414 showed that nearshore depletion occurred during both the cold and hot ‘phase’ of the thermal
415 flow, suggesting that a biological process such as filtration is probably the mechanism
416 responsible for the depletion above the hard-bottom section in the transects rather than
417 advection of planktonic poor water. A similar conclusion was reached by Patten et al. (2011),

418 who showed depleted levels (~40% on average) of microbial cells over a reef with negligible
419 removal over a sandy bottom.

420 Our previous work showed partial grazing resistance of members SAR11 clade to
421 grazing by pelagic and benthic tunicates (Dadon-Pilosof et al. 2017), and in general, lower
422 retention efficiencies on LNA bacteria than other prey types. Selective grazing and
423 specifically low retention on LNA compare to both HNA bacteria and *Synechococcus* was
424 also observed in sponges (Hanson et al. 2009; Jiménez 2011) and preferential retention of
425 *Synechococcus* and eukaryotic algae over other prey types was observed in bivalves (Yahel et
426 al. 2009). The results (although using indirect measurements) of this study suggested
427 inversely to what we expected that members of SAR11 were grazed at higher efficiencies
428 than other available prey (Fig 5,6) somewhat reflecting their overall high abundance. This
429 work strengthens the niche speciation assumption, hinting that some taxa within the hard-
430 bottom community specialize on grazing SAR11. Besides the obvious members of the
431 suspension feeding community, we must consider the heterotrophic nanoflagellates attached
432 to the rocks (Yahel et al. 2006). This cryptic community has a large grazing effect that may
433 be part of the explanation for prey depletion above the hard-bottom. It is known that
434 heterotrophic nanoflagellates are important bacterivores in pelagic waters (Tophøj et al.
435 2018).

436 The apparent lack of grazing evasion by members of SAR11 could be possibly
437 explained by a different mode of interaction to other filter feeders compared to tunicates.
438 Suspension feeders have different filtration organs, using cilia, mucus, or both to capture and
439 process suspended particles. While some are active suspension feeders and specialize in
440 filtration of small particles, others are passive feeders and specialize in filtration of large
441 particles (Gili and Coma 1998). Different grazing strategies of benthic suspension feeders
442 would potentially increase the opportunities for exploitation of available prey by

443 communities of suspension feeders (Gili and Coma 1998). Specialization of different
444 suspension feeders on different prey would explain the homogeneous decrease of different
445 available prey. While removal by benthic filter feeders is a likely biological explanation for
446 the depletion of pico- and nanoplankton we observed, other processes such adsorption to
447 mucus might also have contributed (Decho 1990) although members of SAR11 clade are
448 known as free-living and not particle-associated or biofilm forming bacteria (Giovannoni
449 2017, Haro-Moreno et al. 2020)

450 Evaluation of evasiveness using “in versus out” experiments such those in Dadon-
451 Pilosof et al. (2017) with other taxa, or the measurement of adsorption of different bacteria
452 including cultured *Candidatus Pelagibacter ubique* to transparent exopolymer particles (Long
453 and Azam 1996) with water from different points in the transect remain as possible future
454 experiments to further clarify our results.

455




Figure 7: Schematic illustration of a near-shore thermal flow cycle. (a) during winter, when cooling of nearshore water drives offshore flow at depth and onshore flow at the surface. (b) during spring, when the shallow water near shore warms and water flow offshore at the surface and onshore below.

457 Measurements of removal rates of microbial plankton by benthic organisms at the
458 level of the whole community remain challenging despite of decades of studies of this theme
459 (Sargent and Austin 1949; Odum and Odum 1955; Johannes et al. 1972). More recent studies,
460 assessed spatial gradients of DOC, bacterioplankton and virioplankton concentrations in reef
461 ecosystem resulting in depleted microbial community over the reef compare to negligible
462 removal over sandy bottom nearby (Patten et al. 2011; Nelson et al. 2011). An alternative
463 approach utilizes a control-volume approach either by physically enclosing the community in
464 a bell jar for in situ measurements (e.g., (Hopkinson et al. 1991) or by enclosing and
465 imaginary “box” over the bottom using vertical arrays of samplers and ADCPs to quantify the
466 fluxes of plankton into and out of the control volume (Genin et al. 2002, 2009). A third
467 approach integrates individual rate measurements of dominant benthic suspension feeders
468 with their abundance and size distribution to assess the community flux and its effect on the
469 planktonic community in the overlying water (e.g., Genin et al. 2009; Lucas et al. 2016 and
470 references therein).

471 Further investigation is required to develop and estimate, based on indirect evaluation,
472 cells removal above hard-bottom suspension feeders. Methodology fine tuning is required for
473 estimating the cells removal above the hard-bottom compared to open sea and to discriminate
474 grazing, other biological processes and physical mechanisms. Previous studies showed that
475 SAR11 evade predation of benthic and pelagic ascidians mucus, but this study showed (albeit
476 indirectly) that SAR11 was removed efficiently as other prey cells by hard bottom grazers.
477 Potential grazers might be sponges, bivalves, nanoflagellates attached to the surface of the
478 rocks which filter their prey by using several different mechanisms. Understanding the
479 mechanisms and the variance between the grazers' preferences and/or the microbial cells
480 abilities to evade predation has ramifications on processes affecting the marine food web
481 such as top-down control and, benthic pelagic coupling. Grazing resistance mechanisms are

482 still understudied and should be investigated further to gain knowledge on its effects in the
483 marine ecosystem.

484 [Reference list](#)

- 485 Burnison, B. K. 1980. Modified dimethyl sulfoxide (DMSO) extraction for chlorophyll
486 analysis of phytoplankton. *Can. J. Fish. Aquat. Sci.* **37**: 129–133.
- 487 Calbet, A., and M. R. Landry. 2004. Phytoplankton growth, microzooplankton grazing, and
488 carbon cycling in marine systems. *Limnol. Oceanogr.* **49**: 51–57.
489 doi:10.4319/lo.2004.49.1.0051
- 490 Chesson, J. 1978. Measuring preference in selective predation. *Ecology* 59: 211–215.
491 doi:10.2307/1936364
- 492 Chesson, J. 1983. The estimation and analysis of preference and its Relationship to foraging
493 models. *Ecology* 64: 1297–1304. doi:10.2307/1937838
- 494 Cunningham, A., and G. A. Buonacorsi. 1992. Narrow-angle forward light scattering from
495 individual algal cells: Implications for size and shape discrimination in flow cytometry.
496 *J. Plankton Res.* **14**: 223–234. doi:10.1093/plankt/14.2.223
- 497 Dadon-Pilosof, A., K. R. Conley, Y. Jacobi, and others. 2017. Surface properties of SAR11
498 bacteria facilitate grazing avoidance. *Nat. Microbiol.* doi:10.1038/s41564-017-0030-5
- 499 Dadon-Pilosof A, Lombard F, Genin A, Sutherland KR, Yahel G (2019) Prey taxonomy
500 rather than size determines salp diets. *Limnol Oceanogr* 64(5): 1–
501 15. doi:10.1002/lno.11165
- 502 Decho, A. W. (1990). Microbial exopolymer secretions in ocean environments: their role (s)
503 in food webs and marine processes. *Oceanogr. Mar. Biol. Annu. Rev.*, 28(7), 73-153.
- 504 Eiler, A., D. H. Hayakawa, M. J. Church, D. M. Karl, and M. S. Rappé. 2009. Dynamics of
505 the SAR11 bacterioplankton lineage in relation to environmental conditions in the
506 oligotrophic North Pacific subtropical gyre. *Environ. Microbiol.* **11**: 2291–2300.
507 doi:10.1111/j.1462-2920.2009.01954.x
- 508 Garcia, N. S., J. A. Bonachela, and A. C. Martiny. 2016. Interactions between growth-

509 dependent changes in cell size, nutrient supply and cellular elemental stoichiometry of
510 marine *Synechococcus*. *ISME J.* **10**: 2715–2724. doi:10.1038/ismej.2016.50

511 Genin, A., Yahel, G., Reidenbach, M. A., Monismith, S. G., Koseff, J. R., & Monismith, S.
512 B. (2002). Intense benthic grazing on phytoplankton in coral reefs revealed using the
513 control volume approach. *Oceanography* 90–96.

514 Genin, A., S. G. Monismith, M. A. Reidenbach, G. Yahel, and J. R. Koseff. 2009. Intense
515 benthic grazing of phytoplankton in a coral reef. **54**: 938–951.

516 Gili, J., and R. Coma. 1998. Benthic suspension feeders: their paramount role in littoral
517 marine food webs. **13**: 316–321.

518 Giovannoni, S. J. 2017. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Ann.*
519 *Rev. Mar. Sci.* **9**: annurev-marine-010814-015934. doi:10.1146/annurev-marine-
520 010814-015934

521 Glöckner, F., J. Gasol, N. McDonough, and J. Calewaert. 2012. Marine Microbial Diversity
522 and its role in Ecosystem Functioning and Environmental Change. *Mar. Board-ESF Rep.*
523 84.

524 Gorsky, G., M. J. Chr, J. Blanchot, and I. Palazzoli. 1999. Picoplankton and nanoplankton
525 aggregation by appendicularians- Fecal pellet contents of *Megalocercus huxleyi* in the
526 equatorial. *J. Geophys. Research* **104**: 3381–3390.

527 Hahn, M. W., and M. G. Höfle. 2001. Grazing of protozoa and its effect on populations of
528 aquatic bacteria. *FEMS Microbiol. Ecol.* **35**: 113–121. doi:10.1016/S0168-
529 6496(00)00098-2

530 Hanson, C. E., M. J. Mclaughlin, G. A. Hyndes, and J. Strzelecki. 2009. Estuarine , Coastal
531 and Shelf Science Selective uptake of prokaryotic picoplankton by a marine sponge (
532 *Callyspongia* sp .) within an oligotrophic coastal system. *Estuar. Coast. Shelf Sci.* **84**:
533 289–297. doi:10.1016/j.ecss.2009.05.019

- 534 Haro-Moreno, J. M., Rodriguez-Valera, F., Rosselli, R., Martinez-Hernandez, F., Roda-
535 Garcia, J. J., Gomez, M. L., ... & López-Pérez, M. (2020). Ecogenomics of the SAR11
536 clade. *Environmental Microbiology*, 22(5), 1748-1763.
- 537 Herut, B., A. Almogi-labin, N. Jannink, and I. Gertman. 2000. The seasonal dynamics of
538 nutrient and chlorophyll a concentrations on the SE Mediterranean shelf-slope. **23**: 771–
539 782.
- 540 Hopkinson, C. S., R. D. B. Jansson, and J. P. Schubauer. 1991. Community metabolism and
541 nutrient cycling at ,Gray’s Reef a hard bottom habitat in the Georgia Bight. *Mar. Ecol.*
542 *Prog. Ser.* **73**: 105–120.
- 543 Jiménez, E. 2011. Nutrient fluxes in marine sponges: methodology, geographical variability
544 and the role of associated microorganisms. Universitat Politècnica de Catalunya (UPC).
- 545 Johannes, R. E., J. Alberts, C. D’Elia, R. A. Kinzie, L. R. Pomeroy, W. Sottile, and J. M.
546 Wells. 1972. The Metabolism of Some Coral Reef Communities: A Team Study of
547 Nutrient and Energy Flux at Eniwetok. *Bioscience* **22**: 541–543.
548 doi:<https://doi.org/10.2307/1296314>
- 549 Jones, N. L., J. K. Thompson, K. R. Arrigo, and S. G. Monismith. 2009. Hydrodynamic
550 control of phytoplankton loss to the benthos in an estuarine environment. *Limnol.*
551 *Oceanogr.* **54**: 952–969. doi:10.4319/lo.2009.54.3.0952
- 552 Krom, M. D., and Y. Suari. 2015. Eastern Mediterranean sea: A natural laboratory for
553 studying marine biogeochemical processes. *Ocean Challenge* **21**: 46–52.
- 554 Labiosa, R. G., K. R. Arrigo, S. G. Monismith, and G. Van Dijken. 2003. The interplay
555 between upwelling and deep convective mixing in determining the seasonal
556 phytoplankton dynamics in the Gulf of Aqaba : Evidence from SeaWiFS and MODIS.
557 **48**: 2355–2368.
- 558 Long, R. A., & Azam, F. (1996). Abundant protein-containing particles in the sea. *Aquatic*

- 559 Microbial Ecology, 10(3), 213-221.
- 560 Lucas, L. V, J. E. Cloern, J. K. Thompson, M. T. Stacey, and J. R. Koseff. 2016. Bivalve
561 Grazing Can Shape Phytoplankton Communities. **3**: 1–17.
562 doi:10.3389/fmars.2016.00014
- 563 Marie, D., C. P. D. Brussaard, R. Thyrhaug, and others. 1999. Enumeration of Marine
564 Viruses in Culture and Natural Samples by Flow Cytometry Enumeration of Marine
565 Viruses in Culture and Natural Samples by Flow Cytometry. *Appl. Environmental*
566 *Microbiol.* **65**: 45–52.
- 567 Mary, I. et al. SAR11 dominance among metabolically active low nucleic acid
568 bacterioplankton in surface waters along an Atlantic meridional transect. *Aquat. Microb.*
569 *Ecol.* 45, 107–113 (2006).
- 570 Matz, C., and S. Kjelleberg. 2005. Off the hook - How bacteria survive protozoan grazing.
571 *Trends Microbiol.* **13**: 302–307. doi:10.1016/j.tim.2005.05.009
- 572 Monismith, S. G., A. Genin, M. A. Reidenbach, G. Yahel, and J. R. Koseff. 2006. Thermally
573 Driven Exchanges between a Coral Reef and the Adjoining Ocean. *J. Phys. Oceanogr.*
574 **36**: 1332–1347.
- 575 Monismith, S. G., K. A. Davis, G. G. Shellenbarger, and others. 2010. Flow effects on
576 benthic grazing on phytoplankton by a Caribbean reef. *Limnol. Oceanogr.* **55**: 1881–
577 1892. doi:10.4319/lo.2010.55.5.1881
- 578 Morris, R. M., M. S. Rappé, S. a Connon, K. L. Vergin, W. a Siebold, C. a Carlson, and S. J.
579 Giovannoni. 2002. SAR11 clade dominates ocean surface bacterioplankton
580 communities. *Nature* **420**: 806–810. doi:10.1038/nature01240
- 581 Morris, R. M., C. D. Frazar, and C. A. Carlson. 2012. Basin-scale patterns in the abundance
582 of SAR11 subclades, marine Actinobacteria (OM1), members of the Roseobacter clade
583 and OCS116 in the South Atlantic. *Environ. Microbiol.* **14**: 1133–1144.

- 584 doi:10.1111/j.1462-2920.2011.02694.x
- 585 Nelson, C. E., Alldredge, A. L., McCliment, E. A., Amaral-Zettler, L. A., & Carlson, C. A.
586 (2011). Depleted dissolved organic carbon and distinct bacterial communities in the
587 water column of a rapid-flushing coral reef ecosystem. *The ISME Journal*, 5(8), 1374-
588 1387.
- 589 Odum, H. . ., and E. . Odum. 1955. Trophic Structure and Productivity of a Windward Coral
590 Reef Community on Eniwetok Atoll. *Ecol. Soc. Am.* **25**: 291–320. doi:10.2307/1943285
- 591 Page, E. B. 1963. Ordered Hypotheses for Multiple Treatments : A Significance Test for
592 Linear Ranks. *J. Am.* 37–41.
- 593 Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: assessing
594 small subunit rRNA primers for marine microbiomes with mock communities, time
595 series and global field samples. *Environmental microbiology*, 18(5), 1403-1414.
- 596 Partensky, F., W. R. Hess, and D. Vaultot. 1999. Prochlorococcus, a marine photosynthetic
597 prokaryote of global significance. *Microbiol.Mol Biol.Rev.* **63**: 106–127. doi:doi:1092-
598 2172/99/\$04.00
- 599 Patten, N. L., A. S. J. Wyatt, R. J. Lowe, and A. M. Waite. 2011. Uptake of
600 picophytoplankton, bacterioplankton and virioplankton by a fringing coral reef
601 community (Ningaloo Reef, Australia). *Coral Reefs* **30**: 555–567. doi:10.1007/s00338-
602 011-0777-8
- 603 Pernthaler, J. 2005. Predation on prokaryotes in the water column and its ecological
604 implications. *Nat. Rev. Microbiol.* **3**: 537–546. doi:10.1038/nrmicro1252
- 605 Pomeroy, L., P. leB. Williams, F. Azam, and J. Hobbie. 2007. The Microbial Loop.
606 *Oceanography* **20**: 28–33. doi:10.5670/oceanog.2007.45
- 607 Rappé, M. S., S. a Connon, K. L. Vergin, and S. J. Giovannoni. 2002. Cultivation of the
608 ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**: 630–633.

- 609 doi:10.1038/nature00917
- 610 Raveh, O., N. David, G. Rilov, and E. Rahav. 2015. The temporal dynamics of coastal
611 phytoplankton and bacterioplankton in the eastern mediterranean sea. *PLoS One* **10**.
612 doi:10.1371/journal.pone.0140690
- 613 Riisgard, H. U., and P. S. Larsen. 2001. Minireview: Ciliary filter feeding and bio-fluid
614 mechanics- present understanding and unsolved problems. *Limnitol. Ocean.* **46**: 882–
615 891. doi:10.4319/lo.2001.46.4.0882
- 616 Rilov, G., O. Peleg, E. Yeruham, T. Garval, A. Vichik, and O. Raveh. 2018. Alien turf:
617 Overfishing, overgrazing and invader domination in south-eastern Levant reef
618 ecosystems. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **28**: 351–369. doi:10.1002/aqc.2862
- 619 Rilov, G., O. Peleg, and T. Gan-Haim. 2019. The restructuring of Levant reefs by aliens,
620 ocean warming and overfishing: implications to species interactions and ecosystem
621 functions, *In* S.J. Hawkins, K. Bohn, L.B. Firth, and G.A. Williams [eds.], *Interactions*
622 *in the Marine Benthos Global Patterns and Processes*. Cambridge University press.
- 623 Salter, I., Galand, P.E., Fagervold, S.K., Lebaron, P., Obernosterer, I., Oliver, M.J., et al.
624 (2015) Seasonal dynamics of active SAR11 ecotypes in the oligotrophic Northwest
625 Mediterranean Sea. *ISME J* 9:347–360.
- 626 Sánchez, O., Ferrera, I., Mabrito, I. et al. Seasonal impact of grazing, viral mortality, resource
627 availability and light on the group-specific growth rates of coastal Mediterranean
628 bacterioplankton. *Sci Rep* 10, 19773 (2020). [https://doi.org/10.1038/s41598-020-76590-](https://doi.org/10.1038/s41598-020-76590-5)
629 5
- 630 Sargent, M. C., and T. S. Austin. 1949. Organic productivity of an Atoll. *Eos, Trans. Am.*
631 *Geophys. Union* **30**: 245–249. doi:10.1029/TR030i002p00245
- 632 Simon, N., G. R. Barlow, D. Marie, F. Partensky, and D. Vaultot. 1994. Characterization of
633 oceanic photosynthetic picoeukaryotes by flow cytometry. *J. Phycol.* **30**: 922–935.

- 634 Siokou-Frangou, I., Christaki, U., Mazzocchi, M. G., Montresor, M., Ribera d'Alcalá, M.,
635 Vaqué, D., & Zingone, A. (2010). Plankton in the open Mediterranean Sea: a
636 review. *Biogeosciences*, 7(5), 1543-1586.
- 637 Suari, Y., Dadon-Pilosof, A., Sade, T., Amit, T., Gilboa, M., Gafny, S., ... & Yahel, G.
638 (2019). A long term physical and biogeochemical database of a hyper-eutrophicated
639 Mediterranean micro-estuary. *Data in brief*, 27, 104809.
- 640 Teeling, H., B. M. Fuchs, D. Becher, and others. 2012. Substrate-controlled succession of
641 marine bacterioplankton populations induced by a phytoplankton bloom. *Science* **336**:
642 608–11. doi:10.1126/science.1218344
- 643 Topçu, A., P. Thierry, G. Gérald, and M. Harmelin-vivien. 2010. In situ investigation of
644 *Spongia officinalis* (Demospongiae) particle feeding: Coupling flow cytometry and
645 stable isotope analysis Nur Eda. *J. Exp. Mar. Bio. Ecol.* **389**: 61–69.
- 646 Tophøj, J., Wollenberg, R. D., Sondergaard, T. E., & Eriksen, N. T. (2018). Feeding and
647 growth of the marine heterotrophic nanoflagellates, *Procrystobia sorokini* and
648 *Paraphysomonas imperforata* on a bacterium, *Pseudoalteromonas* sp. with an inducible
649 defence against grazing. *Plos one*, 13(4), e0195935.
- 650 Uysal, Z. 2001. Chroococcoid cyanobacteria *Synechococcus* spp. in the Black Sea: pigments,
651 size, distribution, growth and diurnal variability. *J. Plankton Res.* **23**: 175–190.
- 652 Verity, P. G. 1991. Feeding In Planktonic Protozoans: Evidence For Non-Random
653 Acquisition of Prey. *J. Protozool.* **38**: 69–76. doi:10.1111/j.1550-7408.1991.tb04804.x
- 654 Welschmeyer, N. A., and S. L. Naughton. 1994. Improved chlorophyll a analysis: single
655 fluorometric measurement with no acidification. *Lake Reserv. Manag.* **9**: 123.
- 656 Worden, A. Z., and F. Not. 2008. Ecology and diversity of picoeukaryotes, p. 159–205. *In*
657 *Microbial ecology of the oceans*.
- 658 Yahel, G., D. Marie, A. F. Post, D. Vaultot, K. Fabricius, and A. Genin. 1998. Phytoplankton

- 659 distribution and grazing near coral reefs. *Limnol. Oceanogr.* **43**: 551–563.
660 doi:10.4319/lo.1998.43.4.0551
- 661 Yahel, G., D. I. Eerkes-Medrano, and S. P. Leys. 2006. Size independent selective filtration
662 of ultraplankton by hexactinellid glass sponges. *Aquat. Microb. Ecol.* **45**: 181–194.
663 doi:10.3354/ame045181
- 664 Yahel, G., D. Marie, P. G. Beninger, S. Eckstein, and A. Genin. 2009. In situ evidence for
665 pre-capture qualitative selection in the tropical bivalve *Lithophaga simplex*. *Aquat. Biol.*
666 **6**: 235–246. doi:10.3354/ab00131
- 667 Zubkov, M. V., J. I. Allen, and B. M. Fuchs. 2004. Coexistence of dominant groups in marine
668 bacterioplankton community - A combination of experimental and modelling
669 approaches. *J. Mar. Biol. Assoc. United Kingdom* **84**: 519–529.
670 doi:10.1017/S002531540400952Xh

671

672 [Acknowledgements](#)

673 Research funding was provided by Ruppin Academic center, support provided to
674 A.D.P by the Mediterranean Sea Research Center of Israel. I thank my supervisors Prof.
675 Amatzia Genin and Prof. Gitai Yahel for their advice along the way. Special gratitude to R.
676 Rosenblatt for availability and technical assistance throughout the field experiments.

677 [Author Contributions:](#)

678 **A.D.P.** designed the study and participated in field experiments, data analysis, and
679 manuscript preparation. **K.R.C**, participated in data analysis, and contributed substantially to
680 drafting the manuscript. **M. T.S** designed and performed bioinformatics analyses and
681 contributed substantially to drafting the manuscript. All authors discussed the results and
682 commented on the manuscript during its preparation and approved the submitted version of
683 the manuscript.