



HAL
open science

Thyroid-disrupting chemicals and brain development: an update

Bilal B Mughal, Jean-Baptiste Fini, Barbara A Demeneix

► **To cite this version:**

Bilal B Mughal, Jean-Baptiste Fini, Barbara A Demeneix. Thyroid-disrupting chemicals and brain development: an update. *Endocrine Connections*, 2018, 7 (4), pp.R160 - R186. 10.1530/ec-18-0029 . hal-03919596

HAL Id: hal-03919596

<https://hal.science/hal-03919596>

Submitted on 3 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Thyroid disrupting chemicals and brain development: An update

Bilal B. Mughal, Jean-Baptiste Fini, & Barbara A. Demeneix

CNRS / UMR7221, Muséum national d'Histoire naturelle/ Université Paris-Sorbonne, Paris,
France

Corresponding author and person to whom reprint requests should be addressed:

Prof. Barbara A Demeneix

Département Adaptations du Vivant,

Muséum national d'Histoire naturelle/ Université Paris-Sorbonne

75005, Paris, France

Email: bdem@mnhn.fr, fini@mnhn.fr

Phone: +33 01 40 79 36 07

Fax: +33 01 40 79 36 06

Disclosure statement: BBM and JBF have nothing to disclose. BD is a cofounder of WatchFrog.

Abstract

This review covers recent findings on the main categories of thyroid hormone disrupting chemicals and their effects on brain development. We draw mostly on epidemiological and experimental data published in the last decade. For each chemical class considered, we deal with not only the thyroid hormone disrupting effects but also briefly mention the main mechanisms by which the same chemicals could modify estrogen and/or androgen signalling, thereby exacerbating adverse effects on endocrine-dependent developmental programs. Further, we emphasize recent data showing how maternal thyroid hormone signalling during early pregnancy affects not only offspring IQ, but also neurodevelopmental disease risk. These recent findings add to established knowledge on the crucial importance of iodine and thyroid hormone for optimal brain development. We propose that prenatal exposure to mixtures of thyroid hormone-disrupting chemicals provides a plausible biological mechanism contributing to current increases in incidence of neurodevelopmental disease and IQ loss.

Introduction

Thyroid hormone (TH) is essential for normal brain development where it influences, during specific temporal windows, neurogenesis, neuronal migration, neuronal and glial cell differentiation, myelination, and synaptogenesis. These TH-dependent processes are crucial during early gestation and postnatal development, then continue, albeit at reduced rates, throughout adulthood. During the first 10-12 weeks of gestation, the foetus relies entirely on maternal TH. Hence, severe maternal TH deficiency adversely affects offspring neurodevelopment^{1,2}. Recent epidemiological evidence suggests that even more moderate forms of maternal thyroid dysfunction may affect child cognitive development and increase the risk of neurodevelopmental disorders³⁻⁶. Therefore it is important to gain a better understanding of early thyroid dysfunction on offspring neurodevelopment^{1,7}.

Another major cause of thyroid dysfunction can be the presence of thyroid disrupting chemicals in the maternal and foetal environment. Endocrine disrupting compounds (EDCs) are xenobiotics that modulate hormonal homeostasis thereby inducing adverse effects⁸. Numerous EDCs identified to date contain a halogen group substitution with chlorine and bromine. Interestingly, THs are the only complex halogenated (iodine) molecules produced by and necessary for vertebrate homeostasis, making TH physiology highly vulnerable to EDCS. Halogen substituted phenolic moieties can mimic natural THs and thereby interact with multiple aspects of hormone production, feedback, distribution, entry into cells, intracellular metabolism (deiodination, conjugation) of TH, as well as at the level of receptors, as antagonists or analogues.

The aim of this review is to provide an update on how different chemicals in the environment can disrupt thyroid signalling and thereby affect brain development. A number of previous reviews have addressed certain aspects of this question. Notably, in 1998,

Brucker-Davis and colleagues ⁹ reviewed the different classes of chemicals that could affect thyroid signaling and Zoeller and Crofton ¹⁰ underlined how endocrine disruption affected early brain development. A decade later the situation was updated by Crofton ¹¹ and by Boas and colleagues ¹².

No new major classes of TH-disrupting chemicals have been characterised since the last review appeared. However, within classes certain novel compounds have attracted attention. Most often these new compounds have been introduced to replace a similar chemical for which adverse effects were reported. This has led to many examples of “regrettable substitutions” within classes, cases of which are described below. Thus, our focus remains on perchlorate, phenols, pesticides, polychlorinated-biphenyls (PCBs), poly brominated flame-retardants, perfluorinated compounds (PFCs), and phthalates (Fig. 1). Many of these substances are classed as persistent organic pollutants (POPs) and were banned decades ago yet they remain environmentally relevant due to their previous high production volumes and exceptionally long half-lives.

Perchlorate

Perchlorate is a well-characterised inhibitor of the sodium-iodide symporter (NIS) that is expressed in the basal membrane of thyroid follicular cells and is critical for iodide uptake ¹³. Two other NIS inhibitor classes, nitrates and thiocyanates, are found at significant levels in human fluids, but in molar terms they are respectively 240 and 15 times less active than perchlorate ¹⁴. However, their environmental levels are such that their combined effects should be more often taken into account ^{15,16}.

Given first, its wide-ranging uses as an oxidant in products ranging from in rocket fuel to airbags and second, its high stability, perchlorate contamination is widespread.

Epidemiological data show, that despite its short half-life (<8h in humans) continual exposure means that the chemical is virtually ubiquitous in the US population ¹⁷. Epidemiological evidence showed that perchlorate levels were associated with TSH in women, and this association was stronger in women with <100µg/ L urinary iodine ¹⁸. The relationship was even greater in women who smoke, related to the fact that cigarette smoke is a source of thiocyanate.

Furthermore, recent epidemiological data analysed pregnant women for their thyroid status and perchlorate levels ¹⁹. Offspring born to those women that were both borderline hypothyroid or hypothyroxinemic and had higher perchlorate levels had a higher risk of being in the lowest 10% for IQ scores. The adverse effect of perchlorate was not modified by thyroxine therapy (150µg/day) during pregnancy. However, it is possible that the timing of replacement (after 12 weeks pregnancy) was too late to exert corrective effects. Other recent data also link maternal perchlorate exposure to modified thyroid function during pregnancy ²⁰. Iodine has long been known to be required for thyroid hormone synthesis and both iodine deficiency and maternal hypothyroidism are risk factors for both decreased IQ and neurodevelopmental disease (see ^{21,22}). It is worth noting that in the study cited ²⁰ a large proportion of the women (74%) had urinary iodine levels below the recommended median level (150µg/L) for pregnancy, raising the question of whether iodine deficiency exacerbates the effects of perchlorate (and potentially other TH-disrupting chemicals). This question deserves far more research and needs to be taken into account in both epidemiological and experimental studies. Similarly, the presence of thyroid disrupting chemicals has been identified as a confounder for epidemiological studies assessing effects of iodine supplementation during pregnancy ²³.

Phenols

Two principal phenols are well-characterised TH-disruptors, bisphenol A (BPA) and triclosan (TCS). Both have high production volumes and been so extensively used that they are now virtually ubiquitous contaminants of human fluids²⁴ and the environment²⁵.

Bisphenol A (BPA, 4,4' isopropylidenediphenol)

BPA is an organic synthetic compound, first identified as a synthetic estrogen in 1930s²⁶. Current common uses of BPA are in plastic products such as water bottles and food containers, CDs, DVDs, safety equipment, thermal paper and medical devices. In the US, France and Denmark, BPA is restricted for certain uses, such as baby bottles. More recently, since 2015, France banned the use of BPA in plastic food containers. The same year,(2015) EFSA maintained their opinion delivered in 2013 that BPA poses no health risks but the committee lowered the tolerable daily intake from 50 µg/kg bw/day to 4 µg/kg bw/day²⁷. Despite these recent restrictions, there is still widespread exposure to BPA in human populations²⁸. It is retained in humans and has been found in pregnant women's serum, placenta and breast milk²⁹⁻³⁴. As restrictions were increasingly placed on BPA use, a number of structural BPA analogues such as bisphenol S (BPS), bisphenol F (BPF), and bisphenol B (BPB) were marketed. These analogues are found now in considerable quantities in the human urine³⁵⁻³⁸. These replacement chemicals are often described as “regrettable substitutions” as their EDC-related effects are apparently no less than those of BPA, including effects on TH signalling^{39,40} and ER signalling⁴¹.

As BPA is primarily thought of as an estrogen disruptor but is also a TH modulator. EDC action across endocrine systems is to be expected as crosstalk exists at multiple levels:

from different nuclear receptors⁴² to individual target genes and networks to physiological systems. As BPA can interact with multiple nuclear hormone receptors including ER⁴³, estrogen-related receptors (ERR)⁴⁴, AR^{45,46}, and thyroid hormone receptors (TR)^{39,47,48} potential cross talk needs to be considered at multiple levels.

BPA and ER interaction has been reviewed extensively elsewhere, for both classical and non-classical estrogen receptors^{8,49}. As to TRs, some experimental studies show that BPA does not bind to TR based on a competitive TR binding test⁵⁰⁻⁵², others show T₃-TR mediated agonistic and antagonistic effects of BPA^{48,53,54}. More recently, binding affinities of BPA and its analogues BPF and BPS, with TR were calculated *in silico* and found to be roughly similar⁵⁵. When tested by a spectrum of *in vitro* and *in vivo* methods, all three analogues activated TH signaling in the absence of T₃³⁹. The *in vitro* approaches included competitive binding assays, molecular docking and co-activator binding assays whereas the *in vivo* methodology exploited TH-response gene responses in *Pelophylax nigromaculatus* tadpoles.

In human epidemiology, studies have reported changes in TH parameters as a function of BPA exposure in adults⁵⁶⁻⁵⁹, including in pregnant women⁶⁰⁻⁶². In pregnant women, maternal BPA levels were inversely⁶² or positively⁶⁰ associated with T₄ levels while two studies reported no association^{61,63}. In humans, inverse associations of BPA with TSH have also been reported in both sexes⁶⁰ and in other cases, only in women⁶¹. Yet another study reported a positive association⁶⁴. Such inconsistencies need to be examined notably for methodological differences between studies.

We now focus specifically on BPA and TH and neurodevelopment. Increased prenatal BPA exposure is implicated in several sex-specific changes in child behaviour⁶⁵⁻⁷⁰. Prenatal

BPA exposure is linked to increased internalizing behaviours in boys ^{66,67,70} and increased risk of ADHD related behaviour ⁶⁵. In girls, prenatal BPA exposure has been associated with both internalising and externalising behaviours ^{66,69}, as well as poor executive function ⁶⁸. It is possible that differences in the results is due to varying timing of sample collection, exposure, and assessment among the studies. Studies on post-natal childhood BPA exposure and effects on neurodevelopment are even more inconsistent. Some studies report a positive association of BPA levels with ADHD-linked behaviours in both girls and boys ⁷¹, and anxious, depressive or aggressive behaviours in girls ^{66,67,72}. Others report null association with childhood BPA exposure and neurodevelopmental outcomes ^{68,73}. Pubertal BPA exposure has also been associated with poorer cognitive performance in adolescence ⁷⁴ and adulthood ⁷⁵.

The lack of full endocrine profiles in these epidemiological studies make it hard to pinpoint the exact mechanism linking endocrine disruption and neurodevelopmental outcome. Animal studies however can better define mode of action. Such studies link BPA levels with behavioural outcomes often associated with TH disruption including, hyperactivity (not sex specific) ^{76,77}, anxiety ⁷⁸, and decreased motor activity ⁷⁹. Prenatal BPA exposure in mice also results in mostly sex-specific changes in aggression and cognitive defects ⁸⁰⁻⁸⁵. These sex-specific changes are not surprising due to the role of estrogen in differentiation of sexually dimorphic areas involved in behaviour and cognitive development ⁸⁶. BPA exposure also causes epigenetic changes (methylation) on the *ER- α* gene in the cortex and hypothalamus of male and female mice, and alters mRNA levels of DNA methyltransferases *DNMT1* and *DNMT3A* ^{78,87}. Interestingly, *DNMT3A* is a well-known TH responsive gene, activated by liganded TRs ^{88,89}.

Halogenated BPAs include a bromine (Tetrabromobisphenol-A, TBBPA) or chlorine (Tetrachlorobisphenol-A, TCBPA) substitute on the phenolic ring and are common flame

retardants. TBBPA is currently the flame-retardant with the highest production volume worldwide. It is found in printed electronic circuit boards and in plastics for electrical housings or piping. Due to its high production volume, toxicological effects attributed to TBBPA have been extensively reviewed by governments⁹⁰⁻⁹² and deemed to have no health hazard, risk or concerns to humans. However, Van der Ven and colleagues⁹³ assessing multiple *in vivo* studies on rats concluded, that the margin of exposure for humans was only 2.6 and that TBBPA exposure was a matter of concern for authorities.

What is more, TBBPA has been established *in vitro* as neuro-toxicant that disrupts multiple intracellular pathways including zinc and calcium homeostasis, inducing oxidative stress⁹⁴⁻⁹⁷ as well as acting as a partial GABA_A agonist at 0.1µM⁹⁸. So far, results for *in vivo* developmental toxicity are less consistent. The conclusions of Van der Ven et al (2008) for instance contrast with those of toxicologists^{99,100}. Although Viberg & Eriksson (2011) reported more marked effects for PBDE 99 than TBBPA in neonatal mouse brain, they observed down regulation of a certain classes of nicotinic receptors in the frontal cortex with both chemicals¹⁰¹. In our laboratory, studies on mice showed that gestational exposure to TBBPA decreased TRH receptor and melanocortin 4 receptor basal expression in pups, dramatically affecting T₃ induced repression of these gene expression¹⁰².

Some studies have reported neural defects, including impaired motor function in Zebrafish¹⁰³. Similarly, Nakajima and colleagues reported behavioural effects of TBBPA administration in adult mice and differential accumulation of the chemical according to brain region¹⁰⁴. Similarly, Lilienthal et al. 2008 noted increased latency of hearing responses in a rat one generation study¹⁰⁵. Interestingly, development of the inner ear is known to be a TH-dependent process¹⁰⁶.

Significant reduction in circulating T_4 is the most frequent phenotype seen across rodent studies as a function of TBBPA exposure^{93,107}. T_4 reductions could occur through activation of (UDP-glucuronosyltransferase, UGT), which increases metabolism of T_4 in the liver and subsequent reduction of serum T_4 levels¹⁰⁸. *In vitro*, TBBPA competes with binding of transthyretin (TTR) and interferes with T_3 -dependent cell proliferation¹⁰⁹. In a fluorescent polarization assay, TBBPA was found to modulate both co-activator and co-repressor interactions with TR¹¹⁰. TBBPA also shows thyroid disrupting effects in amphibian models. In *Rana rugosa*, TBBPA displayed inhibitory effects on T_3 -induced tail shortening¹¹¹ and in *Xenopus laevis*, TBBPA exerted antagonistic effects in the presence of high TH levels, but agonistic activity with low TH levels¹¹². TBBPA has also been demonstrated as a TH disruptor using both the amphibian metamorphosis assay^{112,113} and the *Xenopus* embryonic thyroid assay (XETA)¹¹⁴. In the latter study, TBBPA was further found to alter expression of TH target genes implicated in neural stem cell function and differentiation. Whether such effects extend to other proliferative brain regions during development remains to be investigated¹¹⁵.

In rats TBBPA exposure increases estrogen levels¹⁰⁸ and uterine tumours¹¹⁶, effects thought to be related to inhibition of liver estradiol sulfotransferase¹⁰⁹. The combined effects of TBBPA, increasing estrogen and antagonising TH signalling could well interact to modify genetic and cellular responses, as well as inducing longer term adverse physiological responses governing reproduction.

Triclosan (TCS, 2,4,4 - trichloro-hydroxy diphenyl ether)

Triclosan (TCS) is a widely used chlorinated phenolic antimicrobial and antifungal agent. It has been used for over 40 years as an antiseptic, disinfectant, or preservative in medical and personal care products such as hand soaps and shampoos, mouthwash,

toothpaste, and cosmetics. While it has been banned from soaps and body washes in the US, it is still extensively used in skin care products and toothpaste. TCS has a short half-life in humans, i.e. it is rapidly absorbed, metabolized, and eliminated (primarily via urine) with a median excretion half-life of 11 h after oral intake ¹¹⁷. Despite this rapid clearance, the over use of products containing TCS maintain permanent, but varying exposure. The most likely sources in humans are ingestion and skin absorption. TCS has been found in majority of the urine samples obtained via population based studies in North America ^{118,119}. TCS has also been detected in human milk, and pregnant women's urine ¹²⁰⁻¹²². EU has restricted TCS use as a preservative to a maximum concentration of 0.2% in mouthwashes and 0.3% in other categories ¹²³. In the environment, TCS likely accumulates in sediments as it is a lipophilic compound with low aqueous solubility and is commonly found contaminant in solid and water compartments ^{124,125}

Several animal studies have confirmed TCS to act as a TH-disruptive chemical. In pregnant rats, TCS decreases serum T₃ and T₄, disrupts pup sex ratio balance and lowers their body weights ¹²⁶⁻¹²⁸. TH disruption is also evident during weaning rats when their mothers are exposed to triclosan ^{126,127}. In mice, decreased levels T₄ are also observed after a short term oral exposure to triclosan ¹²⁹⁻¹³¹. In amphibian models, the North American bullfrog (*Rana catesbeiana*) and *Xenopus laevis*, TCS exposure results in the disruption of TH-dependent metamorphosis, marked metabolic disorders of the liver and modulation of innate immunity ¹³²⁻¹³⁴.

In addition to TH, numerous studies report adverse effects of TCS exposure on reproductive organ development in male rats i.e., decreased testosterone and sperm production ¹³⁵, and early age of pubertal onset in female mice ¹³⁶. *In vitro* assays have confirmed TCS to act as an estrogen agonist using ER α and ER β reporter gene assays ¹³⁷⁻¹³⁹

stimulate breast and ovarian cancer cell growth in vitro^{140,141} and magnifying the effects of ethinyl estradiol^{136,142}. In rodent models, TCS, like TBBA (see above) inhibits estrogen sulfation by inhibiting sulfotransferases, thus preventing metabolism of estradiol into biologically inactive forms^{143–145} thereby increasing circulating estrogen levels¹⁴³. Similar effects are seen in sheep¹⁴⁴. It is worth noting that these same sulfotransferases metabolise TH as well.

Epidemiological studies have investigated the short term and long-term effect of TCS and TH parameters, with inconsistent findings^{56,146–150}. Many report no significant disruptions in TH levels while some report only most marked effects^{149,151}. Among the effects, some observe a positive association between TCS and total T₃¹⁴⁹ in adolescents, while others report an inverse association between TCS and fT₃¹⁵¹ levels in pregnant women. A prospective study on prenatal TCS exposure recently reported reduced head circumference in boys but not girls¹⁵².

Flavonoids

Flavonoids are phenols that occur as natural food items. Recent work identified the plant extract (*Silybum*) silymarin, and its derivatives silychristin and silybinin, as inhibitors of the membrane TH transporter mct8 (*slc16a2*). Entry of both T₃ and T₄ into target cells is reduced¹⁵³. This feature highlights the possible, and little studied, effects of compounds that interact with membrane TH transporters.

Pesticides

Pesticide usage increased dramatically over the last century, arguably to keep up with the demands of a growing population. However, many studies have shown that pesticide usage is excessive and that yields can be maintained even when halving pesticide use¹⁵⁴.

Many pesticides exert toxicological effects, including on thyroid signalling. Notably, the European Food Safety Authority reported that of 287 pesticide files examined, 101 showed effects indicative of thyroid disruption ¹⁵⁵. Even though many incriminated pesticides have now been banned, many of them are still in use in emerging economies. Further, many of these chemicals are persistent due to their long half-lives and remain in the environment long after their ban. Such pesticides are called legacy pesticides, with many being common environmental contaminants. Here we choose a few examples of this latter category and some others that currently on the market, but are potentially problematic.

Dichlorodiphenyltrichloroethane (DDT)

Dichlorodiphenyltrichloroethane (DDT) is an organochlorine insecticide, first used in World War I to control malaria and typhus. Its initial notoriety arose due to widespread effects on wildlife described by Rachel Carlson in her 1962 book *Silent Spring* ¹⁵⁶, notoriety that led to its ban in the United States by 1972 and worldwide by the Stockholm Convention on Persistent Organic Pollutants later that decade. Despite the ban, it is still used in certain countries to fight against malaria and dengue fever ¹⁵⁷. DDT, and its main metabolite dichlorodiphenyltrichloroethane (DDE), are highly persistent, lipophilic compounds that bioaccumulate and are still found in significant amounts in the environment and in humans including pregnant women ¹⁵⁸⁻¹⁶⁰. Prenatal exposure to p,p'-DDT and p,p'-DDE has been associated with obesity ^{161,162} and a significant reduction in children's psychomotor neurodevelopment ¹⁶²⁻¹⁶⁸, in some cases, in a sex specific manner ^{162,169,170}. The latter is not surprising as DDT binds to and activates estrogen receptors (ERs) in both reproductive and other tissues including the brain ^{171,172}. On the other hand DDE has been shown to inhibit androgens from binding to their receptors ^{173,174}. In adolescent boys, DDE is associated with increased testosterone ¹⁷⁵ and decreased luteinizing hormone ¹⁷⁶ while DDT is associated with

decreases in both luteinizing hormone, and testosterone ¹⁷⁶. In women, *in utero* exposure to DDT, as judged by umbilical cord levels, has been associated with an increased risk of breast cancer later in life ¹⁷⁷.

DDT and its metabolites have also been confirmed as thyroid disrupting chemicals through human epidemiological studies ^{158,178–181}. Studies have found negative association with DDE and total T₃ and T₄ levels ^{182,183} and a positive association with TSH levels ¹⁸³; suggesting an anti-thyroid effect. In contrast, recent studies found a positive association of DDE with total T₃ and T₄ levels, and a non-significant TSH reduction in floriculture workers ^{158,179}. These differences could be due to different levels of exposure and /or exposure to additional chemicals and characteristics of the populations studied, such as iodine or thyroid status and genetic factors.

In experimental studies, rats exposed to DDE exposure exhibit lower free T₄ levels. One target of DDT action on thyroid metabolism may be through the inhibition of TSH-stimulated intracellular accumulation of cyclic adenosine monophosphate (cAMP) by the action of DDT on the TSH receptor ^{184–187}. The highly lipophilic DDT may also interfere indirectly with the TSH receptor by altering the phosphor-lipid composition of the thyroid cell membrane, rendering the TSH receptor unable to internalise and instead be released extracellularly in vesicle forms in the presence of DDT ¹⁸⁸. These vesicles have been suggested to initiate autoimmunity favouring development of Graves' disease ¹⁸⁴. Mice exposed to DDE also exhibit reduced expression of TTR and *Dio2* mRNA which further explains the reduced free T₄ levels observed ^{189,190}. Increased expression of several hepatic enzymes can further contribute to TH degradation ¹⁹⁰. Another study in mice reported that DDT exposure was associated with increased peripheral conversion of T₄ into T₃, reduced TSH levels and morphological changes in the thyroid gland typical of iodine deficiency ¹⁹¹.

Hexachlorobenzene (HCB)

Hexachlorobenzene (HCB) is an organochloride, used primarily as a fungicide for seeds and as a wood preserving agent. It was banned globally in 1979 under the Stockholm Convention on persistent organic pollutants as a pesticide. Its current main source is through the industrial emission as a by-product of the manufacture of chlorinated solvents and pesticides. It is extremely lipophilic and accumulates in the environment. It gained prominence during late 1950s when accidentally over-treated HCB treated seeds were consumed by the general public in Turkey. Affected individuals, primarily children, displayed changed porphyrin metabolism, leading to *porphyria cutanea tarda*, enlarged liver and thyroid gland, and osteoporosis^{192,193}. Similar effects have been observed in HCB exposed rats¹⁹⁴ i.e. hepatic and thyroid neoplasms¹⁹⁵⁻¹⁹⁷, porphyria^{193,198}. Other epidemiological studies have found associations between lower levels of HCB and decreased gestational length¹⁹⁹, poor social competence²⁰⁰, and increased body weight during childhood²⁰¹. Studies on floriculture workers have further revealed an association between HCB with decreased levels total T₄ (TT₄)²⁰², and TT₃²⁰³. In animal studies, HCB is known to disrupt progesterone and estradiol concentrations²⁰⁴⁻²⁰⁶, impair reproductive efficiency^{207,208} and reduce neonatal viability and growth²⁰⁹. It also disrupts levels of T₃ and T₄^{210,211}, leads to goitre²¹², and hypothyroidism²¹³. Disruption of the TH axis may partly be due to HCB's action on the activity and expression of hepatic Dio1 and Dio2 enzymes, respectively²¹⁴. In rats, HCB has been shown to induce apoptosis in the thyroid cells, most likely due to action on mitochondria through oxidative stress^{215,216}. There is also evidence that HCB may competitively inhibit binding of thyroxine to serum carrier proteins^{214,215}. More investigations are required to elucidate the exact mechanisms of HCB on TH signalling.

Chlorpyrifos (CPF, O, O-diethyl O-(3,5,6-trichloro-2-pyridinyl) ester phosphorothioic acid)

Chlorpyrifos is a member of the organophosphate class of insecticides that target the central and peripheral nervous system specifically inhibiting the enzyme activity of acetylcholinesterase ²¹⁷. It is currently one of the most widely used insecticide in US and other countries, being used to manage insect pests on agricultural crop. CPF applications were once particularly heavy in urban areas, where the exposed populations included pregnant women ^{218,219}. Interestingly, its ban in household use in 2001 allowed for a natural experiment within an ongoing birth cohort study at Columbia University. Before the ban, decreases in birth weight and length were observed in association with CPF in newborn cord blood. After the ban, these outcomes disappeared ²²⁰. This result has been observed more recently ²²¹, a result that is not surprising as CPF readily crosses the placenta ²²². Further studies have reported prenatal CPF exposure association with impaired cognition and motor function ²²³, attention deficit hyperactive disorder ²²⁴, deficits in working memory and reduced IQ ²²⁵, and tremors during childhood ²²⁶. While some groups have considered that the levels of CPF in cord blood are too low to induce adverse effects ^{227 228}, one needs to take into account that the half-life of CPF is approximately 27 h ²²⁹. Thus, the possibility of substantially higher *in utero* levels of CPF are considerable. Despite these studies and its recent thorough evaluation by World Health Organization (WHO) and Danish Environmental Protection Agency finding toxicological evidence to be strong and the epidemiological evidence to be of moderate-to-high quality, EPA denied a recent petition for ban calling it “crucial to U.S. agriculture” ²³⁰. Not surprisingly, given the well demonstrated epidemiological data showing negative effects on brain development, this decision has been severely criticised ²³¹.

Given the importance of TH to brain development, the neurological and impaired cognitive outcomes associated with CPF exposure could well have underlying thyroid hormone dependent mechanisms. Two studies based on analysis of NHANES data from the years 1999-2002 describe significant associations between levels of chlorpyrifos metabolite, 3, 5, 6-trichloro-2-pyridinol (TCPY) and thyroid parameters, namely increased TT_4 in both males and females and decreased TSH levels in males, with increased TSH levels in females^{232,233}.

In rat studies, a reduction in brain T_4 levels is seen following prenatal CPF exposure whereas postnatal exposure results in a transient elevation in young adulthood²³⁴. Mice exposed to low dose CPF display reduced serum T_4 levels and display altered thyroid follicular size, with an apparent higher vulnerability in males²³⁵, and anxiety-like behaviour²³⁶. Reduction of T_4 in response to CPF has also been observed in rats²³⁷, whereas exposure to CPF's methyl counterpart (chlorpyrifos-methyl) results in reduced T_4 and increased TSH (hypothyroidism)²³⁸. In our lab, CPF was shown to affect TH signaling using a transgenic reporter. Moreover, a short embryonic exposure impacts mature brain structure (Spirhanzlova et al, manuscript in preparation).

As to the effects on ER and AR signalling, CPF has been found to interfere with the $ER\beta$ mRNA steady state level^{239,240} and exert an $ER\alpha$ -dependent estrogenic effect on cell proliferation *in vitro*²⁴¹ and *in vivo*²⁴². CPF also has anti-androgenic activity as Leydig cells from the rat exposed to CPF *in vitro* exhibit a significant decrease in testosterone biosynthesis²⁴³. More epidemiological and experimental data are urgently needed as this pesticide is being assessed for renewal in the EU from 2019.

Other pesticides

As previously stated, more than a 100 of the 287 pesticides examined by European Food Safety Authority (EFSA), had features indicative of thyroid disruption¹⁵⁵. In the recent review of chemicals that could be used as reference for thyroid disruption screening, Wegner and colleagues identified a number of phytopharmaceuticals or biocides with thyroid disrupting activity²⁴⁴. To cite a few, their list contained different classes of fungicides, organophosphates, along with a number of juvenile hormone analogues. Among the fungicides figure the ethylene bisdithiocarbamates (EBDC) e.g. maneb, mancozeb, ziram, zineb. Their common degradation product, ethylene thiourea (ETU), in addition to being a type IIB carcinogen, interferes with iodide organification by inhibiting thyroid peroxidase (TPO)²⁴⁵ thereby decreasing thyroidal production of T₃ and T₄ in experimental animals^{246,247}. In addition, the animals exhibit increased thyroid/body weight ratio, histopathological changes, and reduced serum protein-bound iodine²⁴⁶. Recent epidemiological data further underlines the importance of iodine status and the effect of ETU on thyroid function²⁴⁸. Another class of fungicides with thyroid disrupting activity are the azoles. This class of fungicide can enhance TH hepatic metabolism through the induction of enzymes uridine diphosphate-glucuronosyl-transferase (UDPGT), thereby increasing biliary elimination of T₃ and T₄²⁴⁹. Reduction of TH level varies considerably depending on the class of azoles²⁵⁰.

The organophosphate insecticide Malathion is an acetylcholinesterase inhibitor, similar to CPF, widely used in North American agriculture and residential landscapes, and in public health pest control programs such as residential mosquito eradication. It has also been characterised as an inhibitor of teleost TPO²⁵¹ and TTR^{252,253}. Other authors have documented increased levels of T₃ and reduced T₄ as a function of malathion exposure^{254,255}. As it is currently an approved insecticide in North America and Europe, it is important to

note that it has been associated with a strongly increased risk of thyroid cancer in spouses of pesticide applicators²⁵⁶.

Pyrethroids are synthetic organic insecticides similar to the natural pyrethrins produced by the flowers of chrysanthemums. Due to their high lipophilicity and persistence, they are prone to bioaccumulation. Further, toxicological studies have demonstrated their the potential to disrupt the endocrine system and exert developmental toxicity²⁵⁷. Permethrin (PM), one of the most heavily used synthetic pyrethroids, exerts estrogenic effect in zebrafish²⁵⁸ while other pyrethroids have been demonstrated as interfering with TRs²⁵⁹, TTR binding²⁶⁰ and Dio1 inhibition^{261,262}. Several rat studies also suggest that pyrethroid insecticides alter serum TH levels, mostly increasing total T₃ levels^{255,263–265}.

Finally, a pyridine-based juvenile hormone analogue pesticide, Pyriproxyfen, has been suggested as a thyroid hormone active substance²⁴⁴. This and other findings led us and colleagues to suspect it could implicated in the increased incidence of Zika induced microcephaly in north eastern Brazil²⁶⁶, especially given its use at high levels in drinking water during the outbreak²⁶⁶

Polychlorinated biphenyls (PCBs)

Polychlorinated biphenyls (PCBs) are a class of organic man-made chemicals that were mass produced globally since the 1920s, until their commercial production ban in the United States in 1979. They were widely used as plasticizers, in hydraulic fluids, heat transfer fluids, lubricants and electrical equipment like capacitors and transformers. A total of 209 possible congeners exists, classed according to the number and position of chlorine atoms carried. PCBs can also be metabolized by hydroxylation to OH-PCBs. Due to their high chemical stability PCBs do not readily break down and are still found in significant quantities

throughout the environment and human fluids²⁶⁷. PCBs and their metabolites are known to efficiently transfer from maternal to foetal blood via the placenta^{268,269} and to nursing children via milk^{270,271}. Prenatal PCB exposure in human has been associated with increased risk of a number of thyroid hormone related disorders including, high BMI²⁷², IQ loss^{273–275}, cognitive defects^{23,273,276–278}, reduced visual recognition memory²⁷⁴, attention and motor deficits^{276,279–281}, increased risk of autism^{282,283} and ADHD^{273,284,285}. In PCB exposed adults, an increased risk of cardiovascular disease has been reported²⁸⁶, as has a slightly increased risk of thyroid autoimmunity in men^{287,288}.

Due to their physiochemical properties, PCBs have long been suspected to act as TH and other steroidal hormone analogues^{2,289}. Numerous publications covering both epidemiological and experimental studies have confirmed the association of hydroxylated and non-hydroxylated PCBs with decreased TH levels, T₄^{290–292} and T₃^{292–295}. In fact serum hypothyroxinemia is the most frequently reported adverse health effect in human populations exposed to PCBs due to displacement of T₄ from TTR and subsequent increase of metabolism²⁹⁶. PCBs in cord blood have also been linked with low thyroid binding globulin (TBG)²⁹³ and high TSH levels^{297,298}. In contrast to the latter study, a recent analysis of three cohorts revealed slightly lower levels of TSH with PCB-153 exposure²⁹⁹.

In experimental studies, PCB and their metabolites demonstrate a clear association with reduced TH levels in animal models^{300–302} and induce long-term effects on behaviour and neurodevelopment^{303,304}. More recent studies have highlighted other possible mechanisms of PCB action on additional TH axis components. TTR disruption may play a role in distribution of hydroxylated PCBs to the placenta and the brain as PCB metabolites are known competitors for TTR's T₄ binding pocket^{296,305}. PCB exposure suppresses NIS expression³⁰¹ though the Akt/FoxO3a/NIS pathway^{306,307}. NIS suppression may also be due

to inflammation by PCB exposure. The PCB induced AhR/JNK pathway stimulates the production of cytokines and thereby suppresses NIS expression³⁰⁸. Hydroxylated PCBs may also inhibit SULT-catalysed THs sulfation³⁰⁹. A more recent study on infants found further an association between PCBs in maternal blood with high T₃ and low rT₃ in cord serum indicating possible action on deiodinases³¹⁰. Finally, iodine status can have a major impact on the effect of PCB exposure²³. A pilot study found that PCB exposure mitigates iodine supplementation during pregnancy in a borderline iodine deficient group and higher PCB levels have a negative impact on the neurocognitive development of the offspring.

The importance of other endocrine systems especially the sex hormones must not be overlooked as many sexually dimorphic changes due to gestational PCB exposure have been reported. Gestationally PCB-exposed females pups display increased birth weight, higher locomotor behaviours, higher corticosterone concentrations while the males display increased anogenital distances^{311,312}. Certain PCB metabolites have been shown to interact with the estrogen receptors acting either as agonists or antagonists^{313,314}. On the other hand, one PCB metabolite (PCB104) exhibits both, AR antagonistic and ER agonist properties³¹⁵. PCBs may also induce estrogenicity indirectly through inhibition of the estrogen sulfotransferase³¹⁶.

Polybrominated flame retardants

Polybrominated diphenyl ethers (PBDEs) are widely used flame retardants being used in furniture, carpets, automobiles, electrical appliances and flame-retardant fabrics. PBDEs are lipophilic in nature and as they are not chemically bound to the substrate, they easily accumulate in the environment. High levels are found in diverse situations from house dust³¹⁷ to river sediments³¹⁸. There are 209 possible congeners of PBDEs, due to the different possible bromine substitutions on the biphenyl backbone. Similar to PCBs, hydroxylated

PBDEs (OH-PBDEs) add to the complexity of chemical interactions and stability. Production and usage of the less brominated PBDEs were banned in Europe in 2004, and more recently extended to BDE-209 (or deca-BDE). As BDE-209, is the most highly brominated compound (10 bromines), it is easily broken down into less brominated congeners³¹⁸. However, general levels of PBDE are increasing despite the ban (see for instance:³¹⁹). In United States, despite similar restrictions and phase out of deca-PBDE at the end of 2013 (except for certain uses) the entire population have detectable levels of at least one PBDE congener in their blood³²⁰.

As PBDEs are persistent, PBDE congeners are still found in significant amounts in the placenta³²¹, foetal blood³²², and breast milk³²³. Early exposure of PBDEs has also been associated with decreased IQ, diminished language and reading abilities, increased problems with hyperactivity and attention, and poorer executive function in children^{324–332}. Among these, two studies further observed sex specific differences. Vuong et al reported significantly poorer executive function among boys with higher concurrent BDE-153 and no associations in girls, while Sagiv et al, observed poorer executive function in girls with higher 4 PBDE (BDE-47, -99, -100, -153) concentrations, but not in boys^{327,333}. These sex-specific differences need to be investigated further as other studies have revealed no statistically significant sex interactions^{328,329}.

The biological mechanism for sex differences in PBDE exposure-related neurotoxicity remains unknown. These behavioural changes are not surprising as PBDEs are well known to pass the blood–brain barrier, accumulate in the central nervous system and induce developmental neurotoxicity³³⁴. Neonatal rodents exposed to PBDEs exhibit behavioural changes^{335–338}, with reduced hippocampal long term potentiation, modified intracellular calcium homeostasis³³⁹, oxidative stress³⁴⁰ and reduced post- synaptic protein levels in the

hippocampus³⁴¹. Poorer attention and executive function suggests that PBDEs may also target the prefrontal cortex region of the brain^{342,343}.

The precise mechanism of PBDE action at a molecular level, still remains to be elucidated. Clearly, one plausible action of PBDEs is through its disruption of TH availability. Several epidemiological studies have reported increased TSH levels, lower total T₄ and, in some studies greater free T₃ levels in humans, including children^{295,344–349}. These TH parameters are not always consistent and the differences in findings may be due to the median levels of PBDEs as demonstrated by a meta-analysis³⁵⁰. High levels of PBDEs were positively correlated with TSH/T₄ levels while low PBDE levels were negatively associated. Decreased levels of T₄ are, however, the most consistently observed adverse effect observed in populations, as a function of PBDE exposure. Decreased circulating TH levels have also been observed in experimental studies on rodents, fish and birds after exposure to perinatal PBDE^{340,351–354}. Further, PBDE congeners have been tested for their agonist and antagonist properties against TRs. Using a reporter gene assay, PBDE congeners, including hydroxylated compounds, inhibited TR-mediated transcription at varying concentrations^{355–359}. The antagonist action of PBDEs on TR is further evident through its effect on Purkinje cell dendrite arborisation and neural progenitor cell differentiation into the oligodendrocyte lineage^{360,361}. In contrast, several hydroxylated PBDEs have been reported to act as agonists on TH-dependent transcription³⁵⁵. Recently, using zebrafish knock down model of TR β , the developmental toxicity of PBDE was demonstrated³⁶².

Several hydroxylated PBDE congeners have been shown to bind to and alter T₄ binding to the two TH distributor proteins, TBG and TTR^{363,364}. The displacement of T₄ from TTR and TBG, may lead to its increased glucuronidation, followed by decrease in circulating T₄ and hence higher TSH. PBDEs and hydroxylated metabolites alter DIO2 activity in

different astrocyte cell lines³⁶⁵. BDE-99 decreased DIO2 activity by up to 80% while 3-OH-BDE-47, 6-OH-BDE-47, and 5-OH-BDE-99 also decreased DIO2 activity by 45–80%. Multiple mechanisms appear to contribute to the decreased DIO2 activity, including weakened expression of *DIO2* mRNA, competitive inhibition of DIO2, and enhanced post-translational degradation of DIO2. As astrocytes produce more than 50 % of T₃ used by the brain Roberts et al. (2015) propose that PBDE exposure could compromise T₃ delivery to the brain³⁶⁵. A possible mechanism for this effect has been investigated *in silico* and thought to be halogen bonding of PBDEs to the active site selenocysteine³⁶⁶. Studies have also found upregulation of *Dio1* and *Dio3*, i.e. inducing local hyperthyroidism, in the periventricular zone of the brain, suggesting another, as yet under-estimated, mechanism impacting neurodevelopment.^{367,368}

Phthalates

Phthalates or phthalate esters are esters of phthalic acid and mainly used as plasticizers and softeners in various commercial products such as furniture, cosmetics, food packaging, and medical equipment such as catheters and perfusion bags/drips. Phthalates are also one of the most volatile EDCs and can be found at high concentrations in enclosed spaces where air conditioning is used such as vehicles³⁶⁹. One of the most widely-used phthalate is di-(2-ethylhexyl) phthalate (DEHP), used as a plasticizer in PVC recycling, but not manufacturing, due to its low cost. This phthalate used in soft medical devices and toys was of major concern due to its high migration rate³⁷⁰ and to the vulnerable population in contact with the compound. Consequently production was banned and manufacturing ceased by BASF in 2002. Other notable common phthalates are dibutyl phthalate (DBP) and its metabolite monobutyl phthalate (MBP), mono-(2-ethylhexyl) phthalate (MEHP), monoethyl phthalate (MEP), benzylbutylphthalate (BBP), diisodecyl phthalate (DIDP), and diisononyl

phthalate (DINP). Alternatives to phthalate such as hexamoll Dinch are now being used in medical devices. A transgenerational study carried out by BASF in 2005, unpublished but described in EFSA 2006³⁷¹ and NICNAS 2012³⁷² reports, showed significant hepatic effects and thyroid hyperplasia induced by Dinch³⁷²

Phthalates do not bio-accumulate in the environment but since they are ubiquitous in our daily life, the potential of consequences of continuous exposure has raised concerns. Current EU legislation focuses on levels in children's toys. Some phthalates (DEHP, BBP and DBP) cannot be used in toys and childcare products. Other phthalates (DINP and DIDP) are banned only from toys that could be potentially placed in mouth. Urinary levels of phthalates serve as a good biomarker and high exposure levels have been found in the general public, including pregnant women³⁷³⁻³⁷⁷.

In humans, DEHP and its metabolites measured in maternal urine have been associated with adverse neurodevelopment and behaviour in offspring³⁷⁸⁻³⁸². Interestingly, the phthalates metabolites, MEHHP, MEOHP and MBP were associated with both mental and behaviour defects in male but not females infants of 6 months³⁷⁸. Prenatal exposure to phthalates has also been associated with "reduced masculine play" among boys of 3- to 6-year³⁸³. In contrast, Téllez-Rojo et al. (2013) reported lower scores on the mental developmental index (Bayley's test) in females but not males at 2-3 years³⁸⁰. These sex specific differences seem to attenuate with age (7-9 age)³⁸⁴. More studies are needed to determine if gender differences are found as a function of phthalate exposure in older children. While these previous studies have focused on cognitive and behavioural defects during early years, other studies have found association between phthalates and more severe neurodevelopmental defects during school ages such a reduced IQ³⁸⁵ and attention deficit disorder (ADD)^{375, 386}.

As for a number of pesticides, these negative effects on brain development can be linked to changes in thyroid hormone levels. In pregnant women, the urinary MBP, MEP and MEHP have been associated with low serum T₄ and fT₄ in these women during the second trimester^{387–392}. These inverse relationships between MEHP and DEHP and T₄ levels have been reported in the adult population including men^{59,393}. In children, there are varying reports of levels of phthalates and its effect on TH levels. Some report inverse relations between the two^{388,394}, others a positive relation⁵⁹, and one that relied on a relatively small cohort found none at all³⁹⁵. This highlights the importance of study design, age group, sample size, and exposure profiles when studying non-persistent chemicals.

Since DBP down-regulates the human NIS promoter³⁹⁶, modulating the transcriptional activity of NIS may be one of the underlying causes of thyroid hyperactivity and decreased circulating T₄ concentrations. DBP also appears to act as a thyroid antagonist when assessed through reporter gene assays³⁹⁷. Zebrafish and male rats exposed to varying concentrations for MEHP and DEHP respectively demonstrate similar low levels of whole body T₄ levels^{398,399}. It is interesting to note that in 2017, ECHA classified DPB, DEHP, BBP and DIBP as substances of high concern due to their endocrine disrupting properties.

Perfluorinated compounds

Perfluorinated chemicals (PFCs) are widely used in the manufacture of fabrics, carpets, surfactants, emulsifiers, Teflon, lubricants, cosmetics, and fire-fighting foams. They are commonly used as surfactants due to their fully fluorinated linear carbon chain attached to a hydrophilic head. Surfactants are compounds that lower the surface tension between two liquids or between a liquid and a solid and therefore are popular in detergents, wetting agents, emulsifiers, foaming agents, and dispersants. They are also highly stable and therefore

extremely persistent in the environment including both wildlife and human populations ⁴⁰⁰. Between the years 2000 and 2002, the main producers discontinued production of certain PFCs, including perfluorohexane sulfonate (PFHxS), perfluorooctanoate (PFOA) and perfluorooctane sulfonate (PFOS). Following its discontinuation, a significant decrease in the serum levels of PFOA and PFOS were observed ⁴⁰¹⁻⁴⁰³. However, another factor that needs to be taken into consideration is their relatively long half-life in humans being approximately 3.8 years for PFOA, 5.4 years for PFOS, and 8.5 years for PFHxS ⁴⁰⁴. Despite the ban, they are still found in significant levels in drinking water. A 2016 study covering 2/3 of drinking water supplies in the United States found unsafe levels of PFCs at the minimum reporting levels required by the EPA ⁴⁰⁵. PFOS has been banned in the EU since 2008 but there are no restrictions on PFOA and PFHxS. In the USA, EPA launched a voluntary campaign in which the companies committed to reduce global facility emissions and product content of PFOA and related chemicals by 2015. The last update for this reduction was in 2013/2014.

Several animal studies have shown low dose exposure of PFCs during neonatal development results in irreversible neurotoxic effects and alterations in spontaneous behaviour, habituation capability, learning and memory (assessed at 4 months) ^{406,407}. PFCs were also shown to alter the levels of synaptophysin and tau proteins in cerebral cortex and the hippocampus. Both proteins are important for the formation and growth of dopaminergic synapses and alterations in the dopamine transporters and receptors are one of the underlying causes of behavioural defects such as ADHD ^{408,409}. Several cross-sectional studies have investigated the potential association between PFC levels in school-age children and ADHD ^{410 411}. Hoffman et al (2010) reported a positive association between levels of PFOS, PFOA and PFHxS with ADHD symptoms while Stein and Savitz (2011) reported an association with only PFHxS. High impulsivity has also been reported in children with high PFC levels ⁴¹² and only high levels of PFOS exposure during pregnancy has been associated with

delayed motor development in the first two years of life^{413,414}. TH dysfunction is a well-established risk for ADHD^{5,415-418}.

PFC exposure and TH disruption have also been reported in humans. A large study of employees in a PFC manufacturer revealed negative association between PFOA and free T₄ levels⁴¹⁹. In the US, women with high levels of PFOA and men with high levels of PFOS are also at increased risk of thyroid diseases⁴²⁰. Low levels of T₄ as a function of PFC exposure have also been confirmed in several animal models. A single dose of PFOS in adult rats resulted in an initial increased fT₄ and decreased TSH levels, followed by decrease in total T₄ and T₃ levels⁴²¹. In other adult rat studies, PFOA exposure resulted in decreased T₄ levels^{422,423}. Perinatal exposure to PFOS also results in decreased levels of T₄ in both the mother and the offspring⁴²⁴⁻⁴²⁷. A test of twenty-four PFCs revealed competitive binding of most PFCs to TTR⁴²⁸ which in turn can explain the dysfunctional levels of T₄ observed in humans and animal models. Of the 24 PFCs, PFHxS that the highest competitive binding followed by PFOA and PFOS equally. PFOS has also been shown to decrease hepatic *Dio1* mRNA while increasing thyroidal *Dio1* mRNA⁴²⁹. Whether this is a direct insult on *Dio1* transcription or a response to levels of T₄, is not yet clear.

Conclusion

The above review covers the main categories of chemicals that affect thyroid signalling. However, we have not reported environmental and human levels of exposure for each chemical class nor effects of mixtures. There are wide variations in exposure to individual chemicals exposure due to geographical location and legislation of the country of residence. Moreover, as we are exposed to multiple chemicals at a given time, it is increasingly important to address the effect of chemicals as a mixture, since synergistic

effects of chemical mixtures without individual effects have been reported ^{430,431}. Our laboratory has shown that exposure to mixtures of common chemicals found in human amniotic fluid, alter thyroid hormone signalling, brain structure and behaviour ^{432,433}. Together, these findings highlight the current impact of EDC exposure on neurodevelopment and argue for rapid public health intervention.

Funding

This work was supported by grants from Centre National de la Recherche Scientifique, Muséum National d'Histoire Naturelle, and European Union Contracts DevCom FP7-People-2013-ITN N°607142, H2020_EDC MIX RISK_GA N°634880 and H2020_HBM4EU_GA n° Contract No. 733032

References

1. Moog, N. K. *et al.* Influence of maternal thyroid hormones during gestation on fetal brain development. *Neuroscience* **342**, 68–100 (2017).
2. Gilbert, M. E., Rovet, J., Chen, Z. & Koibuchi, N. Developmental thyroid hormone disruption: Prevalence, environmental contaminants and neurodevelopmental consequences. *Neurotoxicology* **33**, 842–852 (2012).
3. Henrichs, J. *et al.* Maternal thyroid function during early pregnancy and cognitive functioning in early childhood: The generation R study. *J. Clin. Endocrinol. Metab.* **95**, 4227–4234 (2010).
4. Stagnaro-Green, A. & Pearce, E. Thyroid disorders in pregnancy. *Nat. Rev. Endocrinol.* **8**, 650–658 (2012).
5. Pääkkilä, F. *et al.* The impact of gestational thyroid hormone concentrations on ADHD symptoms of the child. *J. Clin. Endocrinol. Metab.* **99**, E1-8 (2014).
6. Korevaar, T. I. M. *et al.* Association of maternal thyroid function during early pregnancy with offspring IQ and brain morphology in childhood: a population-based prospective cohort study. *Lancet Diabetes Endocrinol.* **4**, 35–43 (2016).
7. Fetene, D. M., Betts, K. S. & Alati, R. Mechanisms in endocrinology maternal thyroid dysfunction during pregnancy and behavioural and psychiatric disorders of children: A systematic review. *Eur. J. Endocrinol.* **177**, R261–R273 (2017).

8. Gore, A. C. *et al.* Executive Summary to EDC-2: The Endocrine Society's second Scientific Statement on endocrine-disrupting chemicals. *Endocrine Reviews* **36**, 593–602 (2015).
9. Brucker-Davis, F. Effects of environmental synthetic chemicals on thyroid function. *Thyroid* **8**, 827–56 (1998).
10. Zoeller, R. T. & Crofton, K. M. Thyroid hormone action in fetal brain development and potential for disruption by environmental chemicals. *Neurotoxicology* **21**, 935–45 (2000).
11. Crofton, K. M. Thyroid disrupting chemicals: Mechanisms and mixtures. *Int. J. Androl.* **31**, 209–222 (2008).
12. Boas, M., Main, K. M. & Feldt-Rasmussen, U. Environmental chemicals and thyroid function: an update. *Curr. Opin. Endocrinol. Diabetes Obes.* **16**, 385–391 (2009).
13. Dohán, O. *et al.* The Na⁺/I symporter (NIS) mediates electroneutral active transport of the environmental pollutant perchlorate. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 20250–20255 (2007).
14. Tonacchera, M. *et al.* Relative Potencies and Additivity of Perchlorate, Thiocyanate, Nitrate, and Iodide on the Inhibition of Radioactive Iodide Uptake by the Human Sodium Iodide Symporter. *Thyroid* **14**, 1012–1019 (2004).
15. Demeneix, B. *Losing Our Minds: How Environmental Pollution Impairs Human Intelligence and Mental Health.* (Oxford University Press, 2014). doi:10.1093/acprof:oso/9780199917518.001.0001
16. Suh, M., Abraham, L., Hixon, J. G. & Proctor, D. M. The effects of perchlorate, nitrate, and thiocyanate on free thyroxine for potentially sensitive subpopulations o. The 2001-2002 and 2007-2008 National Health and Nutrition Examination Surveys. *J. Expo. Sci. Environ. Epidemiol.* **24**, 579–587 (2014).
17. Zoeller, T. R. Environmental chemicals targeting thyroid. *Hormones* **9**, 28–40 (2010).
18. Blount, B. C., Valentin-Blasini, L., Osterloh, J. D., Mauldin, J. P. & Pirkle, J. L. Perchlorate exposure of the US population, 2001-2002. in *Journal of Exposure Science and Environmental Epidemiology* **17**, 400–407 (2007).
19. Taylor, P. N. *et al.* Maternal perchlorate levels in women with borderline thyroid function during pregnancy and the cognitive development of their offspring: Data from the controlled antenatal thyroid

- study. *J. Clin. Endocrinol. Metab.* **99**, 4291–4298 (2014).
20. Steinmaus, C. *et al.* Thyroid hormones and moderate exposure to perchlorate during pregnancy in women in southern California. *Environ. Health Perspect.* **124**, 861–867 (2016).
 21. Rayman, M. P. & Bath, S. C. The new emergence of iodine deficiency in the UK: consequences for child neurodevelopment. *Ann. Clin. Biochem.* **52**, 705–708 (2015).
 22. Andersen, Laurberg, P., Wu, C. S. & Olsen, J. Attention deficit hyperactivity disorder and autism spectrum disorder in children born to mothers with thyroid dysfunction: a Danish nationwide cohort study. *BJOG* **121**, 1365–74 (2014).
 23. Brucker-Davis, F. *et al.* Neurotoxicant exposure during pregnancy is a confounder for assessment of iodine supplementation on neurodevelopment outcome. *Neurotoxicol. Teratol.* **51**, 45–51 (2015).
 24. Provencher, G. *et al.* Determination of bisphenol A, triclosan and their metabolites in human urine using isotope-dilution liquid chromatography–tandem mass spectrometry. *J. Chromatogr. A* **1348**, 97–104 (2014).
 25. Tato, T., Salgueiro-González, N., León, V. M., González, S. & Beiras, R. Ecotoxicological evaluation of the risk posed by bisphenol A, triclosan, and 4-nonylphenol in coastal waters using early life stages of marine organisms (*Isochrysis galbana*, *Mytilus galloprovincialis*, *Paracentrotus lividus*, and *Acartia clausi*). *Environmental Pollution* (2017). doi:10.1016/j.envpol.2017.09.031
 26. Dodds, E. C. & Lawson, W. Synthetic estrogenic agents without the phenanthrene nucleus. *Nature* **137**, 996–996 (1936).
 27. EFSA. Scientific Opinion on the risks to public health related to the presence of bisphenol A (BPA) in foodstuffs. *EFSA J.* **13**, 3978 (2015).
 28. Andra, S. S., Charisiadis, P., Arora, M., Van Vliet-Ostapchouk, J. V. & Makris, K. C. Biomonitoring of human exposures to chlorinated derivatives and structural analogs of bisphenol A. *Environment International* **85**, 352–379 (2015).
 29. Lee, J. *et al.* Bisphenol A distribution in serum, urine, placenta, breast milk, and umbilical cord serum in a birth panel of mother–neonate pairs. *Science of the Total Environment* (2017).

doi:10.1016/j.scitotenv.2017.10.042

30. Cantonwine, D. E. *et al.* Urinary concentrations of bisphenol A and phthalate metabolites measured during pregnancy and risk of preeclampsia. *Environ. Health Perspect.* **124**, 1651–1655 (2016).
31. Troisi, J. *et al.* Placental concentrations of bisphenol A and birth weight from births in the Southeastern U.S. *Placenta* **35**, 947–952 (2014).
32. Vela-Soria, F. *et al.* A multiclass method for endocrine disrupting chemical residue analysis in human placental tissue samples by UHPLC–MS/MS. *Anal. Methods* **3**, 2073 (2011).
33. Mendonca, K., Hauser, R., Calafat, A. M., Arbuckle, T. E. & Duty, S. M. Bisphenol A concentrations in maternal breast milk and infant urine. *Int. Arch. Occup. Environ. Health* **87**, 13–20 (2014).
34. Deceuninck, Y. *et al.* Determination of bisphenol A and related substitutes/analogues in human breast milk using gas chromatography-tandem mass spectrometry. *Anal. Bioanal. Chem.* **407**, 2485–2497 (2015).
35. Ye, X. *et al.* Urinary Concentrations of Bisphenol A and Three Other Bisphenols in Convenience Samples of U.S. Adults during 2000-2014. *Environ. Sci. Technol.* **49**, 11834–11839 (2015).
36. Yang, Y., Guan, J., Yin, J., Shao, B. & Li, H. Urinary levels of bisphenol analogues in residents living near a manufacturing plant in south China. *Chemosphere* **112**, 481–486 (2014).
37. Yang, Y. J., Li, Z. L., Zhang, J., Yang, Y. & Shao, B. Simultaneous determination of bisphenol A, bisphenol AF, tetrachlorobisphenol A, and tetrabromobisphenol A concentrations in water using on-line solid-phase extraction with ultrahigh-pressure liquid chromatography tandem mass spectrometry. *Int. J. Environ. Anal. Chem.* **94**, 16–27 (2014).
38. Thayer, K. A. *et al.* Bisphenol a, bisphenol s, and 4-hydro xyphenyl 4-isopro oxyphenyl sulfone (bpsip) in urine and blood of cashiers. *Environ. Health Perspect.* **124**, 437–444 (2016).
39. Zhang, Y.-F. *et al.* Bisphenol A alternatives bisphenol S and bisphenol F interfere with thyroid hormone signaling pathway in vitro and in vivo. *Environ. Pollut.* (2017). doi:10.1016/j.envpol.2017.11.027
40. Lee, S., Kim, C., Youn, H. & Choi, K. Thyroid hormone disrupting potentials of bisphenol A and its analogues - in vitro comparison study employing rat pituitary (GH3) and thyroid follicular (FRTL-5)

- cells. *Toxicol. Vitro.* **40**, 297–304 (2017).
41. Le Fol, V. *et al.* In vitro and in vivo estrogenic activity of BPA, BPF and BPS in zebrafish-specific assays. *Ecotoxicol. Environ. Saf.* **142**, 150–156 (2017).
 42. MacKay, H. & Abizaid, A. A plurality of molecular targets: The receptor ecosystem for bisphenol-A (BPA). *Hormones and Behavior* (2017). doi:10.1016/j.yhbeh.2017.11.001
 43. Okada, H. *et al.* Direct evidence revealing structural elements essential for the high binding ability of bisphenol a to human estrogen-related receptor-?? *Environ. Health Perspect.* **116**, 32–38 (2008).
 44. Matsushima, A. *et al.* Structural evidence for endocrine disruptor bisphenol A binding to human nuclear receptor ERR γ . *J. Biochem.* **142**, 517–524 (2007).
 45. Xu, L. C. *et al.* Evaluation of androgen receptor transcriptional activities of bisphenol A, octylphenol and nonylphenol in vitro. *Toxicology* **216**, 197–203 (2005).
 46. Teng, C. *et al.* Bisphenol A affects androgen receptor function via multiple mechanisms. *Chem. Biol. Interact.* **203**, 556–564 (2013).
 47. Zoeller, R. T., Bansal, R. & Parris, C. Bisphenol-A, an environmental contaminant that acts as a thyroid hormone receptor antagonist in vitro, increases serum thyroxine, and alters RC3/neurogranin expression in the developing rat brain. *Endocrinology* **146**, 607–12 (2005).
 48. Moriyama, K. *et al.* Thyroid hormone action is disrupted by bisphenol A as an antagonist. *J. Clin. Endocrinol. Metab.* **87**, 5185–5190 (2002).
 49. Alonso-Magdalena, P. *et al.* Bisphenol-A acts as a potent estrogen via non-classical estrogen triggered pathways. *Molecular and Cellular Endocrinology* **355**, 201–207 (2012).
 50. Kitamura, S., Jinno, N., Ohta, S., Kuroki, H. & Fujimoto, N. Thyroid hormonal activity of the flame retardants tetrabromobisphenol A and tetrachlorobisphenol A. *Biochem. Biophys. Res. Commun.* **293**, 554–559 (2002).
 51. Ikezuki, Y., Tsutsumi, O., Takai, Y., Kamei, Y. & Taketani, Y. Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum. Reprod.* **17**, 2839–2841 (2002).

52. Kitamura, S. *et al.* Comparative study of the endocrine-disrupting activity of bisphenol A and 19 related compounds. *Toxicol. Sci.* **84**, 249–259 (2005).
53. Freitas, J. *et al.* Detection of thyroid hormone receptor disruptors by a novel stable in vitro reporter gene assay. *Toxicol. Vitr.* **25**, 257–266 (2011).
54. Hofmann, P. J., Schomburg, L. & Köhrle, J. Interference of endocrine disruptors with thyroid hormone receptor-dependent transactivation. *Toxicol. Sci.* **110**, 125–137 (2009).
55. Goldinger, D. M. *et al.* Endocrine activity of alternatives to BPA found in thermal paper in Switzerland. *Regul. Toxicol. Pharmacol.* **71**, 453–462 (2015).
56. Geens, T. *et al.* Daily intake of bisphenol A and triclosan and their association with anthropometric data, thyroid hormones and weight loss in overweight and obese individuals. *Environ. Int.* **76**, 98–105 (2015).
57. Sriphrapadang, C., Chailurkit, L. O., Aekplakorn, W. & Ongphiphadhanakul, B. Association between bisphenol A and abnormal free thyroxine level in men. *Endocrine* **44**, 441–447 (2013).
58. Wang, F. *et al.* High urinary bisphenol A concentrations in workers and possible laboratory abnormalities. *Occup. Environ. Med.* **69**, 679–84 (2012).
59. Meeker, J. D. & Ferguson, K. K. Relationship between urinary phthalate and bisphenol a concentrations and serum thyroid measures in u.s. adults and adolescents from the national health and nutrition examination survey (NHANES) 2007-2008. *Environ. Health Perspect.* **119**, 1396–1402 (2011).
60. Aung, M. T. *et al.* Thyroid hormone parameters during pregnancy in relation to urinary bisphenol A concentrations: A repeated measures study. *Environ. Int.* **104**, 33–40 (2017).
61. Romano, M. E. *et al.* Gestational urinary bisphenol A and maternal and newborn thyroid hormone concentrations: The HOME Study. *Environ. Res.* **138**, 453–460 (2015).
62. Chevrier, J. *et al.* Maternal urinary bisphenol a during pregnancy and maternal and neonatal thyroid function in the CHAMACOS study. *Environ. Health Perspect.* **121**, 138–144 (2013).
63. Minatoya, M. *et al.* Cord Blood Bisphenol A Levels and Reproductive and Thyroid Hormone Levels of Neonates. *Epidemiology* **28**, S3–S9 (2017).
64. Andrianou, X. D. *et al.* Human exposures to Bisphenol A, Bisphenol F and Chlorinated Bisphenol a

- derivatives and thyroid function. *PLoS One* **11**, (2016).
65. Casas, M. *et al.* Exposure to bisphenol A during pregnancy and child neuropsychological development in the INMA-Sabadell cohort. *Environ. Res.* **142**, 671–679 (2015).
 66. Roen, E. L. *et al.* Bisphenol A exposure and behavioral problems among inner city children at 7-9 years of age. *Environ. Res.* **142**, 739–745 (2015).
 67. Harley, K. G. *et al.* Prenatal and early childhood bisphenol A concentrations and behavior in school-aged children. *Environ. Res.* **126**, 43–50 (2013).
 68. Braun, J. M. *et al.* Impact of Early-Life Bisphenol A Exposure on Behavior and Executive Function in Children. *Pediatrics* **128**, 873–882 (2011).
 69. Braun, J. M. *et al.* Prenatal bisphenol A exposure and early childhood behavior. *Environ. Health Perspect.* **117**, 1945–1952 (2009).
 70. Evans, S. F. *et al.* Prenatal bisphenol A exposure and maternally reported behavior in boys and girls. *Neurotoxicology* **45**, 91–99 (2014).
 71. Tewar, S. *et al.* Association of Bisphenol A exposure and Attention-Deficit/Hyperactivity Disorder in a national sample of U.S. children. *Environ. Res.* **150**, 112–118 (2016).
 72. Perera, F. *et al.* Prenatal bisphenol a exposure and child behavior in an innercity cohort. *Environ. Health Perspect.* **120**, 1190–1194 (2012).
 73. Arbuckle, T. E., Davis, K., Boylan, K., Fisher, M. & Fu, J. Bisphenol A, phthalates and lead and learning and behavioral problems in Canadian children 6-11 years of age: CHMS 2007-2009. *Neurotoxicology* **54**, 89–98 (2016).
 74. Diaz Weinstein, S., Villafane, J. J., Juliano, N. & Bowman, R. E. Adolescent exposure to Bisphenol-A increases anxiety and sucrose preference but impairs spatial memory in rats independent of sex. *Brain Res.* **1529**, 56–65 (2013).
 75. Bowman, R. E. *et al.* Bisphenol-A exposure during adolescence leads to enduring alterations in cognition and dendritic spine density in adult male and female rats. *Horm. Behav.* **69**, 89–97 (2015).
 76. Xu, X. *et al.* Perinatal bisphenol A affects the behavior and SRC-1 expression of male pups but does not

- influence on the thyroid hormone receptors and its responsive gene. *Neurosci. Res.* **58**, 149–155 (2007).
77. Mizuo, K. *et al.* Prenatal and neonatal exposure to bisphenol-A affects the morphine-induced rewarding effect and hyperlocomotion in mice. *Neurosci. Lett.* **356**, 95–98 (2004).
 78. Kundakovic, M. *et al.* Sex-specific epigenetic disruption and behavioral changes following low-dose in utero bisphenol A exposure. *Proc. Natl. Acad. Sci.* **110**, 9956–9961 (2013).
 79. Adriani, W., Della Seta, D., Dessi-Fulgheri, F., Farabollini, F. & Laviola, G. Altered profiles of spontaneous novelty seeking, impulsive behavior, and response to D-amphetamine in rats perinatally exposed to Bisphenol A. *Environmental Health Perspectives* **111**, 395–401 (2003).
 80. Kumar, D. & Kumar Thakur, M. Perinatal exposure to bisphenol-A impairs spatial memory through upregulation of neurexin1 and neuroligin3 expression in male mouse brain. *PLoS One* **9**, (2014).
 81. Jašarević, E. *et al.* Sex and dose-dependent effects of developmental exposure to bisphenol A on anxiety and spatial learning in deer mice (*Peromyscus maniculatus bairdii*) offspring. *Horm. Behav.* **63**, 180–189 (2013).
 82. Kuwahara, R., Kawaguchi, S., Kohara, Y., Cui, H. & Yamashita, K. Perinatal Exposure to Low-Dose Bisphenol A Impairs Spatial Learning and Memory in Male Rats. *J. Pharmacol. Sci.* **123**, 132–139 (2013).
 83. Xu, X., Tian, D., Hong, X., Chen, L. & Xie, L. Sex-specific influence of exposure to bisphenol-A between adolescence and young adulthood on mouse behaviors. *Neuropharmacology* **61**, 565–573 (2011).
 84. Poimenova, A., Markaki, E., Rahiotis, C. & Kitraki, E. Corticosterone-regulated actions in the rat brain are affected by perinatal exposure to low dose of bisphenol A. *Neuroscience* **167**, 741–749 (2010).
 85. Tian, Y.-H., Baek, J.-H., Lee, S.-Y. & Jang, C.-G. Prenatal and postnatal exposure to bisphenol a induces anxiolytic behaviors and cognitive deficits in mice. *Synapse* **64**, 432–439 (2010).
 86. Wolstenholme, J. T., Rissman, E. F. & Connelly, J. J. The role of Bisphenol A in shaping the brain, epigenome and behavior. *Hormones and Behavior* **59**, 296–305 (2011).
 87. Jorgensen, E. M., Alderman, M. H. & Taylor, H. S. Preferential epigenetic programming of estrogen

- response after in utero xenoestrogen (bisphenol-A) exposure. *FASEB J.* **30**, 3194–3201 (2016).
88. Kyono, Y. *et al.* Liganded thyroid hormone receptors transactivate the DNA methyltransferase 3a gene in mouse neuronal cells. *Endocrinology* **157**, 3647–3657 (2016).
89. Kyono, Y., Sachs, L. M., Bilesimo, P., Wen, L. & Denver, R. J. Developmental and thyroid hormone regulation of the DNA methyltransferase 3a gene in xenopus tadpoles. *Endocrinology* **157**, 4961–4972 (2016).
90. Canada. Screening Assessment Report. (2013).
91. Canada. Screening Assessment Report. (2012).
92. EFSA. Scientific Opinion on Tetrabromobisphenol A (TBBPA) and its derivatives in food. *EFSA J.* **9**, 2477 (2011).
93. Van der Ven, L. T. M. *et al.* Endocrine effects of tetrabromobisphenol-A (TBBPA) in Wistar rats as tested in a one-generation reproduction study and a subacute toxicity study. *Toxicology* **245**, 76–89 (2008).
94. Zieminska, E., Ruszczynska, A. & Lazarewicz, J. W. Tetrabromobisphenol A disturbs zinc homeostasis in cultured cerebellar granule cells: A dual role in neurotoxicity. *Food Chem. Toxicol.* **109**, 363–375 (2017).
95. Zieminska, E., Stafiej, A., Toczyłowska, B., Albrecht, J. & Lazarewicz, J. W. Role of Ryanodine and NMDA Receptors in Tetrabromobisphenol A-Induced Calcium Imbalance and Cytotoxicity in Primary Cultures of Rat Cerebellar Granule Cells. *Neurotox. Res.* **28**, 195–208 (2015).
96. Zieminska, E., Stafiej, A., Toczyłowska, B. & Lazarewicz, J. W. Bastadin 12 and ryanodine reveal similarities between thapsigargin- and tetrabromobisphenol A-induced intracellular Ca²⁺ release in cultured cerebellar granule cells. *J. Physiol. Pharmacol.* **65**, 679–686 (2014).
97. Zieminska, E., Lenart, J., Diamandakis, D. & Lazarewicz, J. W. The Role of Ca²⁺ Imbalance in the Induction of Acute Oxidative Stress and Cytotoxicity in Cultured Rat Cerebellar Granule Cells Challenged with Tetrabromobisphenol A. *Neurochem. Res.* **42**, 1–11 (2016).
98. Hendriks, H. S., Van kleef, R. G. D. M., Van den berg, M. & Westerink, R. H. S. Multiple novel modes

- of action involved in the in vitro neurotoxic effects of tetrabromobisphenol-A. *Toxicol. Sci.* **128**, 235–246 (2012).
99. Cope, R. B., Kacew, S. & Dourson, M. A reproductive, developmental and neurobehavioral study following oral exposure of tetrabromobisphenol A on Sprague-Dawley rats. *Toxicology* **329**, 49–59 (2015).
100. Colnot, T., Kacew, S. & Dekant, W. Mammalian toxicology and human exposures to the flame retardant 2,2', 6,6' -tetrabromo-4,4' -isopropylidenediphenol (TBBPA): Implications for risk assessment. *Archives of Toxicology* **88**, 553–573 (2014).
101. Viberg, H. & Eriksson, P. Differences in neonatal neurotoxicity of brominated flame retardants, PBDE 99 and TBBPA, in mice. *Toxicology* **289**, 59–65 (2011).
102. Decherf, S., Seugnet, I., Fini, J. B., Clerget-Froidevaux, M. S. & Demeneix, B. A. Disruption of thyroid hormone-dependent hypothalamic set-points by environmental contaminants. *Mol. Cell. Endocrinol.* **323**, 172–182 (2010).
103. Chen, J. *et al.* TBBPA exposure during a sensitive developmental window produces neurobehavioral changes in larval zebrafish. *Environ. Pollut.* **216**, 53–63 (2016).
104. Nakajima, A. *et al.* Neurobehavioral effects of tetrabromobisphenol A, a brominated flame retardant, in mice. *Toxicol. Lett.* **189**, 78–83 (2009).
105. Lilienthal, H., Verwer, C. M., van der Ven, L. T. M., Piersma, A. H. & Vos, J. G. Exposure to tetrabromobisphenol A (TBBPA) in Wistar rats: Neurobehavioral effects in offspring from a one-generation reproduction study. *Toxicology* **246**, 45–54 (2008).
106. Ng, L., Kelley, M. W. & Forrest, D. Making sense with thyroid hormone-the role of T 3 in auditory development. *Nature Reviews Endocrinology* **9**, 296–307 (2013).
107. NTP-Report. Technical Report on the toxicology studies of tetrabromobisphenol A (CAS NO. 79-94-7) in F344/NTac rats and B6C3F1/N mice and toxicology and carcinogenesis study of tetrabromobisphenol A in Wistar Han [CrI:WI(Han)] rats and B6C3F1/N mice (gavage studies). (2014).
108. Lai, D. Y., Kacew, S. & Dekant, W. Tetrabromobisphenol A (TBBPA): Possible modes of action of

- toxicity and carcinogenicity in rodents. *Food and Chemical Toxicology* **80**, 206–214 (2015).
109. Hamers, T. *et al.* In vitro profiling of the endocrine-disrupting potency of brominated flame retardants. *Toxicol. Sci.* **92**, 157–173 (2006).
110. Lévy-Bimbot, M., Major, G., Courilleau, D., Blondeau, J. P. & Lévi, Y. Tetrabromobisphenol-A disrupts thyroid hormone receptor alpha function in vitro: Use of fluorescence polarization to assay corepressor and coactivator peptide binding. *Chemosphere* **87**, 782–788 (2012).
111. Kitamura, S. *et al.* Anti-thyroid hormonal activity of tetrabromobisphenol A, a flame retardant, and related compounds: Affinity to the mammalian thyroid hormone receptor, and effect on tadpole metamorphosis. *Life Sci.* **76**, 1589–1601 (2005).
112. Zhang, Y. F. *et al.* Tetrabromobisphenol a disrupts vertebrate development via thyroid hormone signaling pathway in a developmental stage-dependent manner. *Environ. Sci. Technol.* **48**, 8227–8234 (2014).
113. Wang, Y., Li, Y., Qin, Z. & Wei, W. Re-evaluation of thyroid hormone signaling antagonism of tetrabromobisphenol A for validating the T3-induced *Xenopus* metamorphosis assay. *J. Environ. Sci. (China)* **52**, 325–332 (2017).
114. Fini, J. B. *et al.* Thyroid hormone signaling in the *Xenopus laevis* embryo is functional and susceptible to endocrine disruption. *Endocrinology* **153**, 5068–81 (2012).
115. Mengeling, B. J. *et al.* A multi-tiered, in vivo, quantitative assay suite for environmental disruptors of thyroid hormone signaling. *Aquat. Toxicol.* **190**, 1–10 (2017).
116. Borghoff, S. J., Wikoff, D., Harvey, S. & Haws, L. Dose- and time-dependent changes in tissue levels of tetrabromobisphenol A (TBBPA) and its sulfate and glucuronide conjugates following repeated administration to female Wistar Han Rats. *Toxicol. Reports* **3**, 190–201 (2016).
117. Yueh, M.-F. & Tukey, R. H. Triclosan: A Widespread Environmental Toxicant with Many Biological Effects. *Annu. Rev. Pharmacol. Toxicol.* **56**, 251–272 (2016).
118. Calafat, A. M., Ye, X., Wong, L. Y., Reidy, J. A. & Needham, L. L. Urinary concentrations of triclosan in the U.S. population: 2003-2004. *Environ. Health Perspect.* **116**, 303–307 (2008).

119. Chapin, R. E. *et al.* NTP-CERHR expert panel report on the reproductive and developmental toxicity of bisphenol A. *Birth Defects Res. Part B - Dev. Reprod. Toxicol.* **83**, 157–395 (2008).
120. Dayan, A. D. Risk assessment of triclosan [Irgasan®] in human breast milk. *Food Chem. Toxicol.* **45**, 125–129 (2007).
121. Arbuckle, T. E. *et al.* Exposure to free and conjugated forms of bisphenol a and triclosan among pregnant women in the MIREC cohort. *Environ. Health Perspect.* **123**, 277–284 (2015).
122. Philippat, C. *et al.* Prenatal exposure to environmental phenols: Concentrations in amniotic fluid and variability in urinary concentrations during pregnancy. *Environ. Health Perspect.* **121**, 1225–1231 (2013).
123. EU_Commission. Regulation (EU) No 358/2014 - of the European Parliament and of the Council on cosmetic products. (2014).
124. Chen, L. *et al.* Accumulation and risk of triclosan in surface sediments near the outfalls of municipal wastewater treatment plants. *Bull. Environ. Contam. Toxicol.* **95**, 525–529 (2015).
125. Diaz-Cruz, M. S. & Barceló, D. *Personal Care Products in the Aquatic Environment. The Handbook of Environmental Chemistry* **36**, (Springer International Publishing, 2015).
126. Axelstad, M., Boberg, J., Vinggaard, A. M., Christiansen, S. & Hass, U. Triclosan exposure reduces thyroxine levels in pregnant and lactating rat dams and in directly exposed offspring. *Food Chem. Toxicol.* **59**, 534–540 (2013).
127. Paul, K. B. *et al.* Developmental triclosan exposure decreases maternal, fetal, and early neonatal thyroxine: A dynamic and kinetic evaluation of a putative mode-of-action. *Toxicology* **300**, 31–45 (2012).
128. Rodríguez, P. E. a & Sanchez, M. S. Maternal exposure to triclosan impairs thyroid homeostasis and female pubertal development in Wistar rat offspring. *J. Toxicol. Environ. Health. A* **73**, 1678–1688 (2010).
129. Crofton, K. M., Paul, K. B., DeVito, M. J. & Hedge, J. M. Short-term in vivo exposure to the water contaminant triclosan: Evidence for disruption of thyroxine. *Environ. Toxicol. Pharmacol.* **24**, 194–197

- (2007).
130. Paul, K. B., Thompson, J. T., Simmons, S. O., Vanden Heuvel, J. P. & Crofton, K. M. Evidence for triclosan-induced activation of human and rodent xenobiotic nuclear receptors. *Toxicol. Vitro*. **27**, 2049–2060 (2013).
 131. Zorrilla, L. M. *et al.* The effects of triclosan on puberty and thyroid hormones in male wistar rats. *Toxicol. Sci.* **107**, 56–64 (2009).
 132. Veldhoen, N. *et al.* Corrigendum to ‘The bactericidal agent triclosan modulates thyroid hormone-associated gene expression and disrupts postembryonic anuran development’ [Aquat. Toxicol. 80 (2006) 217–227] (DOI:10.1016/j.aquatox.2006.08.010). *Aquat. Toxicol.* **83**, 84 (2007).
 133. Fort, D. J. *et al.* Triclosan and thyroid-mediated metamorphosis in anurans: Differentiating growth effects from thyroid-driven metamorphosis in *Xenopus laevis*. *Toxicol. Sci.* **121**, 292–302 (2011).
 134. Regnault, C. *et al.* Metabolic and immune impairments induced by the endocrine disruptors benzo[a]pyrene and triclosan in *Xenopus tropicalis*. *Chemosphere* **155**, 519–527 (2016).
 135. Lan, Z., Kim, T. H., Bi, K. S., Chen, X. H. & Kim, H. S. Triclosan exhibits a tendency to accumulate in the epididymis and shows sperm toxicity in male sprague-dawley rats. *Environ. Toxicol.* **30**, 83–91 (2015).
 136. Stoker, T. E., Gibson, E. K. & Zorrilla, L. M. Triclosan exposure modulates estrogen-dependent responses in the female wistar rat. *Toxicol. Sci.* **117**, 45–53 (2010).
 137. Huang, H. *et al.* The in Vitro estrogenic activities of triclosan and triclocarban. *J. Appl. Toxicol.* **34**, 1060–1067 (2014).
 138. Henry, N. D. & Fair, P. A. Comparison of in vitro cytotoxicity, estrogenicity and anti-estrogenicity of triclosan, perfluorooctane sulfonate and perfluorooctanoic acid. *J. Appl. Toxicol.* **33**, 265–272 (2013).
 139. Gee, R. H., Charles, A., Taylor, N. & Darbre, P. D. Oestrogenic and androgenic activity of triclosan in breast cancer cells. *J. Appl. Toxicol.* **28**, 78–91 (2008).
 140. Kim, J. Y. *et al.* Methoxychlor and triclosan stimulates ovarian cancer growth by regulating cell cycle- and apoptosis-related genes via an estrogen receptor-dependent pathway. *Environ. Toxicol. Pharmacol.*

- 37, 1264–1274 (2014).
141. Lee, H. R., Hwang, K. A., Nam, K. H., Kim, H. C. & Choi, K. C. Progression of breast cancer cells was enhanced by endocrine-disrupting chemicals, triclosan and octylphenol, via an estrogen receptor-dependent signaling pathway in cellular and mouse xenograft models. *Chem. Res. Toxicol.* **27**, 834–842 (2014).
 142. Louis, G. W., Hallinger, D. R. & Stoker, T. E. The effect of triclosan on the uterotrophic response to extended doses of ethinyl estradiol in the weanling rat. *Reprod. Toxicol.* **36**, 71–77 (2013).
 143. Pollock, T., Greville, L. J., Tang, B. & deCatanzaro, D. Triclosan elevates estradiol levels in serum and tissues of cycling and peri-implantation female mice. *Reprod. Toxicol.* **65**, 394–401 (2016).
 144. James, M. O., Li, W., Summerlot, D. P., Rowland-Faux, L. & Wood, C. E. Triclosan is a potent inhibitor of estradiol and estrone sulfonation in sheep placenta. *Environ. Int.* **36**, 942–949 (2010).
 145. Wang, L. Q., Falany, C. N. & James, M. O. Triclosan as a substrate and inhibitor of 3' -phosphoadenosine 5' -phosphosulfate-sulfotransferase and UDP-glucuronosyl transferase in human liver fractions. *Drug Metab. Dispos.* **32**, 1162–1169 (2004).
 146. Aker, A. M. *et al.* Phenols and parabens in relation to reproductive and thyroid hormones in pregnant women. *Environ. Res.* **151**, 30–37 (2016).
 147. Allmyr, M., Panagiotidis, G., Sparve, E., Diczfalusy, U. & Sandborgh-Englund, G. Human exposure to triclosan via toothpaste does not change cyp3a4 activity or plasma concentrations of thyroid hormones. *Basic Clin. Pharmacol. Toxicol.* **105**, 339–344 (2009).
 148. Cullinan, M. P., Palmer, J. E., Carle, A. D., West, M. J. & Seymour, G. J. Long term use of triclosan toothpaste and thyroid function. *Sci. Total Environ.* **416**, 75–79 (2012).
 149. Koeppe, E. S., Ferguson, K. K., Colacino, J. A. & Meeker, J. D. Relationship between urinary triclosan and paraben concentrations and serum thyroid measures in NHANES 2007-2008. *Sci. Total Environ.* **445–446**, 299–305 (2013).
 150. Poole, A. C. *et al.* Crossover Control Study of the Effect of Personal Care Products Containing Triclosan on the Microbiome. *mSphere* **1**, e00056-15 (2016).

151. Wang, X. *et al.* Maternal Urinary Triclosan Concentration in Relation to Maternal and Neonatal Thyroid Hormone Levels : A Prospective Study. *Environ. Health Perspect.* **125**, 67017 (2013).
152. Lassen, T. H. *et al.* Prenatal triclosan exposure and anthropometric measures including anogenital distance in Danish infants. *Environ. Health Perspect.* **124**, 1261–1268 (2016).
153. Johannes, J. *et al.* Silychristin, a Flavonolignan Derived From the Milk Thistle, Is a Potent Inhibitor of the Thyroid Hormone Transporter MCT8. *Endocrinology* **157**, 1694–701 (2016).
154. Hossard, L. *et al.* Effects of halving pesticide use on wheat production. *Sci. Rep.* **4**, (2014).
155. EFSA. Scientific Opinion on the identification of pesticides to be included in cumulative assessment groups on the basis of their toxicological profile. *EFSA J.* **11**, 3293 (2013).
156. Carson, R. Silent Spring. *Foreign Aff.* **76**, 218–219 (1962).
157. Guimarães, R. M., Asmus, C. I. R. F. & Meyer, A. DDT reintroduction for malaria control: the cost-benefit debate for public health. *Cad. Saude Publica* **23**, 2835–2844 (2007).
158. Hernández-Mariano, J. Á. *et al.* Effect of exposure to p,p'-DDE during the first half of pregnancy in the maternal thyroid profile of female residents in a Mexican floriculture area. *Environ. Res.* **156**, 597–604 (2017).
159. Barraza-Vázquez, A., Borja-Aburto, V. H., Bassol-Mayagoitia, S., Monrroy, A. & Recio-Vega, R. Dichlorodiphenyldichloroethylene concentrations in umbilical cord of newborns and determinant maternal factors. *J. Appl. Toxicol.* **28**, 27–34 (2008).
160. Sala, M. *et al.* Levels of hexachlorobenzene and other organochlorine compounds in cord blood: Exposure across placenta. *Chemosphere* **43**, 895–901 (2001).
161. Cano-Sancho, G., Salmon, A. G. & La Merrill, M. A. Association between Exposure to p,p'-DDT and Its Metabolite p,p'-DDE with Obesity: Integrated Systematic Review and Meta-Analysis. *Environ. Health Perspect.* **125**, 96002 (2017).
162. Gaspar, F. W. *et al.* Prenatal DDT and DDE exposure and child IQ in the CHAMACOS cohort. *Environ. Int.* **85**, 206–212 (2015).
163. Zhang, X. *et al.* Transplacental transfer characteristics of organochlorine pesticides in paired maternal

- and cord sera, and placentas and possible influencing factors. *Environ. Pollut.* **233**, 446–454 (2018).
164. Forns, J. *et al.* Novel application of statistical methods for analysis of multiple toxicants identifies DDT as a risk factor for early child behavioral problems. *Environ. Res.* **151**, 91–100 (2016).
165. Kezios, K. L. *et al.* Dichlorodiphenyltrichloroethane (DDT), DDT metabolites and pregnancy outcomes. *Reprod. Toxicol.* **35**, 156–164 (2013).
166. Torres-Sánchez, L. *et al.* Prenatal p,p'-DDE exposure and neurodevelopment among children 3.5-5 years of age. *Environmental Health Perspectives* **121**, 263–268 (2013).
167. Eskenazi, B. *et al.* The pine river statement: Human health consequences of DDT use. *Environmental Health Perspectives* **117**, 1359–1367 (2009).
168. Torres-Sánchez, L. *et al.* In Utero p,p'-DDE exposure and infant neurodevelopment: A perinatal cohort in Mexico. *Environ. Health Perspect.* **115**, 435–439 (2007).
169. Eskenazi, B. *et al.* In utero exposure to dichlorodiphenyltrichloroethane (DDT) and dichlorodiphenyldichloroethylene (DDE) and neurodevelopment among young Mexican American children. *Pediatrics* **118**, 233–241 (2006).
170. Ribas-Fitó, N. *et al.* In utero exposure to background concentrations of DDT and cognitive functioning among preschoolers. *Am. J. Epidemiol.* **164**, 955–962 (2006).
171. Zhuang, S., Zhang, J., Wen, Y., Zhang, C. & Liu, W. Distinct mechanisms of endocrine disruption of DDT-related pesticides toward estrogen receptor α and estrogen-related receptor γ . *Environ. Toxicol. Chem.* **31**, 2597–2605 (2012).
172. Mussi, P. *et al.* Activation of brain estrogen receptors in mice lactating from mothers exposed to DDT. in *Brain Research Bulletin* **65**, 241–247 (2005).
173. Kojima, H., Katsura, E., Takeuchi, S., Niiyama, K. & Kobayashi, K. Screening for estrogen and androgen receptor activities in 200 pesticides by in vitro reporter gene assays using Chinese hamster ovary cells. *Environ. Health Perspect.* **112**, 524–531 (2004).
174. Kelce, W. R. *et al.* Persistent DDT Metabolite p,p'-DDE is a Potent Androgen Receptor Antagonist. *Obstetrical & Gynecological Survey* **51**, 111–114 (1996).

175. Schell, L. M. *et al.* Relationships of polychlorinated biphenyls and dichlorodiphenyldichloroethylene (p,p'-DDE) with testosterone levels in adolescent males. *Environ. Health Perspect.* **122**, 304–309 (2014).
176. Eskenazi, B. *et al.* In utero and childhood DDT, DDE, PBDE and PCBs exposure and sex hormones in adolescent boys: The CHAMACOS study. *Int. J. Hyg. Environ. Health* **220**, 364–372 (2017).
177. Cohn, B. A. *et al.* DDT exposure in utero and breast cancer. *J. Clin. Endocrinol. Metab.* **100**, 2865–2872 (2015).
178. de Cock, M., de Boer, M. R., Lamoree, M., Legler, J. & van de Bor, M. Prenatal exposure to endocrine disrupting chemicals in relation to thyroid hormone levels in infants – a Dutch prospective cohort study. *Environ. Heal.* **13**, 106 (2014).
179. Blanco-Muñoz, J. *et al.* Association between organochlorine pesticide exposure and thyroid hormones in floriculture workers. *Environ. Res.* **150**, 357–363 (2016).
180. Li, C. *et al.* The association between prenatal exposure to organochlorine pesticides and thyroid hormone levels in newborns in Yancheng, China. *Environ. Res.* **129**, 47–51 (2014).
181. Parent, A. S., Naveau, E., Gerard, A., Bourguignon, J. P. & Westbrook, G. L. Early developmental actions of endocrine disruptors on the hypothalamus, hippocampus, and cerebral cortex. *J. Toxicol. Environ. Heal. - Part B Crit. Rev.* **14**, 328–345 (2011).
182. Takser, L. *et al.* Thyroid hormones in pregnancy in relation to environmental exposure to organochlorine compounds and mercury. *Environ. Health Perspect.* **113**, 1039–1045 (2005).
183. Lopez-Espinosa, M. J. *et al.* Association between thyroid hormone levels and 4,4'-DDE concentrations in pregnant women (Valencia, Spain). *Environ. Res.* **109**, 479–485 (2009).
184. Rossi, M., Taddei, A. R., Fasciani, I., Maggio, R. & Giorgi, F. The cell biology of the thyroid-disrupting mechanism of dichlorodiphenyltrichloroethane (DDT). *J. Endocrinol. Invest.* (2017). doi:10.1007/s40618-017-0716-9
185. Rossi, M. *et al.* Ethane Appears to Be an Uncompetitive Inverse Agonist for the Thyrotropin Receptor. *J. Pharmacol. Exp. Ther.* **320**, 465–474 (2007).

186. Rossi, M. *et al.* Presence of a putative steroidal allosteric site on glycoprotein hormone receptors. *Eur. J. Pharmacol.* **623**, 155–159 (2009).
187. Santini, F. *et al.* In vitro assay of thyroid disruptors affecting TSH-stimulated adenylate cyclase activity. *J. Endocrinol. Invest.* **26**, 950–955 (2003).
188. De Gregorio, F. *et al.* The insecticide 1,1,1-trichloro-2,2-bis(p-chlorophenyl) ethane (DDT) alters the membrane raft location of the TSH receptor stably expressed in Chinese hamster ovary cells. *Toxicol. Appl. Pharmacol.* **253**, 121–129 (2011).
189. Liu, C., Shi, Y., Li, H., Wang, Y. & Yang, K. P, p-DDE disturbs the homeostasis of thyroid hormones via thyroid hormone receptors, transthyretin, and hepatic enzymes. *Horm. Metab. Res.* **43**, 391–396 (2011).
190. Liu, C., Ha, M., Li, L. & Yang, K. PCB153 and p,p' -DDE disorder thyroid hormones via thyroglobulin, deiodinase 2, transthyretin, hepatic enzymes and receptors. *Environ. Sci. Pollut. Res.* **21**, 11361–11369 (2014).
191. Yaglova, N. V. & Yaglov, V. V. Changes in thyroid status of rats after prolonged exposure to low dose dichlorodiphenyltrichloroethane. *Bull. Exp. Biol. Med.* **156**, 760–762 (2014).
192. Gocmen, A., Peters, H. A., Cripps, D. J., Bryan, G. T. & Morris, C. R. Hexachlorobenzene episode in Turkey. *Biomed. Environ. Sci.* **2**, 36–43 (1989).
193. Schmid, R. Cutaneous Porphyria in Turkey. *N. Engl. J. Med.* **263**, 397–398 (1960).
194. Chalouati, H., Gamet-Payrastra, L. & Ben Saad, M. Irreversible thyroid disruption induced after subchronic exposure to hexachlorobenzene in male rats. *Toxicol. Ind. Health* **32**, 822–831 (2013).
195. Cabral, J. R. & Shubik, P. Carcinogenic activity of hexachlorobenzene in mice and hamsters. *IARC Sci. Publ.* 411–6 (1986).
196. Cabral, J. R. P., Mollner, T., Raitano, F. & Shubik, P. Carcinogenesis of hexachlorobenzene in mice. *Int. J. Cancer* **23**, 47–51 (1979).
197. Cabral, J. R. P., Shubik, P., MOLLNER, T. & RAITANO, F. Carcinogenic activity of hexachlorobenzene in hamsters. *Nature* **269**, 510–511 (1977).

198. Can, C. & Nigogosyan, G. Acquired toxic porphyria cutanea tarda due to hexachlorobenzene. Report of 348 cases caused by this fungicide. *JAMA* **183**, 88–91 (1963).
199. Eskenazi, B. *et al.* Association of in utero organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ. Health Perspect.* **112**, 1116–1124 (2004).
200. Ribas-Fitó, N. *et al.* Exposure to hexachlorobenzene during pregnancy and children's social behavior at 4 years of age. *Environ. Health Perspect.* **115**, 447–450 (2007).
201. Smink, A. *et al.* Exposure to hexachlorobenzene during pregnancy increases the risk of overweight in children aged 6 years. *Acta Paediatr.* **97**, 1465–1469 (2008).
202. Sala, M., Sunyer, J., Herrero, C., To-Figueras, J. & Grimalt, J. Association between serum concentrations of hexachlorobenzene and polychlorobiphenyls with thyroid hormone and liver enzymes in a sample of the general population. *Occup. Environ. Med.* **58**, 172–7 (2001).
203. Meeker, J. D., Altshul, L. & Hauser, R. Serum PCBs, p,p'-DDE and HCB predict thyroid hormone levels in men. *Environ. Res.* **104**, 296–304 (2007).
204. Foster, W. G., McMahon, A., Villeneuve, D. C. & Jarrell, J. F. Hexachlorobenzene (HCB) suppresses circulating progesterone concentrations during the luteal phase in the cynomolgus monkey. *J. Appl. Toxicol.* **12**, 13–7 (1992).
205. Foster, W. G., McMahon, A., Younglai, E. V., Jarrell, J. F. & Lecavalier, P. Alterations in circulating ovarian steroids in hexachlorobenzene-exposed monkeys. *Reprod. Toxicol.* **9**, 541–548 (1995).
206. Jarrell, J. F. *et al.* Hexachlorobenzene toxicity in the monkey primordial germ cell without induced porphyria. *Reprod. Toxicol.* **7**, 41–7 (1993).
207. Sims, D. E., Singh, A., Donald, A., Jarrell, J. & Villeneuve, D. C. Alteration of primate ovary surface epithelium by exposure to hexachlorobenzene: A quantitative study. *Histol. Histopathol.* **6**, 525–529 (1991).
208. Babineau, K. A., Singh, A., Jarrell, J. F. & Villeneuve, D. C. Surface epithelium of the ovary following oral administration of hexachlorobenzene to the monkey. *J.Submicrosc.Cytol.Pathol.* **23**, 457–464 (1991).

209. Arnold, D. L. *et al.* Long-term toxicity of hexachlorobenzene in the rat and the effect of dietary vitamin A. *Food Chem. Toxicol.* **23**, 779–793 (1985).
210. Hadjab, S., Maurel, D., Cazals, Y. & Siaud, P. Hexachlorobenzene, a dioxin-like compound, disrupts auditory function in rat. *Hear. Res.* **191**, 125–134 (2004).
211. van Raaij, J. A. G. M., Kaptein, E., Visser, T. J. & van den Berg, K. J. Increased glucuronidation of thyroid hormone in hexachlorobenzene-treated rats. *Biochem. Pharmacol.* **45**, 627–631 (1993).
212. Smith, A. G., Dinsdale, D., Cabral, J. R. & Wright, A. L. Goitre and wasting induced in hamsters by hexachlorobenzene. *Arch. Toxicol.* **60**, 343–9 (1987).
213. Van Raaij, J. A. G. M., Frijters, C. M. G. & van den Berg, K. J. Hexachlorobenzene-induced hypothyroidism. Involvement of different mechanisms by parent compound and metabolite. *Biochem. Pharmacol.* **46**, 1385–1391 (1993).
214. Alvarez, L. *et al.* The role of type I and type II 5' deiodinases on hexachlorobenzene-induced alteration of the hormonal thyroid status. *Toxicology* **207**, 349–362 (2005).
215. Chiappini, F., Pontillo, C., Randi, A., Alvarez, L. & Kleiman de Pisarev, D. L. Hexachlorobenzene induces TGF- β 1 expression, which is a regulator of p27 and cyclin D1 modifications. *Toxicol. Lett.* **230**, 1–9 (2014).
216. Chiappini, F., Alvarez, L., Lux-Lantos, V., Randi, A. S. & Kleiman de Pisarev, D. L. Hexachlorobenzene Triggers Apoptosis in Rat Thyroid Follicular Cells. *Toxicol. Sci.* **108**, 301–310 (2009).
217. Muller, M. *et al.* Neurologic dysfunction and genotoxicity induced by low levels of chlorpyrifos. *Neurotoxicology* **45**, 22–30 (2014).
218. Whyatt, R. M. *et al.* Contemporary-use pesticide in personal air samples during pregnancy and blood samples at delivery among urban minority mothers and newborns. *Environ. Health Perspect.* **111**, 749–756 (2003).
219. Perera, F. P. *et al.* The challenge of preventing environmentally related disease in young children: Community-based research in New York City. *Environmental Health Perspectives* **110**, 197–204

- (2002).
220. Whyatt, R. M. *et al.* Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ. Health Perspect.* **112**, 1125–1132 (2004).
 221. Farahat, T., Hala Mohammed, S., Sanad, Z. & NA, F. Organophosphate Pesticide Exposure during Pregnancy and Adverse Perinatal Outcome. *J. Women's Heal. Care* **5**, 1–4 (2016).
 222. Abdel-Rahman, A. a *et al.* Pharmacokinetic profile and placental transfer of a single intravenous injection of [(14)C]chlorpyrifos in pregnant rats. *Arch. Toxicol.* **76**, 452–9 (2002).
 223. Rauh, V. A. *et al.* Impact of Prenatal Chlorpyrifos Exposure on Neurodevelopment in the First 3 Years of Life Among Inner-City Children. *Pediatrics* **118**, e1845–e1859 (2006).
 224. De Cock, M., Maas, Y. G. H. & Van De Bor, M. Does perinatal exposure to endocrine disruptors induce autism spectrum and attention deficit hyperactivity disorders? Review. *Acta Paediatr. Int. J. Paediatr.* **101**, 811–818 (2012).
 225. Rauh, V. *et al.* Seven-year neurodevelopmental scores and prenatal exposure to chlorpyrifos, a common agricultural pesticide. *Environ. Health Perspect.* **119**, 1196–1201 (2011).
 226. Rauh, V. A. *et al.* Prenatal exposure to the organophosphate pesticide chlorpyrifos and childhood tremor. *Neurotoxicology* **51**, 80–86 (2015).
 227. Eaton, D. L. *et al.* Review of the toxicology of chlorpyrifos with an emphasis on human exposure and neurodevelopment. *Critical Reviews in Toxicology* **38**, 1–125 (2008).
 228. Reiss, R., Chang, E. T., Richardson, R. J. & Goodman, M. A review of epidemiologic studies of low-level exposures to organophosphorus insecticides in non-occupational populations. *Critical Reviews in Toxicology* **45**, 531–641 (2015).
 229. Timchalk, C. *et al.* A physiologically based pharmacokinetic and pharmacodynamic (PBPK/PD) model for the organophosphate insecticide chlorpyrifos in rats and humans. *Toxicol. Sci.* **66**, 34–53 (2002).
 230. EPA. EPA Administrator Pruitt Denies Petition to Ban Widely Used Pesticide. (2017).
 231. Trasande, L. When enough data are not enough to enact policy: The failure to ban chlorpyrifos. *PLoS Biol.* **15**, e2003671 (2017).

232. Fortenberry, G. Z., Hu, H., Turyk, M., Barr, D. B. & Meeker, J. D. Association between urinary 3, 5, 6-trichloro-2-pyridinol, a metabolite of chlorpyrifos and chlorpyrifos-methyl, and serum T4 and TSH in NHANES 1999-2002. *Sci. Total Environ.* **424**, 351–355 (2012).
233. Meeker, J. D., Barr, D. B. & Hauser, R. Thyroid hormones in relation to urinary metabolites of non-persistent insecticides in men of reproductive age. *Reprod. Toxicol.* **22**, 437–442 (2006).
234. Slotkin, T. A., Cooper, E. M., Stapleton, H. M. & Seidler, F. J. Does thyroid disruption contribute to the developmental neurotoxicity of chlorpyrifos? *Environ. Toxicol. Pharmacol.* **36**, 284–287 (2013).
235. De Angelis, S. *et al.* Developmental exposure to chlorpyrifos induces alterations in thyroid and thyroid hormone levels without other toxicity signs in CD-1 mice. *Toxicol. Sci.* **108**, 311–319 (2009).
236. Silva, J. G. *et al.* Chlorpyrifos induces anxiety-like behavior in offspring rats exposed during pregnancy. *Neurosci. Lett.* **641**, 94–100 (2017).
237. Ghisari, M. & Bonefeld-Jorgensen, E. C. Impact of environmental chemicals on the thyroid hormone function in pituitary rat GH3 cells. *Mol. Cell. Endocrinol.* **244**, 31–41 (2005).
238. Jeong, S. H., Kim, B. Y., Kang, H. G., Ku, H. O. & Cho, J. H. Effect of chlorpyrifos-methyl on steroid and thyroid hormones in rat F0- and F1-generations. *Toxicology* **220**, 189–202 (2006).
239. Ventura, C. *et al.* Differential mechanisms of action are involved in chlorpyrifos effects in estrogen-dependent or -independent breast cancer cells exposed to low or high concentrations of the pesticide. *Toxicol. Lett.* **213**, 184–193 (2012).
240. Grünfeld, H. T. & Bonefeld-Jorgensen, E. C. Effect of in vitro estrogenic pesticides on human oestrogen receptor α and β mRNA levels. *Toxicol. Lett.* **151**, 467–480 (2004).
241. Yu, K. *et al.* Chlorpyrifos is estrogenic and alters embryonic hatching, cell proliferation and apoptosis in zebrafish. *Chem. Biol. Interact.* **239**, 26–33 (2015).
242. Ventura, C. *et al.* Pesticide chlorpyrifos acts as an endocrine disruptor in adult rats causing changes in mammary gland and hormonal balance. *J. Steroid Biochem. Mol. Biol.* **156**, 1–9 (2016).
243. Viswanath, G. *et al.* Anti-androgenic endocrine disrupting activities of chlorpyrifos and piperophos. *J. Steroid Biochem. Mol. Biol.* **120**, 22–29 (2010).

244. Wegner, S., Browne, P. & Dix, D. Identifying reference chemicals for thyroid bioactivity screening. *Reprod. Toxicol.* **65**, 402–413 (2016).
245. Marinovich, M. *et al.* Thyroid peroxidase as toxicity target for dithiocarbamates. *Arch. Toxicol.* **71**, 508–512 (1997).
246. Mallem, L., Boulakoud, M. S. & Franck, M. Hypothyroidism after medium exposure to the fungicide maneb in the rabbit *Cuniculus lepus*. *Commun. Agric. Appl. Biol. Sci.* **71**, 91–99 (2006).
247. Axelstad, M. *et al.* Exposure to the widely used fungicide mancozeb causes thyroid hormone disruption in rat dams but no behavioral effects in the offspring. *Toxicol. Sci.* **120**, 439–446 (2011).
248. Medda, E. *et al.* Iodine nutritional status and thyroid effects of exposure to ethylenebisdithiocarbamates. *Environ. Res.* **154**, 152–159 (2017).
249. Hester, S. D., Wolf, D. C., Nesnow, S. & Thai, S. F. Transcriptional Profiles in Liver from Rats Treated with Tumorigenic and Non-tumorigenic Triazole Conazole Fungicides: Propiconazole, Triadimefon, and Myclobutanil. *Toxicol. Pathol.* **34**, 879–894 (2006).
250. Campos, É. & Freire, C. Exposure to non-persistent pesticides and thyroid function: A systematic review of epidemiological evidence. *International Journal of Hygiene and Environmental Health* **219**, 481–497 (2016).
251. Yadav, A. K. & Singh, T. P. Pesticide-induced impairment of thyroid physiology in the freshwater catfish, *Heteropneustes fossilis*. *Environ. Pollut.* **43**, 29–38 (1987).
252. Van den Berg, K. J., van Raaij, J. A. G. M., Bragt, P. C. & Notten, W. R. F. Interactions of halogenated industrial chemicals with transthyretin and effects on thyroid hormone levels in vivo. *Arch. Toxicol.* **65**, 15–19 (1991).
253. Ishihara, A., Sawatsubashi, S. & Yamauchi, K. Endocrine disrupting chemicals: Interference of thyroid hormone binding to transthyretins and to thyroid hormone receptors. *Mol. Cell. Endocrinol.* **199**, 105–117 (2003).
254. Sinha, N., Lal, B. & Singh, T. P. Carbaryl-induced thyroid dysfunction in the freshwater catfish *Clarias batrachus*. *Ecotoxicol. Environ. Saf.* **21**, 240–247 (1991).

255. Akhtar, N., Kayani, S. A., Ahmad, M. M. & Shahab, M. Insecticide-induced changes in secretory activity of the thyroid gland in rats. in *Journal of Applied Toxicology* **16**, 397–400 (1996).
256. Lerro, C. C. *et al.* Organophosphate insecticide use and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study. *Occup. Environ. Med.* **72**, 736–744 (2015).
257. Jin, Y. *et al.* Permethrin exposure during puberty has the potential to enantioselectively induce reproductive toxicity in mice. *Environ. Int.* **42**, 144–151 (2012).
258. Jin, Y. *et al.* Enantioselective induction of estrogen-responsive gene expression by permethrin enantiomers in embryo-larval zebrafish. *Chemosphere* **74**, 1238–1244 (2009).
259. Du, G. *et al.* Assessing hormone receptor activities of pyrethroid insecticides and their metabolites in reporter gene assays. *Toxicol. Sci.* **116**, 58–66 (2010).
260. Tu, W. *et al.* Permethrin is a potential thyroid-disrupting chemical: In vivo and in silico evidence. *Aquat. Toxicol.* **175**, 39–46 (2016).
261. Maiti, P. K., Kar, A., Gupta, P. & Chaurasia, S. S. Loss of membrane integrity and inhibition of type-I iodothyronine 5'-monodeiodinase activity by fenvalerate in female mouse. *Biochem. Biophys. Res. Commun.* **214**, 905–909 (1995).
262. Maiti, P. K. & Kar, A. Dimethoate inhibits extrathyroidal 5'-monodeiodination of thyroxine to 3,3',5-triiodothyronine in mice: The possible involvement of the lipid peroxidative process. *Toxicol. Lett.* **91**, 1–6 (1997).
263. Tu, W. *et al.* Acute exposure to synthetic pyrethroids causes bioconcentration and disruption of the hypothalamus-pituitary-thyroid axis in zebrafish embryos. *Sci. Total Environ.* **542**, 876–885 (2016).
264. Kaul, P. P. *et al.* Fenvalerate-induced alterations in circulatory thyroid hormones and calcium stores in rat brain. *Toxicol. Lett.* **89**, 29–33 (1996).
265. Giray, B., Çağlayan, A., Erkekoğlu, P. & Hİncal, F. Fenvalerate exposure alters thyroid hormone status in selenium- and/or iodine-deficient rats. *Biol. Trace Elem. Res.* **135**, 233–241 (2010).
266. Swetox. Pyriproxifen and microcephaly: an investigation of potential ties to the ongoing ' Zika epidemic '. (2016). doi:10.13140/RG.2.1.4853.4166

267. van den Berg, M. *et al.* WHO/UNEP global surveys of PCDDs, PCDFs, PCBs and DDTs in human milk and benefit–risk evaluation of breastfeeding. *Archives of Toxicology* **91**, 83–96 (2017).
268. Meironyté Guvenius, D., Aronsson, A., Ekman-Ordeberg, G., Bergman, Å. & Norén, K. Human prenatal and postnatal exposure to polybrominated diphenyl ethers, polychlorinated biphenyls, polychlorobiphenylols, and pentachlorophenol. *Environmental Health Perspectives* **111**, 1235–1241 (2003).
269. Soechitram, S. D., Athanasiadou, M., Hovander, L., Bergman, Å. & Sauer, P. J. J. Fetal exposure to PCBs and their hydroxylated metabolites in a Dutch cohort. *Environ. Health Perspect.* **112**, 1208–1212 (2004).
270. Fängström, B., Strid, A., Grandjean, P., Weihe, P. & Bergman, Å. A retrospective study of PBDEs and PCBs in human milk from the Faroe Islands. *Environ. Heal.* **4**, 12 (2005).
271. Guvenius, D. M., Hassanzadeh, P., Bergman, A. & Norén, K. Metabolites of polychlorinated biphenyls in human liver and adipose tissue. *Environ. Toxicol. Chem.* **21**, 2264–2269 (2002).
272. Valvi, D. *et al.* Prenatal concentrations of polychlorinated biphenyls, DDE, and DDT and overweight in children: A prospective birth cohort study. *Environ. Health Perspect.* **120**, 451–457 (2012).
273. Caspersen, I. H. *et al.* The influence of maternal dietary exposure to dioxins and PCBs during pregnancy on ADHD symptoms and cognitive functions in Norwegian preschool children. *Environ. Int.* **94**, 649–660 (2016).
274. Stewart, P. W. *et al.* The Relationship between Prenatal PCB Exposure and Intelligence (IQ) in 9-Year-Old Children. *Environ. Health Perspect.* **116**, 1416–1422 (2008).
275. Walkowiak, J. *et al.* Environmental exposure to polychlorinated biphenyls and quality of the home environment: effects on psychodevelopment in early childhood. *Lancet (London, England)* **358**, 1602–1607 (2001).
276. Verner, M. A. *et al.* Prenatal and early-life polychlorinated biphenyl (PCB) levels and behavior in Inuit preschoolers. *Environ. Int.* **78**, 90–94 (2015).
277. Ethier, A. A. *et al.* Assessing new dimensions of attentional functions in children prenatally exposed to

- environmental contaminants using an adapted Posner paradigm. *Neurotoxicol. Teratol.* **51**, 27–34 (2015).
278. Schantz, S. L., Widholm, J. J. & Rice, D. C. Effects of PCB exposure on neuropsychological function in children. *Environ. Health Perspect.* **111**, 357–576 (2003).
279. Behforooz, B., Newman, J., Gallo, M. V. & Schell, L. M. PCBs and measures of attention and impulsivity on a continuous performance task of young adults. *Neurotoxicol. Teratol.* **64**, 29–36 (2017).
280. Šovčíková, E. *et al.* Simple reaction time in 8-9-year old children environmentally exposed to PCBs. *Neurotoxicology* **51**, 138–144 (2015).
281. Berghuis, S. A., Soechitram, S. D., Hitzert, M. M., Sauer, P. J. J. & Bos, A. F. Prenatal exposure to polychlorinated biphenyls and their hydroxylated metabolites is associated with motor development of three-month-old infants. *Neurotoxicology* **38**, 124–130 (2013).
282. Lyall, K. *et al.* Polychlorinated biphenyl and organochlorine pesticide concentrations in maternal mid-pregnancy serum samples: Association with Autism spectrum disorder and intellectual disability. *Environ. Health Perspect.* **125**, 474–480 (2017).
283. Cheslack-Postava, K. *et al.* Maternal serum persistent organic pollutants in the Finnish Prenatal Study of Autism: A pilot study. *Neurotoxicol. Teratol.* **38**, 1–5 (2013).
284. Neugebauer, J. *et al.* The influence of low level pre- and perinatal exposure to PCDD/Fs, PCBs, and lead on attention performance and attention-related behavior among German school-aged children: results from the Duisburg Birth Cohort Study. *Int. J. Hyg. Environ. Health* **218**, 153–162 (2015).
285. Verner, M. A. *et al.* Measured prenatal and estimated postnatal levels of polychlorinated biphenyls (PCBs) and ADHD-related behaviors in 8-year-old children. *Environ. Health Perspect.* **123**, 888–894 (2015).
286. Bergkvist, C., Berglund, M., Glynn, A., Wolk, A. & Åkesson, A. Dietary exposure to polychlorinated biphenyls and risk of myocardial infarction - A population-based prospective cohort study. *Int. J. Cardiol.* **183**, 242–248 (2015).
287. Langer, P. *et al.* Thyroid ultrasound volume, structure and function after long-term high exposure of

- large population to polychlorinated biphenyls, pesticides and dioxin. *Chemosphere* **69**, 118–127 (2007).
288. Langer, P. *et al.* Thyroid volume, iodine intake, autoimmune thyroid disorders, inborn factors, and endocrine disruptors: Twenty-year studies of multiple effects puzzle in slovakia. *Endocr. Regul.* **46**, 191–203 (2012).
289. McKinney, J. D. & Waller, C. L. Polychlorinated biphenyls as hormonally active structural analogues. *Environ. Health Perspect.* **102**, 290–297 (1994).
290. Otake, T. *et al.* Thyroid hormone status of newborns in relation to in utero exposure to PCBs and hydroxylated PCB metabolites. *Environ. Res.* **105**, 240–246 (2007).
291. Herbstman, J. B. *et al.* Birth delivery mode modifies the associations between prenatal polychlorinated biphenyl (PCB) and polybrominated diphenyl ether (PBDE) and neonatal thyroid hormone levels. *Environ. Health Perspect.* **116**, 1376–1382 (2008).
292. Brucker-Davis, F. *et al.* Cord Blood Thyroid Tests in Boys Born With and Without Cryptorchidism: Correlations with Birth Parameters and *In Utero* Xenobiotics Exposure. *Thyroid* **21**, 1133–1141 (2011).
293. Dallaire, R. *et al.* Thyroid hormone levels of pregnant inuit women and their infants exposed to environmental contaminants. *Environ. Health Perspect.* **117**, 1014–1020 (2009).
294. Darnerud, P. O. *et al.* POP levels in breast milk and maternal serum and thyroid hormone levels in mother-child pairs from Uppsala, Sweden. *Environ. Int.* **36**, 180–187 (2010).
295. Leijds, M. M. *et al.* Thyroid hormone metabolism and environmental chemical exposure. *Environ. Heal.* **11**, S10 (2012).
296. Grimm, F. A., Lehmler, H. J., He, X., Robertson, L. W. & Duffel, M. W. Sulfated metabolites of polychlorinated biphenyls are high-affinity ligands for the thyroid hormone transport protein transthyretin. *Environ. Health Perspect.* **121**, 657–662 (2013).
297. Álvarez-Pedrerol, M. *et al.* Thyroid disruption at birth due to prenatal exposure to β -hexachlorocyclohexane. *Environ. Int.* **34**, 737–740 (2008).
298. Hisada, A. *et al.* Associations between levels of hydroxylated PCBs and PCBs in serum of pregnant women and blood thyroid hormone levels and body size of neonates. *Int. J. Hyg. Environ. Health* **217**,

- 546–553 (2014).
299. De Cock, M. *et al.* Thyroid-stimulating hormone levels in newborns and early life exposure to endocrine-disrupting chemicals: Analysis of three European mother-child cohorts. *Pediatr. Res.* **82**, 429–437 (2017).
300. Crofton, K. M. *et al.* Thyroid-hormone-disrupting chemicals: Evidence for dose-dependent additivity or synergism. *Environ. Health Perspect.* **113**, 1549–1554 (2005).
301. Katarzyńska, D., Hrabia, A., Kowalik, K. & Sechman, A. Comparison of the in vitro effects of TCDD, PCB 126 and PCB 153 on thyroid-restricted gene expression and thyroid hormone secretion by the chicken thyroid gland. *Environ. Toxicol. Pharmacol.* **39**, 496–503 (2015).
302. Ahmed, R. G. Early weaning PCB95 exposure alters the neonatal endocrine system: Thyroid adipokine dysfunction. *J. Endocrinol.* **219**, 205–215 (2013).
303. Meerts, I. A. T. M. Developmental Exposure to 4-hydroxy-2,3,3',4',5-pentachlorobiphenyl (4-OH-CB107): Long-Term Effects on Brain Development, Behavior, and Brain Stem Auditory Evoked Potentials in Rats. *Toxicol. Sci.* **82**, 207–218 (2004).
304. Reilly, M. P. *et al.* The effects of prenatal PCBs on adult social behavior in rats. *Horm. Behav.* **73**, 47–55 (2015).
305. Gutleb, A. C. *et al.* In vitro assay shows that PCB metabolites completely saturate thyroid hormone transport capacity in blood of wild polar bears (*Ursus maritimus*). *Environ. Sci. Technol.* **44**, 3149–3154 (2010).
306. Yang, H. *et al.* Molecular mechanisms of 2, 3, 4, 5-pentachlorobiphenyl-induced thyroid dysfunction in FRTL-5 cells. *PLoS One* **10**, (2015).
307. Guo, H. *et al.* Molecular mechanisms of human thyrocyte dysfunction induced by low concentrations of polychlorinated biphenyl 118 through the Akt/FoxO3a/NIS pathway. *J. Appl. Toxicol.* **35**, 992–998 (2015).
308. Xu, B. *et al.* 2,3',4,4',5-pentachlorobiphenyl induces inflammatory responses in the thyroid through JNK and Aryl hydrocarbon receptor-mediated pathway. *Toxicol. Sci.* **149**, 300–311 (2016).

309. Ekuase, E. J., Liu, Y., Lehmler, H. J., Robertson, L. W. & Duffel, M. W. Structure-activity relationships for hydroxylated polychlorinated biphenyls as inhibitors of the sulfation of dehydroepiandrosterone catalyzed by human hydroxysteroid sulfotransferase SULT2A1. *Chem. Res. Toxicol.* **24**, 1720–1728 (2011).
310. Soechitram, S. D., Berghuis, S. A., Visser, T. J. & Sauer, P. J. J. Polychlorinated biphenyl exposure and deiodinase activity in young infants. *Sci. Total Environ.* **574**, 1117–1124 (2017).
311. Dickerson, S. M., Cunningham, S. L., Patisaul, H. B., Woller, M. J. & Gore, A. C. Endocrine disruption of brain sexual differentiation by developmental PCB exposure. *Endocrinology* **152**, 581–594 (2011).
312. Walker, D. M., Goetz, B. M. & Gore, A. C. Dynamic Postnatal Developmental and Sex-Specific Neuroendocrine Effects of Prenatal Polychlorinated Biphenyls in rats. *Mol. Endocrinol.* **28**, 99–115 (2014).
313. deCastro, B. R., Korrick, S. A., Spengler, J. D. & Soto, A. M. Estrogenic Activity of Polychlorinated Biphenyls Present in Human Tissue and the Environment. *Environ. Sci. Technol.* **40**, 2819–2825 (2006).
314. Zhang, Q. *et al.* Characterization of estrogen receptor α activities in polychlorinated biphenyls by in vitro dual-luciferase reporter gene assay. *Environ. Pollut.* **189**, 169–175 (2014).
315. Hamers, T. *et al.* In vitro toxicity profiling of ultrapure non-dioxin-like polychlorinated biphenyl congeners and their relative toxic contribution to PCB mixtures in humans. *Toxicol. Sci.* **121**, 88–100 (2011).
316. Flor, S., He, X., Lehmler, H. J. & Ludewig, G. Estrogenicity and androgenicity screening of PCB sulfate monoesters in human breast cancer MCF-7 cells. *Environ. Sci. Pollut. Res.* **23**, 2186–2200 (2016).
317. Lyche, J. L., Rosseland, C., Berge, G. & Polder, A. Human health risk associated with brominated flame-retardants (BFRs). *Environ. Int.* **74**, 170–180 (2015).
318. Law, R. J. *et al.* Levels and trends of PBDEs and HBCDs in the global environment: Status at the end of 2012. *Environment International* **65**, 147–158 (2014).
319. Fliedner *et al.* Current levels and trends of selected EU Water Framework Directive priority substances in freshwater fish from the German environmental specimen bank. *Environ. Pollut.* **216**, 866–876

- (2016).
320. Sjödin, A. *et al.* Polybrominated diphenyl ethers, polychlorinated biphenyls, and persistent pesticides in serum from the national health and nutrition examination survey: 2003-2008. *Environ. Sci. Technol.* **48**, 753–760 (2014).
 321. Vizcaino, E., Grimalt, J. O., Fernández-Somoano, A. & Tardon, A. Transport of persistent organic pollutants across the human placenta. *Environ. Int.* **65**, 107–115 (2014).
 322. Choi, G. *et al.* Occurrences of major polybrominated diphenyl ethers (PBDEs) in maternal and fetal cord blood sera in Korea. *Sci. Total Environ.* **491–492**, 219–226 (2014).
 323. Gómara, B. *et al.* Distribution of polybrominated diphenyl ethers in human umbilical cord serum, paternal serum, maternal serum, placentas, and breast milk from madrid population, Spain. *Environ. Sci. Technol.* **41**, 6961–6968 (2007).
 324. Zhang, H. *et al.* Prenatal PBDE and PCB exposures and reading, cognition, and externalizing behavior in children. *Environ. Health Perspect.* **125**, 746–752 (2017).
 325. Braun, J. M. *et al.* Prenatal environmental chemical exposures and longitudinal patterns of child neurobehavior. *Neurotoxicology* **62**, 192–199 (2017).
 326. Vuong, A. M. *et al.* Prenatal and postnatal polybrominated diphenyl ether exposure and visual spatial abilities in children. *Environ. Res.* **153**, 83–92 (2017).
 327. Sagiv, S. K. *et al.* Prenatal and childhood polybrominated diphenyl ether (PBDE) exposure and attention and executive function at 9-12 years of age. *Neurotoxicol. Teratol.* **52**, 151–161 (2015).
 328. Chen, A. *et al.* Prenatal polybrominated diphenyl ether exposures and neurodevelopment in U.S. children through 5 years of age: The home study. *Environ. Health Perspect.* **122**, 856–862 (2014).
 329. Eskenazi, B. *et al.* In utero and childhood polybrominated diphenyl ether (PBDE) exposures and neurodevelopment in the CHAMACOS study. *Environmental Health Perspectives* **121**, 257–262 (2013).
 330. Cowell, W. J. *et al.* Prenatal exposure to polybrominated diphenyl ethers and child attention problems at 3-7 years. *Neurotoxicol. Teratol.* **52**, 143–150 (2015).
 331. Ding, G. *et al.* Association between prenatal exposure to polybrominated diphenyl ethers and young

- children's neurodevelopment in China. *Environ. Res.* **142**, 104–111 (2015).
332. Herbstman, J. B. & Mall, J. K. Developmental Exposure to Polybrominated Diphenyl Ethers and Neurodevelopment. *Curr. Environ. Heal. Reports* **1**, 101–112 (2014).
333. Vuong, A. M. *et al.* Childhood polybrominated diphenyl ether (PBDE) exposure and executive function in children in the HOME Study. *International Journal of Hygiene and Environmental Health* (2017). doi:10.1016/j.ijheh.2017.10.006
334. Naert, C., Van Peteghem, C., Kupper, J., Jenni, L. & Naegeli, H. Distribution of polychlorinated biphenyls and polybrominated diphenyl ethers in birds of prey from Switzerland. *Chemosphere* **68**, 977–987 (2007).
335. Zhang, H., Li, X., Nie, J. & Niu, Q. Lactation exposure to BDE-153 damages learning and memory, disrupts spontaneous behavior and induces hippocampus neuron death in adult rats. *Brain Res.* **1517**, 44–56 (2013).
336. Gee, J. R. & Moser, V. C. Acute postnatal exposure to brominated diphenylether 47 delays neuromotor ontogeny and alters motor activity in mice. *Neurotoxicol. Teratol.* **30**, 79–87 (2008).
337. Eriksson, P., Jakobsson, E. & Fredriksson, A. Brominated flame retardants: A novel class of developmental neurotoxicants in our environment? *Environ. Health Perspect.* **109**, 903–908 (2001).
338. Branchi, I., Capone, F., Alleva, E. & Costa, L. G. Polybrominated diphenyl ethers: Neurobehavioral effects following developmental exposure. *Neurotoxicology* **24**, 449–462 (2003).
339. Gassmann, K. *et al.* BDE-47 and 6-OH-BDE-47 modulate calcium homeostasis in primary fetal human neural progenitor cells via ryanodine receptor-independent mechanisms. *Arch. Toxicol.* **88**, 1537–1548 (2014).
340. Costa, L. G., de Laat, R., Tagliaferri, S. & Pellacani, C. A mechanistic view of polybrominated diphenyl ether (PBDE) developmental neurotoxicity. *Toxicol. Lett.* **230**, 282–294 (2014).
341. Dingemans, M. M. L. *et al.* Neonatal exposure to brominated flame retardant BDE-47 reduces long-term potentiation and postsynaptic protein levels in mouse hippocampus. *Environ. Health Perspect.* **115**, 865–870 (2007).

342. Bradner, J. M., Suragh, T. A. & Caudle, W. M. Alterations to the circuitry of the frontal cortex following exposure to the polybrominated diphenyl ether mixture, DE-71. *Toxicology* **312**, 48–55 (2013).
343. Bradner, J. M. *et al.* Exposure to the polybrominated diphenyl ether mixture DE-71 damages the nigrostriatal dopamine system: Role of dopamine handling in neurotoxicity. *Exp. Neurol.* **241**, 138–147 (2013).
344. Jacobson, M. H. *et al.* Serum polybrominated diphenyl ether concentrations and thyroid function in young children. *Environ. Res.* **149**, 222–230 (2016).
345. Xu, X. *et al.* Elevated serum polybrominated diphenyl ethers and alteration of thyroid hormones in children from Guiyu, China. *PLoS One* **9**, (2014).
346. Makey, C. M. *et al.* Polybrominated diphenyl ether exposure and thyroid function tests in North American adults. *Environ. Health Perspect.* **124**, 420–425 (2016).
347. Kiciński, M. *et al.* Neurobehavioral function and low-level exposure to brominated flame retardants in adolescents: a cross-sectional study. *Environ. Heal.* **11**, 86 (2012).
348. GuanGen, H. *et al.* Correlations of PCBs, DIOXIN, and PBDE with TSH in Children's Blood in Areas of Computer E-waste Recycling*. *Biomed Env. Sci* **24**, 112–116 (2011).
349. Abdelouahab, N. *et al.* Maternal and cord-blood thyroid hormone levels and exposure to polybrominated diphenyl ethers and polychlorinated biphenyls during early pregnancy. *Am. J. Epidemiol.* **178**, 701–713 (2013).
350. Zhao, X. *et al.* The correlation between polybrominated diphenyl ethers (PBDES) and thyroid hormones in the general population: A meta-analysis. *PLoS One* **10**, (2015).
351. Lee, E. *et al.* Evaluation of liver and thyroid toxicity in Sprague-Dawley rats after exposure to polybrominated diphenyl ether BDE-209. *J. Toxicol. Sci.* **35**, 535–545 (2010).
352. Miller, V. M., Sanchez-Morrissey, S., Brosch, K. O. & Seegal, R. F. Developmental coexposure to polychlorinated biphenyls and polybrominated diphenyl ethers has additive effects on circulating thyroxine levels in rats. *Toxicol. Sci.* **127**, 76–83 (2012).

353. Kodavanti, P. R. S. *et al.* Developmental exposure to a commercial PBDE Mixture, DE-71: Neurobehavioral, hormonal, and reproductive effects. *Toxicol. Sci.* **116**, 297–312 (2010).
354. Richardson, V. M. *et al.* Possible mechanisms of thyroid hormone disruption in mice by BDE 47, a major polybrominated diphenyl ether congener. *Toxicol. Appl. Pharmacol.* **226**, 244–250 (2008).
355. Ren, X. M., Guo, L. H., Gao, Y., Zhang, B. T. & Wan, B. Hydroxylated polybrominated diphenyl ethers exhibit different activities on thyroid hormone receptors depending on their degree of bromination. *Toxicol. Appl. Pharmacol.* **268**, 256–263 (2013).
356. Li, F. *et al.* Hormone activity of hydroxylated polybrominated diphenyl ethers on human thyroid receptor- β : In vitro and in silico investigations. *Environ. Health Perspect.* **118**, 602–606 (2010).
357. Kitamura, S. *et al.* Thyroid hormone-like and estrogenic activity of hydroxylated PCBs in cell culture. *Toxicology* **208**, 377–387 (2005).
358. Schriks, M., Roessig, J. M., Murk, A. J. & Furlow, J. D. Thyroid hormone receptor isoform selectivity of thyroid hormone disrupting compounds quantified with an in vitro reporter gene assay. *Environ. Toxicol. Pharmacol.* **23**, 302–307 (2007).
359. Kojima, H. *et al.* Nuclear hormone receptor activity of polybrominated diphenyl ethers and their hydroxylated and methoxylated metabolites in transactivation assays using Chinese hamster ovary cells. *Environ. Health Perspect.* **117**, 1210–1218 (2009).
360. Ibhazehiebo, K. *et al.* Disruption of thyroid hormone receptor-mediated transcription and thyroid hormone-induced purkinje cell dendrite arborization by Polybrominated diphenyl ethers. *Environ. Health Perspect.* **119**, 168–175 (2011).
361. Dach, K. *et al.* BDE-99 impairs differentiation of human and mouse NPCs into the oligodendroglial lineage by species-specific modes of action. *Sci. Rep.* **7**, 44861 (2017).
362. Macaulay, L. J. *et al.* Developmental toxicity of the PBDE metabolite 6-OH-BDE-47 in zebrafish and the potential role of thyroid receptor ?? *Aquat. Toxicol.* **168**, 38–47 (2015).
363. Ren, X. M. & Guo, L. H. Assessment of the binding of hydroxylated polybrominated diphenyl ethers to thyroid hormone transport proteins using a site-specific fluorescence probe. *Environ. Sci. Technol.* **46**,

- 4633–4640 (2012).
364. Cao, J. *et al.* Structure-based investigation on the binding interaction of hydroxylated polybrominated diphenyl ethers with thyroxine transport proteins. *Toxicology* **277**, 20–28 (2010).
365. Roberts, S. C., Bianco, A. C. & Stapleton, H. M. Disruption of Type 2 Iodothyronine Deiodinase Activity in Cultured Human Glial Cells by Polybrominated Diphenyl Ethers. *Chem. Res. Toxicol.* **28**, 1265–1274 (2015).
366. Marsan, E. S. & Bayse, C. A. Halogen-Bonding Interactions of Polybrominated Diphenyl Ethers and Thyroid Hormone Derivatives: A Potential Mechanism for the Inhibition of Iodothyronine Deiodinase. *Chem. - A Eur. J.* **23**, 6625–6633 (2017).
367. Bansal, R. *et al.* Polybrominated diphenyl ether (DE-71) interferes with thyroid hormone action independent of effects on circulating levels of thyroid hormone in male rats. *Endocrinology* **155**, 4104–4112 (2014).
368. Dong, W., Macaulay, L. J., Kwok, K. W. H., Hinton, D. E. & Stapleton, H. M. Using whole mount in situ hybridization to examine thyroid hormone deiodinase expression in embryonic and larval zebrafish: A tool for examining OH-BDE toxicity to early life stages. *Aquat. Toxicol.* **132–133**, 190–199 (2013).
369. Geiss, O., Tirendi, S., Barrero-Moreno, J. & Kotzias, D. Investigation of volatile organic compounds and phthalates present in the cabin air of used private cars. *Environ. Int.* **35**, 1188–1195 (2009).
370. Eckert, E. *et al.* Comparative study on the migration of di-2-ethylhexyl phthalate (DEHP) and tri-2-ethylhexyl trimellitate (TOTM) into blood from PVC tubing material of a heart-lung machine. *Chemosphere* **145**, 10–16 (2016).
371. EFSA. Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) on a request from the Commission related to a 5th list of substances for food contact materials. *EFSA J.* **2**, 109 (2004).
372. (STD/1259), N. E. *1,2-Cyclohexanedicarboxylic acid, 1,2-diisononyl ester ('Hexamol DINCH')*. (2012).
373. Blount, B. C. *et al.* Levels of seven urinary phthalate metabolites in a human reference population.

- Environ. Health Perspect.* **108**, 979–982 (2000).
374. Adibi, J. J. *et al.* Prenatal exposures to Phthalates among women in New York and Krakow, Poland. *Environ. Health Perspect.* **111**, 1719–1722 (2003).
375. Latini, G. *et al.* In utero exposure to di-(2-ethylhexyl)phthalate and duration of human pregnancy. *Environ. Health Perspect.* **111**, 1783–1785 (2003).
376. Silva, M. J. *et al.* Detection of phthalate metabolites in human amniotic fluid. *Bull. Environ. Contam. Toxicol.* **72**, 1226–1231 (2004).
377. Calafat, A. M. *et al.* Urinary and amniotic fluid levels of phthalate monoesters in rats after the oral administration of di(2-ethylhexyl) phthalate and di-n-butyl phthalate. *Toxicology* **217**, 22–30 (2006).
378. Kim, Y. *et al.* Prenatal exposure to phthalates and infant development at 6 months: Prospective mothers and children’s environmental health (MOCEH) study. *Environ. Health Perspect.* **119**, 1495–1500 (2011).
379. Polanska, K., Ligocka, D., Sobala, W. & Hanke, W. Phthalate exposure and child development: The Polish Mother and Child Cohort Study. *Early Hum. Dev.* **90**, 477–485 (2014).
380. Téllez-Rojo, M. M. *et al.* Prenatal urinary phthalate metabolites levels and neurodevelopment in children at two and three years of age. *Sci. Total Environ.* **461–462**, 386–390 (2013).
381. Whyatt, R. M. *et al.* Maternal prenatal urinary phthalate metabolite concentrations and child mental, psychomotor, and behavioral development at 3 years of age. *Environ. Health Perspect.* **120**, 290–295 (2012).
382. Minatoya, M. *et al.* Effects of prenatal phthalate exposure on thyroid hormone levels, mental and psychomotor development of infants: The Hokkaido Study on Environment and Children’s Health. *Sci. Total Environ.* **565**, 1037–1043 (2016).
383. Swan, S. H. *et al.* Prenatal phthalate exposure and reduced masculine play in boys. in *International Journal of Andrology* **33**, 259–267 (2010).
384. Miodovnik, A. *et al.* Endocrine disruptors and childhood social impairment. *Neurotoxicology* **32**, 261–267 (2011).

385. Cho, S. C. *et al.* Relationship between environmental phthalate exposure and the intelligence of school-age children. *Environ. Health Perspect.* **118**, 1027–1032 (2010).
386. Chopra, V., Harley, K., Lahiff, M. & Eskenazi, B. Association between phthalates and attention deficit disorder and learning disability in U.S. children, 6-15 years. *Environ. Res.* **128**, 64–69 (2014).
387. Huang, P. C., Kuo, P. L., Guo, Y. L., Liao, P. C. & Lee, C. C. Associations between urinary phthalate monoesters and thyroid hormones in pregnant women. *Hum. Reprod.* **22**, 2715–2722 (2007).
388. Boas, M. *et al.* Childhood exposure to phthalates: Associations with thyroid function, insulin-like growth factor I, and growth. *Environ. Health Perspect.* **118**, 1458–1464 (2010).
389. Huang, P. C. *et al.* Early phthalates exposure in pregnant women is associated with alteration of thyroid hormones. *PLoS One* **11**, (2016).
390. Arbuckle, T. E. *et al.* Phthalate and bisphenol A exposure among pregnant women in Canada - Results from the MIREC study. *Environ. Int.* **68**, 55–65 (2014).
391. Cantonwine, D. E. *et al.* Urinary phthalate metabolite concentrations among pregnant women in Northern Puerto Rico: Distribution, temporal variability, and predictors. *Environ. Int.* **62**, 1–11 (2014).
392. Swan, S. H. *et al.* First trimester phthalate exposure and anogenital distance in newborns. *Hum. Reprod.* **30**, 963–972 (2015).
393. Meeker, J. D., Calafat, A. M. & Hauser, R. Di(2-ethylhexyl) Phthalate metabolites may alter thyroid hormone levels in men. *Environ. Health Perspect.* **115**, 1029–1034 (2007).
394. Wu, M. T. *et al.* Intake of Phthalate-Tainted Foods Alters Thyroid Functions in Taiwanese Children. *PLoS One* **8**, (2013).
395. De Cock, M., De Boer, M. R., Lamoree, M., Legler, J. & Van De Bor, M. Prenatal exposure to endocrine disrupting chemicals (EDCS) in relation to thyroid hormone levels in infants. *Arch. Dis. Child.* **99**, A362 (2014).
396. Breous, E., Wenzel, A. & Loos, U. The promoter of the human sodium/iodide symporter responds to certain phthalate plasticisers. in *Molecular and Cellular Endocrinology* **244**, 75–78 (2005).
397. Shen, O. *et al.* Comparison of in vitro hormone activities of selected phthalates using reporter gene

- assays. *Toxicol. Lett.* **191**, 9–14 (2009).
398. Zhai, W. *et al.* Thyroid endocrine disruption in zebrafish larvae after exposure to mono-(2-ethylhexyl) phthalate (MEHP). *PLoS One* **9**, (2014).
399. Liu, C., Zhao, L., Wei, L. & Li, L. DEHP reduces thyroid hormones via interacting with hormone synthesis-related proteins, deiodinases, transthyretin, receptors, and hepatic enzymes in rats. *Environ. Sci. Pollut. Res.* **22**, 12711–12719 (2015).
400. Houde, M., Martin, J. W., Letcher, R. J., Solomon, K. R. & Muir, D. C. G. Biological monitoring of polyfluoroalkyl substances: A review. *Environmental Science and Technology* **40**, 3463–3473 (2006).
401. Spliethoff, H. M. *et al.* Use of newborn screening program blood spots for exposure assessment: Declining levels of perfluorinated compounds in New York State infants. *Environ. Sci. Technol.* **42**, 5361–5367 (2008).
402. Olsen, G. W. *et al.* Decline in perfluorooctanesulfonate and other polyfluoroalkyl chemicals in American red cross adult blood donors, 2000–2006. *Environ. Sci. Technol.* **42**, 4989–4995 (2008).
403. Kato, K., Wong, L. Y., Jia, L. T., Kuklennyik, Z. & Calafat, A. M. Trends in exposure to polyfluoroalkyl chemicals in the U.S. population: 1999–2008. *Environ. Sci. Technol.* **45**, 8037–8045 (2011).
404. Olsen, G. W. *et al.* Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in retired fluorochemical production workers. *Environ. Health Perspect.* **115**, 1298–1305 (2007).
405. Hu, X. C. *et al.* Detection of Poly- and Perfluoroalkyl Substances (PFASs) in U.S. Drinking Water Linked to Industrial Sites, Military Fire Training Areas, and Wastewater Treatment Plants. *Environ. Sci. Technol. Lett.* **3**, 344–350 (2016).
406. Johansson, N., Fredriksson, A. & Eriksson, P. Neonatal exposure to perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) causes neurobehavioural defects in adult mice. *Neurotoxicology* **29**, 160–169 (2008).
407. Slotkin, T. A., MacKillop, E. A., Meinick, R. L., Thayer, K. A. & Seidler, F. J. Developmental neurotoxicity of perfluorinated chemicals modeled in vitro. *Environ. Health Perspect.* **116**, 716–722

- (2008).
408. Gizer, I. R., Ficks, C. & Waldman, I. D. Candidate gene studies of ADHD: A meta-analytic review. *Hum. Genet.* **126**, 51–90 (2009).
409. Faraone, S. V. *et al.* Molecular genetics of attention-deficit/hyperactivity disorder. *Biol. Psychiatry* **57**, 1313–1323 (2005).
410. Hoffman, K., Webster, T. F., Weisskopf, M. G., Weinberg, J. & Vieira, V. M. Exposure to polyfluoroalkyl chemicals and attention deficit/hyperactivity disorder in U.S. children 12-15 years of age. *Environ. Health Perspect.* **118**, 1762–1767 (2010).
411. Stein, C. R. & Savitz, D. A. Serum perfluorinated compound concentration and attention deficit/hyperactivity disorder in children 5-18 years of age. *Environ. Health Perspect.* **119**, 1466–1471 (2011).
412. Gump, B. B., Wu, Q., Dumas, A. K. & Kannan, K. Perfluorochemical (PFC) exposure in children: Associations with impaired response inhibition. *Environ. Sci. Technol.* **45**, 8151–8159 (2011).
413. Fei, C., McLaughlin, J. K., Lipworth, L. & Olsen, J. Prenatal exposure to perfluorooctanoate (PFOA) and perfluorooctanesulfonate (PFOS) and maternally reported developmental milestones in infancy. *Environ. Health Perspect.* **116**, 1391–1395 (2008).
414. Fei, C. & Olsen, J. Prenatal exposure to perfluorinated chemicals and behavioral or coordination problems at age 7 years. *Environ. Health Perspect.* **119**, 573–578 (2011).
415. Endendijk, J. J., Wijnen, H. A. A., Pop, V. J. M. & van Baar, A. L. Maternal thyroid hormone trajectories during pregnancy and child behavioral problems. *Horm. Behav.* **94**, 84–92 (2017).
416. Modesto, T. *et al.* Maternal Mild Thyroid Hormone Insufficiency in Early Pregnancy and Attention-Deficit/Hyperactivity Disorder Symptoms in Children. *JAMA Pediatr.* **169**, 838–845 (2015).
417. Hauser, P. *et al.* Attention Deficit-Hyperactivity Disorder in People with Generalized Resistance to Thyroid Hormone. *N. Engl. J. Med.* **328**, 997–1001 (1993).
418. Pearce, E. N. Maternal Hypothyroxinemia in Pregnancy Is Associated with Increased Risk for ADHD Symptoms in Children. *Clin. Thyroidol.* **27**, 212–214 (2015).


419. Olsen, G. W. & Zobel, L. R. Assessment of lipid, hepatic, and thyroid parameters with serum perfluorooctanoate (PFOA) concentrations in fluorochemical production workers. *Int. Arch. Occup. Environ. Health* **81**, 231–246 (2007).
420. Melzer, D., Rice, N., Depledge, M. H., Henley, W. E. & Galloway, T. S. Association between serum perfluorooctanoic acid (PFOA) and thyroid disease in the U.S. National Health and Nutrition Examination Survey. *Environ. Health Perspect.* **118**, 686–692 (2010).
421. Chang, S. C. *et al.* Negative bias from analog methods used in the analysis of free thyroxine in rat serum containing perfluorooctanesulfonate (PFOS). *Toxicology* **234**, 21–33 (2007).
422. Martin, M. T. *et al.* Toxicogenomic study of triazole fungicides and perfluoroalkyl acids in rat livers predicts toxicity and categorizes chemicals based on mechanisms of toxicity. *Toxicol. Sci.* **97**, 595–613 (2007).
423. Seacat, A. M. *et al.* Sub-chronic dietary toxicity of potassium perfluorooctanesulfonate in rats. *Toxicology* **183**, 117–131 (2003).
424. Thibodeaux, J. R. *et al.* Exposure to perfluorooctane sulfonate during pregnancy in rat and mouse. I: Maternal and prenatal evaluations. *Toxicological Sciences* **74**, 369–381 (2003).
425. Lau, C. *et al.* Exposure to perfluorooctane sulfonate during pregnancy in rat and mouse. II: Postnatal evaluation. *Toxicological Sciences* **74**, 382–392 (2003).
426. Yu, W. G. *et al.* Prenatal and postnatal impact of perfluorooctane sulfonate (PFOS) on rat development: A cross-foster study on chemical burden and thyroid hormone system. *Environ. Sci. Technol.* **43**, 8416–8422 (2009).
427. Luebker, D. J., York, R. G., Hansen, K. J., Moore, J. A. & Butenhoff, J. L. Neonatal mortality from in utero exposure to perfluorooctanesulfonate (PFOS) in Sprague-Dawley rats: Dose-response, and biochemical and pharmacokinetic parameters. *Toxicology* **215**, 149–169 (2005).
428. Weiss, J. M. *et al.* Competitive binding of poly- and perfluorinated compounds to the thyroid hormone transport protein transthyretin. *Toxicol. Sci.* **109**, 206–216 (2009).
429. Yu, W. G., Liu, W. & Jin, Y. H. Effects of Perfluorooctane Sulfonate on Rat Thyroid Hormone

- Biosynthesis and Metabolism. *Environ. Toxicol. Chem.* **28**, 1 (2008).
430. Delfosse, V. *et al.* Synergistic activation of human pregnane X receptor by binary cocktails of pharmaceutical and environmental compounds. *Nat. Commun.* **6**, 8089 (2015).
431. Silva, E., Rajapakse, N. & Kortenkamp, A. Something from ‘nothing’ - Eight weak estrogenic chemicals combined at concentrations below NOECs produce significant mixture effects. *Environ. Sci. Technol.* **36**, 1751–1756 (2002).
432. Fini, J.-B. *et al.* Human amniotic fluid contaminants alter thyroid hormone signalling and early brain development in *Xenopus* embryos. *Sci. Rep.* **7**, 43786 (2017).
433. Birgersson, L. *et al.* From Cohorts to Molecules: Adverse Impacts of Endocrine Disrupting Mixtures. *bioRxiv* 206664 (2017). doi:10.1101/206664

Figure legend

Figure 1: Endocrine disrupting chemicals (EDCs) action at multiple levels of HPT axis.

Environmental chemicals have the potential to disrupt hypothalamus-pituitary thyroid (HPT) axis, alone or in combination. Given the crucial role for thyroid hormone in brain development, such disruption can have a long lasting functional impact such as IQ loss and increased risk of neurodevelopmental disease. (Note: targets not drawn to scale).



Endocrine disrupting chemicals (EDCs) action at multiple levels of HPT axis. Environmental chemicals have the potential to disrupt hypothalamus-pituitary thyroid (HPT) axis, alone or in combination. Given the crucial role for thyroid hormone in brain development, such disruption can have a long lasting functional impact such as IQ loss and increased risk of neurodevelopmental disease. (Note: targets not drawn to scale).

338x190mm (300 x 300 DPI)