



HAL
open science

Fish metabolome from sub-urban lakes of the Paris area (France) and potential influence of noxious metabolites produced by cyanobacteria

Benjamin Marie, Alison Gallet

► To cite this version:

Benjamin Marie, Alison Gallet. Fish metabolome from sub-urban lakes of the Paris area (France) and potential influence of noxious metabolites produced by cyanobacteria. *Chemosphere*, 2022, 296, pp.134035. 10.1016/j.chemosphere.2022.134035 . hal-03922408

HAL Id: hal-03922408

<https://hal.science/hal-03922408>

Submitted on 4 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Fish metabolome from sub-urban lakes of the Paris area (France)**
2 **and potential influence of noxious metabolites produced by**
3 **cyanobacteria**

4
5 *Benjamin MARIE^{1,*}, Alison GALLET¹*

6
7 ¹ UMR 7245, CNRS/MNHN, Molécules de Communication et Adaptation des Micro-organismes (MCAM), équipe
8 "Cyanobactéries, Cyanotoxines et Environnement", 12 rue Buffon - CP 39, 75231 Paris Cedex 05, France.

9
10 * Corresponding author: benjamin.marie@mnhn.fr

11
12 **Abstract**

13 The recent democratization of high-throughput molecular phenotyping allows the rapid
14 expansion of promising untargeted multi-dimensional approaches (*e.g.* epigenomics,
15 transcriptomics, proteomics, and/or metabolomics). Indeed, these emerging omics tools,
16 processed for ecologically relevant species, may present innovative perspectives for
17 environmental assessments, that could provide early warning of eco(toxico)logical
18 impairments.

19 In a previous pilot study ([Sotton et al., Chemosphere 2019](#)), we explore by ¹H NMR the bio-
20 indicative potential of metabolomics analyses on the liver of 2 sentinel fish species (*Perca*
21 *fluviatilis* and *Lepomis gibbosus*) collected in 8 water bodies of the peri-urban Paris' area
22 (France). In the present study, we further investigate on the same samples the high potential
23 of high-throughput UHPLC-HRMS/MS analyses. We show that the LC-MS metabolome
24 investigation allows a clear separation of individuals according to the species, but also
25 according to their respective sampling lakes. Interestingly, similar variations of *Perca* and
26 *Lepomis* metabolomes occur locally indicating that site-specific environmental constraints
27 drive the metabolome variations which seem to be influenced by the production of noxious
28 molecules by cyanobacterial blooms in certain lakes. Thus, the development of such reliable
29 environmental metabolomics approaches appears to constitute an innovative bio-indicative
30 tool for the assessment of ecological stress, such as toxigenic cyanobacterial blooms, and
31 aim at being further follow up.

32
33 **Keywords:** Untargeted metabolomics, environmental assessment, cyanobacteria,
34 biomarkers, sentinel species

35

36 1. INTRODUCTION

37 In the past decades, the use of aquatic organisms as bio-indicator has been widely adopted
38 to monitor and to manage the quality of water bodies worldwide (Lorenz 2003). In parallel,
39 the difficulties and the rational limits of a direct chemical monitoring for providing sufficient
40 information to adequately assess the risks from anthropogenic chemicals in the natural
41 environment was progressively documented (Reyjol et al., 2014). More recently, the recent
42 democratisation of high-throughput molecular phenotyping has allowed the rapid expansion
43 of promising untargeted multi-dimensional approaches (e.g. epigenomics, transcriptomics,
44 proteomics, metabolomics), that now represent promising perspectives for environmental
45 assessments (Bahamonde et al. 2016; Cordier et al., 2020). It appears now very likely that
46 emerging omics-based analyses, such as metabolomics, developed for ecologically relevant
47 aquatic species, may support the development of new ecological evaluation tools, that could
48 provide early warning of ecotoxicological impairments (Marie 2020).

49 The metabolome is defined as the set of primary and secondary metabolites synthesized at a
50 given time and thus represents its global metabolic chemical picture. This set of metabolites
51 is assumed to represent a final endpoint of the phenotypic response of an organism that can
52 potentially be modified when the ecological and environmental stress conditions change
53 (Bundy et al., 2008). In this way, metabolomic studies have become a relevant approach to
54 investigate physiological processes involved in the homeostatic responses of the organisms
55 encountering environmental stresses from potentially multiple origins (Capello et al., 2016;
56 Good et al., 2020). However, despite its high potential to support the understanding the
57 molecular mechanisms implicated in the ecotoxicological responses of aquatic organisms
58 (Sardans et al., 2011; Viant, 2008), such environmental metabolomics investigations remains
59 still limited. Indeed, numerous studies have compared the metabolite profiles of individuals
60 collecting from a limited set of environments, often including only one contaminated site
61 and a unique reference location (e.g. Capello et al., 2016; Melvin et al., 2019; De Marco et
62 al., 2022). As previously suggested, the selection of a unique “good” reference site may
63 crucially orientate the results of the analyses, and larger efforts should be provided at
64 understanding similarities and differences among various references sites (Simmons et al.,
65 2015; Marjan et al., 2017). Such approaches would gain at investigating larger sets of
66 environments regarding their respective positions within environmental gradients of global
67 ecotoxicological pressures (Jeffries et al., 2010; Davis et al., 2016; Meador et al., 2020).

68
69 Among several aquatic organisms, such as molluscs or arthropods, fishes present remarkable
70 features that make them especially relevant for bio-indication purposes. In fact, they occupy
71 almost any aquatic habitat, being capable of experiencing different and variable
72 environmental conditions (Harris 1995). Fishes are also responsive to numerous abiotic
73 (temperatures, water velocities, sediment loads, hypoxia, ...) or biotic (famine, predation,
74 parasitism, ...) pressures, together with anthropogenic stressors, such as contaminants, that
75 represent additional constraints that fish may experience in disturbed ecosystems
76 (McClanahan 2019). Environmental stressors are known to affect the specific physiological,

77 developmental and reproductive traits, that allow the different fish species to colonize and
78 occupy various ecosystems (Hamilton et al., 2016).
79 Nowadays, cyanobacterial blooms are one of the common sources of stress encountered by
80 organisms living in modern freshwater lentic systems around the world (Harke et al., 2016;
81 Burford et al., 2020). Then, various efforts have been attempted to characterize the
82 potential effect of cyanobacteria proliferation on various fish species according to
83 experimentation performed in the lab, in mesocosms, or from field sampling (Amado et al.,
84 2010; Ferrao-Filho et al., 2011; Sotton et al., 2017a, 2017b; Le Manach et al., 2018). To this
85 end, we initially investigated by ¹H NMR technics the liver metabolome of fish collected from
86 different lakes of the peri-urban Paris area (France), and encountering low or high
87 cyanobacterial proliferation (Sotton et al., 2019). This pilot study leads to the description of
88 putative metabolite concentrations, corresponding to specific NMR signals, that were
89 significantly correlated with higher cyanobacterial occurrence.

90
91 The present study investigates the bio-indicative potential of high-throughput molecular
92 phenotyping supported by LC-HRMS approaches on 2 sentinel fish specimens (*Perca*
93 *fluviatilis* and *Lepomis gibbosus*) that were previously investigated by ¹H NMR (Sotton et al.,
94 2019). We hypothesis that LC-MS-based metabolomics approach may present a higher
95 potential for environmental diagnosis. Indeed, such investigations provide valuable multi-
96 metric characters indicative of biological or ecological conditions, together with a
97 prospective mechanistic under-standing of the cause-effect relationships. With regard to the
98 cyanobacterial occurrence and the associated production of bioactive metabolites in the
99 different lakes, we compare here the metabolome of two different sentinel fish species.
100 Indeed, these organisms could present rather different or comparable sensitivity and
101 responsiveness to local eco(toxico)logical constraints (*i.e.* potentially the production of
102 noxious molecules by cyanobacteria), and may support the general interest for omics-based
103 environmental approach for the characterization of cyanobacterial bloom threat.

104
105

106 **2. MATERIALS AND METHODS**

107 **2.1. Fish sampling**

108 Field sampling campaigns were performed during late summer 2015 (7-10th September), as
109 this “end-of-the-bloom” period appears more appropriate to observe cyanobacterial effects
110 on fishes. Eight peri-urban pounds around Paris’ area (Île-de-France region, France) were
111 selected for their respective eutrophication levels and the presence or the absence of
112 recurrent cyanobacterial blooms: Cergy-Pontoise (Cer), Champs-sur-Marne (Cha), Maurepas
113 (Mau), Rueil (Rue), Verneuil (Ver), Varennes-sur-Seine (Var), Fontenay-sur-Loing (Fon) and
114 Triel (Tri) pounds (Supplementary figure S1) (Maloufi et al., 2016; Sotton et al., 2019). These
115 sites were sampled by the Hydrosphère company (www.hydrosphere.fr) with electric fishing
116 device (FEG 8000, EFKO, Leutkirch, Germany) for capturing live caught fish, that were then
117 promptly sacrificed for liver collection. The investigation of the fish guild indicates that only

118 the perch (*Perca fluviatilis*) and pumpkinseed sunfish (*Lepomis gibbosus*) were presents in all
119 or almost all these pounds ([supplementary table S1](#)) and were further selected as sentinel
120 species for further molecular phenotyping analyses by metabolomics.

121 Briefly, live caught fishes (n=5-10 young-of-the-year per pounds and per species) were
122 briefly euthanized by neck dislocation, directly measured (12.0±4.8 cm), weighed (9.3±2.6
123 g), and then liver of each individual was shortly sampled, flash-frozen in liquid nitrogen and
124 kept at -80°C until analyses, in accordance with European animal ethical concerns and
125 regulations.

126 In every lake, sub-surface chlorophyll-*a* equivalent concentrations attributed to the four-
127 main phytoplankton groups (Chlorophyta, Diatoms, Cyanobacteria and Cryptophyta) were
128 measured with an *in-situ* fluorometer (Fluoroprobe II, Bbe-Moldenke, Germany)
129 ([Supplementary figure S2](#)). Sub-surface water samples filtered on 20- μ m mesh size were also
130 collected for phytoplanktonic community analysis and further metabolomics
131 characterisation, and then kept at -80°C until analysis as previously described ([Sotton et al.,](#)
132 [2019](#)). Physico-chemical parameters of the water column (temperature, conductivity, pH,
133 dissolved O₂ and nitrate concentrations) were monitored with a YSI DSS probe, as previously
134 described ([Sotton et al., 2019](#)).

135

136 **2.2. Liver metabolite extraction and metabolomics analyses**

137 The liver extraction was performed on 132 individuals (comprising 78 *Perca* and 54 *Lepomis*)
138 with methanol/chloroform/water (ratio 2/2/1.8 – 22 mL.g⁻¹ at 4°C) and the polar fraction
139 was analyzed on a 600-MHz NMR spectrometer equipped with a 5-mm cryoprobe (Advance
140 III HD Bruker, Germany) with a noesygppr1d pulse sequence as previously described ([Sotton](#)
141 [et al., 2019](#)). ¹H-NMR spectra were treated with Batman R-package for deconvolution, peak
142 assignment and quantification of 222 putative metabolites ([Hoa et al., 2014](#)).

143 The liver extracted polar phase was additionally injected (2 μ L) on C₁₈ column (Polar
144 Advances II 2.5 pore - Thermo), then eluted at a 300 μ L.min⁻¹ flow rate with a linear gradient
145 of acetonitrile in 0.1% formic acid (5 to 90 % in 21 min) with an ultra-high-performance
146 liquid chromatography (UHPLC) system (ELUTE, Bruker). Consecutively, the individual
147 metabolite contents were analysed using an electrospray ionization hybrid quadrupole time-
148 of-flight (ESI-Qq-TOF) high-resolution mass spectrometer (Compact, Bruker) at 2 Hz speed on
149 positive MS mode on the 50–1500 *m/z* range. The feature peak list was generated from
150 recalibrated MS spectra (< 0.5 ppm for each sample, as an internal calibrant of Na formate
151 was injected at the beginning of each sample analysis) within a 1-15 min window of the LC
152 gradient, with a filtering of 5,000 count of minimal intensity, a minimal occurrence in at least
153 50% of all samples, and combining all charge states and related isotopic forms using
154 MetaboScape 4.0 software (Bruker).

155 Additionally, five pools of six different individuals randomly selected for *Perca* and *Lepomis*
156 (quality check samples) and the phytoplanktonic biomass extracts from the 8 lakes were
157 similarly eluted then analysed on positive autoMS/MS mode at 2-4 Hz on the 50-1500 *m/z*
158 range for further metabolite annotation. Molecular networks were performed with Global

159 Natural Product Social network (<http://gnps.ucsd.edu>) and/or MetGem
160 (<http://metgem.github.io>) softwares, as previously described (Kim Tiam et al., 2019; Le
161 Manach et al., 2019) for cyanobacterial and fish metabolite annotation was assayed using in-
162 house (Le Manach et al., 2019) or CyanoMetDB (Jones et al., 2021), and GNPS, Mona, human
163 metabolome database (HMDB) and Massbank MS/MS spectral libraries. Water samples of
164 each lake, concentrated on a 20- μ m mesh size, were also extracted with 75% methanol (2
165 min sonication, 5 min centrifugation at 15,000 g - 4°C) and then similarly analysed in
166 triplicates on LC-HRMS on autoMS/MS positive mode, as earlier described, and metabolite
167 list annotated with same pipelines.

168

169 **2.4. Data matrix treatment**

170 The resulting the intensity data tables of the 222 metabolites (^1H NMR) and 1252 analytes
171 (LC-MS) were further treated for normalization with Pareto's scaling, inspected data
172 representation by PCA and heatmap with hierarchical clustering (Euclidean distance) and
173 then analysed to investigate the influences of "Species" and "Lakes" parameters on the
174 datasets by PERMANOVA, PLS-DA or multi-block PLS-DA using MixOmics and Mixomics
175 DIABLO R Package (Rohart et al., 2017; Singh et al., 2019), MetaboAnalyst 5 (Chong et al.,
176 2019) and MicrobiomeAnalyst tools (Chong et al., 2020).

177

178

179 **3. RESULTS AND DISCUSSION**

180 **3.1. Distinction of *Perca* and *Lepomis* liver LC-MS metabolome**

181 [Figure 1](#) illustrates the global metabolite profiles analyzed by LC-MS for each fish (comprising
182 *Perca* and *Lepomis*) on a heatmap with hierarchical classifications and individual plot
183 representation of principal component analyses (PCA). A very strong and significant
184 discrimination between the "species" was achieved (PERMANOVA F-value = 94.92; $R^2 = 0.43$;
185 P-value <0.001), when a slighter but still significant discrimination can be also retrieved for
186 "lakes" (PERMANOVA F-value = 4.96; $R^2 = 0.22$; P-value <0.001).

187 In parallel, a molecular network performed with fragmentation data was obtained by LC-
188 MS/MS analyses of *Perca* and *Lepomis* metabolomes. This approach supports the global
189 annotation process of the fish liver metabolites considering both structural identities and
190 similarities (Nothias et al., 2020). The GNPS and the *t*-SNE representation of the molecular
191 network highlights that most, if not all, of the known metabolites (presenting structural
192 identity or analogy hits) are largely shared between the two species (*e.g.* nucleic acids,
193 carnitines, glutathiones, lipids, saccharides, ...). Moreover, numerous species-specific cluster
194 metabolites remain uncharacterized, with relative molecules and clusters presenting no
195 match within the public chemical databases ([supplementary figure S2](#)) (da Silva et al., 2015).
196 Taken together, these analyses illustrate for these two species the various molecular
197 specificities of their respective metabolomes. It also shows the substantial portion of the
198 specific liver molecular metabolism that still remains uncharacterized and that could be

199 related to species-specific nutritional, physiological or toxicological functions (Brusle et al.,
200 2017).

201

202 **3.2. Comparison of *Perca* and *Lepomis* LC-MS metabolome between lakes**

203 In addition, the LC-MS datasets of the two fish species considered separately show even
204 clearer discriminations of the metabolomes of fish originating from the different lakes. The
205 un-supervised PCA and the heatmap with hierarchical classification (Figure 2) present
206 distinct environmental signatures that support reliable discrimination of the lake of sampling
207 for both *Perca* and *Leptomis* (Fig. 2A-B and 2C-D, respectively). These observations strongly
208 suggest that the locality seems to globally influence the local LC-MS metabolome
209 composition for both species. However, the liver metabolomes of individuals from Fontenay-
210 sur-Loing lake (Fon) exhibit quite species-specific relative positions on PCA (Figure 2A and
211 2C). Such differences of the relative position of metabolomes from Fon between the two
212 species suggests that: *i/* for most localities, the specific “lake” signature of the metabolome
213 appears in good agreement between the two species, *ii/* for some specific environments
214 (e.g. Fon), the metabolome signatures are variable from one species to the other one. Thus,
215 the species-specific signature of the *Perca* and *Lepomis* metabolomes in Fon lake may rather
216 traduce the specific responsiveness and/or sensitivity to a local factor occurring at this
217 particular location.

218

219 Interestingly, the two hierarchical classifications performed on LC-MS metabolome datasets
220 of *Perca* and *Lepomis*, respectively, overall show a global similarity of the metabolome of
221 fish originating from the same lake, together with an overall distinction of 3 principal lake
222 clusters. Indeed, these analyses show very similar lake relationships for both species,
223 grouping together all fish from the lakes of ‘Cer and Mau’, from ‘Cha, Fon and Rue’, and from
224 Tri, Var and Ver’ for *Perca*, or ‘Tri and Ver’ for *Lepomis*, as this latter was not retrieved in Var
225 (Figure 2B and 2D). This observation suggests that these localities could somehow present
226 comparable, if not analogous, environmental conditions. This hypothesis was firstly
227 considered regarding NMR metabolomics and potential explicative environmental factors
228 (physico-chemical and biological factors – supplementary table S2) and had suggested a
229 potential influence of cyanobacterial proliferation (BBE_{cya} and pH) (Sotton et al., 2019).
230 Indeed, as initially shown, the Fon, Tri, Var and Ver lakes encounter remarkable high
231 amounts of cyanobacteria belonging to the *Planktothrix*, *Pseudo-Anabaena*, *Aphanizomenon*,
232 *Anabaena* or *Microcystis* genera. Differently, the other lakes (Cer, Mau, Cha and Rue) exhibit
233 phytoplanktonic biomasses principally dominated by green algae, diatoms or cryptophytes
234 (Fig. 3A-B). A molecular network performed with LC-MS/MS data obtained from the
235 phytoplanktonic biomass of these 8 lakes shows that noticeable amounts of cyanopeptides
236 are observed in the water column of the cyanobacterial-rich lakes (Fon, Tri, Var and Ver),
237 with the specific presence of most noxious cyanopeptides (microcystins, cyanopeptolins and
238 anabaenopeptins) being detected in Tri, Var and Ver lakes. In a different manner, Fon water
239 exhibits rather less toxic cyanopeptides (Janssen 2019), such as microginins and aeruginosins

240 (Fig. 3C). Taken together, these observations suggest that the proliferation and the
241 subsequent production of such potentially noxious cyanobacterial metabolites could be also
242 considered as an explicative factor of the local metabolome singularity for the liver of *Perca*
243 and *Lepomis* from Tri, Var and Ver lakes, as previously postulated (Sotton et al., 2019).

244

245 **3.3. Discriminating metabolites between the different lake groups**

246 Thus, we further investigate the fish metabolome distinctions considering together fishes
247 from lakes presenting noxious cyanobacterial metabolites (“Tri/Var/Ver” group for *Perca* and
248 “Tri/Ver” group for *Lepomis*), or not (“Cer/Mau/Cha/Fon/Rue” group). For this purpose, we
249 used PLS-DA to identify metabolites that discriminate the individuals belonging to these two
250 lake groups (Figure 4). Indeed, while unsupervised PCA were first used to evaluate the global
251 dispersion between species and sampling groups, a supervised model such as partial least
252 square differential analysis (PLS-DA) allows to maximize the separation between the
253 different sample classes (e.g. “lake groups”) and to further provide information on
254 discriminating features (variable importance on projection - VIP). For both fish species, the
255 metabolomes present a consistent discriminant analysis scoring (*Perca*: accuracy = 1,
256 performance $R^2 = 0.78$, predictability $Q^2 = 0.72$; *Lepomis*: accuracy = 0.93, $R^2 = 0.59$, $Q^2 =$
257 0.29) and large sets of discriminating variables (*Perca* $VIP_{>1.5}=109$ and *Lepomis* $VIP_{>1.5}=93$)
258 (Figure 4A-B). Among these metabolites, 33 components have been successfully annotated
259 considering their respective molecular mass, the isotopic pattern, together with the
260 fragmentation pattern of all components with GNPS and MetGem algorithms and a manual
261 curation. Overall, the metabolite semi-quantification of the annotated best VIP reveals
262 largely similar variations between *Perca* and *Lepomis* for these two groups of lakes (Figure
263 4C; Supp. fig. S3). Indeed, except for the reduced-glutathione, which present higher
264 concentrations in the livers of *Perca* from “Tri/Var/Ver” lakes, when livers of *Lepomis*
265 exhibits lower concentrations than in the those of fish of the “Cer/Mau/Cha/Fon/Rue”
266 group, all other metabolites present similar quantification patterns between the two
267 species. This discrepancy suggests that glutathione-dependent redox mechanisms could be
268 more extensive in *Perca* than in *Lepomis* livers.

269 Globally, the main differences in the different metabolite concentration between these two
270 lake groups show that fishes collected from the “Cer/Mau/Cha/Fon/Rue” present higher
271 energetic, anti-oxidant/detoxification and lipids reserves (e.g. various nucleic acids,
272 glutathiones, flavonoids and lipo-phosphocholines) than in “Tri/Var/Ver” lakes. In addition,
273 higher amounts of amino acids, di-peptides and acylcarnitines in the liver of fish originating
274 from the “Tri/Var/Ver” lakes could be related to higher protein catabolism and beta-
275 oxidation, that could denote of higher energetic requirements for those specific organisms.
276 Interestingly, the “Tri/Var/Ver” lakes present much more cyanobacteria and related noxious
277 metabolites, that may represent a patent stress source for fishes colonizing those
278 environments. Taken together, these observations are in agreement to the hypothesis of an
279 occurrence of higher stress conditions locally occurring in the Tri, Var and Ver lakes, that
280 could be associated with the local production of potentially noxious metabolites by

281 cyanobacterial proliferation (Sotton et al., 2017a, 2017b; Le Manach et al., 2018).

282

283 In general, various genetic or phenotypic factors (such as development stages,
284 contamination levels, predator/parasite pressures, food availability, ...) could influence and
285 explain the local discrepancy of the metabolome of fishes collected from different
286 environments (reviewed in Marie 2020). However, in the present case, the fact that *Perca*
287 and *Lepomis* metabolomes analyzed in parallel present similar metabolite variations clearly
288 indicates that local environmental constraints may drive such phenotypic co-variations.
289 Then, these present results claim for the high informativeness of LC-MS-based metabolomic
290 imprints for molecular phenotyping on sentinel species and the exploration of
291 environmental stresses (Lohr et al., 2019).

292

293 **3.4. Comparison of ¹H NMR and LC-MS metabolomics for fish livers**

294 Considering the previous ¹H NMR metabolomics datasets retrieved from the same samples
295 on these two fish species (Sotton et al., 2019), the liver metabolome analyzed by NMR
296 shows more limited specificity according to “species” or “lakes” factor than those analyzed
297 by LC-MS. Although the 222 potential metabolites quantified by NMR on 132 fish (78 *Perca*
298 and 54 *Lepomis*) present no obvious structuration of the individual dataset according to
299 “species” or “lakes” variables, when observed on PCA or hierarchical classification, slight but
300 significant relations are however supported by PERMANOVA (F-values = 4.15 and 3.63; R² =
301 0.03 and 0.17; P-values < 0.01 and 0.01, for “species” and “lakes” respectively, Supp. fig. S4).
302 Indeed, according to the relatively lower performances of the hierarchical classifications and
303 PERMANOVA and the PLS-DA (Supp. fig. S4 and S5), the NMR liver metabolomes exhibit a
304 limited discrimination on “lakes” for both species, when compared to LC-MS metabolomics
305 performances.

306 Overall, regarding both hierarchical classification, individual plots of PCA or PERMANOVA
307 performances, the dataset obtained from the LC-MS metabolome analysis presents a clearer
308 discriminant pattern between the two species or the 8 lakes, than the NMR dataset
309 generated on the same samples (with the same extraction procedure) (Fig. 1 and 2; Supp.
310 fig. S4 and S5). This observation is further confirmed by direct comparison of the NMR and
311 LC-MS metabolome dataset using MixOmics Diabolo tool for multi-block PLS-DA (Supp. fig. S6
312 and S7). Indeed, although the two datasets appear mostly congruent, the individual plot
313 distribution carries much more variance on the PLS-DA projection for the LC-MS analysis
314 than for the NMR, considering both fish species.

315

316 On one side, the ¹H NMR metabolomics allows the reliable quantification of main liver
317 metabolites (belonging principally to amino acids, sugars, TCA metabolites, ...), by detecting
318 characteristic chemical groups, with regards to their respective position within the
319 molecules (Martin et al., 2015). These primary metabolites are key contributors to the
320 cellular metabolism and are supposed to be responsive to variable physiological conditions
321 of fish encountering peculiar biological outcomes and/or environmental constraints (Roques

322 [et al., 2020](#)). However, this approach remains focused on few principal metabolites involved
323 in main cellular pathways which measured cellular contents (expressed in *ppm*) can be
324 globally involved in multiple regulation processes, and not being specific of subtle molecular
325 regulation processes. On the other hand, ¹H NMR presents only low sensitivity and an overall
326 limited discrimination capability for molecules presenting similar chemical structures, such
327 as lipids, that exhibits an aggregated and unspecific signal on NMR spectra ([Emwas et al.,](#)
328 [2019](#)). For this reason, it is supposed to present lower discriminating power than more
329 selective and less qualitative, but more sensitive, high-resolution mass spectrometry-based
330 metabolomics approaches.

331 On the other side, although LC-MS-based metabolomic remains a largely selective analytical
332 approach (according to both LC separation and ionisation respective properties of each
333 molecule), it presents un-precedent analytical capabilities in terms of sensitivity, dynamic
334 range and number of characterized components and offer now unprecedented perspectives
335 for the investigation for molecular phenotyping applied to environmental sciences ([Beyoglu](#)
336 [and Idle, 2020](#)). We assume, that a single LC-MS fingerprint analyse does not embed the
337 whole metabolite picture of the biological compartment (the fish liver, in the present case),
338 especially because of its selectivity performances ([Gika et al., 2019](#)). Indeed, this noticeable
339 selectivity can have multifactorial origin along the analytical procedures and be related to
340 both the extraction solvent, the selected chromatographic gradient and column, together
341 with the ionisation principle and mode. However, it provided specific and precise
342 measurements of a large number of components that can endorse high-throughput and in-
343 depth phenotyping. For comparison, ¹H NMR metabolomics allow the global quantification
344 of the CH₂-CH₂ or CH₂-CH=CH fatty acid bounds ([Marchand et al., 2018](#)), when LC-MS
345 metabolomics can discriminate and characterize hundreds of different lipids belonging to
346 various sub-classes ([Dreier et al., 2020](#)), providing undoubtedly more informative and
347 discriminant features. For these reasons, recent efforts are still been performed in order to
348 compared the inputs of both NMR- and LC-MS-based metabolomics in order to explore the
349 respective advantages and caveats of both approaches for environmental investigations
350 ([Ameline et al., 2019](#); [Hayden et al., 2019](#); [Labine et al., 2020](#)). In addition, multi-omics
351 integration tools, such as MixOmics DIABLO ([Singh et al., 2019](#)), also provides innovative
352 solutions for the comparison of heterogeneous dataset joint matrices obtained from a single
353 set of samples or individuals ([supplementary figures S6-S7](#)).

354 However, the robustness and the repeatability of the operational analytical pipeline
355 constitute key requirements of first order for environmental metabolomics, as recently
356 experienced by clinic development of environmental toxicology ([Silva et al., 2020](#)).
357 Nowadays, it also appears even more crucial to carefully take into account the socio-
358 economic and the environmental costs when considering the development of different
359 pipelines for operational and sustainable metabolomic applications ([Jones et al., 2013](#)).

360

361

362 **3.5. Use of LC-MS metabolomics fingerprint for environmental assessment?**

363 Although the value of LC-MS metabolomics for investigating the impacts of environmental
364 stressors or contaminants, and their respective mode of action has been well-explored in
365 medical sciences (Beyoğlu et al., 2020; Zhang et al., 2020) or in ecotoxicology laboratory-
366 based studies on aquatic models (Le Manach et al., 2018; Huang et al., 2017; Ekman et al.,
367 2015; Gil-Solsona et al., 2017), such methods have been used only faintly in field research so
368 far (Bundy et al., 2009; Lankadurai et al., 2013). Apart from a limited number of evidence of
369 the utility of NMR-based metabolomics in environmental fish studies (Capello et al., 2016;
370 Sotton et al., 2019; Wei et al., 2018), few other examples indicate that field-based LC-MS
371 metabolomics constitutes an emerging and powerful, but still underused, approach for
372 increasing our understanding of *in situ* biological, physiological, ecological or ecotoxicological
373 processes (Meador et al., 2020; Reverter et al., 2017; Goode et al., 2020).

374 So far, aquatic ecotoxicology has been widely using fishes as reference models to investigate
375 both physical or chemical contaminants (Li et al., 2021; Mallik et al., 2021), and their
376 deleterious effects on diverse biological mechanisms including their behaviour (Almroth et
377 al., 2021) or specific enzymatic activities (Villeneuve et al., 2021), on short or longer terms
378 (Thoré et al., 2021). Among the large list of the characters of interest on which
379 ecotoxicological studies aim at being dedicated, the metabolome appears as one of the main
380 promising functional traits (Martynuik et al., 2016). Indeed, metabolomics seem to present
381 many key advantages for environmental studies, such as its immediate connexion with the
382 global physiology of the organisms (Rosato et al., 2018), its pertinence for all model or non-
383 model organisms (Hillyer et al., 2022; Reverter et al., 2017), the multi-variable character of
384 the dataset and its statistical strength (Chu et al., 2019; Mendez et al., 2019), and its
385 appliance for non-lethal and minimal-stress approaches (Colette et al., 2016). However,
386 applying metabolomics to ecotoxicological questions is still currently limited by a lack of
387 conceptual, technical and data frameworks, which require to be tailored to reach its full
388 potential for applied environmental purposes (Walker et al., 2022).

389
390 Our analysis constitutes one of the first attempts to push forward the potential of high-
391 throughput molecular phenotyping, and especially through LC-MS-based metabolomics, for
392 environmental assessment. Indeed, this organism molecular phenotyping supported by
393 multi-variable chemometric investigation offers rich biological information that serves at
394 describing specific phenotypic plasticity. Moreover, this molecular variability/responsiveness
395 can further be confronted to local environmental factors (such as noxious cyanobacterial
396 metabolites in the present case) in order to demonstrate correlation or causality
397 relationships.

398
399 As described by Pompfret and co-workers (2019), environmental metabolomics exhibit very
400 promising perspectives for operational bio-monitoring applications, according to its
401 reliability, its reproducibility, and its high predictive potential. However, these authors also
402 point out that the responsiveness and robustness of the bio-indicative object, which is
403 characterized through the analytical prism of the metabolomics, remains crucial and have to

404 be carefully evaluated and tested with an appropriate experimental design (Pompfret et al.,
405 2019). However, for ethical concerns, fish bio-monitoring would also gain at been less
406 invasive and deleterious for the organisms. To this end, non-lethal mucus sampling has been
407 investigated by LC-MS and have demonstrated the remarkable informativeness of this
408 approach for environmental studies (Reverter et al. 2017), but these efforts still remain
409 explorative and require additional investigations.

410
411

412 **4. Conclusions**

413 The main challenge facing us remains to be able to develop more integrative approaches to
414 connect chemical, biological and ecological assessment, in the context of anthropized
415 natural environments confronting multi-stressor pressures. A major caveat of the use of fish
416 environmental bio-indication by metabolomics remains maybe the lack of standardized
417 methodology and, overall, of dedicated databases (Viant et al., 2019) fulfilled by studies
418 considering together different species, populations, development stages, seasons and
419 environments. These data could be used to provide baseline reference values that would
420 further support machine learning or artificial neural network tools for the training of
421 decision-making models (Mendez et al., 2019).

422
423

424 **Author contributions**

425 B.M. designed experiment, and performed sample analysis. B.M. and A.G. analysed data.
426 B.M. wrote the manuscript. All authors have given approval to the final version of the
427 manuscript.

428

429 **Notes**

430 The authors declare not conflict of interest

431

432 **ACKNOWLEDGMENTS**

433 We would like to thanks B. Sotton for his collaboration on NMR data acquisition and
434 treatment. We are also grateful to A. Paris for its support and prolific discussions. This work
435 was supported by grants from CNRS (Défi ENVIROMICS "Toxycyfish" project). The NMR and
436 the MS spectra were respectively acquired at the Plateau technique de Résonance
437 Magnétique Nucleaire and the Plateau technique de spectrométrie de masse bio-organique,
438 Muséum National d'Histoire Naturelle, Paris, France. This work benefitted from the French
439 GDR "Aquatic Ecotoxicology" framework which aims at fostering stimulating scientific
440 discussions and collaborations for more integrative approaches.

441

442

443 **REFERENCES**

444 Almroth, B. C., Cartine, J., Jönander, C., Karlsson, M., Langlois, J., Lindström, M., ... & Sturve,
445 J. (2021). Assessing the effects of textile leachates in fish using multiple testing
446 methods: From gene expression to behavior. *Ecotoxicology and Environmental*
447 *Safety*, 207, 111523.

448 Amado, L., Monserrat, J., 2010. Oxidative stress generation by microcystins in aquatic
449 animals: Why and how. *Environment International* 36, 226–235.

450 Ameline, A., Garnier, D., Gheddar, L., Richeval, C., Gaulier, J. M., Raul, J. S., & Kintz, P. (2019).
451 Identification and analytical characterization of seven NPS, by combination of 1H
452 NMR spectroscopy, GC–MS and UPLC–MS/MS®, to resolve a complex toxicological
453 fatal case. *Forensic science international*, 298, 140-148.

454 Bahamonde, P. A.; Feswick, A.; Isaacs, M. A.; Munkittrick, K. R.; Martyniuk, C. J. (2016)
455 Defining the role of omics in assessing ecosystem health: Perspectives from the
456 Canadian environmental monitoring program. *Environmental toxicology and*
457 *chemistry* 35(1), 20-35.

458 Beyoğlu, D.; Idle, J. R. (2020). Metabolomic and lipidomic biomarkers for premalignant liver
459 disease diagnosis and therapy. *Metabolites* 10(2), 50.

460 Brusle, J., & i Anadon, G. G. (2017). The structure and function of fish liver. In *Fish*
461 *morphology* (pp. 77-93). Routledge.

462 Bundy, J. G.; Davey, M. P.; Viant, M. R. (2009) Environmental metabolomics: a critical review
463 and future perspectives. *Metabolomics* 5(1), 3.

464 Burford, M. A., Carey, C. C., Hamilton, D. P., Huisman, J., Paerl, H. W., Wood, S. A., & Wulff,
465 A. (2020) Perspective: Advancing the research agenda for improving understanding of
466 cyanobacteria in a future of global change. *Harmful Algae*, 91, 101601.

467 Cappello, T.; Brandão, F.; Guilherme, S.; Santos, M. A.; Maisano, M.; Mauceri, A.; ... Pereira,
468 P. (2016) Insights into the mechanisms underlying mercury-induced oxidative stress
469 in gills of wild fish (*Liza aurata*) combining ¹H NMR metabolomics and conventional
470 biochemical assays. *Science of the Total Environment* 548, 13-24.

471 Chong, J.; Wishart, D. S.; Xia, J. (2019) Using metaboanalyst 4.0 for comprehensive and
472 integrative metabolomics data analysis. *Current protocols in bioinformatics* 68(1):e86.

473 Chong, J.; Liu, P.; Zhou, G.; Xia, J. (2020) Using MicrobiomeAnalyst for comprehensive
474 statistical, functional, and meta-analysis of microbiome data. *Nature Protocols* 15,
475 799-821.

476 Chu, S. H., Huang, M., Kelly, R. S., Benedetti, E., Siddiqui, J. K., Zeleznik, O. A., ... &
477 Consortium of Metabolomics Studies Statistics Working Group. (2019). Integration of
478 metabolomic and other omics data in population-based study designs: An
479 epidemiological perspective. *Metabolites*, 9(6), 117.

480 Collette, T. W., Skelton, D. M., Davis, J. M., Cavallin, J. E., Jensen, K. M., Kahl, M. D., ... &
481 Ekman, D. R. (2016). Metabolite profiles of repeatedly sampled urine from male
482 fathead minnows (*Pimephales promelas*) contain unique lipid signatures following
483 exposure to anti-androgens. *Comparative Biochemistry and Physiology Part D:*
484 *Genomics and Proteomics*, 19, 190-198.

485 Cordier T., Alonso-Sáez L., Apothéloz-Perret-Gentil L., Aylagas E.; Bohan D. A., Bouchez A.,...
486 Keeley, N. (2021) Ecosystems monitoring powered by environmental genomics: a
487 review of current strategies with an implementation roadmap. *Molecular Ecology* 30,
488 2937-58.

489 da Silva, R. R.; Dorrestein, P. C.; Quinn, R. A. (2015) Illuminating the dark matter in
490 metabolomics. *Proceedings of the National Academy of Sciences* 112(41), 12549-
491 12550.

492 Davis, J. M., Ekman, D. R., Teng, Q., Ankley, G. T., Berninger, J. P., Cavallin, J. E., ... & Collette,
493 T. W. (2016). Linking field-based metabolomics and chemical analyses to prioritize
494 contaminants of emerging concern in the Great Lakes basin. *Environmental*
495 *toxicology and chemistry*, 35(10), 2493-2502.

496 De Marco, G., Brandão, F., Pereira, P., Pacheco, M., & Cappello, T. (2022). Organ-Specific
497 Metabolome Deciphering Cell Pathways to Cope with Mercury in Wild Fish (Golden
498 Grey Mullet *Chelon auratus*). *Animals*, 12(1), 79.

499 Dreier, D. A.; Bowden, J. A.; Aristizabal-Henao, J. J.; Denslow, N. D.; Martyniuk, C. J. (2020)
500 Ecotoxicolipidomics: An emerging concept to understand chemical-metabolic
501 relationships in comparative fish models. *Comparative Biochemistry and Physiology*
502 *Part D: Genomics and Proteomics* 36, 100742.

503 Ekman, D. R.; Skelton, D. M.; Davis, J. M.; Villeneuve, D. L.; Cavallin, J. E.; Schroeder, A.; ...
504 Collette, T. W. (2015) Metabolite profiling of fish skin mucus: a novel approach for
505 minimally-invasive environmental exposure monitoring and surveillance.
506 *Environmental Science & Technology* 49(5), 3091-3100.

507 Emwas, A. H.; Roy, R.; McKay, R. T.; Tenori, L.; Saccenti, E.; Gowda, G. A.; ... Wishart, D. S.
508 (2019) NMR spectroscopy for metabolomics research. *Metabolites* 9(7), 123.

509 Ferrão-Filho, A. da S., Kozlowsky-Suzuki, B., 2011. Cyanotoxins: bioaccumulation and effects
510 on aquatic animals. *Marine drugs* 9(12), 2729–2772.

511 Gika, H.; Virgiliou, C.; Theodoridis, G.; Plumb, R. S.; Wilson, I. D. (2019) Untargeted LC/MS-
512 based metabolic phenotyping (metabonomics/metabolomics): The state of the art.
513 *Journal of Chromatography B* 1117, 136-147.

514 Gil-Solsona, R., Nacher-Mestre, J., Lacalle-Bergeron, L., Sancho, J. V., Calduch-Giner, J. A.,
515 Hernández, F., & Pérez-Sánchez, J. (2017) Untargeted metabolomics approach for
516 unraveling robust biomarkers of nutritional status in fasted gilthead sea bream
517 (*Sparus aurata*). *PeerJ* 5, e2920.

518 Goode, K. L.; Dunphy, B. J.; Parsons, D. M. (2020) Environmental metabolomics as an
519 ecological indicator: Metabolite profiles in juvenile fish discriminate sites with
520 different nursery habitat qualities. *Ecological Indicators* 115, 106361.

521 Hamilton, P.; Cowx, I.; et al.; Tyler, C. R. (2016) Population-level consequences for wild fish
522 exposed to sublethal concentrations of chemicals—a critical review. *Fish and Fisheries*
523 17(3), 545-566.

524 Hao, J.; Liebeke, M.; Astle, W.; De Iorio, M.; Bundy, J. G.; Ebbels, T. M. D. (2014) Bayesian
525 deconvolution and quantification of metabolites in complex ¹D NMR spectra using

526 BATMAN. *Nature protocols* 9(6), 1416–1427.

527 Harke, M. J., Steffen, M. M., Gobler, C. J., Otten, T. G., Wilhelm, S. W., Wood, S. A., & Paerl,
528 H. W. (2016). A review of the global ecology, genomics, and biogeography of the toxic
529 cyanobacterium, *Microcystis* spp. *Harmful algae*, 54, 4-20.

530 Harris, J. H. (1995) The use of fish in ecological assessments. *Australian Journal of Ecology*
531 20(1), 65-80.

532 Hayden, H. L., Rochfort, S. J., Ezernieks, V., Savin, K. W., & Mele, P. M. (2019). Metabolomics
533 approaches for the discrimination of disease suppressive soils for *Rhizoctonia solani*
534 AG8 in cereal crops using 1H NMR and LC-MS. *Science of the Total Environment*, 651,
535 1627-1638.

536 Hillyer, K. E., Raes, E., Karsh, K., Holmes, B., Bissett, A., & Beale, D. J. (2022). Metabolomics as
537 a tool for in situ study of chronic metal exposure in estuarine invertebrates.
538 *Environmental Pollution*, 292, 118408.

539 Huang, S. S.; Benskin, J. P.; Veldhoen, N.; Chandramouli, B.; Butler, H.; Helbing, C. C.;
540 Cosgrove, J. R. (2017) A multi-omic approach to elucidate low-dose effects of
541 xenobiotics in zebrafish (*Danio rerio*) larvae. *Aquatic toxicology* 182, 102-112.

542 Janssen, E. M. L. (2019) Cyanobacterial peptides beyond microcystins—A review on co-
543 occurrence, toxicity, and challenges for risk assessment. *Water Research*, 151, 488-
544 499.

545 Jeffries, K. M., Jackson, L. J., Ikonomou, M. G., & Habibi, H. R. (2010). Presence of natural and
546 anthropogenic organic contaminants and potential fish health impacts along two
547 river gradients in Alberta, Canada. *Environmental Toxicology and Chemistry*, 29(10),
548 2379-2387.

549 Jones, O. A., Maguire, M. L., Griffin, J. L., Dias, D. A., Spurgeon, D. J., & Svendsen, C. (2013).
550 Metabolomics and its use in ecology. *Austral Ecology*, 38(6), 713-720.

551 Jones, M. R., Pinto, E., Torres, M. A., Dörr, F., Mazur-Marzec, H., Szubert, K., ... & Janssen, E.
552 M. L. (2021). CyanoMetDB, a comprehensive public database of secondary
553 metabolites from cyanobacteria. *Water Research*, 196, 117017.

554 Jousse, C., Dalle, C., Abila, A., Traïkia, M., Diogon, M., Lyan, B., ... & Delbac, F. (2020). A
555 combined LC-MS and NMR approach to reveal metabolic changes in the hemolymph
556 of honeybees infected by the gut parasite *Nosema ceranae*. *Journal of Invertebrate*
557 *Pathology*, 176, 107478.

558 Kim Tiam, S.; Gugger, M.; Demay, J.; Le Manach, S.; Duval, C.; Bernard, C.; Marie, B. (2019)
559 Insights into the Diversity of Secondary Metabolites of *Planktothrix* Using a Biphasic
560 Approach Combining Global Genomics and Metabolomics. *Toxins* 11(9), 498.

561 Labine, L. M., & Simpson, M. J. (2020). The use of nuclear magnetic resonance (NMR) and
562 mass spectrometry (MS)–based metabolomics in environmental exposure
563 assessment. *Current Opinion in Environmental Science & Health*, 15, 7-15.

564 Lankadurai, B. P.; Nagato, E. G.; Simpson, M. J. (2013) Environmental metabolomics: an
565 emerging approach to study organism responses to environmental stressors.
566 *Environmental Reviews* 21(3), 180-205.

567 Le Manach, S.; Sotton, B.; Huet, H.; Duval, C.; Paris, A.; Marie, A.; ... ; Marie, B. (2018)
568 Physiological effects caused by microcystin-producing and non-microcystin producing
569 *Microcystis aeruginosa* on medaka fish: A proteomic and metabolomic study on liver.
570 *Environmental pollution* 234, 523-537.

571 Le Manach, S., Duval, C., Marie, A., Djediat, C., Catherine, A., Edery, M., ... & Marie, B. (2019)
572 Global metabolomic characterizations of *Microcystis* spp. highlights clonal diversity in
573 natural bloom-forming populations and expands metabolite structural diversity.
574 *Frontiers in microbiology*, 10, 791.

575 Li, H., Jing, T., Li, T., Huang, X., Gao, Y., Zhu, J., ... & Mu, W. (2021). Ecotoxicological effects of
576 pyraclostrobin on tilapia (*Oreochromis niloticus*) via various exposure routes.
577 *Environmental Pollution*, 285, 117188.

578 Lohr, K. E.; Khattri, R. B.; Guingab-Cagmat, J.; Camp, E. F.; Merritt, M. E.; Garrett, T. J.,
579 Patterson, J. T. (2019) Metabolomic profiles differ among unique genotypes of a
580 threatened Caribbean coral. *Scientific reports* 9(1), 1-11.

581 Lorenz, C. M. (2003) Bioindicators for ecosystem management, with special reference to
582 freshwater systems. In *Trace Metals and other Contaminants in the Environment* (Vol.
583 6, pp. 123-152). Elsevier.

584 Mallik, A., Xavier, K. M., Naidu, B. C., & Nayak, B. B. (2021). Ecotoxicological and
585 physiological risks of microplastics on fish and their possible mitigation measures.
586 *Science of The Total Environment*, 779, 146433.

587 Maloufi, S.; Catherine, A.; Mouillot, D.; Louvard, C.; Couté, A.; Bernard, C.; Troussellier, M.
588 (2017) Environmental heterogeneity among lakes promotes hyper β -diversity across
589 phytoplankton communities. *Freshwater Biology*, 61(5), 633-645.

590 Marchand, J.; Martineau, E.; Guitton, Y.; Le Bizec, B.; Dervilly-Pinel, G.; Giraudeau, P. A
591 (2018) multidimensional ^1H NMR lipidomics workflow to address chemical food
592 safety issues. *Metabolomics* 14(5), 60.

593 Marie, B. (2020) Disentangling of the ecotoxicological signal using “omics” analyses, a lesson
594 from the survey of the impact of cyanobacterial proliferations on fishes. *Science of*
595 *The Total Environment* 736, 139701.

596 Marjan, P., Bragg, L. M., MacLatchy, D. L., Servos, M. R., & Martyniuk, C. J. (2017). How does
597 reference site selection influence interpretation of omics data?: Evaluating liver
598 transcriptome responses in male rainbow darter (*Etheostoma caeruleum*) across an
599 urban environment. *Environmental science & technology*, 51(11), 6470-6479.

600 Martin, J. C.; Maillot, M.; Mazerolles, G.; Verdu, A.; Lyan, B.; Migne, C.; ... ; Manach, C. (2015)
601 Can we trust untargeted metabolomics? Results of the metabo-ring initiative, a large-
602 scale, multi-instrument inter-laboratory study. *Metabolomics* 11(4), 807-821.

603 Martyniuk, C. J., & Simmons, D. B. (2016). Spotlight on environmental omics and toxicology:
604 a long way in a short time. *Comparative Biochemistry and Physiology Part D:*
605 *Genomics and Proteomics*, 19, 97-101.

606 McClanahan, T. R.; Schroeder, R. E.; Friedlander, A. M.; Vigliola, L.; Wantiez, L.; Caselle, J. E.;
607 ...; Oddenyo, R. M. (2019) Global baselines and benchmarks for fish biomass:

608 comparing remote reefs and fisheries closures. *Marine Ecology Progress Series*, 612,
609 167-192.

610 Meador, J. P.; Bettcher, L. F.; Ellenberger, M. C.; Senn, T. D. (2020) Metabolomic profiling for
611 juvenile Chinook salmon exposed to contaminants of emerging concern. *Science of*
612 *The Total Environment* 747, 141097.

613 Melvin, S. D., Lanctôt, C. M., Doriean, N. J., Bennett, W. W., & Carroll, A. R. (2019). NMR-
614 based lipidomics of fish from a metal (loid) contaminated wetland show differences
615 consistent with effects on cellular membranes and energy storage. *Science of the*
616 *Total Environment*, 654, 284-291.

617 Mendez, K. M., Reinke, S. N., & Broadhurst, D. I. (2019). A comparative evaluation of the
618 generalised predictive ability of eight machine learning algorithms across ten clinical
619 metabolomics data sets for binary classification. *Metabolomics*, 15(12), 1-15.

620 Nothias, LF., Petras, D., Schmid, R. *et al.* (2020) Feature-based molecular networking in the
621 GNPS analysis environment. *Nat Methods* 17, 905–908.

622 Pomfret, S. M.; Brua, R. B.; Izral, N. M.; Yates, A. G. (2020) Metabolomics for biomonitoring:
623 an evaluation of the metabolome as an indicator of aquatic ecosystem health.
624 *Environmental Reviews*, 28(1), 89-98.

625 Reverter, M.; Sasal, P.; Banaigs, B.; Lecchini, D.; Lecellier, G.; Tapissier-Bontemps, N. (2017)
626 Fish mucus metabolome reveals fish life-history traits. *Coral Reefs* 36(2), 463-475.

627 Rohart, F.; Gautier, B.; Singh, A.; Lê Cao, K. A. (2017) mixOmics: An R package for 'omics
628 feature selection and multiple data integration. *PLoS computational biology* 13(11),
629 e1005752.

630 Roques, S.; Deborde, C.; Richard, N., Marchand; Y., Larroquet, L.; Prigent, S.; ...; Fauconneau,
631 B. (2020) Proton-NMR Metabolomics of Rainbow Trout Fed a Plant-Based Diet
632 Supplemented with Graded Levels of a Protein-Rich Yeast Fraction Reveal Several
633 Metabolic Processes Involved in Growth. *J. Nutrition* 150(9), 2268-2277.

634 Rosato, A., Tenori, L., Cascante, M., Carulla, P. R. D. A., Dos Santos, V. A. M., & Saccenti, E.
635 (2018). From correlation to causation: analysis of metabolomics data using systems
636 biology approaches. *Metabolomics*, 14(4), 1-20.

637 Sardans, J., Peguelas, J., Rivas-Ubach, A., 2011 Ecological metabolomics: Overview of current
638 developments and future challenges. *Chemoecology* 21(4), 191–225.

639 Silva, R. A., Pereira, T. C., Souza, A. R., & Ribeiro, P. R. (2020). ¹H NMR-based metabolite
640 profiling for biomarker identification. *Clinica Chimica Acta*, 502, 269-279.

641 Simmons, D. B., Benskin, J. P., Cosgrove, J. R., Duncker, B. P., Ekman, D. R., Martyniuk, C. J., &
642 Sherry, J. P. (2015). Omics for aquatic ecotoxicology: Control of extraneous variability
643 to enhance the analysis of environmental effects. *Environmental toxicology and*
644 *chemistry*, 34(8), 1693-1704.

645 Singh, A.; Shannon, C. P.; Gautier, B.; Rohart, F.; Vacher, M.; Tebbutt, S. J.; Lê Cao, K. A.
646 (2019) DIABLO: an integrative approach for identifying key molecular drivers from
647 multi-omics assays. *Bioinformatics* 35(17), 3055-3062.

648 Sotton, B., Paris, A., Le Manach, S., Blond, A., Lacroix, G., Millot, A., ... & Marie, B. (2017a)
649 Metabolic changes in Medaka fish induced by cyanobacterial exposures in
650 mesocosms: an integrative approach combining proteomic and metabolomic
651 analyses. *Scientific reports*, 7(1), 1-13.

652 Sotton, B., Paris, A., Le Manach, S., Blond, A., Lacroix, G., Millot, A., ... & Marie, B. (2017b)
653 Global metabolome changes induced by cyanobacterial blooms in three
654 representative fish species. *Science of the Total Environment*, 590, 333-342.

655 Sotton, B.; Paris, A.; Le Manach, S.; Blond, A.; Duval, C.; Qiao, Q.; ...; Marie, B. (2019)
656 Specificity of the metabolic signatures of fish from cyanobacteria rich lakes.
657 *Chemosphere* 226, 183-191.

658 Thoré, E. S., Philippe, C., Brendonck, L., & Pinceel, T. (2021). Towards improved fish tests in
659 ecotoxicology-efficient chronic and multi-generational testing with the killifish
660 *Nothobranchius furzeri*. *Chemosphere*, 129697.

661 Viant, M. R.; Ebbels, T. M.; Beger, R. D.; Ekman, D. R.; Epps, D. J.; Kamp, H.; ...; Rocca-Serra,
662 P. (2019) Use cases, best practice and reporting standards for metabolomics in
663 regulatory toxicology. *Nature communications* 10(1), 1-10.

664 Villeneuve, D. L., Blackwell, B. R., Cavallin, J. E., Cheng, W. Y., Feifarek, D. J., Jensen, K. M., ...
665 & Ankley, G. T. (2021). Case Study in 21st Century Ecotoxicology: Using In Vitro
666 Aromatase Inhibition Data to Predict Short-Term In Vivo Responses in Adult Female
667 Fish. *Environmental Toxicology and Chemistry*, 40(4), 1155-1170.

668 Walker, T. W., Alexander, J. M., Allard, P. M., Baines, O., Baldy, V., Bardgett, R. D., ... &
669 Salguero-Gómez, R. (2022). Functional Traits 2.0: The power of the metabolome for
670 ecology. *Journal of Ecology*, 110(1), 4-20.

671 Wei, F.; Sakata, K.; Asakura, T.; Date, Y.; Kikuchi, J. (2018) Systemic homeostasis in
672 metabolome, ionome, and microbiome of wild yellowfin goby in estuarine
673 ecosystem. *Scientific reports* 8(1), 1-12.

674 Zhang, X. W.; Li, Q. H.; Dou, J. J. (2020) Mass spectrometry-based metabolomics in health
675 and medical science: a systematic review. *RSC Advances* 10(6), 3092-3104.

676 **Legends of figures**

677

678 **Figure 1.** Visualization of the dataset structuration for LC-MS metabolomics (comprising 132
679 fish and 1252 analytes) on PCA (A) and heatmap with hierarchical clustering (B) for *Perca*
680 ($n=78$) and *Lepomis* ($n=54$) fish collected during the 7-10th of September 2015 within 8
681 pounds of the peri-urban Paris' area.

682

683 **Figure 2.** Individual plot of PCA (A and C) and heatmap with hierarchical classification (B and
684 D) of LC-MS liver metabolomes of *Perca* ($n=78$; A-B) and *Lepomis* ($n=54$; C-D) collected
685 during the 7-10th of September 2015 within 8 pounds of the peri-urban Paris' area.

686


687 **Figure 3.** Phytoplankton composition estimated by BBE measurement of the 8 lake sub-
688 surface water (A), corresponding cyanobacteria relative composition for Fon, Tri, Var and
689 Ver (B), and molecular networking of metabolites extracted from the filtered biomass of the
690 respective water of the 8 lakes generated with *t*-SNE algorithm, with cyanobacteria peptide
691 clusters indicated in bold and potentially more noxious family clusters (microcystin,
692 cyanopeptolmin and anabaenopeptin) indicated in red (C).

693


694 **Figure 4.** Individual plots of PLS-DA, testing the difference between the different lake groups
695 presenting, or not, high content of potentially noxious cyanobacterial metabolites
696 (Cha/Cer/Fon/Mau/Rue and Tri/Var/Ver, respectively) according to the LC-MS metabolomes of *Perca*
697 (A) and *Lepomis* (B), and corresponding list of annotated best VIP list (metabolite score > 1.5) of
698 *Perca* and *Lepomis* metabolomes (C). Accuracy, predictability and quality performances and number
699 of VIP metabolite score > 1.5) for sampling lake discrimination are indicated for each species. Dots
700 between brackets indicate when the metabolite concentration is more important in the
701 Cha/Cer/Fon/Mau/Rue (green) or Tri/Var/Ver (red) lake group. * indicates that this metabolite was
702 also observed as being negatively correlated with cyanobacteria occurrence within the different lake
703 according to NMR analysis (Sotton et al., 2019).


704

A





B










- Lakes
- Cergy
 - △ Champs
 - + Fontenay
 - × Maurepas
 - ◇ Rueil
 - ▽ Triel
 - Varennes
 - * Verneuil



- Lakes
- Cergy
 - △ Champs
 - + Fontenay
 - × Maurepas
 - ◇ Rueil
 - ▽ Triel
 - Varennes
 - * Verneuil







C

MW	Annotation	Comp. 1_Perca	Comp. 1_Lepomis
267.09601	Adenosine	9.33 (●<●)	14.3 (●<●)
347.06212	Adenosine 5'-mono-phosphate	2.02 (●<●)	7.64 (●<●)
307.08355	Glutathione (reduced)	7.49 (●<●)	5.28 (●<●)
335.11488	Trp-Met	5.64 (●<●)	0.17
203.11734	Acetyl-carnitine	5.27 (●<●)	2.25 (●<●)
284.13698	Analog: Isoflavonoid	4.84 (●<●)	1.53 (●<●)
463.07428	Adenylosuccinic acid	4.06 (●<●)	0.07
612.15076	Glutathione (oxidized)	3.82 (●<●)	0.05
245.16244	Valeryl-carnitine*	3.76 (●<●)	1.64 (●<●)
463.07419	Adenylosuccinic acid	3.46 (●<●)	1.99 (●<●)
383.10736	Analog: Cyanidin arabonosine	3.41 (●<●)	0.41
567.33206	Lysophosphatidylcholine 22:6	3.34 (●<●)	1.60 (●<●)
626.16765	Analog: N-acetylglutamic acid	1.51 (●<●)	3.08 (●<●)
181.07481	Tyr	1.55 (●<●)	2.67 (●<●)
250.02948	Analog: N-acetylglutamic acid	2.57 (●<●)	0.03
300.13247	Analog: Hydroxyflavanone	2.50 (●<●)	1.03
559.07138	Adenosine 5'-diphosphoribose	2.49 (●<●)	0.59
404.00264	Uridine 5'-diphosphate	2.40 (●<●)	0.55
135.05505	Adenine	1.77 (●<●)	2.38 (●<●)
607.08199	Uridine-5-diphosphoacetylgalactosamine	2.37 (●<●)	0.14
363.05757	Guanosine 5-monophosphate	2.33 (●<●)	1.59 (●<●)
324.10545	Analog: 6-O-α-D-glucopyranosyl-D-fructose	1.89 (●<●)	2.30 (●<●)
392.16974	Trp-Trp	0.02	2.29 (●<●)
321.06967	Cytidine triphosphate	2.28 (●<●)	0.19
278.15223	Tyr-Pro	2.21 (●<●)	0.02
521.19585	sn-glycero-3-phosphocholine	1.69 (●<●)	2.21 (●<●)
665.12496	1,4-Dihydrnicotinamide adenine dinucleotide	1.54 (●<●)	2.16 (●<●)
204.07959	Trp	2.13 (●<●)	0.10
242.06799	Thymidine	2.12 (●<●)	0.36
283.09143	Guanosine	0.44	2.08 (●<●)
429.14971	Analog: Adenosine 5'-monophosphate	1.54 (●<●)	2.06 (●<●)
257.10277	sn-glycero-3-phosphocholine	2.04 (●<●)	0.60
325.37028	N-Oleylethanolamine	2.03 (●<●)	0.77