



**HAL**  
open science

## Lithospheric strength control over depositional environments and foreland accommodation in the Western USA Basin during the Early Triassic

Gwénaél Caravaca, Arnaud Brayard, Emmanuelle Vennin, Michel Guiraud, Nicolas Olivier, Christophe Thomazo, Emmanuel Fara, Gilles Escarguel, Kevin G. Bylund, James F. Jenks, et al.

### ► To cite this version:

Gwénaél Caravaca, Arnaud Brayard, Emmanuelle Vennin, Michel Guiraud, Nicolas Olivier, et al.. Lithospheric strength control over depositional environments and foreland accommodation in the Western USA Basin during the Early Triassic. 15ème Congrès Français de Sédimentologie, Association des Sédimentologistes Français, Oct 2015, Chambéry, France. hal-03922918

**HAL Id: hal-03922918**

**<https://hal.science/hal-03922918>**

Submitted on 27 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# Lithospheric strength control over depositional environments and foreland accommodation in the Western USA Basin during the Early Triassic

Gwénaél Caravaca\*, Arnaud Brayard, Emmanuelle Vennin, Michel Guiraud, Nicolas Olivier, Christophe Thomazo, Emmanuel Fara, Gilles Escarguel, Kevin Bylund, James Jenks, Daniel Stephen

\* [gwenael.caravaca@u-bourgogne.fr](mailto:gwenael.caravaca@u-bourgogne.fr)




## Context of the study




P/T Boundary-Smithian interval (~1,5 ma) in the Sonoma Foreland Basin (western US)

- ⇒ Basin formed due to Golconda Allochthon emplacement around the PTB (Sonoma Orogeny, ~252 Ma)
- ⇒ Excellent fossil and sedimentary record of the aftermath of the PTB-crisis
- ⇒ Paleogeography of the region still poorly constrained

# Context of the study



Brayard et al., 2013



Empiric observations => **Sedimentary record varies in the basin:**

- Transgressive trends recorded in the basin with marine incursions from the North
- **Different facies associations:** from terrigenous and dolomitic red beds, to mixed bioclastic limestones and terrigenous sediments
- **Variation in sedimentary thickness**

*Aim of this work*



*What is the nature of the main controlling factor(s) over the sedimentation in the Sonoma Foreland Basin during the PTU-Smithian interval?*

*Role of the foreland nature of the basin in the aforementioned sedimentary distribution?*


# Methods

## Database:

- 43 selected sections after literature and field work
- Study of the PTU-Smithian interval
- Biostratigraphic framework (after Brayard *et al.*, 2013)

## Palinspastic reconstructions:

- Necessary to obtain an unbiased view of the basin
- Original location of the sections




## Results: thickness variations in the basin

**Isopach map** of the PTU-Smithian interval  
⇒ **Spatial distribution of sedimentary thickness**

2 different parts in the basin:

- **Northern high thicknesses zone** (>300 up to ~550m thick)
- **Southern low thicknesses zone** (~10 up to 250m thick)




# Results: thickness variations in the basin


East-West pattern:

- Thickening trend towards West
- Lower thicknesses in westernmost parts

⇒ **Coherent with Foreland Basin System**  
(sensu DeCelles & Giles, 1996)



DeCelles & Giles, 1996



Caravaca et al., in prep.


## Results: thickness variations in the basin

Sedimentary accumulation caused by **foreland flexuration**

**2 different parts** in the basin(N-S pattern):

- Northern high thicknesses zone (>300 up to ~550m thick)
- Southern low thicknesses zone (~10 up to 250m thick)

⇒ **Why 2 different zones in this single basin as the result of Golconda Allochthon emplacement?**



## Results: terranes map of the basin


Composition of the **basement of the basin**?

⇒ Construction of a new **terrane map** from:

- U/Pb dating (literature data)
- Bouguer gravity anomaly map (Kucks, 1999)

**Five terranes identified, with different ages:**

- Archean Wyoming Terrane
- Archean Grouse Creek Block
- Paleoproterozoic Mojave Terrane
- Mesoproterozoic Yavapai Terrane
- Mesoproterozoic Farmington Terrane



## Results: terranes map of the basin

⇒ Lithospheric nature is key control to rheological behavior

Oldest lithospheres are **colder, more rigid, and show weaker density**

⇒ **Strong lithospheres**

- Wyoming Terrane (Archean)
- Grouse Creek Block (Archean)
- Mojave Terrane (Paleoproterozoic)


Conversely, juvenile lithospheres show **lesser rigidity**

⇒ **Weak uplifted lithosphere**

- Yavapai Terrane (Mesoproterozoic) with positive relief due to ARM orogeny

⇒ **Weak attenuated lithosphere**

- Farmington Terrane (Mesoproterozoic Mobile Belt)




# Discussion

Northern part of the basin:

- **Weak lithosphere** (Farmington Terrane)
- **High sedimentary thicknesses**
- Lower thicknesses on strong lithospheres (Wyoming Terrane and Grouse Creek Block)

Southern part of the basin:

- **Strong lithosphere** (Mojave Terrane)
- **Low sedimentary thicknesses**




Caravaca et al., in prep.

# Perspectives: depositional settings

Depositional settings distributions between the *Owenites* and *Anasibirites* ammonoids biozones (middle Smithian)


Striking differences between southern and northern parts of the basin:

- ⇒ Differences between paleontological and sedimentological record
- ⇒ Paleogeography to be reconstructed using this integrative set of data
- ⇒ Impact of lithospheric control



Depositional settings:

- Orange: Deep subtidal
- Yellow: Shallow subtidal
- Blue: Subtidal to intertidal (microbial carb.)
- Brown: Transitional/coastal (red beds)



## *Conclusion*





- Sonoma Basin acts as a **foreland system** during the P/T Boundary-Smithian interval (~1,5ma)
- The **main controlling factor** over sedimentary accumulation and thus depositional settings distribution **is of lithospheric origin**

*Thank you for paying attention!*





Retrodeformation method





Oldest lithospheres are:

- Thicker and colder than juvenile counterparts (due to R\* elements depletion)
  - More buoyant ( $d^{\circ} < d^{\circ}$  juvenile lithos.)
- ⇒ More rigid and resistant to flexuration






Fig. 2. Lithospheric thermal thickness versus geologic age of the continental lithosphere. The Archean lithosphere has bimodal thickness distribution centered at  $\sim 350$  and  $\sim 220$  km. Gray area shows the lithospheric thickness estimates derived from thermal data (Artemieva and Mooney, 2001). Key: Ar—Archean; ePt, mPt, lPt—early, middle and late Proterozoic, respectively; Pz—Paleozoic; Cz—Cenozoic.

Carpathians



Magallanes basin



**Figure 12.** Conceptual cross sections depicting the major structural and stratigraphic differences between foreland basin systems that develop atop (a) continental and cratonic lithosphere (modified from Horton and DeCelles [1997]) and (b) attenuated rift and back-arc basin lithosphere in successor retroarc foreland basins (this study; with aspects from Romans et al. [2011]). In particular, topographic loading from a mafic thrust belt, a lateral gradient in flexural rigidity, and sediment loading of the foredeep promote a broad and deep region of subsidence.