



# The determinants of saccade targeting strategy in neurodevelopmental disorders: the influence of suboptimal reading experience

Marie Vernet, Stéphanie Bellocchi, Jérémy Danna, Delphine Massendari,  
Marianne Jover, Yves Chaix, Stéphanie Ducrot

## ► To cite this version:

Marie Vernet, Stéphanie Bellocchi, Jérémy Danna, Delphine Massendari, Marianne Jover, et al.. The determinants of saccade targeting strategy in neurodevelopmental disorders: the influence of suboptimal reading experience. *Vision Research*, 2023, 204, 10.1016/j.visres.2022.108162 . hal-03924190

**HAL Id: hal-03924190**

**<https://hal.science/hal-03924190>**

Submitted on 5 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**The determinants of saccade targeting strategy in neurodevelopmental disorders: the influence of suboptimal reading experience**

Marie Vernet<sup>a,b,e</sup>, Stéphanie Bellocchi<sup>d</sup>, Jérémy Danna<sup>e</sup>, Delphine Massendari<sup>b</sup>, Marianne Jover<sup>c</sup>, Yves Chaix<sup>a,f</sup> & Stéphanie Ducrot<sup>b</sup>

<sup>a</sup>ToNIC, Toulouse NeuroImaging Center, University of Toulouse, Inserm, UPS, France

<sup>b</sup>Aix Marseille Univ, CNRS, LPL, Aix-en-Provence, France

<sup>c</sup>Aix Marseille Univ, PSYCLE, Aix-en-Provence, France

<sup>d</sup>Univ. Paul Valéry Montpellier 3, EPSYLON UR 4556, F34000, Montpellier, France

<sup>e</sup>CLLE, Université de Toulouse, CNRS, Toulouse, France

<sup>f</sup>Children's Hospital, Toulouse-Purpan University Hospital, Toulouse, France

**SHORT RUNNING TITLE:** Saccade targeting strategy in pathological reading

**Correspondence**

Marie Vernet

Toulouse Neuro-Imaging Center

Pavillon Baudot – CHU Purpan

Place du Dr Joseph Baylac

31024 Toulouse, France

E-mail: marie.vernet@inserm.fr

## ABSTRACT

Whether eye-movements deficits are causal in reading disorders (RD) or rather a consequence of linguistic processing difficulty experienced by disabled readers has been extensively debated. Since RD are frequently comorbid with the Neurofibromatosis type1 (NF1), children with NF1 were used as a comparison group for children with dyslexia in this study. Eye movements were recorded while 21 dyslexic, 20 NF1, and 20 typically developing children performed an oculomotor lateralized bisection task. In this experiment, we manipulated the type of stimulus - discrete (words and strings of hashes) versus continuous (solid lines) - and the visual field where the stimulus was displayed (left vs. right). The results showed that (1) only proficient readers (TD and NF1 without RD) showed fully developed oculomotor mechanisms for efficient reading, with a clear preferred viewing location located to the left of the word's centre in both visual fields, and fine-tuned saccade targeting guided by the between-character space information and (2) NF1 poor readers mirrored the dyslexic eye movement behaviour, with less accuracy and more variability in saccadic programming, no sensitivity to the discreteness of the stimuli, particularly in the left visual field. We concluded that disruption to oculomotor behaviour reflects the fact that many of the processes involved in reading are not yet automatized for children with RD, independently of NF1. This suggests that the differences in saccade targeting strategy between children with and without RD would be secondary consequences of their reduced reading experience.

**KEYWORDS.** Saccadic computation; Reading experience; Developmental dyslexia; Neurofibromatosis type 1; Parafoveal processing.

48    **ABBREVIATIONS.**

49	DD	Children with developmental dyslexia
50	ILP	Initial landing position
51	LVF	Left visual field
52	NF1	Neurofibromatosis type 1
53	NF1RD	Children with neurofibromatosis type 1 with reading disorders
54	NF1noRD	Children with neurofibromatosis type 1 without reading disorders
55	PVL	Preferred viewing location
56	RAN	Rapid Automatized Naming
57	RD	Reading disorders
58	RVF	Right visual field
59	TD	Typically developing children
60	VP	Viewing position

## 1. INTRODUCTION

As reported by several studies, many adults with dyslexia or poor reading skills experienced early reading difficulties (Eloranta et al., 2018; Smart et al., 2017). Indeed, learning to read can be a hard task for some beginners, requiring becoming more efficient in linguistic processing and to gain cognitive control of saccadic eye movements. Understanding the mechanisms underlying reading difficulties has therefore been the focus of ongoing research, with particular interest in the role of eye movements in the learning-to-read process.

Much research has focused on linguistic processing abilities as determinants of individual differences in reading acquisition (e.g., Kim & Pallante, 2012; Landerl et al., 2019; Leppänen et al., 2004). If phonological processing is a necessary (but not sufficient) condition for the development of adequate word recognition skill, our ability to read depends also on visuo-attentional processes and eye movement control (e.g., Bellocchi et al., 2017; Besner et al., 2005; Facoetti et al., 2010; Plaza & Cohen, 2007; for reviews, see Hung, 2021; Leibnitz et al., 2017; Premeti et al., 2022). Reading does require children to focus selectively on words in a left-to-right attentional scanning. The reader's eyes have to learn to land in the best position within the words to be read, extract the relevant information, and program the next saccade (Ducrot et al., 2013; O'Regan & Lévy-Schoen, 1987; Rayner, 1986).

As reading proficiency increases, a child becomes familiar with the printed form of words and changes in eye movement behaviour are observed: the span of effective vision increases (extending asymmetrically in the direction of reading, even in beginning readers; Häikiö et al., 2009; Rayner, 1986), the saccades size lengthens, the fixations duration and the number of refixations decrease (Blythe, 2014; Blythe & Joseph, 2011; Rayner, 2009). Moreover, the role of attention in reading evolves with expertise, i.e., expert readers need less attention to process words than beginning readers (Bellocchi & Leclercq, 2021; LaBerge & Brown, 1989; Leclercq & Siéoff, 2016).

86       The present study further investigates the perceptual and attentional factors that  
87 determine the eyes' landing position in children and whether or not the pattern of eye  
88 movements (and in particular the primary saccade landing sites) can provide useful  
89 information to help characterize the reading skills in neurodevelopmental disorders. Word  
90 length, that is provided by the visually salient extra spacing between words, in most languages  
91 that use an alphabetic script, is a primary source of information in order to guide the eyes  
92 during text reading and determine where the eyes first land in a word (e.g., Inhoff et al., 2003;  
93 Rayner et al., 1998). Rayner (1979) first demonstrated that the landing position in a word  
94 during continuous reading, referred to as the "preferred viewing location" (PVL), is usually  
95 located halfway between the beginning and the middle of the word for languages read from  
96 left-to-right (McConkie et al., 1988; Radach & Kempe, 1993; Rayner, 1979; Vitu et al.,  
97 1990). It is thought that the PVL reflects a strategy whereby readers aim to fixate the centre of  
98 the word (e.g., McConkie et al., 1988; Reichle et al., 2003; but see Vitu, 2003) and fall short  
99 of that optimal viewing position (OVP) (i.e., saccades would undershoot their goal), owing to  
100 the properties of the oculomotor system (Engbert & Krügel, 2010; Joseph et al., 2009;  
101 McConkie et al., 1988; Nuthmann et al., 2005; O'Regan & Lévy-Schoen, 1987).  
102 Alternatively, a visual strategy could be developed to send the eyes to a location that  
103 optimizes information uptake from the fixated word (Legge et al., 1997; O'Regan & Lévy-  
104 Schoen, 1987). The evidence strongly suggests that in addition to length information, salient  
105 orthographic information can be extracted from the parafovea and influence saccade  
106 computation (e.g., orthographically unfamiliar initial trigrams, Hyönä, 1995; White &  
107 Liversedge, 2006). Furthermore, developmental research suggests that beginning readers do  
108 learn to optimize saccade targeting strategies, with average initial landing positions (ILPs)  
109 gradually shifted towards the word centre during the first years of formal reading instruction  
110 (Ducrot et al., 2013; Huestegge et al., 2009; Joseph et al., 2009; Vorstius et al., 2014). Based

on the dual-route cascaded model of Coltheart et al. (2001), the child would rapidly switch from a grapho-phonological decoding mode (indirect route), with a letter-by-letter processing, to an orthographic treatment mode (direct route) with an OVP for word processing. Similarly, the attention deployed on words would be adapted to the preferred global processing mode (Ans et al., 1998) and would be related to the efficiency of the oculomotor parameters (Prado et al., 2007). Thus, during the first year of learning, oculomotor control and especially the PVL will be adjusted in parallel to the development of an optimal lexical and visual-attentional processing strategy for word reading.

Of particular interest here is the study by Ducrot and Pynte (2002) who used an oculomotor lateralized bisection task and demonstrated that the presence of between-character space information influences saccade programming. According to this view, PVL “is due to attentional processes that develop for any type of discrete stimulus, whether or not reading is actually required” (Ducrot & Pynte, 2002, p. 1142). These authors argued that “linguistic-like” stimuli (i.e., with inter-character spaces) encourage a saccade targeting strategy that takes the direction of visual exploration into account and aims to land left of the centre of targets (for languages read from left to right), in preparation for subsequent left-to-right attentional scanning. Note that this oculomotor strategy is modulated according to the visual field (VF) in which the stimulus is displayed, resulting in a left-right asymmetry in saccade extent which is larger in the left visual field (LVF) than in the right visual field (RVF). In the same vein, Bellocchi et al. (2019) found that the average landing positions of fifth grade typical reading children were left of centre for stimuli with inter-character spaces (words and strings of hash marks) in both the left and right VFs. Interestingly, beginning readers and children with DD were only influenced by the linguistic nature of stimuli such that ILPs were shifted more toward the beginning of the strings only when the stimulus was a word.

Naturally, a reasonable question arises: do children with different levels of reading ability differ in their saccade computation strategies when processing parafoveally presented stimuli? In the present study, we used the oculomotor lateralized bisection task as Ducrot and Pynte (2002) to further investigate saccade programming in typical reading children and children with reading difficulties, under conditions that modulate the direction of visual exploration and minimize the influence of linguistic factors.

A huge amount of data has reported differences between the eye movement patterns of dyslexic readers and those of skilled readers. Poor readers and individuals diagnosed as having dyslexia exhibit inefficient oculomotor behaviour while reading, with longer saccades, shorter fixations, and more refixations than typical readers (Ashby et al., 2005; Franzen et al., 2021; Gagl et al., 2014; Gangl et al., 2018; Hawelka et al., 2010; Kirkby et al., 2022; Lefton et al., 1979; Prado et al., 2007; for review, see Rayner, 1998). Also relevant for the present study are experiments reporting a missing parafoveal preview benefit in poor readers or dyslexic children (Silva et al., 2016). The difficulty of pre-processing the word to the right of fixation may affect both saccade-planning and its execution towards the PVL. In that sense, dyslexic readers tend to initially fixate words closer to their beginning than good readers (Gagl et al., 2014; Gangl et al., 2018; Hawelka et al., 2010; Kirkby et al., 2022; McDonald et al., 2006; McKeben et al., 2004). However, there is still some unsettled disputes on the role of these atypical eye movement patterns as a causal factor in dyslexia (see Blythe et al., 2018, asserting that instead they might primarily reflect an underlying difficulty in written word recognition or decoding). Indeed, the attempts to replicate differences in eye-movement patterns of dyslexic, poor and control readers when performing nonreading tasks have been essentially unsuccessful, suggesting that children “with delayed reading skills (identified by an assessment of age referenced reading ability rather than a diagnosis of dyslexia possess the



capacity to execute eye movements equivalent to those with good reading skills” (Hindmarsh et al., 2021, p. 1135).

The point of agreement is that in a reading task, children with DD exhibited atypical visual-processing skills. Among them, the deployment of visual attention seems to be impaired in DD individuals. As mentioned above, several studies have shown that in left-to-right reading systems, the fixation point of skilled readers was often located toward the left of the strings, when they processed multiple characters at once, under the influence of reading habits and experience (e.g., Ducrot & Pynte, 2002; Sireteanu et al., 2005), suggesting a left dominance of attention distribution in the process. DD children have a deficit in RVF attentional inhibition, and their attentional resources can be excessively interfered by RVF information (Li et al., 2021). It results in the ignorance of LVF information (see Facoetti et al., 2006; Hari & Renvall, 2001), which leads to no significant difference between LVF and RVF, showing an unbiased distribution pattern of attention resources (Li et al., 2021). In that vein, Ducrot and colleagues found a symmetric OVP curve for dyslexic readers in foveal vision, contrary to control who demonstrated a typical OVP effect with a left-right asymmetric J-shape curve (Bellocchi & Ducrot, 2021; Ducrot et al., 2003, see also Bellocchi et al., 2019, for a less pronounced left-right asymmetry in parafoveal vision). In addition, children with DD are more affected by visual crowding (even with non-linguistic stimuli, Joo et al., 2018; Spinelli et al., 2002) and benefit more from increased text spacing (Perea et al., 2012; Zorzi et al., 2012; for reviews, see Bellocchi et al., 2013; Gori & Facoetti, 2015). Finally, the atypical eye-movement patterns observed in dyslexic children have been linked to impaired visuo-attentional processing, and poor visual attentional span abilities, resulting in fewer letters simultaneously processed and more rightwards fixations (Prado et al., 2007). As suggested by Facoetti (2012), a possible neurobiological substrate of visuo-spatial attention deficits in DD could be a weakened or abnormal magnocellular input to the dorsal visual

stream. Magnocellular–dorsal deficits could lead to reading difficulties through impaired serial attentional orienting (Facoetti et al., 2010; Franceschini et al., 2012; Vidyasagar & Pammer, 2010) or poor eye movement control (Stein, 2001; but see Goswami, 2015; Hutzler et al., 2006, for a different point of view).

One of the main motivations of the study was to examine whether deviant eye movements exhibited by dyslexic readers are a symptom of dyslexia or constitute a more causal impairment. Few studies have explored eye movement behaviour in children with a range of reading abilities that goes beyond dyslexia, using eye tracking during a task preserving the perceptual and oculomotor demands of reading (but removing the higher order processing). Could the inefficient eye movement patterns reported in DD be transposed to a reading deficit in another vulnerable population? To find answers to this question, we focused on neurofibromatosis type 1 (NF1), that is a genetic disease in which a high prevalence of reading deficits has been shown (between 30 % to 80 %; Arnold et al., 2020; Chaix et al., 2017; Hyman et al., 2005), with a profound impact on their academic achievement (Krab et al., 2008). In this neurodevelopmental disorder, visual-processing deficits were also demonstrated (e.g., Hyman et al., 2005; Ribeiro et al., 2012; Vernet et al., 2022) and a first eye-tracking study showed atypicality in the saccadic system (Lasker et al., 2003). Some studies have suggested a delay in the maturation of low-level vision processes in children with NF1 (Lasker et al., 2003; Ribeiro et al., 2012). To our knowledge, no study has investigated the involvement of eye-movement control in the reading difficulties observed in NF1 children.

Summing up, reading is primary a visual task requiring good oculomotor and visuo-attentional skills and in which the initial fixation position of a word plays a crucial early role. With expertise, a left-right asymmetry in VF emerges and the discreteness nature of the word becomes a factor of considerable importance when computing the incoming saccade. Some

findings suggest that in reading disorders, such as DD, there is atypical eye movement control and specifically, the implementation of an inefficient saccade targeting strategy. To our knowledge, very few studies have examined the factors that lead to this ineffective strategy in children with severe reading difficulties, either in DD or in NF1. Thus, we assume that it is crucial to better understand the determinants of eyes' landing position in pathological reading in each of these populations. Here, we examined whether children with reading disorders (RD) differed from typical developing readers in saccade targeting strategies, and whether lateralized presentation would differently impact saccade programming in children with and without RD. We hypothesized that if a detrimental effect of LVF presentation is observed, this effect may occur especially for children who have not yet fully developed oculomotor mechanisms for efficient reading (i.e., DD and NF1 children; Bellocchi et al., 2019; Ducrot & Grainger, 2007; Nazir, 2000). Moreover, according to Ducrot and Pynte (2002), we predicted that if landing positions are an indicator of the development of reading skills, ILPs would differ according to stimulus type and reading experience, with only experienced readers being able to take account of the presence of between-character space information to compute saccades. To test this hypothesis, we manipulated the type (linguistic vs. non-linguistic) as well as the discreteness (stimuli with inter-character spaces vs. continuous lines) of the stimuli.

We conducted a three-step analysis. With the first analysis, we investigated the extent to which attentional factors, stimulus features, and reading ability influence eye behaviour, and more specifically the PVL effect in parafoveal vision. At the same time, we explored possible visual and oculomotor deficits in children with DD. The second analysis aimed to quantify the use of different saccade targeting strategies and closer understand their relationship to reading abilities, by comparing NF1 children with and without RD. Finally, a third analysis including all children was carried out, in order to further investigate the relation between individual

landing position and reading skills. We wanted to know if independently of the clinical diagnosis, reading ability only can reveal different saccade targeting strategies.

## **2. METHOD**

### **2.1. Participants**

Twenty typically developing children (TD, 10 females), 21 children with DD (8 females), and 20 children with NF1 (13 females) participated in the present study.

The inclusion criteria were the same as those described in the study by Vernet et al. (2022). All children were between 8 and 12 years old and attended school from grades 2 through 6. They were all French native speakers, right-handed and presented normal or corrected-to-normal vision. There were no significant differences between groups for chronological age (TD mean age = 118.30 months, SD = 11.48; DD mean age = 120.90 months, SD = 14.18; NF1: mean age = 121.57 months, SD = 18.68). All children underwent a complete medical and neuropsychological screening. Children were excluded from the study if they had Attention Deficit Hyperactivity Disorder (i.e., more than 6 symptoms of Hyperactivity and/or Inattention in a parent rating on the DSM-5 diagnosis criteria), intellectual disability (i.e., WISC total IQ below 70 and/or standard score below 7 on the Similarities and/or Picture Concepts subtests), neurological or psychiatric disorder (e.g., epilepsy, brain tumor, or autism), or a known hearing impairment.

For the DD children, the reading level was controlled to ensure that children exhibited reading disorders (i.e., scores at least 1 SDs below the mean on the reading speed and accuracy indices of the Alouette test (Lefavrais, 2005) and at least 1.5 SDs below the mean on the ODEDYS-2 test (Jacquier-Roux et al., 2002)). The reading level was also assessed in the TD group to verify that they were all typical readers (i.e., scores at least superior to -0.5 SDs

on the Alouette-test (Lefavrais, 2005) and on the ODEDYS-2 test (Jacquier-Roux et al., 2002)). As expected, statistical analyses revealed that the DD group displayed significantly poorer performances than the TD group for the three reading indices with large effect sizes [reading age:  $t(39) = -10.022$ ,  $p < .001$ ,  $d = -3.131$ ; reading accuracy:  $t(39) = -5.703$ ,  $p < .001$ ,  $d = -1.782$ ; reading fluency:  $t(39) = -11.277$ ,  $p < .001$ ,  $d = -3.523$ ].

For the NF1 children, participation was offered to all children who met the clinical diagnostic criteria for NF1, according to the Neurofibromatosis Conference Statement (National Institutes of Health, 1988). The reading level was systematically assessed but was not used as an inclusion criterion for this specific subject.

The French ethics committee Review Board approved the study (2014-A01239-38 and 2014-A01960-47), which was conducted according to the guidelines of the Declaration of Helsinki (World Health Organisation, 2008). The written informed consent was obtained from every participant and their legal tutors prior to inclusion in the study. Note that this study included participants from a larger study examining procedural learning and memory<sup>1</sup>.

## 2.2. Material

*Oculomotor lateralized bisection task.* The material consists of 3 lists of stimuli: solid lines, strings of hashes and words. Each list is composed of 20 items of 5 or 6 characters long. We selected a set of words from the first-grade lemma lexicon of the Manulex database (Lété et al., 2004) that provides reliable information on grade-level word frequency. Half of the targets were high frequency words, with average printed frequency of 419 occurrences per million and the other half were low frequency words (printed frequency of occurrence of 16 occurrences per million). Stimuli were displayed in either the LVF or RVF relative to the

---

<sup>1</sup> DYSTAC-MAP cohort (ANR-13-APPR-0010)

initial fixation point. The distance between the item and the fixation cross was set at 1.25° from the end of the stimulus when displayed in the LVF or from the beginning of the stimulus when displayed in the RVF. To sum up, two factors were manipulated: (1) the type of stimulus with 3 conditions (i.e., words, strings of hashes and solid lines) and (2) the visual field with 2 conditions (i.e., LVF and RVF).

*Visual-processing skills assessment.* The *Developmental Eye Movement* (DEM) test (Garzia et al., 1990) is commonly used to assess visual-processing skills. It consists of reading aloud vertically and horizontally organized digits printed on three different sheets of paper (Test A, B and C). Children were instructed to read aloud the digits as fast and as accurately as possible. This test provides three main indices: (1) the vertical reading time (VT, in seconds) that represents the sum of the time spent on naming the vertically organized digits of Test A and B; (2) the adjusted horizontal time (HT, in seconds) that represents the adjusted time required for reading the horizontally organized digits presented in the Test C; and (3) the total number of errors, which gives the accuracy on the execution of Test C.

*Reading skills assessment.* One of the most widely used French reading tests for assessing reading efficiency in children is the *Alouette test* (Lefavrais, 1967, 2005). In this test, the child is asked to read a 265-word text passage aloud as quickly and accurately as possible, within 3 minutes. The text consists of real words in syntactically correct but semantically poor sentences. The Alouette test provides three main indices: reading age, reading fluency, and reading accuracy.

### **2.3. Apparatus and procedure**

All children were recruited at both the Hospital of Aix-en-Provence and the Timone University Hospital (Marseille). They were seen individually during two sessions of approximately 1 hour and a half each. The first one was conducted directly at the hospital

and was intended to verify the inclusion criteria. This session included also neuropsychological assessment with the evaluation of reading level. The second session was carried out at the laboratory to perform the oculomotor bisection task. All children were tested individually in a quiet, separate room. The lighting in the room was adjusted to a comfortable level for each participant and each task.

Some of the following detail is taken from Ducrot et al. (2013). For the oculomotor lateralized bisection task, the eye movements were recorded by a mobile infrared, head-mounted eye tracker (Eyelink 2, SR Research Ltd., Canada). The recording was based on infrared-light reflection from the pupil and cornea of the right eye at a sampling rate of 250 Hz and a spatial resolution of less than  $0.04^\circ$ . A chin-and-forehead rest was used to minimize head movements.

Before the experiment, a calibration step of the eye tracker was performed according to a 9-point grid extended to the whole computer screen. The eye tracker was connected to a Dell D-type docking station and a Dell Latitude D600 laptop computer for data recording and stimulus delivery. The stimuli were displayed in white lowercase letters against a black background in 22-point Courier New font, using a 14-inch colour monitor, at a resolution of  $1400 \times 1050$  pixels. Participants were seated in front of the screen at 60 cm. At this distance, one letter subtended a visual angle of  $0.38^\circ$  and  $1^\circ$  is equal to 0.95 cm. The space between letters was 1 mm that corresponds to 2.8 pt or  $0.09^\circ$ .

Each trial consisted of the following sequence of events (Figure 1). First, children were required to maintain fixation on the cross in the centre of the screen. Note that the experimenter repeatedly reminded the child to fixate the cross and not to move their eyes away from this point. Five-hundred ms after, the central fixation point was replaced by a parafoveal target displayed in the RVF or the LVF of the fixation point. Therefore, the children were instructed to “move their eyes, as quickly and accurately as possible, to the

position they considered to be the middle of the item” and then validate this position by pressing a button. After the response, the screen was cleared for the next trial that started 500 ms later. Participants performed 12 practice trials beforehand.

**[Insert Figure 1 About here]**

## **2.4. Data analysis**

The Emaa software package (Ducrot et al., 2006) was used to analyze the eye-tracking data. Trials were excluded from the analysis when an eye-tracker sampling error or a blink occurred during stimulus presentation. We also excluded trials when the saccade latency was shorter than 80 ms or longer than 800 ms (4.3%), the average eye position before saccade onset deviated from the fixation cross by more than  $\pm 0.5$  characters (2.9%), or the participant took more than 5000 ms before pressing the button (6.8%).

We used the same eye-movement measures as Ducrot et al. (2013): (1) the saccade latency which is the time between the onset of the target and the beginning of the saccade, (2) the saccade size which is the size of the initial saccade, and finally (3) the refixation probability which is the probability of making an additional fixation within the word before leaving it. To study PVL effects, we also computed (4) the initial landing position in the stimulus. This last parameter is used to determine the landing frequency at each position of the item. Saccade latency was reported in milliseconds, whereas saccade size and initial landing position were expressed in number of characters. For all the variables measured, means were calculated for each participant in each condition.

## **2.5. Statistical analysis**

A three-step analysis was conducted. A first analysis was conducted to examine saccade targeting strategies of children with and without DD in parafoveal vision. ANOVA were



performed with a 2 Visual Field  $\times$  3 Type of stimulus  $\times$  2 Groups of children model. All factors except the group of children were within-participant variables. Further analyses were carried out in each group independently with statistical a priori comparisons.

In a second analysis, the same ANOVA were performed within the NF1 group, to better understand the factors that determine the saccade targeting strategy implemented by children with RD. Based on their performance on the Alouette test, NF1 children who presented reading disorders (i.e., NF1RD; at least 12 months delay in reading age on the Alouette test compared to chronological age) were compared to NF1 children without reading disorders (i.e., NF1noRD). Note that 11 children were included in the NF1RD group and 9 in the NF1noRD group. The two groups differed significantly in terms of reading age [ $t(18) = -4.435$ ,  $p < .001$ ], reading accuracy [ $t(18) = -4.144$ ,  $p < .001$ ] and reading speed [ $t(18) = -5.218$ ,  $p < .001$ ]<sup>2</sup>.

Finally, a final analysis including all children was carried out to further investigate the relation between individual ILPs and reading skills. Each child was assigned an ILP index (their initial-landing position for words in the RVF) and a reading level index (their reading age). The correlations between these two measures were tested. Then, cluster analyses were conducted. The ILPs were analysed by interactive partitioning (K-means), minimizing the within-cluster variability and maximizing the between-cluster variability, with a three-group solution.

### 3. RESULTS

#### 3.1. Comparisons between children with and without DD

---

<sup>2</sup> There were no significant differences between the two groups for the chronological age [ $t(18) = -5.316$ ,  $p > .05$ ] and the handedness ratio ( $\chi^2 = 0.669$ ,  $p > .05$ ).

[Insert Table 1 about here]

### 3.1.1. Saccade Size

The median amplitude of left-to-right saccades in TD children was 3.46 characters, while that of dyslexics was 3.49. For right-to-left saccades, however, we observed a shift of the saccade's distribution toward shorter amplitudes for DD children (as compared to TD children, 3.7 and 4.1 for DD and TD children, respectively). The analysis of incoming saccade size yielded a significant interaction between stimulus type and side of presentation [ $F(2,76) = 12.505, p < .001, \eta_p^2 = 0.247$ ] and three-way interaction between the group (TD vs. DYS children), the type of stimulus (words, strings of hashes and solid lines), and the presentation side (LVF vs. RVF) [ $F(2, 76) = 3.423, p = .038, \eta_p^2 = 0.080$ ]. Separate analyses for TD and DYS children indicated that the interaction between the stimulus type and the presentation side was significant in the two groups [ $F(2,38) = 7.128, p = .002, \eta_p^2 = 0.273$  and  $F(2,38) = 8.509, p < .001, \eta_p^2 = 0.309$ , for the TD and DD children respectively].

As can be seen in Table 1, for TD children, initial-saccade extent was equivalent in size for all stimuli composed of discrete elements (words and strings of hashes), but was different for continuous solid lines. Pairwise comparisons revealed that the difference between words and strings of hashes was nonsignificant [ $F(1,38) = 1.089, ns$ ] and that the stimulus-type by presentation-side interaction result from the strong opposition between discrete stimuli (words and strings of hashes) and continuous stimuli (solid lines) [ $F(1,38) = 8.413, p < .01$ ], thus confirming the influence of stimulus discreteness vs. continuousness in determining ILPs (see Ducrot et al., 2002, for similar conclusions).

For DD children, pairwise comparisons revealed that the stimulus-type by presentation-side interaction was entirely explained by the strong opposition between linguistic stimuli (words) and non-linguistic stimuli (strings hashes and solid lines) [ $F(1,38) = 10.208, p < .01$ ], and no difference between strings of hashes and solid lines [ $F$

< 1]. As Table 1 shows, DD children do tend to initially fixate *words* at a point somewhere between the beginning and the centre of the target when the saccade was rightward and closer to the word middle for left presentations. For non-linguistic stimuli, however, the mean ILPS were approximately symmetrical in the left and right VFs.

### 3.1.2. Latency

As first raised by Radach & McConkie (1998), this variable is “of particular interest, because it provides a link to temporal aspects of eye movement control that may well significantly modulate spatial saccade parameters” (p. 83). The ANOVA yielded a main effect of type of stimulus [ $F(2,78) = 92.602$ ,  $p < .001$ ,  $\eta_p^2 = .704$ ], regardless of the presentation side (left or right). Pairwise comparisons revealed that the interaction was due to the difference between discrete and continuous stimuli [183 ms for discrete stimuli vs. 220 ms for continuous stimuli,  $F(1,78) = 578.436$ ,  $p < .001$ ]. The difference between words and strings of hashes was nonsignificant [ $F(1,78) = 1.35$ , ns]. Note that strings of hashes are letter-like and similar in visual complexity to letters, so processing these symbols might activate a decoding process similar to processing text. Strings of hashes also differ substantially from solid lines in shape, size and brightness. A similar tendency in expert readers has already been observed by Ducrot and Pynte (2002), with shorter latencies in the discrete condition than in continuous condition (173 vs. 193 ms,  $p = .06$ , Experiment 3). There was also a main effect of group [ $F(1,39) = 4.839$ ,  $p = .034$ ,  $\eta_p^2 = .110$ ] and a stimulus type  $\times$  group interaction [ $F(2,78) = 6.274$ ,  $p = .003$ ,  $\eta_p^2 = .139$ ], showing that latencies were longer for DD children (207 ms,  $SD = 31$ ) than for TD children (196 ms,  $SD = 20.5$ ) (see Bucci et al., 2008; Fischer & Weber, 1990; Pirozzolo & Rayner, 1979, for similar results), in particular for discrete stimuli (193 vs. 173 ms for DD and TD respectively, Table 1).

In order to examine whether symmetrical patterns were associated with short latencies (saccades aim to target the centre of the word but undershoot their goal) and asymmetrical patterns with long latencies (the eyes land at the location that enables maximum information intake), we calculated (1) an asymmetry index (the difference between the mean amplitude of a child initial saccades in the left and right VFs) and (2) a rapidity index (his or her mean saccade latency, for leftward and rightward saccades) for each participant and each type of target. The correlation between these two indices was not significant with a lack of correlation for words [ $r(39) = .05$ ,  $p > .10$ ], strings of hashes [ $r(39) = -.03$ ,  $p > .10$ ] and solid lines [ $r(39) = .07$ ,  $p > .10$ ]. The same tendency was found when separate analyses were made for TD and DD children [ $r(18) = -.19$ ,  $p > .10$ , and  $r(19) = .40$ ,  $p > .05$ , respectively].

### 3.1.3. Refixation probability

We found a significant effect of group on refixation probability, [ $F(1,38) = 6.626$ ,  $p = .014$ ,  $\eta_p^2 = .148$ ], with surprisingly DD children making less refixations (57.64%) than TD children (68.09%). There was also an interaction between presentation side and group [ $F(1,38) = 9.442$ ,  $p = .004$ ,  $\eta_p^2 = .199$ ], with an “adverse” effect of fixating the RVF among the DD children (as compared to the TD group). Interestingly, the analyses revealed three-way interaction between the group, the presentation side and the type of stimulus [ $F(2,76) = 3.632$ ,  $p = .031$ ,  $\eta_p^2 = .087$ ]. Separate analyses were done for TD and DD children and revealed that the interaction between presentation side and stimulus type was only significant in TD children [ $F(2,38) = 3.333$ ,  $p = .046$ ,  $\eta_p^2 = .149$  and  $F < 1$ , for TD and DD children respectively]. For the DD group analysis, we only found a main effect of the side of presentation [ $F(1,19) = 9.005$ ,  $p < .007$ ,  $\eta_p^2 = .322$ ], with children making more refixations on the RVF (64%) than on the LVF (51%). For the TD group, pairwise comparisons revealed a significant difference between RVF and LVF for words only [ $F(1,38) = 8.725$ ,  $p < .01$  for words, and  $F_s < 1$  for strings of hashes and solid lines], thus suggesting an advantage for the

RVF limited to word recognition. Note that, even if no reading was required, the bisection task elicited comparable ILPs patterns and a RVF advantage for words in TD children, thus supports the idea that some of the factors responsible for the VF and ILPs asymmetries in natural reading are also at work in this task.

To sum up, saccades toward stimuli located in the RVF were not significantly affected by stimulus type or reading disorder, all children being able to land at the PVL. In contrast, leftward saccades were influenced by stimulus type and reading ability. In particular, the left-right asymmetry of the incoming saccade was influenced by the inter-character spacing in TD children whereas the asymmetry was restricted to words in dyslexics. Moreover, the TD readers fixated closer to the PVL of the discrete stimuli in the LVF compared to the dyslexic children, with leftward saccades about one character longer. These results are consistent with those of Ducrot & Pynte (2002) who showed that expert readers land to the left of the centre in all discrete stimuli (strings of hashes and words) for both presentation sides (see Bellocchi, Mancini, et al., 2013, for similar results in 5<sup>th</sup> graders). These results suggest that, as a result of reading experience, the children tested in this experiment had already developed automatized routines for programming right-going saccades. The fact that saccades from right-to-left are in the opposite direction of reading would lead to less automatized routines for saccade computation, making them more vulnerable to the influence of reading level and VF. It must be emphasized that many children with dyslexia do not read as much as individuals with typical reading skills, and so, this non-optimal saccade targeting strategy in dyslexia may also results from poor reading experience (see Huettig et al., 2018, for a discussion).

### **3.2. Comparisons of NF1 children with and without RD**

[Insert Table 2 about here]

### 3.2.1. Saccade size

In clinical populations, as suggested by Ablinger et al. (2014), a “standard way to report on the spatial aspect of eye movement control is to plot the individual distributions of the landing positions of incoming saccades over all letter positions within a target word” (p. 652; see also Radach & Kempe, 1993; Radach & McConkie, 1998).

Zooming in on the frequency distributions of ILPs (Figure 2) reveals different distribution profiles between NF1 children who presented reading disorders (NF1RD) and those without reading disorders (NF1noRD). We found a significant correlation between the reading age of the NF1 children and the frequency of their initial landing position at P2 [ $r(18) = .47, p < .05$ ]. Whereas a PVL was distinctly observed in NF1noRD children between the beginning and the centre of the word (i.e., P2), no PVL emerged in NF1RD children [ $t(39) = -3.241, p < .01$ ].

[Insert Figure 2 about here]

More precisely, for both groups, there was an interaction between stimulus type and presentation side [ $F(2,20) = 7.236, p = .004, \eta_p^2 = 0.420$  and  $F(2,16) = 3.593, p = .05, \eta_p^2 = 0.313$ , for NF1RD and NF1noRD respectively]. For the NF1RD group, pairwise comparisons were used to reveal a difference in initial-saccade size between linguistic stimuli (words) and non-linguistic stimuli (strings hashes and solid lines) [ $F(1,20) = 14.323, p < .005$ ]. The difference between strings of hashes and solid lines was non-significant [ $F < 1$ ]. For the NF1noRD group, the amplitude of the initial saccade was similar for words and strings of hashes, but different for solid lines, thus confirming the importance of the discreteness of the stimuli for unimpaired readers [ $F(1,16) = 4.49, p = .05$ ], and [ $F < 1$ ], for (words and strings of hashes) vs. solid lines, and strings of hashes vs. words, respectively].

Of particular interest is that NF1RD readers mirrored the dyslexic eye movement behaviour, with more variability in saccadic programming, and no sensibility to the discreteness of the stimuli (see Figure 3). The data showed no main effect of group [ $F < 1$ ] nor double interaction between presentation side, stimulus-type and group [ $F < 1$ ], but a presentation side  $\times$  stimulus-type interaction [ $F(2,58) = 13.732$ ,  $p < .001$ ,  $\eta_p^2 = .321$ ], mainly explained by the sensitivity to the linguistic nature of the stimuli [ $F(2,58) = 58.81$ ,  $p < .001$ , for the opposition between linguistic vs. non-linguistic stimuli, and  $F < 1$ , for the difference between strings of hashes vs. solid lines].

**[Insert Figure 3 about here]**

The children with NF1noRD showed fully developed oculomotor mechanisms for efficient reading similar to those of TD children [ $F < 1$ ], with a clear PVL located to the left of the word's centre in both visual fields for the 2 groups [ $F < 1$ , for the group  $\times$  presentation side  $\times$  stimulus type interaction], and fine-tuned saccade targeting guided by the between-character space information. The stimulus-type by presentation-side interaction [ $F(2, 54) = 8.088$ ,  $p < .001$ ,  $\eta_p^2 = 0.231$ ] was explained by the strong difference between ILPs distributions in discrete stimuli (words and strings of hashes) vs. continuous stimuli (solid lines) [ $F(1,54) = 10.08$ ,  $p < .01$ ], the difference between words and strings of hashes being non-significant [ $F < 1$ ].

### **3.2.2. Latency**

As detailed in Table 2, for both groups of NF1, there was a main effect of stimulus type [ $F(2,20) = 19.360$ ,  $p < .001$ ,  $\eta_p^2 = .638$  and  $F(2, 16) = 50.890$ ,  $p < .001$ ,  $\eta_p^2 = .864$ , for NF1RD and NF1noRD respectively]. This difference was mainly explained by shorter

latencies for discrete stimuli than for continuous stimuli [ $F(1,20) = 38.620$ ,  $p < .001$  and  $F(1,16) = 101.705$ ,  $p < .001$ , for NF1RD and NF1noRD respectively].

Consistent with what had previously been observed for saccade size, the saccade latencies of NF1RD children were similar to those of DD children. There was a main effect of stimulus type [ $F(2,62) = 40.205$ ,  $p < .001$ ,  $\eta_p^2 = .565$ ] regardless of the group [ $F < 1$ ]. Similarly, the saccade latencies in NF1noRD were similar to those found in TD children, with shorter latencies when targeting discrete stimuli and longer saccade latencies for continuous stimuli [ $F(2,54) = 119.607$ ,  $p < .001$ ,  $\eta_p^2 = .816$ ]. There was no main effect of group [ $F < 1$ ] nor interaction between stimulus-type and group [ $F < 1$ ]. These results seem to suggest that it is less time-consuming to target the PVL in a word (or in a linguistic-like stimulus) than to target the middle of a solid line.

### 3.2.3. Refixation probability

We found a significant interaction between stimulus type and presentation side, for both groups of NF1 [ $F(2,20) = 3.569$ ,  $p = .047$ ,  $\eta_p^2 = 0.263$  and  $F(2,16) = 3.639$ ,  $p = .050$ ,  $\eta_p^2 = .313$  for NF1RD and NF1noRD respectively]. In line with what had previously been observed for TD children, pairwise comparisons in the NF1noRD group revealed a significant effect of the VF for words on the refixation probability [ $F(1,16) = 10.156$ ,  $p < .005$  for words, and no VF difference for strings of hashes and solid lines,  $F_s < 1$ ]. For the NF1RD group, pairwise comparisons yielded an opposite effect, with children performing more refixation on words in the RVF [ $F(1,20) = 9.028$ ,  $p < .01$  for words and  $F_s < 1$  for strings of hashes and solid lines].

When pooling the data of NF1noRD and TD children, this interaction remained [ $F(2,54) = 4.666$ ,  $p = .014$ ,  $\eta_p^2 = 0.153$ ], and could be entirely explained by a larger VF asymmetry for words [ $F(1,54) = 20.643$ ,  $p < .001$  for words, and  $F_s < 1$  for strings of hashes and solid lines]. Note that this larger VF asymmetry for words also resulted in a larger “linguistic” effect in



RVF than in LVF. Comparing NF1RD performance with that of DD children, the ANOVA showed no main effect of participant group [ $F < 1$ ] nor double interaction between presentation side, stimulus-type and group [ $F < 1$ ]. We found, however, a main effect of presentation side [ $F(1,29) = 7.448, p = .011, \eta_p^2 = .204$ ] and a presentation side  $\times$  stimulus-type interaction [ $F(2,58) = 4.055, p = .022, \eta_p^2 = 0.123$ ], mainly explained by the cost for word recognition in the RVF [ $F(1,58) = 37.383, p < .001$  for words, and  $F_s < 1$  for the strings of hashes and solid lines]. Here again, the difference between words and non-linguistic stimuli was larger in the RVF than in the LVF (but in reverse). The opposite effects in the VF asymmetries found for children with and without RD suggests that the parafoveal words were processed (even if no linguistic processing was required in the bisection task), resulting in a classical RVF advantage for children without RD and a cost for the children with RD.

To sum up, the eye movements of NF1RD children during the ocular bisection task are different from those of NF1noRD children, but similar to those of DD children. In particular, our results showed a less pronounced left-right asymmetry in the saccade amplitude for RD children than for TD and NF1noRD children. This pattern of result reflects, in the context of reading deficits, less stable ILPs and a not yet automatized saccade targeting strategy. This is consistent with the data obtained on the VP effect in foveal vision, showing reduced left-right asymmetry in the RD children's VP-curve (Bellocchi & Ducrot, 2021; Ducrot et al., 2003). The fact that NF1RD and children with dyslexia showed similar saccade targeting strategy, is consistent with the proposition of Huettig et al. (2018) that "this at least partly reflects their common reduced reading experience rather than a causal impairment due to a reading disorder" (p. 339).

### 3.3. Cross-population results

Each child was assigned an ILP index (their initial-landing position for words in the RVF) and a reading level index (their reading age<sup>3</sup>). The correlations between these two measures are displayed in Figure 4. We found a significant correlation between reading age and the initial landing position at P2 [ $r(59) = .493$ ,  $p < .001$ ], with a higher reading level being associated with a higher probability to land first at the PVL (i.e., P2). A significant negative correlation was also found between reading age and the initial landing position at P1 [ $r(59) = -.266$ ,  $p < .05$ ], suggesting that poor readers are more likely to land at the beginning of the word. Note that no significant correlation was shown for landing positions at P3 [ $r(59) = -.136$ ,  $p > .05$ ].

**[Insert Figure 4. about here]**

Cluster analyses with a three-group solution yielded a relatively distinct separation of participants into clusters with very little overlap. Figure 5 shows the three clusters of participants for the distribution of the ILP frequency at each position and the relationship between reading level and ILP. This allowed for an evaluation of the ways in which our readers are similar or different in reading skills and eye movement behaviour. This analysis strategy is supported by the idea that, as word identification becomes more efficient, children shift from a letter-by-letter sequential decoding strategy (characterised by ILPs close to the word beginning to be able to scan it from left to right) to more parallel processing (with ILPs shifted toward the word centre (see also, Ablinger et al., 2013; Schattka et al., 2010). Cluster 2 comprised almost half of the participants (46%, mainly children without RD) and corresponds to an expert reader strategy with an ILP most of the time at the PVL (i.e., P2).

---

<sup>3</sup> We chose to use the reading age measure since it is frequently used in eye-tracking and dyslexia studies (e.g., Bellocchi et al., 2019; Liu et al., 2022; Prado et al., 2007). Note that this measure was strongly correlated in the present study with the other two reading indices of the Alouette-test [reading accuracy,  $r(59) = .684$ ,  $p < .001$ ; reading fluency  $r(59) = .911$ ,  $p < .001$ ].

The two remaining clusters differed from the cluster of “typical” readers. Cluster 1 (12%, mainly DD) showed a more careful eye-movement strategy (i.e., fixated closer to the beginning of the stimuli, P1), as usually found for beginning readers. It comprised readers with the lowest reading level scores, and this group might stick to serial reading, suggesting an immaturity of the oculomotor control associated with a sequential processing strategy. Members of Cluster 3 (42% of the participants, mainly RD children) presented an intermediate reading level and a flattened curve with an ILP covering a wider area around the centre of the word. This pattern could be the reflect of an oculomotor instability. In that sense, this subgroup of readers also presented severe visual-processing deficits at the DEM-test (-1.27 SD on the VT index and -1.97 SD on the HTaj index, with 41% of the children below -1.5 SD on the VT index, and 59% below -1.5 SD on the HTaj index)<sup>4</sup>.

We want to draw attention to the fact that the 3 clusters differed<sup>5</sup> significantly in terms of reading level [ $F(2,58) = 3.877$ ,  $p < .05$ ,  $\eta_p^2 = .118$ ]. The pairwise comparisons showed a significant higher reading age in the cluster 2 (Mean = 115.61) compared with the cluster 1 (Mean = 94.43;  $p < .05$ ) and 3 (Mean = 100.81;  $p < .05$ ), respectively. However, no significant difference emerged between clusters 1 and 3 in terms of reading age [ $p > .05$ ] and between the 3 clusters in terms of chronological age [ $F(2,58) = 0.647$ ,  $p > .05$ ,  $\eta_p^2 = .022$ ].

To sum up, taken together, the correlation and cluster analyses revealed a strong link between reading expertise and ILP. Cluster analyses also pointed to individual differences, with three subgroups of readers using different saccade targeting strategies, irrespective of whether they have been classified as neurotypical, dyslexic or NF1. In our view, the present

---

<sup>4</sup> Note that the cluster 1 group had significantly poor performance at the DEM-test, as well, probably linked to their slowness in their responses and their low reading age (-2.19 SD on the VT index and -1.65 SD on the HTaj index). No such visual-processing deficits was observed in cluster 2 ( $-1 < SD < 0$ , for the VT and HTaj index).

<sup>5</sup> For these statistical analyses, Kruskal–Wallis tests followed by Dunn–Bonferroni method for pairwise comparisons were carried out.

results strongly suggest that the reading ability and the mode of processing used (shift from a sequential reading to a more parallel orthographic processing) are sufficient to account for the different patterns of eye movement behaviour.

**[Insert Figure 5 about here]**

## **4. DISCUSSION**

The aim of the present study was to investigate the extent to which attentional factors, stimulus features, and reading ability influence the position where the eyes first land in a target (word or non-linguistic stimulus) displayed in parafoveal vision. A secondary objective was to better understand the relationship between saccade eye movements and reading difficulties and to examine the possible modulation of clinical diagnosis (DD and NF1) and reading skills (good and poor readers) on saccade targeting strategies as expressed by ILPs.

### **4.1. What determines where children first fixate?**

As previously demonstrated by Ducrot et al. (2013), our data showed that all children tend to initially fixate words at a point somewhere between the beginning and the centre of the word while executing rightward saccades, regardless of their reading level (with or without RD) or clinical diagnosis (dyslexics or NF1 group). In other words, even with a minimum experience with reading, children show an adult-like PVL pattern in the RVF (for similar results, see Bellocchi et al., 2019; Ducrot et al., 2013; Joseph et al., 2009). Note that rightward saccades represent the most frequent saccades in left-to-right writing systems, leading to early development of automatized procedures for saccade computation.

One aspect that warrants particular attention is the left-right asymmetry in the saccade size launched toward parafoveally located stimuli. As stated before, “linguistic-like” stimuli encourage a saccade targeting strategy that aims to land at the PVL in both VFs, with greater difficulty in reaching this goal for the LVF due to the greater distance and unusual reading direction (Ducrot & Pynte, 2002; Vitu et al., 1995). Our results showed a more pronounced left-right asymmetry for TD and NF1noRD children than for DD and NF1RD readers (see Bellocchi et al., 2019, for similar results with DD). One possible explanation to account for this increase in asymmetry for TD readers was suggested by Huang et al. (2019), as reading experience seems to “increase the ratio between recognition in the centre and recognition away from the centre of fixation, as reflected in the changes of the shape of the so-called Form Resolving Field (which maps the distribution of recognition accuracy onto eccentricity) from symmetrical and reversed-V-shaped to slightly narrower on the side of the reading direction” (see also, Lorusso et al., 2004; Zagarra-Moran & Geiger, 1993). Moreover, children without RD used a saccade targeting strategy that aims to land left of centre (PVL) for all discrete stimuli (i.e., with inter-character spaces), even in the LVF, as already observed in adults (Ducrot & Pynte, 2002). In contrast, for children with RD, the asymmetry was less pronounced and restricted to words only. NF1RD and DD children were influenced by the linguistic nature of stimuli, such that they do tend to initially fixate words closer to their beginning (as compared to the strings of hashes).

The results, therefore, suggest that the pre-processing mechanism responsible for the direction of attentional scanning (left-to-right) proposed in Ducrot and Pynte (2002) is well automatized in children without RD, which “anticipated doing left-to-right attentional scanning of discrete stimuli and took the direction of this upcoming attentional scanning into account when computing the incoming saccade” (Ducrot & Pynte, 2002, p. 1142). Note that the fact that children used different saccade targeting strategies depending on the presence of

inter-character spaces or the linguistic nature of the stimulus suggests that the left–right asymmetry in the PVL effect may not result *only* from oculomotor constraints associated with saccade execution.

#### **4.2. How reading ability affects eye movement behaviour?**

A major aim of the present study was to determine whether reading level and changes in written language experience over the course of DD, NF1 and TD children's developmental trajectory are associated with changes in reading strategies, as indexed by saccade targeting strategies. Firstly, the individual distributions of first fixation positions were modulated by reading skills. Cluster analyses confirmed that the landing position of eye movement saccades on words is a sensitive index of reading efficiency (Ducrot et al., 2013; Joseph et al., 2009; Juhasz et al., 2008), with three subgroups of readers using different saccade targeting strategies, irrespective of whether they have been classified as neurotypical, dyslexic or NF1. Good readers (cluster 2) made their first fixation most of the time at the PVL whereas, in children with poor reading skills, ILPs either occurred at the beginning of the word (cluster 1) or covered a wider area around the centre of the word with high variability (cluster 3). A convincing explanation for these saccade targeting differences is based on the dual-route model of reading (Coltheart et al., 2001). As experience with written language grows, children shift from a letter-by-letter sublexical decoding strategy to more parallel lexical processing (Frith, 1985). In that sense, Tydgat and Grainger (2009) suggested that as children develop their capacity to perform parallel letter processing, their ILPs shift from being close to the beginning of words toward a position closer to the centre of words (see also Huestegge et al., 2009). Meanwhile, visuo-attentional abilities would gradually develop with reading experience (Bosse & Valdois, 2009; Hawelka & Wimmer, 2005). In that vein, children show a smaller span of effective vision than adults (Häikiö et al., 2009; Rayner, 1986), the size of

children's perceptual span increasing with reading proficiency (Sperlich et al., 2016). It may be that in very fluent reading, the visuo-attentional capacity approaches visual acuity constraints for letter encoding (Hautala et al., 2022).

We also explored the impact of the reading level on the VF effects. Children with RD made shorter saccades in words than did TD readers, with ILPs located left of centre for both left and right VF only for the TD children. One possibility is that children with RD experience difficulties in oculomotor control. Shifted ILPs may arise simply because children with RD, as beginning readers, exhibit shorter average saccade amplitude (cluster 1) and/or increased variability (cluster 3) in these saccade amplitudes (see also Ducrot et al., 2013, for a similar finding with a lateralized lexical decision task). In line with this proposal is the fact that an eye-tracking training can improve saccadic eye movements in beginning children, with accurately aiming for the OVP of each word (Lehtimäki & Reilly, 2005). It is also possible that oculomotor behaviour and saccade targeting strategy depend on the efficiency of linguistic processing (Rayner, 1998). Proficient readers with high-quality lexical representations and efficient lexical processing will aim to optimize information uptake from words by initially fixating the OVP. In contrast, beginning or poor readers, relying on a letter-by-letter phonological decoding strategy, will prefer to initially fixate words closer to their beginning. In that sense, shorter incoming saccade amplitudes and increased refixation rates have been reported in poor, dyslexic, beginning, and slow readers (e.g., Ducrot et al., 2013; Hawelka et al., 2010; Huestegge et al., 2009; McKeen et al., 2004; Rayner et al., 2010, but see also Valdois et al., 2004, for the hypothesis that dyslexic readers suffer from a narrowed visual attentional window). However, if the second account was the sole explanation, then why would DD and NF1RD children aim to land to the right of the middle of a word displayed in the LVF? Because acuity drops rapidly with increasing eccentricity, word recognition is assumed to be better when the initial fixation is left rather than right of the

centre of the word. Rather, these results suggest that oculomotor behaviour change as linguistic processing becomes more efficient, thus confirming that poor and proficient readers face different oculomotor, perceptual, and cognitive constraints when reading.

As a whole, the current results support the hypothesis that saccade targeting strategies are an acquired reading skill that develops with reading experience. Conversely, the children's reading strategies (i.e., whether they use lexical or sublexical route) can be understood from the characteristics of saccadic eye movement.

#### **4.3. A Causal Role for Inefficient Eye Movements in Reading Disability?**

Everyone agrees that DD readers' eye movements are different from typical readers. These eye movement differences have been found when reading isolated words/pseudowords, sentences or texts, and were reported in different languages with different orthographic depths (e.g., Hutzler & Wimmer, 2004; Hutzler et al., 2006). It has been suggested that they might result from poor oculomotor control associated with dyslexia (Pavlidis, 1981). However, evidence in support of a low-level oculomotor account of the reading behaviour of dyslexics are highly controversial (e.g., Eden et al., 1994; Stanley et al., 1983; Stein et al., 1988, for reviews, see Rayner, 1998).

Understanding the source of such differences in eye movements has been the focus of this work, a sample of children with dyslexia while also examining another neurodevelopmental disorder, known to present high comorbidity with RD. In line with previous observations, we found different saccadic programming in children with RD. DD children and NF1 with RD presented longer latencies than TD children, in particular for discrete stimuli (see also Bucci et al., 2008). Differences in the left-right asymmetry in the saccade size were also shown between groups, as previously discussed above. RD readers



were able to accurately perceive and use parafoveal visual information in order to guide their eyes during text reading, at least in the usual reading direction. However, they revealed a greater difficulty in reaching the centre of targets presented in the LVF due to the greater distance and an unusual reading direction. McKeben et al. (2004) have previously shown that “a mechanism for adapting the gain of reactive, stimulus-triggered saccades (Deubel, 1995) was active during reading in dyslexics, but that it was quantitatively less developed” (p. 396). Moreover, RD children didn’t show the classical asymmetrical landing position pattern for “linguistic-like” stimuli (with inter-character spaces). This suggests that in poor readers, the preprocessing mechanism that detects the presence or absence of spaces between characters and responsible for the direction of attentional scanning is not entirely operational, in particular for the LVF. This result could be interpreted as a deficient allocation of visual attention in RD children hampering the exact planning of fine-tuned leftward saccades (e.g., Facoetti, 2012; Facoetti et al., 2010; Lobier et al., 2012; Valdois et al., 2004; see Gavril et al., 2021 for a meta-analysis). In that sense, it was reported that the orienting visuospatial attention is impaired in dyslexia with a LVF “mini-neglect” and a RVF over-distractibility (Facoetti, 2001; Hari, 2001; Hari & Renvall, 2001). The similar pattern of results for RD and beginning readers (i.e., ILPs determined by the linguistic nature of stimuli instead of the presence of inter-character spaces; Bellocchi, et al., 2013) speaks against this view. We rather suggest that the lack of sensibility to the discreteness of the stimuli we observed in RD children in the saccade computation, reflects their reading difficulties and/or a lack of experience with written language (see also Hawelka et al., 2006). The cluster analyses support this idea with children of cluster 1 (with the lowest reading level scores) showing an immature saccade targeting strategy (characterised by ILPs near the words’ beginning and sequential reading), which mirrors their poor reading skills. That goes in line with the above findings on the asymmetry of the saccade extent and the data of Franzen et al. (2021) showing “a

laborious and more effortful reading strategy in DD, resembling a pattern observed in beginning and poorer readers” (Franzen et al., 2021, p. 9, see also Ashby et al., 2005; Rayner, 1998). These findings can be accounted for by assuming that visual attention distribution (as expressed by ILP) adapts to the differences in reading experience. Therefore, an important result to retain from our study is that eye movements per se are not a cause of dyslexia.

What about the children of cluster 3? They presented an intermediate reading level and exhibits an increased variance in the saccade amplitudes resulting in a flattened and diffuse LP-curve. According to Snell and Theeuwes (2020), a clean oculomotor behaviour is reflected in a narrower spread of ILPs, around the PVL (as observed in experienced readers of the cluster 2). ILP spreads in cluster 3 readers widened upon an increased number of positions. Note that the cluster 3 comprised 70% of reading-disabled children (mainly DD). Interestingly, it has been reported that individuals with DD demonstrate a more distributed/diffused mode of attention and have difficulty in narrowing their focus of attention (Facoetti et al., 2000, 2003; Franceschini et al., 2012). Such detrimental effect of the increased variability in the ILPs received support from two recent studies (Franzen et al., 2021; Rima & Schmid, 2021). Franzen et al. (2021) found an increased number of unexpected/atypical saccades in readers with dyslexia, thus suggesting that their eyes occasionally move to seemingly random places. Regarding the distribution of the children’s ILPs in the cluster 3, it is conceivable that an increased oculomotor instability would result in more spatial variability in ILPs. Taken together with the performance at the DEM-test, this pattern of results may reveal possible occasional oculomotor control deficits in RD, as already reported in previous studies (e.g., Freedman et al., 2017; Raghuram et al., 2018; Tiadi et al., 2016; Vagge et al., 2015), which might arise from underlying visual-processing deficits, such as visuo-attentional deficits (Facoetti et al., 2003, 2019; Franceschini et al., 2012, 2022; Valdois et al., 2004; Vernet et al., 2022).

The point of particular interest is that poor NF1 readers mirrored the dyslexic eye movement behaviour, with less accuracy and more variability in saccadic programming, no sensibility to the discreteness of the stimuli, particularly in the LVF. Proficient NF1 readers showed fully automatized oculomotor mechanisms for proficient reading, as TD children do, using a saccade targeting strategy that aims to land left of the centre for all discrete stimuli so as to be able to scan the stimulus from left to right, thus left-shifting PLP for both left and right VF. Note that it was the first study investigating eye movement control related to reading context in NF1. We provided key evidence showing a difference in the saccadic targeting strategies used by NF1 children depending on the presence of a RD associated. In addition, the atypical saccade targeting strategies used by NF1RD children (timid eye-movement strategy in cluster 1 and inaccurate ILPs in cluster 3) were not specific to this disease. Finally, the fact that NF1RD and DD children showed similar saccade targeting strategies gives clear empirical support to the idea that this at least partly reflects their common poor reading skills (probably related to their limited reading experience, see Cunningham & Stanovich, 1997; Goswami, 2015) rather than an impairment causally related to a reading disorder.

There are some limitations to the present study. The first limitation regards the generalizability of the findings with the small sample size of the two NF1 groups due to the prevalence of this rare disease and our strict inclusion and exclusion criteria. For instance, we have made the choice not to include NF1 children with comorbid ADHD although the large proportion of ADHD in this population (i.e., 40%, Hyman et al., 2005). But our priority was to ensure that our results accurately reflect saccade targeting strategies associated with a RD without possible modulation related to inattentive or impulsive behaviour. Another limitation comes from the absence of a direct measure of print exposure to measure the amount of children's reading and be able to identify the impoverished reading experience

of RD children as a causal claim. One way could be to systematically include in our future studies traditional self-report questionnaires and print exposure checklists assessing parents' familiarity with children's book titles (see Mol & Bus, 2011, for a meta-analysis). In addition, the heated debate on the causes of dyslexia needs more longitudinal studies starting in kindergarteners to increase the likelihood of being able to reliably identify explicit causal links.

What should be taken from the individual analysis is that most of the DD readers of our sample do not exhibit oculomotor deficits (see Hutzler et al., 2006; Prado et al., 2007, for similar conclusions with non-linguistic tasks). Furthermore, many of the impairments displayed by RD children are also shown by beginning readers or illiterate individuals who have received very little reading instruction, thus suggesting that eye movements are functional and perform the same function in DD readers that they do in proficient readers. We suggest that such differences in performance between children with and without DD are, to a substantial extent, the consequence of their poor experience with written language. Specifically, disruption to oculomotor behaviour simply reflects the fact that many of the component processes involved in learning to read have not yet become fully automatized (LaBerge & Samuels, 1974). For DD readers who exhibited deviant saccade targeting strategies, the possibility of occasional oculomotor control deficits cannot be entirely ruled out supporting the notion that non-linguistic processes could serve as an additional source to explain their impaired reading. Although these low-level oculomotor deficits in dyslexia are difficult to quantify, it would be important to propose a careful assessment of the visual-processing and visuo-attentional skills underlying reading to ensure diagnostic and targeted intervention adapted to their deficiencies.

830

## 831 ACKNOWLEDGEMENTS

832         We are very thankful to all the children and their parents who agreed to contribute to  
833 this study. The authors would also like to thank all the research members of the DYSTAC-  
834 MAP project, especially Dr F. Audic and Dr M.-O. Livet for the inclusion of the children in  
835 the Aix-Marseille hospitals. Finally, we are very grateful for the support and financial  
836 contributions of the “Association Neurofibromatoses et Recklinghausen” and the “Fondation  
837 de France”.

838

## 839 FUNDING

840         This work was supported by the French National Research Agency (ANR-13-APPR-  
841 0010) and by a PhD Fellowship for the first author awarded by the “Fondation de France” and  
842 the “Association Neurofibromatoses et Recklinghausen” (VISALECT\_NF1\_00099576).

843

## 844 DECLARATION OF COMPETING INTEREST

845         The authors have no conflicts of interest to disclose.

- 847 Ablinger, I., Huber, W., Schattka, K. I., & Radach, R. (2013). Recovery in a letter-by-letter  
848 reader: more efficiency at the expense of normal reading strategy. *Neurocase*, 19(3),  
849 236–255. <https://doi.org/10.1080/13554794.2012.667119>
- 850 Ans, B., Carbonnel, S., & Valdois, S. (1998). A connectionist multiple-trace memory model  
851 for polysyllabic word reading. *Psychological Review*, 105(4), 678–723.  
852 <https://doi.org/10.1037/0033-295X.105.4.678-723>
- 853 Arnold, S. S., Payne, J. M., McArthur, G., North, K. N., & Barton, B. (2020). Profiling the  
854 Word Reading Abilities of School-Age Children with Neurofibromatosis Type 1.  
855 *Journal of the International Neuropsychological Society*, 1–13.  
856 <https://doi.org/10.1017/S135561772000106X>
- 857 Ashby, J., Rayner, K., & Clifton, C. (2005). Eye movements of highly skilled and average  
858 readers: Differential effects of frequency and predictability. *Quarterly Journal of*  
859 *Experimental Psychology Section A: Human Experimental Psychology*, 58(6), 1065–  
860 1086. <https://doi.org/10.1080/02724980443000476>
- 861 Bellocchi, S., & Ducrot, S. (2021). “Same, same but different”: The optimal viewing position  
862 effect in developmental dyslexia, developmental coordination disorder and comorbid  
863 disorders. *Dyslexia*, 27(3), 294–311. <https://doi.org/10.1002/dys.1688>
- 864 Bellocchi, S., & Leclercq, V. (2021). Exploring the Moderation Effect of Educational Stage  
865 on Visual Magnocellular Functioning Linked to Reading: A Study in French Primary  
866 School Children. *Children*, 8(2), 68. <https://doi.org/10.3390/children8020068>
- 867 Bellocchi, S., Mancini, J., Jover, M., Huau, A., Ghio, A., André, C., & Ducrot, S. (2013).  
868 Dyslexic readers and saccade computation : effects of reading exposure and visuo-  
869 perceptual constraints [poster communication]. *XVIIIth ESCoP Conference, Budapest,*  
870 *Hungary*. <https://hal.archives-ouvertes.fr/hal-01510227>
- 871 Bellocchi, S., Massendari, D., Grainger, J., & Ducrot, S. (2019). Effects of inter-character  
872 spacing on saccade programming in beginning readers and dyslexics. *Child*  
873 *Neuropsychology*, 25(4), 482–506. <https://doi.org/10.1080/09297049.2018.1504907>
- 874 Bellocchi, S., Muneaux, M., Bastien-Toniazio, M., & Ducrot, S. (2013). I can read it in your  
875 eyes: What eye movements tell us about visuo-attentional processes in developmental  
876 dyslexia. *Research in Developmental Disabilities*, 34(1), 452–460.  
877 <https://doi.org/10.1016/j.ridd.2012.09.002>
- 878 Bellocchi, S., Muneaux, M., Huau, A., Lévêque, Y., Jover, M., & Ducrot, S. (2017).  
879 Exploring the Link between Visual Perception, Visual–Motor Integration, and Reading  
880 in Normal Developing and Impaired Children using DTVP-2. *Dyslexia*, 23(3), 296–315.  
881 <https://doi.org/10.1002/dys.1561>
- 882 Besner, D., Risko, E. F., & Sklair, N. (2005). Spatial Attention as a Necessary Preliminary to  
883 Early Processes in Reading. *Canadian Journal of Experimental Psychology/Revue*  
884 *Canadienne de Psychologie Expérimentale*, 59(2), 99–108.  
885 <https://doi.org/10.1037/h0087465>
- 886 Blythe, H. I. (2014). Developmental Changes in Eye Movements and Visual Information  
887 Encoding Associated With Learning to Read. *Current Directions in Psychological*  
888 *Science*, 23(3), 201–207. <https://doi.org/10.1177/0963721414530145>

- 889 Blythe, H. I., & Joseph, H. S. S. L. (2011). Children's eye movements during reading. In S. P.  
890 Liversedge, I. Gilchrist, & S. Everling (Eds.), *The Oxford Handbook of Eye Movements*  
891 (pp. 634–662). Oxford University Press.  
892 <https://doi.org/10.1093/oxfordhb/9780199539789.013.0036>
- 893 Blythe, H. I., Kirkby, J., & Liversedge, S. (2018). Comments on: “What Is Developmental  
894 Dyslexia?” *Brain Sci.* 2018, 8, 26. The Relationship between Eye Movements and  
895 Reading Difficulties. *Brain Sciences*, 8(6), 100. <https://doi.org/10.3390/brainsci8060100>
- 896 Bosse, M., & Valdois, S. (2009). Influence of the visual attention span on child reading  
897 performance: a cross-sectional study. *Journal of Research in Reading*, 32(2), 230–253.  
898 <https://doi.org/10.1111/j.1467-9817.2008.01387.x>
- 899 Brysbaert, M., & Nazir, T. A. (2005). Visual constraints in written word recognition: evidence  
900 from the optimal viewing-position effect. *Journal of Research in Reading*, 28(3), 216–  
901 228. <https://doi.org/10.1111/j.1467-9817.2005.00266.x>
- 902 Bucci, M. P., Brémond-Gignac, D., & Kapoula, Z. (2008). Poor binocular coordination of  
903 saccades in dyslexic children. *Graefes' Archive for Clinical and Experimental*  
904 *Ophthalmology*, 246(3), 417–428. <https://doi.org/10.1007/s00417-007-0723-1>
- 905 Chaix, Y., Lion-François, L., Kemlin, I., Faure-Marie, N., Castelnau, P., Barbarot, S.,  
906 Preclaire, E., Lelong, S., Gentil, C., Lauwers-Cancès, V., Rivier, F., Rodriguez, D.,  
907 Carniero, M., Iannuzzi, S., Dorison, N., & Schweitzer, E. (2017). Deficit in phonological  
908 processes: a characteristic of the neuropsychological profile of children with NF1. *Child*  
909 *Neuropsychology*, 24(4), 558–574. <https://doi.org/10.1080/09297049.2017.1313970>
- 910 Coltheart, Max, Rastle, K., Perry, C., Langdon, R., & Ziegler, J. (2001). DRC: A dual route  
911 cascaded model of visual word recognition and reading aloud. *Psychological Review*,  
912 108(1), 204–256. <https://doi.org/10.1037/0033-295X.108.1.204>
- 913 Cunningham, A. E., & Stanovich, K. E. (1997). Early reading acquisition and its relation to  
914 reading experience and ability 10 years later. *Developmental Psychology*, 33(6), 934–  
915 945. <https://doi.org/10.1037/0012-1649.33.6.934>
- 916 Deubel, H. (1995). Separate adaptive mechanisms for the control of reactive and volitional  
917 saccadic eye movements. *Vision Research*, 35(23–24), 3529–3540.  
918 [https://doi.org/10.1016/0042-6989\(95\)00058-M](https://doi.org/10.1016/0042-6989(95)00058-M)
- 919 Ducrot, S., & Grainger, J. (2007). Deployment of spatial attention to words in central and  
920 peripheral vision. *Perception and Psychophysics*, 69(4), 578–590.  
921 <https://doi.org/10.3758/BF03193915>
- 922 Ducrot, S., Lété, B., Descottes, C., Muneaux, M., & Ghio, A. (2006). *The Emaa*  
923 *(EyeMovement Acquisition and Analysis) software package*. Unpublished Technical  
924 Report (67 p.), University of Provence.
- 925 Ducrot, S., Lété, B., Sprenger-Charolles, L., Pynte, J., & Billard, C. (2003). The Optimal  
926 Viewing Position Effect in Beginning and Dyslexic Readers. *Current Psychology*  
927 *Letters. Behaviour, Brain & Cognition*, 1(10).
- 928 Ducrot, S., & Pynte, J. (2002). What determines the eyes' landing position in words?  
929 *Perception and Psychophysics*, 64(7), 1130–1144. <https://doi.org/10.3758/BF03194762>
- 930 Ducrot, S., Pynte, J., Ghio, A., & Lété, B. (2013). Visual and linguistic determinants of the  
931 eyes' initial fixation position in reading development. *Acta Psychologica*, 142(3), 287–

298. <https://doi.org/10.1016/j.actpsy.2013.01.013>
- Eden, G. F., Stein, J. F., Wood, H. M., & Wood, F. B. (1994). Differences in eye movements and reading problems in dyslexic and normal children. *Vision Research*, 34(10), 1345–1358. [https://doi.org/10.1016/0042-6989\(94\)90209-7](https://doi.org/10.1016/0042-6989(94)90209-7)
- Eloranta, A., Närhi, V. M., Eklund, K. M., Ahonen, T. P. S., & Aro, T. I. (2018). Resolving reading disability—Childhood predictors and adult- age outcomes. *Dyslexia*, 25(1), dys.1605. <https://doi.org/10.1002/dys.1605>
- Engbert, R., & Krügel, A. (2010). Readers Use Bayesian Estimation for Eye Movement Control. *Psychological Science*, 21(3), 366–371. <https://doi.org/10.1177/0956797610362060>
- Facoetti, A. (2001). Facilitation and inhibition mechanisms of human visuospatial attention in a non-search task. *Neuroscience Letters*, 298(1), 45–48. [https://doi.org/10.1016/S0304-3940\(00\)01719-5](https://doi.org/10.1016/S0304-3940(00)01719-5)
- Facoetti, A. (2012). Spatial attention disorders in developmental dyslexia: towards the prevention of reading acquisition deficits. *Visual Aspect of Dyslexia*, 123–136.
- Facoetti, A., Corradi, N., Ruffino, M., Gori, S., & Zorzi, M. (2010). Visual spatial attention and speech segmentation are both impaired in preschoolers at familial risk for developmental dyslexia. *Dyslexia*, 16(3), 226–239. <https://doi.org/10.1002/dys.413>
- Facoetti, A., Franceschini, S., & Gori, S. (2019). Role of Visual Attention in Developmental Dyslexia. In *Developmental Dyslexia across Languages and Writing Systems* (pp. 307–326). Cambridge University Press. <https://doi.org/10.1017/9781108553377.014>
- Facoetti, A., Luisa Lorusso, M., Paganoni, P., Umiltà, C., & Gastone Mascetti, G. (2003). The role of visuospatial attention in developmental dyslexia: evidence from a rehabilitation study. *Cognitive Brain Research*, 15(2), 154–164. [https://doi.org/10.1016/S0926-6410\(02\)00148-9](https://doi.org/10.1016/S0926-6410(02)00148-9)
- Facoetti, A., Paganoni, P., & Lorusso, M. L. (2000). The spatial distribution of visual attention in developmental dyslexia. *Experimental Brain Research*, 132(4), 531–538. <https://doi.org/10.1007/s002219900330>
- Facoetti, A., Zorzi, M., Cestnick, L., Lorusso, M. L., Molteni, M., Paganoni, P., Umiltà, C., & Mascetti, G. G. (2006). The relationship between visuo-spatial attention and nonword reading in developmental dyslexia. *Cognitive Neuropsychology*, 23(6), 841–855. <https://doi.org/10.1080/02643290500483090>
- Fischer, B., & Weber, H. (1990). Saccadic reaction times of dyslexic and age-matched normal subjects. *Perception*, 19(6), 805–818. <https://doi.org/10.1068/p190805>
- Franceschini, S., Bertoni, S., & Puccio, G. (2022). *Hidden in the Labyrinth : Visuo-spatial Attention Deficit in Children with Developmental Dyslexia*. 1–18.
- Franceschini, S., Gori, S., Ruffino, M., Pedrolli, K., & Facoetti, A. (2012). A causal link between visual spatial attention and reading acquisition. *Current Biology*, 22(9), 814–819. <https://doi.org/10.1016/j.cub.2012.03.013>
- Franzen, L., Stark, Z., & Johnson, A. P. (2021). Individuals with dyslexia use a different visual sampling strategy to read text. *Scientific Reports*, 11(1), 6449. <https://doi.org/10.1038/s41598-021-84945-9>



974 Freedman, E. G., Molholm, S., Gray, M. J., Belyusar, D., & Foxe, J. J. (2017). Saccade  
975 adaptation deficits in developmental dyslexia suggest disruption of cerebellar-dependent  
976 learning. *Journal of Neurodevelopmental Disorders*, 9(1), 36.  
977 <https://doi.org/10.1186/s11689-017-9218-5>

978 Frith, U. (1985). Beneath the surface of developmental dyslexia. In K. E. Patterson, J. C.  
979 Marshall, & M. Coltheart (Eds.), *Surface dyslexia* (Vol. 32, Issue 1).

980 Gagl, B., Hawelka, S., & Hutzler, F. (2014). A similar correction mechanism in slow and  
981 fluent readers after suboptimal landing positions. *Frontiers in Human Neuroscience*,  
982 8(JUNE), 1–10. <https://doi.org/10.3389/fnhum.2014.00355>

983 Gangl, M., Moll, K., Jones, M. W., Banfi, C., Schulte-Körne, G., & Landerl, K. (2018).  
984 Lexical Reading in Dysfluent Readers of German. *Scientific Studies of Reading*, 22(1),  
985 24–40. <https://doi.org/10.1080/10888438.2017.1339709>

986 Garzia, R. P., Richman, J. E., Nicholson, S. B., & Gaines, C. S. (1990). A new visual-verbal  
987 saccade test: the development eye movement test (DEM). *Journal of the American*  
988 *Optometric Association*, 61(2), 124–135.

989 Gavril, L., Roşan, A., & Szamosközi, Ştefan. (2021). The role of visual-spatial attention in  
990 reading development: a meta-analysis. *Cognitive Neuropsychology*, 38(6), 387–407.  
991 <https://doi.org/10.1080/02643294.2022.2043839>

992 Gori, S., & Facoetti, A. (2015). How the visual aspects can be crucial in reading acquisition:  
993 The intriguing case of crowding and developmental dyslexia. *Journal of Vision*, 15(1),  
994 1–20. <https://doi.org/10.1167/15.1.8>

995 Goswami, U. (2015). Sensory theories of developmental dyslexia: three challenges for  
996 research. *Nature Reviews Neuroscience*, 16(1), 43–54. <https://doi.org/10.1038/nrn3836>

997 Häikiö, T., Bertram, R., Hyönä, J., & Niemi, P. (2009). Development of the letter identity  
998 span in reading: Evidence from the eye movement moving window paradigm. *Journal of*  
999 *Experimental Child Psychology*, 102(2), 167–181.  
1000 <https://doi.org/10.1016/j.jecp.2008.04.002>

1001 Hari, R. (2001). Left minineglect in dyslexic adults. *Brain*, 124(7), 1373–1380.  
1002 <https://doi.org/10.1093/brain/124.7.1373>

1003 Hari, R., & Renvall, H. (2001). Impaired processing of rapid stimulus sequences in dyslexia.  
1004 *Trends in Cognitive Sciences*, 5(12), 525–532. [https://doi.org/10.1016/S1364-](https://doi.org/10.1016/S1364-6613(00)01801-5)  
1005 [6613\(00\)01801-5](https://doi.org/10.1016/S1364-6613(00)01801-5)

1006 Hautala, J., Hawelka, S., Loberg, O., & Leppänen, P. H. T. (2022). A dynamic adjustment  
1007 model of saccade lengths in reading for word-spaced orthographies: evidence from  
1008 simulations and invisible boundary experiments. *Journal of Cognitive Psychology*, 34(4),  
1009 435–453. <https://doi.org/10.1080/20445911.2021.2011895>

1010 Hawelka, S., Gagl, B., & Wimmer, H. (2010). A dual-route perspective on eye movements of  
1011 dyslexic readers. *Cognition*, 115(3), 367–379.  
1012 <https://doi.org/10.1016/j.cognition.2009.11.004>

1013 Hawelka, S., Huber, C., & Wimmer, H. (2006). Impaired visual processing of letter and digit  
1014 strings in adult dyslexic readers. *Vision Research*, 46(5), 718–723.  
1015 <https://doi.org/10.1016/j.visres.2005.09.017>

- 1016 Hawelka, S., & Wimmer, H. (2005). Impaired visual processing of multi-element arrays is  
1017 associated with increased number of eye movements in dyslexic reading. *Vision*  
1018 *Research*, 45(7), 855–863. <https://doi.org/10.1016/j.visres.2004.10.007>
- 1019 Hindmarsh, G. P., Black, A. A., White, S. L. J., Hopkins, S., & Wood, J. M. (2021). Eye  
1020 movement patterns and reading ability in children. *Ophthalmic and Physiological Optics*,  
1021 41(5), 1134–1143. <https://doi.org/10.1111/opo.12854>
- 1022 Huang, C., Lorusso, M. L., Luo, Z., & Zhao, J. (2019). Developmental Differences in the  
1023 Relationship Between Visual Attention Span and Chinese Reading Fluency. *Frontiers in*  
1024 *Psychology*, 10. <https://doi.org/10.3389/fpsyg.2019.024>
- 1025 Huestegge, L., Radach, R., Corbic, D., & Huestegge, S. M. (2009). Oculomotor and linguistic  
1026 determinants of reading development: A longitudinal study. *Vision Research*, 49(24),  
1027 2948–2959. <https://doi.org/10.1016/j.visres.2009.09.012>
- 1028 Huettig, F., Lachmann, T., Reis, A., & Petersson, K. M. (2018). Distinguishing cause from  
1029 effect – many deficits associated with developmental dyslexia may be a consequence of  
1030 reduced and suboptimal reading experience. *Language, Cognition and Neuroscience*,  
1031 33(3), 333–350. <https://doi.org/10.1080/23273798.2017.1348528>
- 1032 Hung, Y.-N. (2021). The Science of Reading: The Eyes Cannot Lie. *International Journal of*  
1033 *Education and Literacy Studies*, 9(4), 26. <https://doi.org/10.7575/aiac.ijels.v.9n.4p.26>
- 1034 Hutzler, F., Kronbichler, M., Jacobs, A. M., & Wimmer, H. (2006). Perhaps correlational but  
1035 not causal: No effect of dyslexic readers’ magnocellular system on their eye movements  
1036 during reading. *Neuropsychologia*, 44(4), 637–648.  
1037 <https://doi.org/10.1016/j.neuropsychologia.2005.06.006>
- 1038 Hutzler, F., & Wimmer, H. (2004). Eye movements of dyslexic children when reading in a  
1039 regular orthography. *Brain and Language*, 89(1), 235–242.  
1040 [https://doi.org/10.1016/S0093-934X\(03\)00401-2](https://doi.org/10.1016/S0093-934X(03)00401-2)
- 1041 Hyman, S. L., Shores, A., & North, K. N. (2005). The nature and frequency of cognitive  
1042 deficits in children with neurofibromatosis type 1. *Neurology*, 65(7), 1037–1044.  
1043 <https://doi.org/10.1212/01.wnl.0000179303.72345.ce>
- 1044 Hyönä, J. (1995). An eye movement analysis of topic-shift effect during repeated reading.  
1045 *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 21(5), 1365–  
1046 1373. <https://doi.org/10.1037/0278-7393.21.5.A>
- 1047 Inhoff, A. W., Radach, R., Eiter, B. M., & Juhasz, B. (2003). Distinct Subsystems for the  
1048 Parafoveal Processing of Spatial and Linguistic Information during Eye Fixations in  
1049 Reading. *The Quarterly Journal of Experimental Psychology Section A*, 56(5), 803–827.  
1050 <https://doi.org/10.1080/02724980244000639>
- 1051 Jacquier-Roux, M., Valdois, S., & Zorman, M. (2002). *Outil de dépistage des dyslexies*  
1052 *[Dyslexia screening tool]*. Laboratoire de Cogni-Sciences IUFM, Grenoble, France  
1053 (2002).
- 1054 Joo, S. J., White, A. L., Strodtman, D. J., & Yeatman, J. D. (2018). Optimizing text for an  
1055 individual’s visual system: The contribution of visual crowding to reading difficulties.  
1056 *Cortex*, 103, 291–301. <https://doi.org/10.1016/j.cortex.2018.03.013>
- 1057 Joseph, H. S. S. L., Liversedge, S. P., Blythe, H. I., White, S. J., & Rayner, K. (2009). Word  
1058 length and landing position effects during reading in children and adults. *Vision*

- 1059 *Research*, 49(16), 2078–2086. <https://doi.org/10.1016/j.visres.2009.05.015>
- 1060 Juhasz, B. J., White, S. J., Liversedge, S. P., & Rayner, K. (2008). Eye movements and the  
1061 use of parafoveal word length information in reading. *Journal of Experimental*  
1062 *Psychology: Human Perception and Performance*, 34(6), 1560–1579.  
1063 <https://doi.org/10.1037/a0012319>
- 1064 Kim, Y.-S., & Pallante, D. (2012). Predictors of reading skills for kindergartners and first  
1065 grade students in Spanish: a longitudinal study. *Reading and Writing*, 25(1), 1–22.  
1066 <https://doi.org/10.1007/s11145-010-9244-0>
- 1067 Kirkby, J. A., Barrington, R. S., Drieghe, D., & Liversedge, S. P. (2022). Parafoveal  
1068 processing and transposed- letter effects in dyslexic reading. *Dyslexia*, 28(3), 359–374.  
1069 <https://doi.org/10.1002/dys.1721>
- 1070 Krab, L. C., Aarsen, F. K., de Goede-Bolder, A., Catsman-Berrevoets, C. E., Arts, W. F.,  
1071 Moll, H. A., & Elgersma, Y. (2008). Impact of neurofibromatosis type 1 on school  
1072 performance. *J Child Neurol*, 23(9), 1002–1010. <https://doi.org/0883073808316366>  
1073 [pii]10.1177/0883073808316366 [doi]
- 1074 LaBerge, D., & Brown, V. (1989). Theory of attentional operations in shape identification.  
1075 *Psychological Review*, 96(1), 101–124. <https://doi.org/10.1037/0033-295X.96.1.101>
- 1076 LaBerge, D., & Samuels, S. J. (1974). Toward a theory of automatic information processing  
1077 in reading. *Cognitive Psychology*, 6(2), 293–323. [https://doi.org/10.1016/0010-](https://doi.org/10.1016/0010-0285(74)90015-2)  
1078 [0285\(74\)90015-2](https://doi.org/10.1016/0010-0285(74)90015-2)
- 1079 Landerl, K., Freudenthaler, H. H., Heene, M., De Jong, P. F., Desrochers, A., Manolitsis, G.,  
1080 Parrila, R., & Georgiou, G. K. (2019). Phonological Awareness and Rapid Automatized  
1081 Naming as Longitudinal Predictors of Reading in Five Alphabetic Orthographies with  
1082 Varying Degrees of Consistency. *Scientific Studies of Reading*, 23(3), 220–234.  
1083 <https://doi.org/10.1080/10888438.2018.1510936>
- 1084 Lasker, a G., Denckla, M. B., & Zee, D. S. (2003). Ocular motor behavior of children with  
1085 neurofibromatosis 1. *Journal of Child Neurology*, 18(5), 348–355.  
1086 <https://doi.org/10.1177/08830738030180050301>
- 1087 Leclercq, V., & Siéoff, E. (2016). Attentional Processing of Letter Strings by Children. *Child*  
1088 *Neuropsychology*, 22(1), 110–132. <https://doi.org/10.1080/09297049.2014.977242>
- 1089 Lefavrais, P. (1967). *Test de l'alouette: manuel*. Les éditions du centre de psychologie  
1090 appliquée.
- 1091 Lefavrais, P. (2005). *Alouette-R*. Les éditions du centre de psychologie appliquée.
- 1092 Lefton, L. A., Nagle, R. J., Johnson, G., & Fisher, D. F. (1979). Eye Movement Dynamics of  
1093 Good and Poor Readers: Then and Now. *Journal of Reading Behavior*, 11(4), 319–328.  
1094 <https://doi.org/10.1080/10862967909547338>
- 1095 Legge, G. E., Klitz, T. S., & Tjan, B. S. (1997). Mr. Chips: An ideal-observer model of  
1096 reading. *Psychological Review*, 104(3), 524–553. [https://doi.org/10.1037/0033-](https://doi.org/10.1037/0033-295X.104.3.524)  
1097 [295X.104.3.524](https://doi.org/10.1037/0033-295X.104.3.524)
- 1098 Lehtimäki, T. M., & Reilly, R. G. (2005). Improving eye movement control in young readers.  
1099 *Artificial Intelligence Review*, 24(3–4), 477–488. [https://doi.org/10.1007/s10462-005-](https://doi.org/10.1007/s10462-005-9010-x)  
1100 [9010-x](https://doi.org/10.1007/s10462-005-9010-x)

- 1101 Leibnitz, L., Grainger, J., Muneaux, M., & Ducrot, S. (2017). Processus visuo-attentionnels et  
1102 lecture : une synthèse. *L'Année Psychologique*, 116(4), 597–622.  
1103 <https://doi.org/10.4074/S0003503316000403>
- 1104 Leppänen, U., Niemi, P., Aunola, K., & Nurmi, J.-E. (2004). Development of reading skills  
1105 among preschool and primary school pupils. *Reading Research Quarterly*, 39(1), 72–93.  
1106 <https://doi.org/10.1598/RRQ.39.1.5>
- 1107 Lété, B., Sprenger-Charolles, L., & Colé, P. (2004). MANULEX: A grade-level lexical  
1108 database from French elementary school readers. *Behavior Research Methods,*  
1109 *Instruments, & Computers*, 36(1), 156–166. <https://doi.org/10.3758/BF03195560>
- 1110 Li, J., Yang, Y., & Zhao, J. (2021). The development of visual simultaneous processing skill  
1111 subcomponents of Chinese children with developmental dyslexia and the relationship  
1112 with reading. *Acta Psychologica Sinica*, 53(8), 821.  
1113 <https://doi.org/10.3724/SP.J.1041.2021.00821>
- 1114 Liu, T., Thiebaut de Schotten, M., Altarelli, I., Ramus, F., & Zhao, J. (2022). Neural  
1115 dissociation of visual attention span and phonological deficits in developmental dyslexia:  
1116 A hub- based white matter network analysis. *Human Brain Mapping, November 2021*,  
1117 1–10. <https://doi.org/10.1002/hbm.25997>
- 1118 Lobier, M., Zoubrinetzky, R., & Valdois, S. (2012). The visual attention span deficit in  
1119 dyslexia is visual and not verbal. *Cortex*, 48(6), 768–773.  
1120 <https://doi.org/10.1016/j.cortex.2011.09.003>
- 1121 Lorusso, M. L., Facoetti, A., Pesenti, S., Cattaneo, C., Molteni, M., & Geiger, G. (2004).  
1122 Wider recognition in peripheral vision common to different subtypes of dyslexia. *Vision*  
1123 *Research*, 44(20), 2413–2424. <https://doi.org/10.1016/j.visres.2004.05.001>
- 1124 McConkie, G. W., Kerr, P. W., Reddix, M. D., & Zola, D. (1988). Eye movement control  
1125 during reading: I. The location of initial eye fixations on words. *Vision Research*, 28(10),  
1126 1107–1118. [https://doi.org/10.1016/0042-6989\(88\)90137-X](https://doi.org/10.1016/0042-6989(88)90137-X)
- 1127 McDonald, R., Lerman, K., & Pereira, F. (2006). Multilingual dependency analysis with a  
1128 two-stage discriminative parser. *CoNLL 2006 - Proceedings of the 10th Conference on*  
1129 *Computational Natural Language Learning, June*, 216–220.
- 1130 McKeen, M., Trauzettel-Klosinski, S., Reinhardt, J., Duerrwaechter, U., Adler, M., &  
1131 Klosinski, G. (2004). Eye movement control during single-word reading in dyslexics.  
1132 *Journal of Vision*, 4(5), 388–402. <https://doi.org/10.1167/4.5.4>
- 1133 Mol, S. E., & Bus, A. G. (2011). To read or not to read: A meta-analysis of print exposure  
1134 from infancy to early adulthood. *Psychological Bulletin*, 137(2), 267–296.  
1135 <https://doi.org/10.1037/a0021890>
- 1136 Nazir, T. A. (2000). Traces of Print Along the Visual Pathway. In *Reading as a Perceptual*  
1137 *Process* (pp. 3–22). Elsevier. <https://doi.org/10.1016/B978-008043642-5/50003-6>
- 1138 Nazir, T. A., Ben-Boutayab, N., Decoppet, N., Deutsch, A., & Frost, R. (2004). Reading  
1139 habits, perceptual learning, and recognition of printed words. *Brain and Language*,  
1140 88(3), 294–311. [https://doi.org/10.1016/S0093-934X\(03\)00168-8](https://doi.org/10.1016/S0093-934X(03)00168-8)
- 1141 Nuthmann, A., Engbert, R., & Kliegl, R. (2005). Mislocated fixations during reading and the  
1142 inverted optimal viewing position effect. *Vision Research*, 45(17), 2201–2217.  
1143 <https://doi.org/10.1016/j.visres.2005.02.014>

- 1144 O'Regan, J. K., & Lévy-Schoen, A. (1987). Eye-movement strategy and tactics in word  
1145 recognition and reading. In M. Coltheart (Ed.), *Attention and performance 12: The*  
1146 *psychology of reading* (pp. 363–383). Lawrence Erlbaum Associates, Inc.
- 1147 Pavlidis, G. T. (1981). Do eye movements hold the key to dyslexia? *Neuropsychologia*, 19(1),  
1148 57–64. [https://doi.org/10.1016/0028-3932\(81\)90044-0](https://doi.org/10.1016/0028-3932(81)90044-0)
- 1149 Perea, M., Panadero, V., Moret-Tatay, C., & Gómez, P. (2012). The effects of inter-letter  
1150 spacing in visual-word recognition: Evidence with young normal readers and  
1151 developmental dyslexics. *Learning and Instruction*, 22(6), 420–430.  
1152 <https://doi.org/10.1016/j.learninstruc.2012.04.001>
- 1153 Pirozzolo, F. J., & Rayner, K. (1979). The Neural Control of Eye Movements in Acquired and  
1154 Developmental Reading Disorders. In H. Whitaker & H. A. Whitaker (Eds.), *Studies in*  
1155 *Neurolinguistics* (pp. 97–123). Elsevier. [https://doi.org/10.1016/B978-0-12-746304-](https://doi.org/10.1016/B978-0-12-746304-9.50009-4)  
1156 [9.50009-4](https://doi.org/10.1016/B978-0-12-746304-9.50009-4)
- 1157 Plaza, M., & Cohen, H. (2007). The contribution of phonological awareness and visual  
1158 attention in early reading and spelling. *Dyslexia*, 239(13), 234–239.  
1159 <https://doi.org/10.1002/dys>
- 1160 Prado, C., Dubois, M., & Valdois, S. (2007). The eye movements of dyslexic children during  
1161 reading and visual search: Impact of the visual attention span. *Vision Research*, 47(19),  
1162 2521–2530. <https://doi.org/10.1016/j.visres.2007.06.001>
- 1163 Premeti, A., Bucci, M. P., & Isel, F. (2022). Evidence from ERP and Eye Movements as  
1164 Markers of Language Dysfunction in Dyslexia. *Brain Sciences*, 12(1), 73.  
1165 <https://doi.org/10.3390/brainsci12010073>
- 1166 Radach, R., & Kempe, V. (1993). An individual analysis of initial fixation positions in  
1167 reading. In *Perception and cognition: Advances in eye movement research*. (pp. 213–  
1168 225). North-Holland/Elsevier Science Publishers.
- 1169 Radach, R., & McConkie, G. W. (1998). Determinants of Fixation Positions in Words During  
1170 Reading. In *Eye Guidance in Reading and Scene Perception* (pp. 77–100). Elsevier.  
1171 <https://doi.org/10.1016/B978-008043361-5/50005-5>
- 1172 Raghuram, A., Gowrisankaran, S., Swanson, E., Zurakowski, D., Hunter, D. G., & Waber, D.  
1173 P. (2018). Frequency of Visual Deficits in Children With Developmental Dyslexia.  
1174 *JAMA Ophthalmology*, 136(10), 1089.  
1175 <https://doi.org/10.1001/jamaophthalmol.2018.2797>
- 1176 Rayner, K. (1975). Parafoveal identification during a fixation in reading. *Acta Psychologica*,  
1177 39(4), 271–281. [https://doi.org/10.1016/0001-6918\(75\)90011-6](https://doi.org/10.1016/0001-6918(75)90011-6)
- 1178 Rayner, K. (1979). Eye guidance in reading: Fixation locations within words. *Perception*,  
1179 8(6), 21–30.
- 1180 Rayner, K. (1986). Eye movements and the perceptual span in beginning and skilled readers.  
1181 *Journal of Experimental Child Psychology*, 41(2), 211–236.  
1182 [https://doi.org/10.1016/0022-0965\(86\)90037-8](https://doi.org/10.1016/0022-0965(86)90037-8)
- 1183 Rayner, K. (1998). Eye movements in reading and information processing: 20 years of  
1184 research. *Psychological Bulletin*, 124(3), 372–422. [https://doi.org/10.1037/0033-](https://doi.org/10.1037/0033-2909.124.3.372)  
1185 [2909.124.3.372](https://doi.org/10.1037/0033-2909.124.3.372)

- 1186 Rayner, K. (2009). Eye Movements in Reading: Models and Data. *Journal of Eye Movement*  
1187 *Research*, 2(5), 1–10. <https://doi.org/10.16910/jemr.2.5.2>
- 1188 Rayner, K., Fischer, M. H., & Pollatsek, A. (1998). Unspaced text interferes with both word  
1189 identification and eye movement control. *Vision Research*, 38(8), 1129–1144.  
1190 [https://doi.org/10.1016/S0042-6989\(97\)00274-5](https://doi.org/10.1016/S0042-6989(97)00274-5)
- 1191 Rayner, K., Slattery, T. J., & Bélanger, N. N. (2010). Eye movements, the perceptual span,  
1192 and reading speed. *Psychonomic Bulletin & Review*, 17(6), 834–839.  
1193 <https://doi.org/10.3758/PBR.17.6.834>
- 1194 Rayner, K., White, S. J., Kambe, G., Miller, B., & Liversedge, S. (2003). On the processing of  
1195 meaning from parafoveal vision during eye fixations in reading. In J. Hyönä, R. Radach,  
1196 & H. Deubel (Eds). *The Mind's Eye: Cognitive and Applied Aspects of Eye Movement*  
1197 *Research*, 213–234.
- 1198 Reichle, E. D., Rayner, K., & Pollatsek, A. (2003). The E-Z Reader model of eye-movement  
1199 control in reading: Comparisons to other models. *Behavioral and Brain Sciences*, 26(4),  
1200 445–476. <https://doi.org/10.1017/S0140525X03000104>
- 1201 Ribeiro, M. J., Violante, I. R., Ribeiro, I., Bernardino, I., Ramos, F., Saraiva, J., Reviriego, P.,  
1202 Upadhyaya, M., Silva, E. D., & Castelo-Branco, M. (2012). Abnormal achromatic and  
1203 chromatic contrast sensitivity in neurofibromatosis type 1. *Investigative Ophthalmology*  
1204 *and Visual Science*, 53(1), 287–293. <https://doi.org/10.1167/iovs.11-8225>
- 1205 Rima, S., & Schmid, M. C. (2021). Reading Specific Small Saccades Predict Individual  
1206 Phonemic Awareness and Reading Speed. *Frontiers in Neuroscience*, 15(December), 1–  
1207 13. <https://doi.org/10.3389/fnins.2021.663242>
- 1208 Schattka, K. I., Radach, R., & Huber, W. (2010). Eye movement correlates of acquired central  
1209 dyslexia. *Neuropsychologia*, 48(10), 2959–2973.  
1210 <https://doi.org/10.1016/j.neuropsychologia.2010.06.005>
- 1211 Silva, S., Faisca, L., Araújo, S., Casaca, L., Carvalho, L., Petersson, K. M., & Reis, A. (2016).  
1212 Too little or too much? Parafoveal preview benefits and parafoveal load costs in dyslexic  
1213 adults. *Annals of Dyslexia*, 66(2), 187–201. <https://doi.org/10.1007/s11881-015-0113-z>
- 1214 Sireteanu, R., Goertz, R., Bachert, I., & Wandert, T. (2005). Children with developmental  
1215 dyslexia show a left visual “minineglect.” *Vision Research*, 45(25–26), 3075–3082.  
1216 <https://doi.org/10.1016/j.visres.2005.07.030>
- 1217 Smart, D., Youssef, G. J., Sanson, A., Prior, M., Toumbourou, J. W., & Olsson, C. A. (2017).  
1218 Consequences of childhood reading difficulties and behaviour problems for educational  
1219 achievement and employment in early adulthood. *British Journal of Educational*  
1220 *Psychology*, 87(2), 288–308. <https://doi.org/10.1111/bjep.12150>
- 1221 Snell, J., & Theeuwes, J. (2020). A story about statistical learning in a story: Regularities  
1222 impact eye movements during book reading. *Journal of Memory and Language*,  
1223 113(August 2019), 104127. <https://doi.org/10.1016/j.jml.2020.104127>
- 1224 Sperlich, A., Meixner, J., & Laubrock, J. (2016). Development of the perceptual span in  
1225 reading: A longitudinal study. *Journal of Experimental Child Psychology*, 146, 181–201.  
1226 <https://doi.org/10.1016/j.jecp.2016.02.007>
- 1227 Spinelli, D., De Luca, M., Judica, A., & Zoccolotti, P. (2002). Crowding effects on word  
1228 identification in developmental dyslexia. *Cortex*, 38(2), 179–200.

- 1229 [https://doi.org/10.1016/S0010-9452\(08\)70649-X](https://doi.org/10.1016/S0010-9452(08)70649-X)
- 1230 Stanley, G., Smith, G. A., & Howell, E. A. (1983). Eye-movements and sequential tracking in  
1231 dyslexic and control children. *British Journal of Psychology*, 74(2), 181–187.  
1232 <https://doi.org/10.1111/j.2044-8295.1983.tb01852.x>
- 1233 Stein, J. F. (2001). The magnocellular theory of developmental dyslexia. *Dyslexia*, 7(1), 12–  
1234 36. <https://doi.org/10.1002/dys.186>
- 1235 Stein, J. F., Riddell, P. M., & Fowler, S. (1988). Disordered vergence control in dyslexic  
1236 children. *British Journal of Ophthalmology*, 72(3), 162–166.  
1237 <https://doi.org/10.1136/bjo.72.3.162>
- 1238 Tiadi, A., Gérard, C.-L., Peyre, H., Bui-Quoc, E., & Bucci, M. P. (2016). Immaturity of  
1239 Visual Fixations in Dyslexic Children. *Frontiers in Human Neuroscience*, 10(FEB2016),  
1240 1–7. <https://doi.org/10.3389/fnhum.2016.00058>
- 1241 Tydgate, I., & Grainger, J. (2009). Serial Position Effects in the Identification of Letters,  
1242 Digits, and Symbols. *Journal of Experimental Psychology: Human Perception and*  
1243 *Performance*, 35(2), 480–498. <https://doi.org/10.1037/a0013027>
- 1244 Vagge, A., Cavanna, M., Traverso, C. E., & Iester, M. (2015). Evaluation of ocular  
1245 movements in patients with dyslexia. *Annals of Dyslexia*, 65(1), 24–32.  
1246 <https://doi.org/10.1007/s11881-015-0098-7>
- 1247 Valdois, S., Bosse, M.-L., & Tainturier, M.-J. (2004). The cognitive deficits responsible for  
1248 developmental dyslexia: Review of evidence for a selective visual attentional disorder.  
1249 *Dyslexia*, 10(4), 339–363. <https://doi.org/10.1002/dys.284>
- 1250 Vernet, M., Jover, M., Bellocchi, S., Maziero, S., Jucla, M., Tallet, J., Danna, J., Chaix, Y., &  
1251 Ducrot, S. (2022). Visual-processing deficits in children with neurofibromatosis type 1:  
1252 A clinical marker of reading difficulties. *European Journal of Paediatric Neurology*, 38,  
1253 25–32. <https://doi.org/10.1016/j.ejpn.2022.03.009>
- 1254 Vidyasagar, T. R., & Pammer, K. (2010). Dyslexia: a deficit in visuo-spatial attention, not in  
1255 phonological processing. *Trends in Cognitive Sciences*, 14(2), 57–63.  
1256 <https://doi.org/10.1016/j.tics.2009.12.003>
- 1257 Vitu, F. (2003). The basic assumptions of E-Z Reader are not well-founded. *Behavioral and*  
1258 *Brain Sciences*, 26(4), 506–507. <https://doi.org/10.1017/S0140525X0351010X>
- 1259 Vitu, F., O'Regan, J. K., Inhoff, A. W., & Topolski, R. (1995). Mindless reading: Eye-  
1260 movement characteristics are similar in scanning letter strings and reading texts.  
1261 *Perception & Psychophysics*, 57(3), 352–364. <https://doi.org/10.3758/BF03213060>
- 1262 Vitu, F., O'Regan, J. K., & Mittau, M. (1990). Optimal landing position in reading isolated  
1263 words and continuous text. *Perception & Psychophysics*, 47(6), 583–600.  
1264 <https://doi.org/10.3758/BF03203111>
- 1265 Vorstius, C., Radach, R., & Lonigan, C. J. (2014). Eye movements in developing readers: A  
1266 comparison of silent and oral sentence reading. *Visual Cognition*, 22(3–4), 458–485.  
1267 <https://doi.org/10.1080/13506285.2014.881445>
- 1268 White, S. J., & Liversedge, S. P. (2006). Foveal processing difficulty does not modulate non-  
1269 foveal orthographic influences on fixation positions. *Vision Research*, 46(3), 426–437.  
1270 <https://doi.org/10.1016/j.visres.2005.07.006>

- 1271 World Health Organisation. (2008). *59th general assembly* (A. WM (ed.)). WHO.
- 1272 Zegarra-Moran, O., & Geiger, G. (1993). Visual Recognition in the Peripheral Field: Letters  
1273 versus Symbols and Adults versus Children. *Perception*, 22(1), 77–90.  
1274 <https://doi.org/10.1068/p220077>
- 1275 Zorzi, M., Barbiero, C., Facoetti, A., Lonciari, I., Carrozzi, M., Montico, M., Bravar, L.,  
1276 George, F., Pech-Georgel, C., & Ziegler, J. C. (2012). Extra-large letter spacing  
1277 improves reading in dyslexia. *Proceedings of the National Academy of Sciences of the*  
1278 *United States of America*, 109(28), 11455–11459.  
1279 <https://doi.org/10.1073/pnas.1205566109>
- 1280



1281 Table 1. Mean of initial saccade size (in characters), saccade latency (in milliseconds) and  
1282 refixation probability (in percent) as a function of stimulus type, visual field and group in  
1283 Experiment 1.

	Initial saccade size		Saccade latency		Refixation probability	
	Mean (SD)		Mean (SD)		Mean (SD)	
	TD	DD	TD	DD	TD	DD
Left Visual Field						
Words	4.33 (0.74)	3.95 (1.30)	170.55 (13.86)	194.94 (28.02)	74.32 (19.69)	55.75 (24.52)
Strings of hashes	4.34 (0.93)	3.72 (0.64)	173.59 (19.96)	193.64 (25.79)	69.24 (24.72)	49.20 (22.42)
Solid lines	3.59 (0.84)	3.64 (0.99)	217.92 (28.47)	220.19 (29.70)	69.02 (19.53)	48.93 (21.98)
Right Visual Field						
Words	3.40 (0.78)	3.27 (0.87)	173.85 (21.51)	191.36 (33.15)	53.64 (28.45)	72.34 (24.84)
Strings of hashes	3.49 (0.54)	3.63 (0.56)	173.91 (21.84)	192.18 (38.67)	70.88 (22.71)	59.44 (22.77)
Solid lines	3.54 (0.59)	3.60 (0.57)	221.18 (17.23)	220.63 (24.91)	71.43 (24.44)	60.17 (26.62)

1284 *Notes.* Saccade size was measured according to the central fixation point. TD: typically  
1285 developing children; DD: children with developmental dyslexia.

1286

1287 Table 2. Mean of initial saccade size (in characters), saccade latency (in milliseconds) and  
1288 refixation probability (in percent) as a function of stimulus type, visual field, and group in  
1289 Experiment 2.

	Initial saccade size		Saccade latency		Refixation probability	
	Mean (SD)		Mean (SD)		Mean (SD)	
	NF1noRD	NF1RD	NF1noRD	NF1RD	NF1noRD	NF1RD
Left Visual Field						
Words	4.46 (0.71)	4.22 (0.52)	173.91 (8.81)	188.88 (12.88)	74.96 (25.10)	47.79 (26.49)
Strings of hashes	4.43 (0.26)	3.55 (0.55)	175.38 (11.29)	191.10 (19.02)	71.03 (23.59)	51.89 (23.82)
Solid lines	3.67 (0.96)	3.64 (1.22)	219.34 (32.24)	222.05 (35.39)	69.05 (24.85)	45.98 (23.06)
Right Visual Field						
Words	3.43 (0.57)	3.34 (0.63)	173.47 (15.93)	182.89 (12.20)	53.92 (33.42)	72.41 (21.02)
Strings of hashes	3.50 (0.90)	3.53 (0.70)	174.61 (14.54)	184.78 (17.63)	70.24 (25.97)	47.05 (29.55)
Solid lines	3.67 (1.18)	3.52 (0.65)	220.33 (22.14)	222.51 (25.85)	71.11 (30.09)	47.65 (23.47)

1290 *Notes.* Initial saccade size was measured according to the central fixation point. NF1RD: NF1  
1291 children with reading disorders; NF1noRD: NF1 children without reading disorders.





Figure 1. Procedure used in the oculomotor lateralized bisection task. The different display conditions of the stimuli are represented in this figure with the 3 types of stimuli (words, hashes and lines) and the 2 visual fields of presentation (LVF and RVF). When the target item appeared, participants were instructed to move their eyes to look at what they considered to be the middle of the item (see arrows and dashed vertical lines in the figure).



1298

1299 Figure 2. Individual distribution of first fixation position specifically for words presented in  
 1300 the RVF in NF1noRD (i.e., left part of the figure) and NF1RD (i.e., right part of the figure)  
 1301 children.

1302 *Notes.* This representation was made only for the words presented in the RVF to observe the  
 1303 oculomotor behaviour in the most reading-like situation. Each line represents the individual  
 1304 performance of a different participant. NF1RD: NF1 children with reading disorders;  
 1305 NF1noRD: NF1 children without reading disorders.





Figure 3. Comparison of the initial landing position's frequency for words in the RVF between the 4 groups of children in Experiment 1 and 2.


*Notes.* This representation was made only for the words presented in the RVF to observe the oculomotor behaviour in the most reading-like situation. TD: Typically developing children; DD: children with developmental dyslexia; NF1RD: NF1 children with reading disorders; NF1noRD: NF1 children without reading disorders.



1313

1314 Figure 4. Frequency of the initial landing of the first saccade (in percentage) at each position  
 1315 of words according to reading age (in months) and groups.

1316 Notes. These representations were made only for the words presented in the RVF to observe  
 1317 the oculomotor behaviour in the most reading-like situation. TD: Typically developing  
 1318 children; DD: children with developmental dyslexia; NF1RD: NF1 children with reading  
 1319 disorders; NF1noRD: NF1 children without reading disorders.



1320

1321 Figure 5. Distributions of the initial landing position (in percentage) according to the 3  
 1322 clusters found for all groups combined.

1323 *Notes.* This figure was made only for the words presented in the RVF to observe the  
 1324 oculomotor behaviour in the most reading-like situation.