

# The determinants of saccade targeting strategy in neurodevelopmental disorders: the influence of suboptimal reading experience

Marie Vernet, Stéphanie Bellocchi, Jérémy Danna, Delphine Massendari,

Marianne Jover, Yves Chaix, Stéphanie Ducrot

# ▶ To cite this version:

Marie Vernet, Stéphanie Bellocchi, Jérémy Danna, Delphine Massendari, Marianne Jover, et al.. The determinants of saccade targeting strategy in neurodevelopmental disorders: the influence of suboptimal reading experience. Vision Research, 2023, 204, pp.108162. 10.1016/j.visres.2022.108162 . hal-03924190

# HAL Id: hal-03924190 https://hal.science/hal-03924190v1

Submitted on 5 Jan 2023  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	The determinants of saccade targeting strategy in neurodevelopmental disorders: the
2	influence of suboptimal reading experience
3	Marie Vernet <sup>a,b,e</sup> , Stéphanie Bellocchi <sup>d</sup> , Jérémy Danna <sup>e</sup> , Delphine Massendari <sup>b</sup> , Marianne
4	Jover <sup>c</sup> , Yves Chaix <sup>a,f</sup> & Stéphanie Ducrot <sup>b</sup>
5	<sup>a</sup> ToNIC, Toulouse NeuroImaging Center, University of Toulouse, Inserm, UPS, France
6	<sup>b</sup> Aix Marseille Univ, CNRS, LPL, Aix-en-Provence, France
7	<sup>c</sup> Aix Marseille Univ, PSYCLE, Aix-en-Provence, France
8	<sup>d</sup> Univ. Paul Valéry Montpellier 3, EPSYLON UR 4556, F34000, Montpellier, France
9	<sup>e</sup> CLLE, Université de Toulouse, CNRS, Toulouse, France
10	<sup>f</sup> Children's Hospital, Toulouse-Purpan University Hospital, Toulouse, France
11	
12	
13 14	SHORT RUNNING TITLE: Saccade targeting strategy in pathological reading
14	SHOKI KONNING IIILE. Saccade targeting strategy in pathological reading
16	
17	
18	
19	Correspondence
20	Marie Vernet
21	Toulouse Neuro-Imaging Center
22	Pavillon Baudot – CHU Purpan
23	Place du Dr Joseph Baylac
24	31024 Toulouse, France
25	E-mail: marie.vernet@inserm.fr

#### ABSTRACT

27 Whether eye-movements deficits are causal in reading disorders (RD) or rather a consequence of linguistic processing difficulty experienced by disabled readers has been 28 extensively debated. Since RD are frequently comorbid with the Neurofibromatosis type1 29 (NF1), children with NF1 were used as a comparison group for children with dyslexia in this 30 study. Eye movements were recorded while 21 dyslexic, 20 NF1, and 20 typically developing 31 children performed an oculomotor lateralized bisection task. In this experiment, we 32 manipulated the type of stimulus - discrete (words and strings of hashes) versus continuous 33 (solid lines) - and the visual field where the stimulus was displayed (left vs. right). The results 34 35 showed that (1) only proficient readers (TD and NF1 without RD) showed fully developed 36 oculomotor mechanisms for efficient reading, with a clear preferred viewing location located to the left of the word's centre in both visual fields, and fine-tuned saccade targeting guided 37 by the between-character space information and (2) NF1 poor readers mirrored the dyslexic 38 eye movement behaviour, with less accuracy and more variability in saccadic programming, 39 no sensitivity to the discreteness of the stimuli, particularly in the left visual field. We 40 concluded that disruption to oculomotor behaviour reflects the fact that many of the processes 41 involved in reading are not yet automatized for children with RD, independently of NF1. This 42 suggests that the differences in saccade targeting strategy between children with and without 43 RD would be secondary consequences of their reduced reading experience. 44

45

46 KEYWORDS. Saccadic computation; Reading experience; Developmental dyslexia;
47 Neurofibromatosis type 1; Parafoveal processing.

# 48 **ABBREVIATIONS**.

- 49 DD Children with developmental dyslexia
- 50 ILP Initial landing position
- 51 LVF Left visual field
- 52 NF1 Neurofibromatosis type 1
- 53 NF1RD Children with neurofibromatosis type 1 with reading disorders
- 54 NF1noRD Children with neurofibromatosis type 1 without reading disorders
- 55 PVL Preferred viewing location
- 56 RAN Rapid Automatized Naming
- 57 RD Reading disorders
- 58RVFRight visual field
- 59 TD Typically developing children
- 60 VP Viewing position

#### 61 **1. INTRODUCTION**

As reported by several studies, many adults with dyslexia or poor reading skills experienced early reading difficulties (Eloranta et al., 2018; Smart et al., 2017). Indeed, learning to read can be a hard task for some beginners, requiring becoming more efficient in linguistic processing and to gain cognitive control of saccadic eye movements. Understanding the mechanisms underlying reading difficulties has therefore been the focus of ongoing research, with particular interest in the role of eye movements in the learning-to-read process.

Much research has focused on linguistic processing abilities as determinants of 68 individual differences in reading acquisition (e.g., Kim & Pallante, 2012; Landerl et al., 2019; 69 Leppänen et al., 2004). If phonological processing is a necessary (but not sufficient) condition 70 for the development of adequate word recognition skill, our ability to read depends also on 71 visuo-attentional processes and eye movement control (e.g., Bellocchi et al., 2017; Besner et 72 73 al., 2005; Facoetti et al., 2010; Plaza & Cohen, 2007; for reviews, see Hung, 2021; Leibnitz et al., 2017; Premeti et al., 2022). Reading does require children to focus selectively on words in 74 75 a left-to-right attentional scanning. The reader's eyes have to learn to land in the best position within the words to be read, extract the relevant information, and program the next saccade 76 (Ducrot et al., 2013; O'Regan & Lévy-Schoen, 1987; Rayner, 1986). 77

As reading proficiency increases, a child becomes familiar with the printed form of 78 words and changes in eye movement behaviour are observed: the span of effective vision 79 increases (extending asymmetrically in the direction of reading, even in beginning readers; 80 Häikiö et al., 2009; Rayner, 1986), the saccades size lengthens, the fixations duration and the 81 number of refixations decrease (Blythe, 2014; Blythe & Joseph, 2011; Rayner, 2009). 82 Moreover, the role of attention in reading evolves with expertise, i.e., expert readers need less 83 attention to process words than beginning readers (Bellocchi & Leclercq, 2021; LaBerge & 84 85 Brown, 1989; Leclercq & Siéroff, 2016).

The present study further investigates the perceptual and attentional factors that 86 determine the eyes' landing position in children and whether or not the pattern of eye 87 movements (and in particular the primary saccade landing sites) can provide useful 88 information to help characterize the reading skills in neurodevelopmental disorders. Word 89 length, that is provided by the visually salient extra spacing between words, in most languages 90 that use an alphabetic script, is a primary source of information in order to guide the eyes 91 during text reading and determine where the eyes first land in a word (e.g., Inhoff et al., 2003; 92 Rayner et al., 1998). Rayner (1979) first demonstrated that the landing position in a word 93 during continuous reading, referred to as the "preferred viewing location" (PVL), is usually 94 located halfway between the beginning and the middle of the word for languages read from 95 left-to-right (McConkie et al., 1988; Radach & Kempe, 1993; Rayner, 1979; Vitu et al., 96 1990). It is thought that the PVL reflects a strategy whereby readers aim to fixate the centre of 97 98 the word (e.g., McConkie et al., 1988; Reichle et al., 2003; but see Vitu, 2003) and fall short of that optimal viewing position (OVP) (i.e., saccades would undershoot their goal), owing to 99 100 the properties of the oculomotor system (Engbert & Krügel, 2010; Joseph et al., 2009; 101 McConkie et al., 1988; Nuthmann et al., 2005; O'Regan & Lévy-Schoen, 1987). Alternatively, a visual strategy could be developed to send the eyes to a location that 102 optimizes information uptake from the fixated word (Legge et al., 1997; O'Regan & Lévy-103 104 Schoen, 1987). The evidence strongly suggests that in addition to length information, salient orthographic information can be extracted from the parafovea and influence saccade 105 computation (e.g., orthographically unfamiliar initial trigrams, Hyönä, 1995; White & 106 107 Liversedge, 2006). Furthermore, developmental research suggests that beginning readers do learn to optimize saccade targeting strategies, with average initial landing positions (ILPs) 108 109 gradually shifted towards the word centre during the first years of formal reading instruction (Ducrot et al., 2013; Huestegge et al., 2009; Joseph et al., 2009; Vorstius et al., 2014). Based 110

on the dual-route cascaded model of Coltheart et al. (2001), the child would rapidly switch 111 from a grapho-phonological decoding mode (indirect route), with a letter-by-letter processing, 112 to an orthographic treatment mode (direct route) with an OVP for word processing. Similarly, 113 114 the attention deployed on words would be adapted to the preferred global processing mode (Ans et al., 1998) and would be related to the efficiency of the oculomotor parameters (Prado 115 et al., 2007). Thus, during the first year of learning, oculomotor control and especially the 116 PVL will be adjusted in parallel to the development of an optimal lexical and visual-117 attentional processing strategy for word reading. 118

119 Of particular interest here is the study by Ducrot and Pynte (2002) who used an oculomotor lateralized bisection task and demonstrated that the presence of between-character 120 space information influences saccade programming. According to this view, PVL "is due to 121 122 attentional processes that develop for any type of discrete stimulus, whether or not reading is actually required" (Ducrot & Pynte, 2002, p. 1142). These authors argued that "linguistic-123 124 like" stimuli (i.e., with inter-character spaces) encourage a saccade targeting strategy that takes the direction of visual exploration into account and aims to land left of the centre of 125 targets (for languages read from left to right), in preparation for subsequent left-to-right 126 127 attentional scanning. Note that this oculomotor strategy is modulated according to the visual field (VF) in which the stimulus is displayed, resulting in a left-right asymmetry in saccade 128 extent which is larger in the left visual field (LVF) than in the right visual field (RVF). In the 129 same vein, Bellocchi et al. (2019) found that the average landing positions of fifth grade 130 typical reading children were left of centre for stimuli with inter-character spaces (words and 131 strings of hash marks) in both the left and right VFs. Interestingly, beginning readers and 132 children with DD were only influenced by the linguistic nature of stimuli such that ILPs were 133 shifted more toward the beginning of the strings only when the stimulus was a word. 134

Naturally, a reasonable question arises: do children with different levels of reading ability differ in their saccade computation strategies when processing parafoveally presented stimuli? In the present study, we used the oculomotor lateralized bisection task as Ducrot and Pynte (2002) to further investigate saccade programming in typical reading children and children with reading difficulties, under conditions that modulate the direction of visual exploration and minimize the influence of linguistic factors.

A huge amount of data has reported differences between the eve movement patterns of 141 dyslexic readers and those of skilled readers. Poor readers and individuals diagnosed as 142 having dyslexia exhibit inefficient oculomotor behaviour while reading, with longer saccades, 143 shorter fixations, and more refixations than typical readers (Ashby et al., 2005; Franzen et al., 144 2021; Gagl et al., 2014; Gangl et al., 2018; Hawelka et al., 2010; Kirkby et al., 2022; Lefton 145 146 et al., 1979; Prado et al., 2007; for review, see Rayner, 1998). Also relevant for the present study are experiments reporting a missing parafoveal preview benefit in poor readers or 147 148 dyslexic children (Silva et al., 2016). The difficulty of pre-processing the word to the right of fixation may affect both saccade-planning and its execution towards the PVL. In that sense, 149 dyslexic readers tend to initially fixate words closer to their beginning than good readers 150 151 (Gagl et al., 2014; Gangl et al., 2018; Hawelka et al., 2010; Kirkby et al., 2022; McDonald et al., 2006; McKeben et al., 2004). However, there is still some unsettled disputes on the role of 152 153 these atypical eye movement patterns as a causal factor in dyslexia (see Blythe et al., 2018, asserting that instead they might primarily reflect an underlying difficulty in written word 154 recognition or decoding). Indeed, the attempts to replicate differences in eye-movement 155 patterns of dyslexic, poor and control readers when performing nonreading tasks have been 156 essentially unsuccessful, suggesting that children "with delayed reading skills (identified by 157 an assessment of age referenced reading ability rather than a diagnosis of dyslexia possess the 158

159 capacity to execute eye movements equivalent to those with good reading skills" (Hindmarsh160 et al., 2021, p. 1135).

161 The point of agreement is that in a reading task, children with DD exhibited atypical visual-processing skills. Among them, the deployment of visual attention seems to be 162 impaired in DD individuals. As mentioned above, several studies have shown that in left-to-163 right reading systems, the fixation point of skilled readers was often located toward the left of 164 the strings, when they processed multiple characters at once, under the influence of reading 165 habits and experience (e.g., Ducrot & Pynte, 2002; Sireteanu et al., 2005), suggesting a left 166 167 dominance of attention distribution in the process. DD children have a deficit in RVF attentional inhibition, and their attentional resources can be excessively interfered by RVF 168 information (Li et al., 2021). It results in the ignorance of LVF information (see Facoetti et 169 170 al., 2006; Hari & Renvall, 2001), which leads to no significant difference between LVF and RVF, showing an unbiased distribution pattern of attention resources (Li et al., 2021). In that 171 172 vein, Ducrot and colleagues found a symmetric OVP curve for dyslexic readers in foveal vision, contrary to control who demonstrated a typical OVP effect with a left-right 173 asymmetric J-shape curve (Bellocchi & Ducrot, 2021; Ducrot et al., 2003, see also Bellocchi 174 et al., 2019, for a less pronounced left-right asymmetry in parafoveal vision). In addition, 175 children with DD are more affected by visual crowding (even with non-linguistic stimuli, Joo 176 et al., 2018; Spinelli et al., 2002) and benefit more from increased text spacing (Perea et al., 177 178 2012; Zorzi et al., 2012; for reviews, see Bellocchi et al., 2013; Gori & Facoetti, 2015). Finally, the atypical eye-movement patterns observed in dyslexic children have been linked to 179 impaired visuo-attentional processing, and poor visual attentional span abilities, resulting in 180 fewer letters simultaneously processed and more rightwards fixations (Prado et al., 2007). As 181 suggested by Facoetti (2012), a possible neurobiological substrate of visuo-spatial attention 182 183 deficits in DD could be a weakened or abnormal magnocellular input to the dorsal visual

stream. Magnocellular–dorsal deficits could lead to reading difficulties through impaired
serial attentional orienting (Facoetti et al., 2010; Franceschini et al., 2012; Vidyasagar &
Pammer, 2010) or poor eye movement control (Stein, 2001; but see Goswami, 2015; Hutzler
et al., 2006, for a different point of view).

One of the main motivations of the study was to examine whether deviant eye 188 movements exhibited by dyslexic readers are a symptom of dyslexia or constitute a more 189 causal impairment. Few studies have explored eye movement behaviour in children with a 190 range of reading abilities that goes beyond dyslexia, using eye tracking during a task 191 192 preserving the perceptual and oculomotor demands of reading (but removing the higher order processing). Could the inefficient eye movement patterns reported in DD be transposed to a 193 reading deficit in another vulnerable population? To find answers to this question, we focused 194 195 on neurofibromatosis type 1 (NF1), that is a genetic disease in which a high prevalence of reading deficits has been shown (between 30 % to 80 %; Arnold et al., 2020; Chaix et al., 196 197 2017; Hyman et al., 2005), with a profound impact on their academic achievement (Krab et al., 2008). In this neurodevelopmental disorder, visual-processing deficits were also 198 demonstrated (e.g., Hyman et al., 2005; Ribeiro et al., 2012; Vernet et al., 2022) and a first 199 200 eye-tracking study showed atypicality in the saccadic system (Lasker et al., 2003). Some studies have suggested a delay in the maturation of low-level vision processes in children with 201 NF1 (Lasker et al., 2003; Ribeiro et al., 2012). To our knowledge, no study has investigated 202 203 the involvement of eye-movement control in the reading difficulties observed in NF1 children. 204

Summing up, reading is primary a visual task requiring good oculomotor and visuoattentional skills and in which the initial fixation position of a word plays a crucial early role. With expertise, a left-right asymmetry in VF emerges and the discreteness nature of the word becomes a factor of considerable importance when computing the incoming saccade. Some

findings suggest that in reading disorders, such as DD, there is atypical eye movement control 209 and specifically, the implementation of an inefficient saccade targeting strategy. To our 210 knowledge, very few studies have examined the factors that lead to this ineffective strategy in 211 212 children with severe reading difficulties, either in DD or in NF1. Thus, we assume that it is crucial to better understand the determinants of eyes' landing position in pathological reading 213 in each of these populations. Here, we examined whether children with reading disorders 214 215 (RD) differed from typical developing readers in saccade targeting strategies, and whether 216 lateralized presentation would differently impact saccade programming in children with and without RD. We hypothesized that if a detrimental effect of LVF presentation is observed, 217 218 this effect may occur especially for children who have not yet fully developed oculomotor mechanisms for efficient reading (i.e., DD and NF1 children; Bellocchi et al., 2019; Ducrot & 219 Grainger, 2007; Nazir, 2000). Moreover, according to Ducrot and Pynte (2002), we predicted 220 221 that if landing positions are an indicator of the development of reading skills, ILPs would differ according to stimulus type and reading experience, with only experienced readers being 222 223 able to take account of the presence of between-character space information to compute 224 saccades. To test this hypothesis, we manipulated the type (linguistic vs. non-linguistic) as well as the discreteness (stimuli with inter-character spaces vs. continuous lines) of the 225 226 stimuli.

We conducted a three-step analysis. With the first analysis, we investigated the extent to which attentional factors, stimulus features, and reading ability influence eye behaviour, and more specifically the PVL effect in parafoveal vision. At the same time, we explored possible visual and oculomotor deficits in children with DD. The second analysis aimed to quantify the use of different saccade targeting strategies and closer understand their relationship to reading abilities, by comparing NF1 children with and without RD. Finally, a third analysis including all children was carried out, in order to further investigate the relation between individual landing position and reading skills. We wanted to know if independently of the clinicaldiagnosis, reading ability only can reveal different saccade targeting strategies.

236

### 237 **2. METHOD**

## 238 **2.1. Participants**

Twenty typically developing children (TD, 10 females), 21 children with DD (8
females), and 20 children with NF1 (13 females) participated in the present study.

The inclusion criteria were the same as those described in the study by Vernet et al. 241 (2022). All children were between 8 and 12 years old and attended school from grades 2 242 through 6. They were all French native speakers, right-handed and presented normal or 243 corrected-to-normal vision. There were no significant differences between groups for 244 245 chronological age (TD mean age = 118.30 months, SD = 11.48; DD mean age = 120.90246 months, SD = 14.18; NF1: mean age = 121.57 months, SD = 18.68). All children underwent a complete medical and neuropsychological screening. Children were excluded from the study 247 if they had Attention Deficit Hyperactivity Disorder (i.e., more than 6 symptoms of 248 Hyperactivity and/or Inattention in a parent rating on the DSM-5 diagnosis criteria), 249 intellectual disability (i.e., WISC total IQ below 70 and/or standard score below 7 on the 250 Similarities and/or Picture Concepts subtests), neurological or psychiatric disorder (e.g., 251 epilepsy, brain tumor, or autism), or a known hearing impairment. 252

For the DD children, the reading level was controlled to ensure that children exhibited reading disorders (i.e., scores at least 1 SDs below the mean on the reading speed and accuracy indices of the Alouette test (Lefavrais, 2005) and at least 1.5 SDs below the mean on the ODEDYS-2 test (Jacquier-Roux et al., 2002)). The reading level was also assessed in the TD group to verify that they were all typical readers (i.e., scores at least superior to -0.5 SDs

on the Alouette-test (Lefavrais, 2005) and on the ODEDYS-2 test (Jacquier-Roux et al., 2002)). As expected, statistical analyses revealed that the DD group displayed significantly poorer performances than the TD group for the three reading indices with large effect sizes [reading age: t(39) = -10.022, p < .001, d = -3.131; reading accuracy: t(39) = -5.703, p < .001, d = -1.782; reading fluency: t(39) = -11.277, p < .001, d = -3.523].

For the NF1 children, participation was offered to all children who met the clinical 263 NF1. Neurofibromatosis diagnostic criteria for according to the Conference 264 Statement (National Institutes of Health, 1988). The reading level was systematically assessed 265 but was not used as an inclusion criterion for this specific subject. 266

The French ethics committee Review Board approved the study (2014-A01239-38 and 2014-A01960-47), which was conducted according to the guidelines of the Declaration of Helsinki (World Health Organisation, 2008). The written informed consent was obtained from every participant and their legal tutors prior to inclusion in the study. Note that this study included participants from a larger study examining procedural learning and memory<sup>1</sup>.

272

#### **273 2.2. Material**

Oculomotor lateralized bisection task. The material consists of 3 lists of stimuli: solid
lines, strings of hashes and words. Each list is composed of 20 items of 5 or 6 characters long.
We selected a set of words from the first-grade lemma lexicon of the Manulex database (Lété
et al., 2004) that provides reliable information on grade-level word frequency. Half of the
targets were high frequency words, with average printed frequency of 419 occurrences per
million and the other half were low frequency words (printed frequency of occurrence of 16
occurrences per million). Stimuli were displayed in either the LVF or RVF relative to the

<sup>&</sup>lt;sup>1</sup> DYSTAC-MAP cohort (ANR-13-APPR-0010)

initial fixation point. The distance between the item and the fixation cross was set at 1.25°
from the end of the stimulus when displayed in the LVF or from the beginning of the stimulus
when displayed in the RVF. To sum up, two factors were manipulated: (1) the type of
stimulus with 3 conditions (i.e., words, strings of hashes and solid lines) and (2) the visual
field with 2 conditions s (i.e., LVF and RVF).

Visual-processing skills assessment. The Developmental Eye Movement (DEM) test 286 (Garzia et al., 1990) is commonly used to assess visual-processing skills. It consists of reading 287 aloud vertically and horizontally organized digits printed on three different sheets of paper 288 (Test A, B and C). Children were instructed to read aloud the digits as fast and as accurately 289 290 as possible. This test provides three main indices: (1) the vertical reading time (VT, in 291 seconds) that represents the sum of the time spent on naming the vertically organized digits of 292 Test A and B; (2) the adjusted horizontal time (HT, in seconds) that represents the adjusted time required for reading the horizontally organized digits presented in the Test C; and (3) the 293 total number of errors, which gives the accuracy on the execution of Test C. 294

*Reading skills assessment.* One of the most widely used French reading tests for assessing reading efficiency in children is the *Alouette test* (Lefavrais, 1967, 2005). In this test, the child is asked to read a 265-word text passage aloud as quickly and accurately as possible, within 3 minutes. The text consists of real words in syntactically correct but semantically poor sentences. The Alouette test provides three main indices: reading age, reading fluency, and reading accuracy.

301

#### 2.3. Apparatus and procedure

All children were recruited at both the Hospital of Aix-en-Provence and the Timone University Hospital (Marseille). They were seen individually during two sessions of approximatively 1 hour and a half each. The first one was conducted directly at the hospital and was intended to verify the inclusion criteria. This session included also neuropsychological assessment with the evaluation of reading level. The second session was carried out at the laboratory to perform the oculomotor bisection task. All children were tested individually in a quiet, separate room. The lighting in the room was adjusted to a comfortable level for each participant and each task.

Some of the following detail is taken from Ducrot et al. (2013). For the oculomotor lateralized bisection task, the eye movements were recorded by a mobile infrared, headmounted eye tracker (Eyelink 2, SR Research Ltd., Canada). The recording was based on infrared-light reflection from the pupil and cornea of the right eye at a sampling rate of 250 Hz and a spatial resolution of less than 0.04°. A chin-and-forehead rest was used to minimize head movements.

316 Before the experiment, a calibration step of the eye tracker was performed according to a 9-point grid extended to the whole computer screen. The eye tracker was connected to a 317 Dell D-type docking station and a Dell Latitude D600 laptop computer for data recording and 318 stimulus delivery. The stimuli were displayed in white lowercase letters against a black 319 background in 22-point Courier New font, using a 14-inch colour monitor, at a resolution of 320 1400×1050 pixels. Participants were seated in front of the screen at 60 cm. At this distance, 321 one letter subtended a visual angle of 0.38° and 1° is equal to 0.95 cm. The space between 322 letters was 1 mm that corresponds to 2.8 pt or  $0.09^{\circ}$ . 323

Each trial consisted of the following sequence of events (Figure 1). First, children were required to maintain fixation on the cross in the centre of the screen. Note that the experimenter repeatedly reminded the child to fixate the cross and not to move their eyes away from this point. Five-hundred ms after, the central fixation point was replaced by a parafoveal target displayed in the RVF or the LVF of the fixation point. Therefore, the children were instructed to "move their eyes, as quickly and accurately as possible, to the

position they considered to be the middle of the item" and then validate this position by
pressing a button. After the response, the screen was cleared for the next trial that started 500
ms later. Participants performed 12 practice trials beforehand.
[Insert Figure 1 About here]
2.4. Data analysis
The Emaa software package (Ducrot et al., 2006) was used to analyze the eye-tracking
data. Trials were excluded from the analysis when an eye-tracker sampling error or a blink
occurred during stimulus presentation. We also excluded trials when the saccade latency was
shorter than 80 ms or longer than 800 ms (4.3%), the average eye position before saccade
onset deviated from the fixation cross by more than $\pm$ 0.5 characters (2.9%), or the participant
took more than 5000 ms before pressing the button (6.8%).
We used the same eye-movement measures as Ducrot et al. (2013): (1) the saccade
latency which is the time between the onset of the target and the beginning of the saccade, (2)
the saccade size which is the size of the initial saccade, and finally (3) the refixation
probability which is the probability of making an additional fixation within the word before
leaving it. To study PVL effects, we also computed (4) the initial landing position in the
stimulus. This last parameter is used to determine the landing frequency at each position of
the item. Saccade latency was reported in milliseconds, whereas saccade size and initial
landing position were expressed in number of characters. For all the variables measured,

351 **2.5. Statistical analysis** 

350

A three-step analysis was conducted. A first analysis was conducted to examine saccade targeting strategies of children with and without DD in parafoveal vision. ANOVA were

means were calculated for each participant in each condition.

performed with a 2 Visual Field  $\times$  3 Type of stimulus  $\times$  2 Groups of children model. All factors except the group of children were within-participant variables. Further analyses were carried out in each group independently with statistical a priori comparisons.

In a second analysis, the same ANOVA were performed within the NF1 group, to better 357 understand the factors that determine the saccade targeting strategy implemented by children 358 with RD. Based on their performance on the Alouette test, NF1 children who presented 359 reading disorders (i.e., NF1RD; at least 12 months delay in reading age on the Alouette test 360 compared to chronological age) were compared to NF1 children without reading disorders 361 (i.e., NF1noRD). Note that 11 children were included in the NF1RD group and 9 in the 362 363 NF1noRD group. The two groups differed significantly in terms of reading age [t(18) = -364 4.435, p < .001], reading accuracy [t(18) = -4.144, p < .001] and reading speed [t(18) = -4.144, p < .001]5.218, p < .001<sup>2</sup>. 365

Finally, a final analysis including all children was carried out to further investigate the relation between individual ILPs and reading skills. Each child was assigned an ILP index (their initial-landing position for words in the RVF) and a reading level index (their reading age). The correlations between these two measures were tested. Then, cluster analyses were conducted. The ILPs were analysed by interactive partitioning (K-means), minimizing the within-cluster variability and maximizing the between-cluster variability, with a three-group solution.

373

## **374 3. RESULTS**

#### 375 **3.1. Comparisons between children with and without DD**

<sup>&</sup>lt;sup>2</sup> There were no significant differences between the two groups for the chronological age [t(18) = -5.316, p > .05] and the handedness ratio ( $\chi 2 = 0.669$ , p > .05).

[Insert Table 1 about here]

376

377

## **3.1.1.** Saccade Size

The median amplitude of left-to-right saccades in TD children was 3.46 characters, 378 while that of dyslexics was 3.49. For right-to-left saccades, however, we observed a shift of 379 the saccade's distribution toward shorter amplitudes for DD children (as compared to TD 380 children, (3.7 and 4.1 for DD and TD children, respectively). The analysis of incoming 381 saccade size yielded a significant interaction between stimulus type and side of presentation 382  $[F(2,76) = 12.505, p < .001, \eta_p^2 = 0.247]$  and three-way interaction between the group (TD vs. 383 DYS children), the type of stimulus (words, strings of hashes and solid lines), and the 384 presentation side (LVF vs. RVF) [F(2, 76) = 3.423, p = .038,  $\eta_p^2 = 0.080$ ]. Separate analyses 385 for TD and DYS children indicated that the interaction between the stimulus type and the 386 presentation side was significant in the two groups [F(2,38) = 7.128, p = .002,  $\eta_p^2 = 0.273$  and 387 F(2,38) = 8.509, p <.001,  $\eta_p^2 = 0.309$ , for the TD and DD children respectively]. 388

As can be seen in Table 1, for TD children, initial-saccade extent was equivalent in 389 size for all stimuli composed of discrete elements (words and strings of hashes), but was 390 different for continuous solid lines. Pairwise comparisons revealed that the difference 391 between words and strings of hashes was nonsignificant [F(1,38) = 1.089, ns] and that the 392 stimulus-type by presentation-side interaction result from the strong opposition between 393 discrete stimuli (words and strings of hashes) and continuous stimuli (solid lines) 394 [F(1,38) = 8.413, p < .01], thus confirming the influence of stimulus discreteness vs. 395 continuousness in determining ILPs (see Ducrot et al., 2002, for similar conclusions). 396

For DD children, pairwise comparisons revealed that the stimulus-type by presentation-side interaction was entirely explained by the strong opposition between linguistic stimuli (words) and non-linguistic stimuli (strings hashes and solid lines) [F(1,38) = 10.208, p < .01], and no difference between strings of hashes and solid lines [F <1]. As Table 1 shows, DD children do tend to initially fixate *words* at a point somewhere
between the beginning and the centre of the target when the saccade was rightward and closer
to the word middle for left presentations. For non-linguistic stimuli, however, the mean ILPS
were approximately symmetrical in the left and right VFs.

405

# 3.1.2. Latency

As first raised by Radach & McConkie (1998), this variable is "of particular interest, 406 407 because it provides a link to temporal aspects of eye movement control that may well significantly modulate spatial saccade parameters" (p. 83). The ANOVA yielded a main effect 408 of type of stimulus [F(2,78) = 92.602, p < .001,  $\eta_p^2 = .704$ ], regardless of the presentation side 409 (left or right). Pairwise comparisons revealed that the interaction was due to the difference 410 between discrete and continuous stimuli [183 ms for discrete stimuli vs. 220 ms for 411 continuous stimuli, F(1,78) = 578.436, p < .001]. The difference between words and strings of 412 413 hashes was nonsignificant [F(1,78) = 1.35, ns]. Note that strings of hashes are letter-like and similar in visual complexity to letters, so processing these symbols might activate a decoding 414 process similar to processing text. Strings of hashes also differ substantially from solid lines 415 in shape, size and brightness. A similar tendency in expert readers has already been observed 416 by Ducrot and Pynte (2002), with shorter latencies in the discrete condition than in continuous 417 condition (173 vs. 193 ms, p = .06, Experiment 3). There was also a main effect of group 418  $[F(1,39) = 4.839, p = .034, \eta_p^2 = .110]$  and a stimulus type × group interaction [F(2,78) =419 6.274, p = .003,  $\eta_p^2$  = .139], showing that latencies were longer for DD children (207 ms, SD 420 = 31) than for TD children (196 ms, SD = 20.5) (see Bucci et al., 2008; Fischer & Weber, 421 422 1990; Pirozzolo & Rayner, 1979, for similar results), in particular for discrete stimuli (193 vs. 173 ms for DD and TD respectively, Table 1). 423

In order to examine whether symmetrical patterns were associated with short latencies 424 425 (saccades aim to target the centre of the word but undershoot their goal) and asymmetrical patterns with long latencies (the eyes land at the location that enables maximum information 426 427 intake), we calculated (1) an asymmetry index (the difference between the mean amplitude of a child initial saccades in the left and right VFs) and (2) a rapidity index (his or her mean 428 saccade latency, for leftward and rightward saccades) for each participant and each type of 429 target. The correlation between these two indices was not significant with a lack of correlation 430 for words [r(39) = .05, p > .10], strings of hashes [r(39) = -.03, p > .10] and solid lines [r(39)431 = .07, p > .10]. The same tendency was found when separate analyses were made for TD and 432 DD children [r(18) = -.19, p > .10, and r(19) = .40, p > .05, respectively]. 433

434

#### 3.1.3. Refixation probability

We found a significant effect of group on refixation probability, [F(1,38) = 6.626, p =435 .014,  $\eta_p^2 = .148$ ], with surprisingly DD children making less refixations (57.64%) than TD 436 children (68.09%). There was also an interaction between presentation side and group 437  $[F(1,38) = 9.442, p = .004, \eta_p^2 = .199]$ , with an "adverse" effect of fixating the RVF among 438 the DD children (as compared to the TD group). Interestingly, the analyses revealed three-439 way interaction between the group, the presentation side and the type of stimulus [F(2,76) =440 3.632, p = .031,  $\eta_p^2$  = .087]. Separate analyses were done for TD and DD children and 441 442 revealed that the interaction between presentation side and stimulus type was only significant in TD children [F(2,38) = 3.333, p = .046,  $\eta_p^2$  = .149 and F < 1, for TD and DD children 443 respectively]. For the DD group analysis, we only found a main effect of the side of 444 presentation [F(1,19) = 9.005, p <.007,  $\eta_p^2$  = .322], with children making more refixations on 445 the RVF (64%) than on the LVF (51%). For the TD group, pairwise comparisons revealed a 446 significant difference between RVF and LVF for words only [F(1,38) = 8.725, p < .01 for 447 words, and Fs < 1 for strings of hashes and solid lines], thus suggesting an advantage for the 448

RVF limited to word recognition. Note that, even if no reading was required, the bisection task elicited comparable ILPs patterns and a RVF advantage for words in TD children, thus supports the idea that some of the factors responsible for the VF and ILPs asymmetries in natural reading are also at work in this task.

453

To sum up, saccades toward stimuli located in the RVF were not significantly affected 454 by stimulus type or reading disorder, all children being able to land at the PVL. In contrast, 455 leftward saccades were influenced by stimulus type and reading ability. In particular, the left-456 right asymmetry of the incoming saccade was influenced by the inter-character spacing in TD 457 458 children whereas the asymmetry was restricted to words in dyslexics. Moreover, the TD 459 readers fixated closer to the PVL of the discrete stimuli in the LVF compared to the dyslexic children, with leftward saccades about one character longer. These results are consistent with 460 those of Ducrot & Pynte (2002) who showed that expert readers land to the left of the centre 461 in all discrete stimuli (strings of hashes and words) for both presentation sides (see Bellocchi, 462 Mancini, et al., 2013, for similar results in 5<sup>th</sup> graders). These results suggest that, as a result 463 of reading experience, the children tested in this experiment had already developed 464 automatized routines for programming right-going saccades. The fact that saccades from 465 right-to-left are in the opposite direction of reading would lead to less automatized routines 466 for saccade computation, making them more vulnerable to the influence of reading level and 467 VF. It must be emphasized that many children with dyslexia do not read as much as 468 individuals with typical reading skills, and so, this non-optimal saccade targeting strategy in 469 dyslexia may also results from poor reading experience (see Huettig et al., 2018, for a 470 discussion). 471

472

#### 473 **3.2.** Comparisons of NF1 children with and without RD

[Insert Table 2 about here]

#### 475 **3.2.1.** Saccade size

In clinical populations, as suggested by Ablinger et al. (2014), a "standard way to report on the spatial aspect of eye movement control is to plot the individual distributions of the landing positions of incoming saccades over all letter positions within a target word" (p. 652; see also Radach & Kempe, 1993; Radach & McConkie, 1998).

Zooming in on the frequency distributions of ILPs (Figure 2) reveals different distribution profiles between NF1 children who presented reading disorders (NF1RD) and those without reading disorders (NF1noRD). We found a significant correlation between the reading age of the NF1 children and the frequency of their initial landing position at P2 [r(18) = .47, p < .05]. Whereas a PVL was distinctly observed in NF1noRD children between the beginning and the centre of the word (i.e., P2), no PVL emerged in NF1RD children [t(39) = -3.241, p < .01].

487

#### [Insert Figure 2 about here]

More precisely, for both groups, there was an interaction between stimulus type and 488 presentation side [F(2,20) = 7.236, p = .004,  $\eta_p^2 = 0.420$  and F(2,16) = 3.593, p = .05,  $\eta_p^2 =$ 489 0.313, for NF1RD and NF1noRD respectively]. For the NF1RD group, pairwise comparisons 490 were used to reveal a difference in initial-saccade size between linguistic stimuli (words) and 491 non-linguistic stimuli (strings hashes and solid lines) [F(1,20) = 14.323, p < .005]. The 492 difference between strings of hashes and solid lines was non-significant [F < 1]. For the 493 NF1noRD group, the amplitude of the initial saccade was similar for words and strings of 494 hashes, but different for solid lines, thus confirming the importance of the discreteness of the 495 stimuli for unimpaired readers [[F(1,16) = 4.49, p = .05], and [F < 1], for (words and strings 496 of hashes) vs. solid lines, and strings of hashes vs. words, respectively]. 497

Of particular interest that NF1RD readers mirrored dyslexic 498 is the eye movement behaviour, with more variability in saccadic programming, and no sensibility to 499 the discreteness of the stimuli (see Figure 3). The data showed no main effect of group [F < 1]500 nor double interaction between presentation side, stimulus-type and group [F < 1], but a 501 presentation side × stimulus-type interaction [F(2,58) = 13.732, p < .001,  $\eta_p^2$  = .321], mainly 502 explained by the sensitivity to the linguistic nature of the stimuli [F(2,58) = 58.81, p < .001,503 for the opposition between linguistic vs. non-linguistic stimuli, and F < 1, for the difference 504 between strings of hashes vs. solid lines]. 505

506

## [Insert Figure 3 about here]

507 The children with NF1noRD showed fully developed oculomotor mechanisms for efficient reading similar to those of TD children [F < 1], with a clear PVL located to the left 508 of the word's centre in both visual fields for the 2 groups  $[F < 1, for the group \times presentation]$ 509 510 side  $\times$  stimulus type interaction], and fine-tuned saccade targeting guided by the betweencharacter space information. The stimulus-type by presentation-side interaction [F(2, 54) =511 8.088, p < .001,  $\eta_p^2 = 0.231$ ] was explained by the strong difference between ILPs 512 distributions in discrete stimuli (words and strings of hashes) vs. continuous stimuli (solid 513 lines) [F(1,54) = 10.08, p < .01], the difference between words and strings of hashes being 514 non-significant [F<1]. 515

516 **3.2.2.** Latency

As detailed in Table 2, for both groups of NF1, there was a main effect of stimulus type  $[F(2,20) = 19.360, p < .001, \eta_p^2 = .638 \text{ and } F(2, 16) = 50.890, p < .001, \eta_p^2 = .864, \text{ for}$ NF1RD and NF1noRD respectively]. This difference was mainly explained by shorter 520 latencies for discrete stimuli than for continuous stimuli [F(1,20) = 38.620, p < .001] and 521 F(1,16) = 101.705, p < .001, for NF1RD and NF1noRD respectively].

Consistent with what had previously been observed for saccade size, the saccade 522 latencies of NF1RD children were similar to those of DD children. There was a main effect of 523 stimulus type [F(2,62) = 40.205, p < .001,  $\eta_p^2$  = .565] regardless of the group [F < 1]. 524 Similarly, the saccade latencies in NF1noRD were similar to those found in TD children, with 525 shorter latencies when targeting discrete stimuli and longer saccade latencies for continuous 526 stimuli [F(2,54) = 119.607, p < .001,  $\eta_p^2$  = .816]. There was no main effect of group [F < 1] 527 nor interaction between stimulus-type and group [F < 1]. These results seem to suggest that it 528 529 is less time-consuming to target the PVL in a word (or in a linguistic-like stimulus) than to 530 target the middle of a solid line.

531

### **3.2.3.** Refixation probability

532 We found a significant interaction between stimulus type and presentation side, for both groups of NF1 [F(2,20) = 3.569, p = .047,  $\eta_p^2 = 0.263$  and F(2,16) = 3.639, p = .050,  $\eta_p^2 =$ 533 .313 for NF1RD and NF1noRD respectively]. In line with what had previously been observed 534 for TD children, pairwise comparisons in the NF1noRD group revealed a significant effect of 535 the VF for words on the refixation probability [F(1,16) = 10.156, p < .005 for words, and no536 VF difference for strings of hashes and solid lines, Fs < 1]. For the NF1RD group, pairwise 537 comparisons yielded an opposite effect, with children performing more refixation on words in 538 the RVF [F(1,20) = 9.028, p < .01 for words and Fs < 1 for strings of hashes and solid lines]. 539

540 When pooling the data of NF1noRD and TD children, this interaction remained [F(2,54) 541 = 4.666, p = .014,  $\eta_p^2 = 0.153$ ], and could be entirely explained by a larger VF asymmetry for 542 words [F(1,54) = 20.643, p < .001 for words, and Fs < 1 for strings of hashes and solid lines]. 543 Note that this larger VF asymmetry for words also resulted in a larger "linguistic" effect in

RVF than in LVF. Comparing NF1RD performance with that of DD children, the ANOVA 544 showed no main effect of participant group [F < 1] nor double interaction between 545 presentation side, stimulus-type and group [F < 1]. We found, however, a main effect of 546 presentation side [F(1,29) = 7.448, p = .011,  $\eta_p^2$  = .204] and a presentation side × stimulus-547 type interaction [F(2,58) = 4.055, p = .022,  $\eta_p^2 = 0.123$ ], mainly explained by the cost for word 548 recognition in the RVF [F(1,58) = 37.383, p < .001for words, and Fs < 1 for the strings of 549 hashes and solid lines]. Here again, the difference between words and non-linguistic stimuli 550 was larger in the RVF than in the LVF (but in reverse). The opposite effects in the VF 551 asymmetries found for children with and without RD suggests that the parafoveal words were 552 processed (even if no linguistic processing was required in the bisection task), resulting in a 553 classical RVF advantage for children without RD and a cost for the children with RD. 554

To sum up, the eye movements of NF1RD children during the ocular bisection task are 555 different from those of NF1noRD children, but similar to those of DD children. In particular, 556 our results showed a less pronounced left-right asymmetry in the saccade amplitude for RD 557 children than for TD and NF1noRD children. This pattern of result reflects, in the context of 558 reading deficits, less stable ILPs and a not yet automatized saccade targeting strategy. This is 559 consistent with the data obtained on the VP effect in foveal vision, showing reduced left-right 560 asymmetry in the RD children's VP-curve (Bellocchi & Ducrot, 2021; Ducrot et al., 2003). 561 The fact that NF1RD and children with dyslexia showed similar saccade targeting strategy, is 562 consistent with the proposition of Huettig et al. (2018) that "this at least partly reflects their 563 common reduced reading experience rather than a causal impairment due to a reading 564 disorder" (p. 339). 565

#### 567

#### **3.3.** Cross-population results

Each child was assigned an ILP index (their initial-landing position for words in the 568 RVF) and a reading level index (their reading age<sup>3</sup>). The correlations between these two 569 measures are displayed in Figure 4. We found a significant correlation between reading age 570 and the initial landing position at P2 [r(59) = .493, p < .001], with a higher reading level being 571 associated with a higher probability to land first at the PVL (i.e., P2). A significant negative 572 correlation was also found between reading age and the initial landing position at P1 [r(59) = -573 .266, p < .05], suggesting that poor readers are more likely to land at the beginning of the 574 word. Note that no significant correlation was shown for landing positions at P3 [r(59) = -575 .136, p > .05]. 576

577

### [Insert Figure 4. about here]

Cluster analyses with a three-group solution yielded a relatively distinct separation of 578 579 participants into clusters with very little overlap. Figure 5 shows the three clusters of participants for the distribution of the ILP frequency at each position and the relationship 580 between reading level and ILP. This allowed for an evaluation of the ways in which our 581 readers are similar or different in reading skills and eye movement behaviour. This analysis 582 strategy is supported by the idea that, as word identification becomes more efficient, children 583 shift from a letter-by-letter sequential decoding strategy (characterised by ILPs close to the 584 word beginning to be able to scan it from left to right) to more parallel processing (with ILPs 585 shifted toward the word centre (see also, Ablinger et al., 2013; Schattka et al., 2010). Cluster 586 2 comprised almost half of the participants (46%, mainly children without RD) and 587 corresponds to an expert reader strategy with an ILP most of the time at the PVL (i.e., P2). 588

<sup>&</sup>lt;sup>3</sup> We chose to use the reading age measure since it is frequently used in eye-tracking and dyslexia studies (e.g., Bellocchi et al., 2019; Liu et al., 2022; Prado et al., 2007). Note that this measure was strongly correlated in the present study with the other two reading indices of the Alouette-test [reading accuracy, r(59) = .684, p < .001; reading fluency r(59) = .911, p < .001].

The two remaining clusters differed from the cluster of "typical" readers. Cluster 1 (12%, 589 mainly DD) showed a more careful eye-movement strategy (i.e., fixated closer to the 590 beginning of the stimuli, P1), as usually found for beginning readers. It comprised readers 591 with the lowest reading level scores, and this group might stick to serial reading, suggesting 592 an immaturity of the oculomotor control associated with a sequential processing strategy. 593 Members of Cluster 3 (42% of the participants, mainly RD children) presented an 594 intermediate reading level and a flattened curve with an ILP covering a wider area around the 595 centre of the word. This pattern could be the reflect of an oculomotor instability. In that sense, 596 this subgroup of readers also presented severe visual-processing deficits at the DEM-test (-597 1.27 SD on the VT index and -1.97 SD on the HTaj index, with 41% of the children below -598 1.5 SD on the VT index, and 59% below -1.5 SD on the HTaj index)<sup>4</sup>. 599

We want to draw attention to the fact that the 3 clusters differed<sup>5</sup> significantly in terms of reading level [F(2,58) = 3.877, p < .05,  $\eta_p^2$  = .118]. The pairwise comparisons showed a significant higher reading age in the cluster 2 (Mean = 115.61) compared with the cluster 1 (Mean = 94.43; p <.05) and 3 (Mean = 100.81; p < .05), respectively. However, no significant difference emerged between clusters 1 and 3 in terms of reading age [p >.05] and between the 3 clusters in terms of chronological age [F(2,58) = 0.647, p > .05,  $\eta_p^2$  = .022].

To sum up, taken together, the correlation and cluster analyses revealed a strong link between reading expertise and ILP. Cluster analyses also pointed to individual differences, with three subgroups of readers using different saccade targeting strategies, irrespective of whether they have been classified as neurotypical, dyslexic or NF1. In our view, the present

<sup>&</sup>lt;sup>4</sup> Note that the cluster 1 group had significantly poor performance at the DEM-test, as well, probably linked to their slowness in their responses and their low reading age (-2.19 SD on the VT index and -1.65 SD on the HTaj index). No such visual-processing deficits was observed in cluster 2 (-1 $\leq$ SD $\leq$ 0, for the VT and HTaj index).

<sup>&</sup>lt;sup>5</sup> For these statistical analyses, Kruskal–Wallis tests followed by Dunn–Bonferroni method for pairwise comparisons were carried out.

610 results strongly suggest that the reading ability and the mode of processing used (shift from a 611 sequential reading to a more parallel orthographic processing) are sufficient to account for the 612 different patterns of eye movement behaviour.

613

## [Insert Figure 5 about here]

614

#### 615 **4. DISCUSSION**

The aim of the present study was to investigate the extent to which attentional factors, stimulus features, and reading ability influence the position where the eyes first land in a target (word or non-linguistic stimulus) displayed in parafoveal vision. A secondary objective was to better understand the relationship between saccade eye movements and reading difficulties and to examine the possible modulation of clinical diagnosis (DD and NF1) and reading skills (good and poor readers) on saccade targeting strategies as expressed by ILPs.

622

623

#### 4.1. What determines where children first fixate?

As previously demonstrated by Ducrot et al. (2013), our data showed that all children 624 tend to initially fixate words at a point somewhere between the beginning and the centre of 625 the word while executing rightward saccades, regardless of their reading level (with or 626 without RD) or clinical diagnosis (dyslexics or NF1 group). In other words, even with 627 a minimum experience with reading, children show an adult-like PVL pattern in the RVF (for 628 629 similar results, see Bellocchi et al., 2019; Ducrot et al., 2013; Joseph et al., 2009). Note that rightward saccades represent the most frequent saccades in left-to-right writing systems, 630 631 leading to early development of automatized procedures for saccade computation.

One aspect that warrants particular attention is the left-right asymmetry in the saccade 632 size launched toward parafoveally located stimuli. As stated before, "linguistic-like" stimuli 633 encourage a saccade targeting strategy that aims to land at the PVL in both VFs, with greater 634 635 difficulty in reaching this goal for the LVF due to the greater distance and unusual reading direction (Ducrot & Pynte, 2002; Vitu et al., 1995). Our results showed a more pronounced 636 left-right asymmetry for TD and NF1noRD children than for DD and NF1RD readers (see 637 Bellocchi et al., 2019, for similar results with DD). One possible explanation to account for 638 this increase in asymmetry for TD readers was suggested by Huang et al. (2019), as reading 639 experience seems to "increase the ratio between recognition in the centre and recognition 640 641 away from the centre of fixation, as reflected in the changes of the shape of the so-called Form Resolving Field (which maps the distribution of recognition accuracy onto eccentricity) 642 643 from symmetrical and reversed-V-shaped to slightly narrower on the side of the reading 644 direction" (see also, Lorusso et al., 2004; Zegarra-Moran & Geiger, 1993). Moreover, children without RD used a saccade targeting strategy that aims to land left of centre (PVL) 645 646 for all discrete stimuli (i.e., with inter-character spaces), even in the LVF, as already observed 647 in adults (Ducrot & Pynte, 2002). In contrast, for children with RD, the asymmetry was less pronounced and restricted to words only. NF1RD and DD children were influenced by the 648 linguistic nature of stimuli, such that they do tend to initially fixate words closer to their 649 beginning (as compared to the strings of hashes). 650

The results, therefore, suggest that the pre-processing mechanism responsible for the direction of attentional scanning (left-to-right) proposed in Ducrot and Pynte (2002) is well automatized in children without RD, which "anticipated doing left-to-right attentional scanning of discrete stimuli and took the direction of this upcoming attentional scanning into account when computing the incoming saccade" (Ducrot & Pynte, 2002, p. 1142). Note that the fact that children used different saccade targeting strategies depending on the presence of inter-character spaces or the linguistic nature of the stimulus suggests that the left–right
asymmetry in the PVL effect may not result *only* from oculomotor constraints associated with
saccade execution.

660

661

## 4.2. How reading ability affects eye movement behaviour?

A major aim of the present study was to determine whether reading level and changes in 662 written language experience over the course of DD, NF1 and TD children's developmental 663 trajectory are associated with changes in reading strategies, as indexed by saccade targeting 664 strategies. Firstly, the individual distributions of first fixation positions were modulated by 665 666 reading skills. Cluster analyses confirmed that the landing position of eve movement saccades on words is a sensitive index of reading efficiency (Ducrot et al., 2013; Joseph et al., 2009; 667 Juhasz et al., 2008), with three subgroups of readers using different saccade targeting 668 669 strategies, irrespective of whether they have been classified as neurotypical, dyslexic or NF1. Good readers (cluster 2) made their first fixation most of the time at the PVL whereas, in 670 children with poor reading skills, ILPs either occurred at the beginning of the word (cluster 1) 671 or covered a wider area around the centre of the word with high variability (cluster 3). A 672 convincing explanation for these saccade targeting differences is based on the dual-route 673 model of reading (Coltheart et al., 2001). As experience with written language grows, children 674 shift from a letter-by-letter sublexical decoding strategy to more parallel lexical processing 675 676 (Frith, 1985). In that sense, Tydgat and Grainger (2009) suggested that as children develop 677 their capacity to perform parallel letter processing, their ILPs shift from being close to the beginning of words toward a position closer to the centre of words (see also Huestegge et al., 678 2009). Meanwhile, visuo-attentional abilities would gradually develop with reading 679 680 experience (Bosse & Valdois, 2009; Hawelka & Wimmer, 2005). In that vein, children show a smaller span of effective vision than adults (Häikiö et al., 2009; Rayner, 1986), the size of 681

children's perceptual span increasing with reading proficiency (Sperlich et al., 2016). It may
be that in very fluent reading, the visuo-attentional capacity approaches visual acuity
constraints for letter encoding (Hautala et al., 2022).

We also explored the impact of the reading level on the VF effects. Children with RD 685 made shorter saccades in words than did TD readers, with ILPs located left of centre for both 686 left and right VF only for the TD children. One possibility is that children with RD experience 687 difficulties in oculomotor control. Shifted ILPs may arise simply because children with RD, 688 as beginning readers, exhibit shorter average saccade amplitude (cluster 1) and/or increased 689 variability (cluster 3) in these saccade amplitudes (see also Ducrot et al., 2013, for a similar 690 691 finding with a lateralized lexical decision task). In line with this proposal is the fact that an 692 eye-tracking training can improve saccadic eye movements in beginning children, with accurately aiming for the OVP of each word (Lehtimäki & Reilly, 2005). It is also possible 693 694 that oculomotor behaviour and saccade targeting strategy depend on the efficiency of linguistic processing (Rayner, 1998). Proficient readers with high-quality lexical 695 696 representations and efficient lexical processing will aim to optimize information uptake from words by initially fixating the OVP. In contrast, beginning or poor readers, relying on a letter-697 by-letter phonological decoding strategy, will prefer to initially fixate words closer to their 698 699 beginning. In that sense, shorter incoming saccade amplitudes and increased refixation rates 700 have been reported in poor, dyslexic, beginning, and slow readers (e.g., Ducrot et al., 2013; Hawelka et al., 2010; Huestegge et al., 2009; McKeben et al., 2004; Rayner et al., 2010, but 701 702 see also Valdois et al., 2004, for the hypothesis that dyslexic readers suffer from a narrowed 703 visual attentional window). However, if the second account was the sole explanation, then why would DD and NF1RD children aim to land to the right of the middle of a word 704 705 displayed in the LVF? Because acuity drops rapidly with increasing eccentricity, word 706 recognition is assumed to be better when the initial fixation is left rather than right of the

centre of the word. Rather, these results suggest that oculomotor behaviour change as
linguistic processing becomes more efficient, thus confirming that poor and proficient readers
face different oculomotor, perceptual, and cognitive constraints when reading.

As a whole, the current results support the hypothesis that saccade targeting strategies are an acquired reading skill that develops with reading experience. Conversely, the children's reading strategies (i.e., whether they use lexical or sublexical route) can be understood from the characteristics of saccadic eye movement.

714

### 715 **4.3.** A Causal Role for Inefficient Eye Movements in Reading Disability?

716 Everyone agrees that DD readers' eye movements are different from typical readers. These eye movement differences have been found when reading isolated words/pseudowords, 717 sentences or texts, and were reported in different languages with different orthographic depths 718 719 (e.g., Hutzler & Wimmer, 2004; Hutzler et al., 2006). It has been suggested that they might 720 result from poor oculomotor control associated with dyslexia (Pavlidis, 1981). However, 721 evidence in support of a low-level oculomotor account of the reading behaviour of dyslexics 722 are highly controversial (e.g., Eden et al., 1994; Stanley et al., 1983; Stein et al., 1988, for reviews, see Rayner, 1998). 723

Understanding the source of such differences in eye movements has been the focus of this work, a sample of children with dyslexia while also examining another neurodevelopmental disorder, known to present high comorbidity with RD. In line with previous observations, we found different saccadic programming in children with RD. DD children and NF1 with RD presented longer latencies than TD children, in particular for discrete stimuli (see also Bucci et al., 2008). Differences in the left-right asymmetry in the saccade size were also shown between groups, as previously discussed above. RD readers

were able to accurately perceive and use parafoveal visual information in order to guide their 731 732 eyes during text reading, at least in the usual reading direction. However, they revealed a greater difficulty in reaching the centre of targets presented in the LVF due to the greater 733 734 distance and an unusual reading direction. McKeben et al. (2004) have previously shown that "a mechanism for adapting the gain of reactive, stimulus-triggered saccades (Deubel, 1995) 735 was active during reading in dyslexics, but that it was quantitatively less developed" (p. 396). 736 737 Moreover, RD children didn't show the classical asymmetrical landing position pattern for "linguistic-like" stimuli (with inter-character spaces). This suggests that in poor readers, the 738 preprocessing mechanism that detects the presence or absence of spaces between characters 739 740 and responsible for the direction of attentional scanning is not entirely operational, in particular for the LVF. This result could be interpreted as a deficient allocation of visual 741 742 attention in RD children hampering the exact planning of fine-tuned leftward saccades (e.g., 743 Facoetti, 2012; Facoetti et al., 2010; Lobier et al., 2012; Valdois et al., 2004; see Gavril et al., 2021 for a meta-analysis). In that sense, it was reported that the orienting visuospatial 744 745 attention is impaired in dyslexia with a LVF "mini-neglect" and a RVF over-distractibility 746 (Facoetti, 2001; Hari, 2001; Hari & Renvall, 2001). The similar pattern of results for RD and beginning readers (i.e., ILPs determined by the linguistic nature of stimuli instead of the 747 presence of inter-character spaces; Bellocchi, et al., 2013) speaks against this view. We rather 748 suggest that the lack of sensibility to the discreteness of the stimuli we observed in RD 749 children in the saccade computation, reflects their reading difficulties and/or a lack of 750 experience with written language (see also Hawelka et al., 2006). The cluster analyses support 751 this idea with children of cluster 1 (with the lowest reading level scores) showing an immature 752 saccade targeting strategy (characterised by ILPs near the words' beginning and sequential 753 754 reading), which mirrors their poor reading skills. That goes in line with the above findings on the asymmetry of the saccade extent and the data of Franzen et al. (2021) showing "a 755

Iaborious and more effortful reading strategy in DD, resembling a pattern observed in beginning and poorer readers" (Franzen et al., 2021, p. 9, see also Ashby et al., 2005; Rayner, 1998). These findings can be accounted for by assuming that visual attention distribution (as expressed by ILP) adapts to the differences in reading experience. Therefore, an important result to retain from our study is that eye movements per se are not a cause of dyslexia.

What about the children of cluster 3? They presented an intermediate reading level and 761 exhibits an increased variance in the saccade amplitudes resulting in a flattened and diffuse 762 LP-curve. According to Snell and Theeuwes (2020), a clean oculomotor behaviour is reflected 763 in a narrower spread of ILPs, around the PVL (as observed in experienced readers of the 764 765 cluster 2). ILP spreads in cluster 3 readers widened upon an increased number of positions. 766 Note that the cluster 3 comprised 70% of reading-disabled children (mainly DD). Interestingly, it has been reported that individuals with DD demonstrate a more 767 768 distributed/diffused mode of attention and have difficulty in narrowing their focus of attention (Facoetti et al., 2000, 2003; Franceschini et al., 2012). Such detrimental effect of the 769 increased variability in the ILPs received support from two recent studies (Franzen et al., 770 2021; Rima & Schmid, 2021). Franzen et al. (2021) found an increased number of 771 unexpected/atypical saccades in readers with dyslexia, thus suggesting that their eyes 772 773 occasionally move to seemingly random places. Regarding the distribution of the children's 774 ILPs in the cluster 3, it is conceivable that an increased oculomotor instability would result in more spatial variability in ILPs. Taken together with the performance at the DEM-test, this 775 776 pattern of results may reveal possible occasional oculomotor control deficits in RD, as already reported in previous studies (e.g., Freedman et al., 2017; Raghuram et al., 2018; Tiadi et al., 777 2016; Vagge et al., 2015), which might arise from underlying visual-processing deficits, such 778 as visuo-attentional deficits (Facoetti et al., 2003, 2019; Franceschini et al., 2012, 2022; 779 Valdois et al., 2004; Vernet et al., 2022). 780

The point of particular interest is that poor NF1 readers mirrored the dyslexic eye 781 movement behaviour, with less accuracy and more variability in saccadic programming, no 782 sensibility to the discreteness of the stimuli, particularly in the LVF. Proficient NF1 readers 783 784 showed fully automatized oculomotor mechanisms for proficient reading, as TD children do, using a saccade targeting strategy that aims to land left of the centre for all discrete stimuli so 785 as to be able to scan the stimulus from left to right, thus left-shifting PLP for both left and 786 787 right VF. Note that it was the first study investigating eye movement control related to 788 reading context in NF1. We provided key evidence showing a difference in the saccadic targeting strategies used by NF1 children depending on the presence of a RD associated. In 789 addition, the atypical saccade targeting strategies used by NF1RD children (timid eye-790 movement strategy in cluster 1 and inaccurate ILPs in cluster 3) were not specific to this 791 disease. Finally, the fact that NF1RD and DD children showed similar saccade targeting 792 793 strategies gives clear empirical support to the idea that this at least partly reflects their 794 common poor reading skills (probably related to their limited reading experience, see 795 Cunningham & Stanovich, 1997; Goswami, 2015) rather than an impairment causally related 796 to a reading disorder.

797 There are some limitations to the present study. The first limitation regards the 798 generalizability of the findings with the small sample size of the two NF1 groups due to the 799 prevalence of this rare disease and our strict inclusion and exclusion criteria. For instance, we have made the choice not to include NF1 children with comorbid ADHD although the large 800 proportion of ADHD in this population (i.e., 40%, Hyman et al., 2005). But our priority 801 was to ensure that our results accurately reflect saccade targeting strategies associated with a 802 RD without possible modulation related to inattentive or impulsive behaviour. Another 803 limitation comes from the absence of a direct measure of print exposure to measure 804 the amount of children's reading and be able to identify the impoverished reading experience 805

of RD children as a causal claim. One way could be to systematically include in our future
studies traditional self-report questionnaires and print exposure checklists assessing parents'
familiarity with children's book titles (see Mol & Bus, 2011, for a meta-analysis). In addition,
the heated debate on the causes of dyslexia needs more longitudinal studies starting in
kindergarteners to increase the likelihood of being able to reliably identify explicit causal
links.

What should be taken from the individual analysis is that most of the DD readers of our 812 sample do not exhibit oculomotor deficits (see Hutzler et al., 2006; Prado et al., 2007, for 813 similar conclusions with non-linguistic tasks). Furthermore, many of the impairments 814 815 displayed by RD children are also shown by beginning readers or illiterate individuals who 816 have received very little reading instruction, thus suggesting that eye movements are functional and perform the same function in DD readers that they do in proficient readers. We 817 818 suggest that such differences in performance between children with and without DD are, to a substantial extent, the consequence of their poor experience with written language. 819 Specifically, disruption to oculomotor behaviour simply reflects the fact that many of the 820 component processes involved in learning to read have not yet become fully automatized 821 (LaBerge & Samuels, 1974). For DD readers who exhibited deviant saccade targeting 822 strategies, the possibility of occasional oculomotor control deficits cannot be entirely ruled 823 out supporting the notion that non-linguistic processes could serve as an additional source to 824 explain their impaired reading. Although these low-level oculomotor deficits in dyslexia are 825 difficult to quantify, it would be important to propose a careful assessment of the visual-826 processing and visuo-attentional skills underlying reading to ensure diagnostic and targeted 827 intervention adapted to their deficiencies. 828

830

#### 831 ACKNOWLEDGEMENTS

We are very thankful to all the children and their parents who agreed to contribute to this study. The authors would also like to thank all the research members of the DYSTAC-MAP project, especially Dr F. Audic and Dr M.-O. Livet for the inclusion of the children in the Aix-Marseille hospitals. Finally, we are very grateful for the support and financial contributions of the "Association Neurofibromatoses et Recklinghausen" and the "Fondation de France".

838

### 839 FUNDING

This work was supported by the French National Research Agency (ANR-13-APPR-0010) and by a PhD Fellowship for the first author awarded by the "Fondation de France" and the "Association Neurofibromatoses et Recklinghausen" (VISALECT\_NF1\_00099576).

843

### 844 DECLARATION OF COMPETING INTEREST

845 The authors have no conflicts of interest to disclose.

### REFERENCES

- Ablinger, I., Huber, W., Schattka, K. I., & Radach, R. (2013). Recovery in a letter-by-letter
  reader: more efficiency at the expense of normal reading strategy. *Neurocase*, *19*(3),
  236–255. https://doi.org/10.1080/13554794.2012.667119
- Ans, B., Carbonnel, S., & Valdois, S. (1998). A connectionist multiple-trace memory model
  for polysyllabic word reading. *Psychological Review*, *105*(4), 678–723.
  https://doi.org/10.1037/0033-295X.105.4.678-723
- Arnold, S. S., Payne, J. M., Mcarthur, G., North, K. N., & Barton, B. (2020). Profiling the
  Word Reading Abilities of School-Age Children with Neurofibromatosis Type 1. *Journal of the International Neuropsychological Society*, 1–13.
  https://doi.org/10.1017/S135561772000106X
- Ashby, J., Rayner, K., & Clifton, C. (2005). Eye movements of highly skilled and average
  readers: Differential effects of frequency and predictability. *Quarterly Journal of Experimental Psychology Section A: Human Experimental Psychology*, 58(6), 1065–
  1086. https://doi.org/10.1080/02724980443000476
- Bellocchi, S., & Ducrot, S. (2021). "Same, same but different": The optimal viewing position
  effect in developmental dyslexia, developmental coordination disorder and comorbid
  disorders. *Dyslexia*, 27(3), 294–311. https://doi.org/10.1002/dys.1688
- Bellocchi, S., & Leclercq, V. (2021). Exploring the Moderation Effect of Educational Stage
  on Visual Magnocellular Functioning Linked to Reading: A Study in French Primary
  School Children, 8(2), 68. https://doi.org/10.3390/children8020068
- Bellocchi, S., Mancini, J., Jover, M., Huau, A., Ghio, A., André, C., & Ducrot, S. (2013).
  Dyslexic readers and saccade computation : effects of reading exposure and visuoperceptual constraints [poster communication]. *XVIIIth ESCoP Conference, Budapest, Hungary*. https://hal.archives-ouvertes.fr/hal-01510227
- Bellocchi, S., Massendari, D., Grainger, J., & Ducrot, S. (2019). Effects of inter-character
  spacing on saccade programming in beginning readers and dyslexics. *Child Neuropsychology*, 25(4), 482–506. https://doi.org/10.1080/09297049.2018.1504907
- Bellocchi, S., Muneaux, M., Bastien-Toniazzo, M., & Ducrot, S. (2013). I can read it in your
  eyes: What eye movements tell us about visuo-attentional processes in developmental
  dyslexia. *Research in Developmental Disabilities*, 34(1), 452–460.
- 877 https://doi.org/10.1016/j.ridd.2012.09.002
- Bellocchi, S., Muneaux, M., Huau, A., Lévêque, Y., Jover, M., & Ducrot, S. (2017).
  Exploring the Link between Visual Perception, Visual–Motor Integration, and Reading
  in Normal Developing and Impaired Children using DTVP-2. *Dyslexia*, 23(3), 296–315.
  https://doi.org/10.1002/dys.1561
- Besner, D., Risko, E. F., & Sklair, N. (2005). Spatial Attention as a Necessary Preliminary to
  Early Processes in Reading. *Canadian Journal of Experimental Psychology/Revue Canadienne de Psychologie Expérimentale*, 59(2), 99–108.
- 885 https://doi.org/10.1037/h0087465
- Blythe, H. I. (2014). Developmental Changes in Eye Movements and Visual Information
   Encoding Associated With Learning to Read. *Current Directions in Psychological Science*, 23(3), 201–207. https://doi.org/10.1177/0963721414530145

- Blythe, H. I., & Joseph, H. S. S. L. (2011). Children's eye movements during reading. In S. P.
  Liversedge, I. Gilchrist, & S. Everling (Eds.), *The Oxford Handbook of Eye Movements*
- 891 (pp. 634–662). Oxford University Press.
- 892 https://doi.org/10.1093/oxfordhb/9780199539789.013.0036
- Blythe, H. I., Kirkby, J., & Liversedge, S. (2018). Comments on: "What Is Developmental
  Dyslexia?" Brain Sci. 2018, 8, 26. The Relationship between Eye Movements and
  Reading Difficulties. *Brain Sciences*, 8(6), 100. https://doi.org/10.3390/brainsci8060100
- Bosse, M., & Valdois, S. (2009). Influence of the visual attention span on child reading
  performance: a cross-sectional study. *Journal of Research in Reading*, *32*(2), 230–253.
  https://doi.org/10.1111/j.1467-9817.2008.01387.x
- Brysbaert, M., & Nazir, T. A. (2005). Visual constraints in written word recognition: evidence
  from the optimal viewing-position effect. *Journal of Research in Reading*, 28(3), 216–
  228. https://doi.org/10.1111/j.1467-9817.2005.00266.x
- Bucci, M. P., Brémond-Gignac, D., & Kapoula, Z. (2008). Poor binocular coordination of
  saccades in dyslexic children. *Graefe's Archive for Clinical and Experimental Ophthalmology*, 246(3), 417–428. https://doi.org/10.1007/s00417-007-0723-1
- Chaix, Y., Lion-François, L., Kemlin, I., Faure-Marie, N., Castelnau, P., Barbarot, S.,
  Preclaire, E., Lelong, S., Gentil, C., Lauwers-Cancès, V., Rivier, F., Rodriguez, D.,
  Carniero, M., Iannuzzi, S., Dorison, N., & Schweitzer, E. (2017). Deficit in phonological
  processes: a characteristic of the neuropsychological profile of children with NF1. *Child Neuropsychology*, 24(4), 558–574. https://doi.org/10.1080/09297049.2017.1313970
- Coltheart, Max, Rastle, K., Perry, C., Langdon, R., & Ziegler, J. (2001). DRC: A dual route
  cascaded model of visual word recognition and reading aloud. *Psychological Review*, *108*(1), 204–256. https://doi.org/10.1037/0033-295X.108.1.204
- Cunningham, A. E., & Stanovich, K. E. (1997). Early reading acquisition and its relation to
  reading experience and ability 10 years later. *Developmental Psychology*, *33*(6), 934–
  945. https://doi.org/10.1037/0012-1649.33.6.934
- Deubel, H. (1995). Separate adaptive mechanisms for the control of reactive and volitional
  saccadic eye movements. *Vision Research*, *35*(23–24), 3529–3540.
  https://doi.org/10.1016/0042-6989(95)00058-M
- Ducrot, S., & Grainger, J. (2007). Deployment of spatial attention to words in central and
  peripheral vision. *Perception and Psychophysics*, 69(4), 578–590.
  https://doi.org/10.3758/BF03193915
- Ducrot, S., Lété, B., Descottes, C., Muneaux, M., & Ghio, A. (2006). *The Emaa*(*EyeMovement Acquisition and Analysis*) software package. Unpublished Technical
  Report (67 p.), University of Provence.
- Ducrot, S., Lété, B., Sprenger-Charolles, L., Pynte, J., & Billard, C. (2003). The Optimal
  Viewing Position Effect in Beginning and Dyslexic Readers. *Current Psychology Letters. Behaviour, Brain & Cognition*, 1(10).
- Ducrot, S., & Pynte, J. (2002). What determines the eyes' landing position in words?
   *Perception and Psychophysics*, 64(7), 1130–1144. https://doi.org/10.3758/BF03194762
- Ducrot, S., Pynte, J., Ghio, A., & Lété, B. (2013). Visual and linguistic determinants of the
  eyes' initial fixation position in reading development. *Acta Psychologica*, 142(3), 287–

- 932 298. https://doi.org/10.1016/j.actpsy.2013.01.013
- Eden, G. F., Stein, J. F., Wood, H. M., & Wood, F. B. (1994). Differences in eye movements
  and reading problems in dyslexic and normal children. *Vision Research*, *34*(10), 1345–
  1358. https://doi.org/10.1016/0042-6989(94)90209-7
- Eloranta, A., Närhi, V. M., Eklund, K. M., Ahonen, T. P. S., & Aro, T. I. (2018). Resolving
  reading disability—Childhood predictors and adult- age outcomes. *Dyslexia*, 25(1),
  dys.1605. https://doi.org/10.1002/dys.1605
- Bayesian Estimation for Eye Movement
  Control. *Psychological Science*, *21*(3), 366–371.
  https://doi.org/10.1177/0956797610362060
- Facoetti, A. (2001). Facilitation and inhibition mechanisms of human visuospatial attention in
  a non-search task. *Neuroscience Letters*, 298(1), 45–48. https://doi.org/10.1016/S03043940(00)01719-5
- Facoetti, A. (2012). Spatial attention disorders in developmental dyslexia: towards the
  prevention of reading acquisition deficits. *Visual Aspect of Dyslexia*, 123–136.
- Facoetti, A., Corradi, N., Ruffino, M., Gori, S., & Zorzi, M. (2010). Visual spatial attention
  and speech segmentation are both impaired in preschoolers at familial risk for
  developmental dyslexia. *Dyslexia*, *16*(3), 226–239. https://doi.org/10.1002/dys.413
- Facoetti, A., Franceschini, S., & Gori, S. (2019). Role of Visual Attention in Developmental
  Dyslexia. In *Developmental Dyslexia across Languages and Writing Systems* (pp. 307– 326). Cambridge University Press. https://doi.org/10.1017/9781108553377.014
- Facoetti, A., Luisa Lorusso, M., Paganoni, P., Umiltà, C., & Gastone Mascetti, G. (2003). The
  role of visuospatial attention in developmental dyslexia: evidence from a rehabilitation
  study. *Cognitive Brain Research*, *15*(2), 154–164. https://doi.org/10.1016/S09266410(02)00148-9
- Facoetti, A., Paganoni, P., & Lorusso, M. L. (2000). The spatial distribution of visual
  attention in developmental dyslexia. *Experimental Brain Research*, *132*(4), 531–538.
  https://doi.org/10.1007/s002219900330
- Facoetti, A., Zorzi, M., Cestnick, L., Lorusso, M. L., Molteni, M., Paganoni, P., Umiltà, C., &
  Mascetti, G. G. (2006). The relationship between visuo-spatial attention and nonword
  reading in developmental dyslexia. *Cognitive Neuropsychology*, 23(6), 841–855.
  https://doi.org/10.1080/02643290500483090
- Fischer, B., & Weber, H. (1990). Saccadic reaction times of dyslexic and age-matched normal
   subjects. *Perception*, *19*(6), 805–818. https://doi.org/10.1068/p190805
- Franceschini, S., Bertoni, S., & Puccio, G. (2022). *Hidden in the Labyrinth : Visuo-spatial Attention Deficit in Children with Developmental Dyslexia*. 1–18.
- Franceschini, S., Gori, S., Ruffino, M., Pedrolli, K., & Facoetti, A. (2012). A causal link
  between visual spatial attention and reading acquisition. *Current Biology*, 22(9), 814–
  819. https://doi.org/10.1016/j.cub.2012.03.013
- 971 Franzen, L., Stark, Z., & Johnson, A. P. (2021). Individuals with dyslexia use a different
  972 visual sampling strategy to read text. *Scientific Reports*, *11*(1), 6449.
  973 https://doi.org/10.1038/s41598-021-84945-9

- Freedman, E. G., Molholm, S., Gray, M. J., Belyusar, D., & Foxe, J. J. (2017). Saccade
  adaptation deficits in developmental dyslexia suggest disruption of cerebellar-dependent
  learning. *Journal of Neurodevelopmental Disorders*, 9(1), 36.
  https://doi.org/10.1186/s11689-017-9218-5
- Frith, U. (1985). Beneath the surface of developmental dyslexia. In K. E. Patterson, J. C.
  Marshall, & M. Coltheart (Eds.), *Surface dyslexia* (Vol. 32, Issue 1).
- Gagl, B., Hawelka, S., & Hutzler, F. (2014). A similar correction mechanism in slow and
  fluent readers after suboptimal landing positions. *Frontiers in Human Neuroscience*,
  8(JUNE), 1–10. https://doi.org/10.3389/fnhum.2014.00355
- Gangl, M., Moll, K., Jones, M. W., Banfi, C., Schulte-Körne, G., & Landerl, K. (2018).
  Lexical Reading in Dysfluent Readers of German. *Scientific Studies of Reading*, 22(1),
  24–40. https://doi.org/10.1080/10888438.2017.1339709
- Garzia, R. P., Richman, J. E., Nicholson, S. B., & Gaines, C. S. (1990). A new visual-verbal
  saccade test: the development eye movement test (DEM). *Journal of the American Optometric Association*, 61(2), 124–135.
- Gavril, L., Roşan, A., & Szamosközi, Ştefan. (2021). The role of visual-spatial attention in
   reading development: a meta-analysis. *Cognitive Neuropsychology*, *38*(6), 387–407.
   https://doi.org/10.1080/02643294.2022.2043839
- Gori, S., & Facoetti, A. (2015). How the visual aspects can be crucial in reading acquisition:
  The intriguing case of crowding and developmental dyslexia. *Journal of Vision*, 15(1),
  1–20. https://doi.org/10.1167/15.1.8
- Goswami, U. (2015). Sensory theories of developmental dyslexia: three challenges for
  research. *Nature Reviews Neuroscience*, *16*(1), 43–54. https://doi.org/10.1038/nrn3836
- Häikiö, T., Bertram, R., Hyönä, J., & Niemi, P. (2009). Development of the letter identity
  span in reading: Evidence from the eye movement moving window paradigm. *Journal of Experimental Child Psychology*, *102*(2), 167–181.
  https://doi.org/10.1016/j.jecp.2008.04.002
- Hari, R. (2001). Left minineglect in dyslexic adults. *Brain*, *124*(7), 1373–1380.
   https://doi.org/10.1093/brain/124.7.1373
- Hari, R., & Renvall, H. (2001). Impaired processing of rapid stimulus sequences in dyslexia.
   *Trends in Cognitive Sciences*, 5(12), 525–532. https://doi.org/10.1016/S1364-6613(00)01801-5
- Hautala, J., Hawelka, S., Loberg, O., & Leppänen, P. H. T. (2022). A dynamic adjustment
  model of saccade lengths in reading for word-spaced orthographies: evidence from
  simulations and invisible boundary experiments. *Journal of Cognitive Psychology*, *34*(4),
  435–453. https://doi.org/10.1080/20445911.2021.2011895
- Hawelka, S., Gagl, B., & Wimmer, H. (2010). A dual-route perspective on eye movements of
  dyslexic readers. *Cognition*, 115(3), 367–379.
- 1012 https://doi.org/10.1016/j.cognition.2009.11.004
- Hawelka, S., Huber, C., & Wimmer, H. (2006). Impaired visual processing of letter and digit
  strings in adult dyslexic readers. *Vision Research*, 46(5), 718–723.
  https://doi.org/10.1016/j.visres.2005.09.017

- Hawelka, S., & Wimmer, H. (2005). Impaired visual processing of multi-element arrays is
  associated with increased number of eye movements in dyslexic reading. *Vision Research*, 45(7), 855–863. https://doi.org/10.1016/j.visres.2004.10.007
- Hindmarsh, G. P., Black, A. A., White, S. L. J., Hopkins, S., & Wood, J. M. (2021). Eye
  movement patterns and reading ability in children. Ophthalmic and Physiological Optics,
  41(5), 1134–1143. https://doi.org/10.1111/opo.12854
- Huang, C., Lorusso, M. L., Luo, Z., & Zhao, J. (2019). Developmental Differences in the
   Relationship Between Visual Attention Span and Chinese Reading Fluency. *Frontiers in Psychology*, 10. https://doi.org/10.3389/fpsyg.2019.024
- Huestegge, L., Radach, R., Corbic, D., & Huestegge, S. M. (2009). Oculomotor and linguistic
  determinants of reading development: A longitudinal study. *Vision Research*, 49(24),
  2948–2959. https://doi.org/10.1016/j.visres.2009.09.012
- Huettig, F., Lachmann, T., Reis, A., & Petersson, K. M. (2018). Distinguishing cause from
  effect many deficits associated with developmental dyslexia may be a consequence of
  reduced and suboptimal reading experience. *Language, Cognition and Neuroscience, 33*(3), 333–350. https://doi.org/10.1080/23273798.2017.1348528
- Hung, Y.-N. (2021). The Science of Reading: The Eyes Cannot Lie. *International Journal of Education and Literacy Studies*, 9(4), 26. https://doi.org/10.7575/aiac.ijels.v.9n.4p.26
- Hutzler, F., Kronbichler, M., Jacobs, A. M., & Wimmer, H. (2006). Perhaps correlational but
  not causal: No effect of dyslexic readers' magnocellular system on their eye movements
  during reading. Neuropsychologia, 44(4), 637–648.
  https://doi.org/10.1016/j.neuropsychologia.2005.06.006
- Hutzler, F., & Wimmer, H. (2004). Eye movements of dyslexic children when reading in a
  regular orthography. *Brain and Language*, 89(1), 235–242.
  https://doi.org/10.1016/S0093-934X(03)00401-2
- Hyman, S. L., Shores, A., & North, K. N. (2005). The nature and frequency of cognitive
  deficits in children with neurofibromatosis type 1. *Neurology*, 65(7), 1037–1044.
  https://doi.org/10.1212/01.wnl.0000179303.72345.ce
- Hyönä, J. (1995). An eye movement analysis of topic-shift effect during repeated reading. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 21(5), 1365–
  1373. https://doi.org/10.1037/0278-7393.21.5.A
- Inhoff, A. W., Radach, R., Eiter, B. M., & Juhasz, B. (2003). Distinct Subsystems for the
  Parafoveal Processing of Spatial and Linguistic Information during Eye Fixations in
  Reading. *The Quarterly Journal of Experimental Psychology Section A*, 56(5), 803–827.
  https://doi.org/10.1080/02724980244000639
- Jacquier-Roux, M., Valdois, S., & Zorman, M. (2002). *Outil de dépistage des dyslexies [Dyslexia screening tool]*. Laboratoire de Cogni-Sciences IUFM, Grenoble, France
   (2002).
- Joo, S. J., White, A. L., Strodtman, D. J., & Yeatman, J. D. (2018). Optimizing text for an
  individual's visual system: The contribution of visual crowding to reading difficulties. *Cortex*, 103, 291–301. https://doi.org/10.1016/j.cortex.2018.03.013
- Joseph, H. S. S. L., Liversedge, S. P., Blythe, H. I., White, S. J., & Rayner, K. (2009). Word
   length and landing position effects during reading in children and adults. *Vision*

- 1059 *Research*, 49(16), 2078–2086. https://doi.org/10.1016/j.visres.2009.05.015
- Juhasz, B. J., White, S. J., Liversedge, S. P., & Rayner, K. (2008). Eye movements and the
  use of parafoveal word length information in reading. *Journal of Experimental Psychology: Human Perception and Performance*, *34*(6), 1560–1579.
  https://doi.org/10.1037/a0012319
- Kim, Y.-S., & Pallante, D. (2012). Predictors of reading skills for kindergartners and first
  grade students in Spanish: a longitudinal study. *Reading and Writing*, 25(1), 1–22.
  https://doi.org/10.1007/s11145-010-9244-0
- Kirkby, J. A., Barrington, R. S., Drieghe, D., & Liversedge, S. P. (2022). Parafoveal
  processing and transposed- letter effects in dyslexic reading. *Dyslexia*, 28(3), 359–374.
  https://doi.org/10.1002/dys.1721
- Krab, L. C., Aarsen, F. K., de Goede-Bolder, A., Catsman-Berrevoets, C. E., Arts, W. F.,
  Moll, H. A., & Elgersma, Y. (2008). Impact of neurofibromatosis type 1 on school
  performance. *J Child Neurol*, 23(9), 1002–1010. https://doi.org/0883073808316366
  [pii]10.1177/0883073808316366 [doi]
- LaBerge, D., & Brown, V. (1989). Theory of attentional operations in shape identification.
   *Psychological Review*, 96(1), 101–124. https://doi.org/10.1037/0033-295X.96.1.101
- LaBerge, D., & Samuels, S. J. (1974). Toward a theory of automatic information processing
  in reading. *Cognitive Psychology*, 6(2), 293–323. https://doi.org/10.1016/00100285(74)90015-2
- Landerl, K., Freudenthaler, H. H., Heene, M., De Jong, P. F., Desrochers, A., Manolitsis, G.,
  Parrila, R., & Georgiou, G. K. (2019). Phonological Awareness and Rapid Automatized
  Naming as Longitudinal Predictors of Reading in Five Alphabetic Orthographies with
  Varying Degrees of Consistency. *Scientific Studies of Reading*, 23(3), 220–234.
  https://doi.org/10.1080/10888438.2018.1510936
- Lasker, a G., Denckla, M. B., & Zee, D. S. (2003). Ocular motor behavior of children with
  neurofibromatosis 1. *Journal of Child Neurology*, *18*(5), 348–355.
  https://doi.org/10.1177/08830738030180050301
- Leclercq, V., & Siéroff, E. (2016). Attentional Processing of Letter Strings by Children. *Child Neuropsychology*, 22(1), 110–132. https://doi.org/10.1080/09297049.2014.977242
- Lefavrais, P. (1967). *Test de l'alouette: manuel*. Les éditions du centre de psychologie
   appliquée.
- 1091 Lefavrais, P. (2005). *Alouette-R*. Les éditions du centre de psychologie appliquée.
- Lefton, L. A., Nagle, R. J., Johnson, G., & Fisher, D. F. (1979). Eye Movement Dynamics of
  Good and Poor Readers: Then and Now. *Journal of Reading Behavior*, *11*(4), 319–328.
  https://doi.org/10.1080/10862967909547338
- Legge, G. E., Klitz, T. S., & Tjan, B. S. (1997). Mr. Chips: An ideal-observer model of
   reading. *Psychological Review*, *104*(3), 524–553. https://doi.org/10.1037/0033 295X.104.3.524
- Lehtimäki, T. M., & Reilly, R. G. (2005). Improving eye movement control in young readers.
   *Artificial Intelligence Review*, 24(3–4), 477–488. https://doi.org/10.1007/s10462-005 9010-x

- Leibnitz, L., Grainger, J., Muneaux, M., & Ducrot, S. (2017). Processus visuo-attentionnels et
  lecture : une synthèse. *L'Année Psychologique*, *116*(4), 597–622.
  https://doi.org/10.4074/S0003503316000403
- Leppänen, U., Niemi, P., Aunola, K., & Nurmi, J.-E. (2004). Development of reading skills
  among preschool and primary school pupils. *Reading Research Quarterly*, *39*(1), 72–93.
  https://doi.org/10.1598/RRQ.39.1.5
- 1107 Lété, B., Sprenger-Charolles, L., & Colé, P. (2004). MANULEX: A grade-level lexical
  1108 database from French elementary school readers. *Behavior Research Methods*,
  1109 *Instruments, & Computers*, *36*(1), 156–166. https://doi.org/10.3758/BF03195560
- Li, J., Yang, Y., & Zhao, J. (2021). The development of visual simultaneous processing skill
  subcomponents of Chinese children with developmental dyslexia and the relationship
  with reading. *Acta Psychologica Sinica*, 53(8), 821.
  https://doi.org/10.3724/SP.J.1041.2021.00821
- Liu, T., Thiebaut de Schotten, M., Altarelli, I., Ramus, F., & Zhao, J. (2022). Neural dissociation of visual attention span and phonological deficits in developmental dyslexia:
  A hub- based white matter network analysis. *Human Brain Mapping, November 2021*, 1–10. https://doi.org/10.1002/hbm.25997
- Lobier, M., Zoubrinetzky, R., & Valdois, S. (2012). The visual attention span deficit in
  dyslexia is visual and not verbal. *Cortex*, 48(6), 768–773.
  https://doi.org/10.1016/j.cortex.2011.09.003
- Lorusso, M. L., Facoetti, A., Pesenti, S., Cattaneo, C., Molteni, M., & Geiger, G. (2004).
  Wider recognition in peripheral vision common to different subtypes of dyslexia. *Vision Research*, 44(20), 2413–2424. https://doi.org/10.1016/j.visres.2004.05.001
- McConkie, G. W., Kerr, P. W., Reddix, M. D., & Zola, D. (1988). Eye movement control
  during reading: I. The location of initial eye fixations on words. *Vision Research*, 28(10),
  1107–1118. https://doi.org/10.1016/0042-6989(88)90137-X
- McDonald, R., Lerman, K., & Pereira, F. (2006). Multilingual dependency analysis with a
   two-stage discriminative parser. *CoNLL 2006 Proceedings of the 10th Conference on Computational Natural Language Learning, June*, 216–220.
- McKeben, M., Trauzettel-Klosinski, S., Reinhanrd, J., Duerrwaechter, U., Adler, M., &
  Klosinski, G. (2004). Eye movement control during single-word reading in dyslexics. *Journal of Vision*, 4(5), 388–402. https://doi.org/10.1167/4.5.4
- Mol, S. E., & Bus, A. G. (2011). To read or not to read: A meta-analysis of print exposure
  from infancy to early adulthood. *Psychological Bulletin*, *137*(2), 267–296.
  https://doi.org/10.1037/a0021890
- Nazir, T. A. (2000). Traces of Print Along the Visual Pathway. In *Reading as a Perceptual Process* (pp. 3–22). Elsevier. https://doi.org/10.1016/B978-008043642-5/50003-6
- Nazir, T. A., Ben-Boutayab, N., Decoppet, N., Deutsch, A., & Frost, R. (2004). Reading
  habits, perceptual learning, and recognition of printed words. *Brain and Language*,
  88(3), 294–311. https://doi.org/10.1016/S0093-934X(03)00168-8
- Nuthmann, A., Engbert, R., & Kliegl, R. (2005). Mislocated fixations during reading and the
  inverted optimal viewing position effect. *Vision Research*, 45(17), 2201–2217.
  https://doi.org/10.1016/j.visres.2005.02.014

- O'Regan, J. K., & Lévy-Schoen, A. (1987). Eye-movement strategy and tactics in word 1144 recognition and reading. In M. Coltheart (Ed.), Attention and performance 12: The 1145 1146 psychology of reading (pp. 363–383). Lawrence Erlbaum Associates, Inc.
- Pavlidis, G. T. (1981). Do eye movements hold the key to dyslexia? *Neuropsychologia*, 19(1), 1147 57-64. https://doi.org/10.1016/0028-3932(81)90044-0 1148
- Perea, M., Panadero, V., Moret-Tatay, C., & Gómez, P. (2012). The effects of inter-letter 1149 spacing in visual-word recognition: Evidence with young normal readers and 1150 developmental dyslexics. Learning and Instruction, 22(6), 420-430. 1151 https://doi.org/10.1016/j.learninstruc.2012.04.001
- 1152
- 1153 Pirozzolo, F. J., & Rayner, K. (1979). The Neural Control of Eye Movements in Acquired and Developmental Reading Disorders. In H. Whitaker & H. A. Whitaker (Eds.), Studies in 1154 Neurolinguistics (pp. 97-123). Elsevier. https://doi.org/10.1016/B978-0-12-746304-1155 1156 9.50009-4
- 1157 Plaza, M., & Cohen, H. (2007). The contribution of phonological awarness and visual attention in early reading and spelling. Dyslexia, 239(13), 234–239. 1158 https://doi.org/10.1002/dys 1159
- Prado, C., Dubois, M., & Valdois, S. (2007). The eye movements of dyslexic children during 1160 1161 reading and visual search: Impact of the visual attention span. Vision Research, 47(19), 2521-2530. https://doi.org/10.1016/j.visres.2007.06.001 1162
- 1163 Premeti, A., Bucci, M. P., & Isel, F. (2022). Evidence from ERP and Eye Movements as 1164 Markers of Language Dysfunction in Dyslexia. Brain Sciences, 12(1), 73. https://doi.org/10.3390/brainsci12010073 1165
- Radach, R., & Kempe, V. (1993). An individual analysis of initial fixation positions in 1166 reading. In Perception and cognition: Advances in eye movement research. (pp. 213– 1167 225). North-Holland/Elsevier Science Publishers. 1168
- Radach, R., & McConkie, G. W. (1998). Determinants of Fixation Positions in Words During 1169 Reading. In Eye Guidance in Reading and Scene Perception (pp. 77–100). Elsevier. 1170 https://doi.org/10.1016/B978-008043361-5/50005-5 1171
- 1172 Raghuram, A., Gowrisankaran, S., Swanson, E., Zurakowski, D., Hunter, D. G., & Waber, D. P. (2018). Frequency of Visual Deficits in Children With Developmental Dyslexia. 1173 JAMA Ophthalmology, 136(10), 1089. 1174
- https://doi.org/10.1001/jamaophthalmol.2018.2797 1175
- Rayner, K. (1975). Parafoveal identification during a fixation in reading. Acta Psychologica, 1176 39(4), 271-281. https://doi.org/10.1016/0001-6918(75)90011-6 1177
- Rayner, K. (1979). Eye guidance in reading: Fixation locations within words. Perception, 1178 1179 8(6), 21–30.
- Rayner, K. (1986). Eye movements and the perceptual span in beginning and skilled readers. 1180 Journal of Experimental Child Psychology, 41(2), 211–236. 1181 https://doi.org/10.1016/0022-0965(86)90037-8 1182
- 1183 Rayner, K. (1998). Eye movements in reading and information processing: 20 years of research. Psychological Bulletin, 124(3), 372-422. https://doi.org/10.1037/0033-1184 2909.124.3.372 1185

- Rayner, K. (2009). Eye Movements in Reading: Models and Data. *Journal of Eye Movement Research*, 2(5), 1–10. https://doi.org/10.16910/jemr.2.5.2
- 1188 Rayner, K., Fischer, M. H., & Pollatsek, A. (1998). Unspaced text interferes with both word
  1189 identification and eye movement control. *Vision Research*, *38*(8), 1129–1144.
  1190 https://doi.org/10.1016/S0042-6989(97)00274-5
- 1191 Rayner, K., Slattery, T. J., & Bélanger, N. N. (2010). Eye movements, the perceptual span,
  1192 and reading speed. *Psychonomic Bulletin & Review*, *17*(6), 834–839.
  1193 https://doi.org/10.3758/PBR.17.6.834
- Rayner, K., White, S. J., Kambe, G., Miller, B., & Liversedge, S. (2003). On the processing of
  meaning from parafoveal vision during eye fixations in reading. In J. Hyönä, R. Radach,
  & H. Deubel (Eds). *The Mind's Eye: Cognitive and Applied Aspects of Eye Movement Research*, 213–234.
- Reichle, E. D., Rayner, K., & Pollatsek, A. (2003). The E-Z Reader model of eye-movement
  control in reading: Comparisons to other models. *Behavioral and Brain Sciences*, 26(4),
  445–476. https://doi.org/10.1017/S0140525X03000104
- Ribeiro, M. J., Violante, I. R., Ribeiro, I., Bernardino, I., Ramos, F., Saraiva, J., Reviriego, P.,
  Upadhyaya, M., Silva, E. D., & Castelo-Branco, M. (2012). Abnormal achromatic and
  chromatic contrast sensitivity in neurofibromatosis type 1. *Investigative Ophthalmology and Visual Science*, 53(1), 287–293. https://doi.org/10.1167/iovs.11-8225
- Rima, S., & Schmid, M. C. (2021). Reading Specific Small Saccades Predict Individual
   Phonemic Awareness and Reading Speed. *Frontiers in Neuroscience*, 15(December), 1–
   13. https://doi.org/10.3389/fnins.2021.663242
- Schattka, K. I., Radach, R., & Huber, W. (2010). Eye movement correlates of acquired central dyslexia. *Neuropsychologia*, 48(10), 2959–2973.
  https://doi.org/10.1016/j.neuropsychologia.2010.06.005
- Silva, S., Faísca, L., Araújo, S., Casaca, L., Carvalho, L., Petersson, K. M., & Reis, A. (2016).
  Too little or too much? Parafoveal preview benefits and parafoveal load costs in dyslexic adults. *Annals of Dyslexia*, 66(2), 187–201. https://doi.org/10.1007/s11881-015-0113-z
- Sireteanu, R., Goertz, R., Bachert, I., & Wandert, T. (2005). Children with developmental dyslexia show a left visual "minineglect." *Vision Research*, 45(25–26), 3075–3082.
  https://doi.org/10.1016/j.visres.2005.07.030
- Smart, D., Youssef, G. J., Sanson, A., Prior, M., Toumbourou, J. W., & Olsson, C. A. (2017).
  Consequences of childhood reading difficulties and behaviour problems for educational
  achievement and employment in early adulthood. *British Journal of Educational Psychology*, 87(2), 288–308. https://doi.org/10.1111/bjep.12150
- Snell, J., & Theeuwes, J. (2020). A story about statistical learning in a story: Regularities
  impact eye movements during book reading. *Journal of Memory and Language*, *113*(August 2019), 104127. https://doi.org/10.1016/j.jml.2020.104127
- Sperlich, A., Meixner, J., & Laubrock, J. (2016). Development of the perceptual span in
   reading: A longitudinal study. *Journal of Experimental Child Psychology*, *146*, 181–201.
   https://doi.org/10.1016/j.jecp.2016.02.007
- Spinelli, D., De Luca, M., Judica, A., & Zoccolotti, P. (2002). Crowding effects on word
   identification in developmental dyslexia. *Cortex*, *38*(2), 179–200.

- 1229 https://doi.org/10.1016/S0010-9452(08)70649-X
- Stanley, G., Smith, G. A., & Howell, E. A. (1983). Eye-movements and sequential tracking in
  dyslexic and control children. *British Journal of Psychology*, 74(2), 181–187.
  https://doi.org/10.1111/j.2044-8295.1983.tb01852.x
- Stein, J. F. (2001). The magnocellular theory of developmental dyslexia. *Dyslexia*, 7(1), 12–
   36. https://doi.org/10.1002/dys.186
- Stein, J. F., Riddell, P. M., & Fowler, S. (1988). Disordered vergence control in dyslexic
  children. *British Journal of Ophthalmology*, 72(3), 162–166.
  https://doi.org/10.1136/bjo.72.3.162
- Tiadi, A., Gérard, C.-L., Peyre, H., Bui-Quoc, E., & Bucci, M. P. (2016). Immaturity of
  Visual Fixations in Dyslexic Children. *Frontiers in Human Neuroscience*, 10(FEB2016),
  1–7. https://doi.org/10.3389/fnhum.2016.00058
- Tydgat, I., & Grainger, J. (2009). Serial Position Effects in the Identification of Letters,
  Digits, and Symbols. *Journal of Experimental Psychology: Human Perception and Performance*, *35*(2), 480–498. https://doi.org/10.1037/a0013027
- Vagge, A., Cavanna, M., Traverso, C. E., & Iester, M. (2015). Evaluation of ocular
  movements in patients with dyslexia. *Annals of Dyslexia*, 65(1), 24–32.
  https://doi.org/10.1007/s11881-015-0098-7
- Valdois, S., Bosse, M.-L., & Tainturier, M.-J. (2004). The cognitive deficits responsible for
   developmental dyslexia: Review of evidence for a selective visual attentional disorder.
   *Dyslexia*, 10(4), 339–363. https://doi.org/10.1002/dys.284
- Vernet, M., Jover, M., Bellocchi, S., Maziero, S., Jucla, M., Tallet, J., Danna, J., Chaix, Y., &
  Ducrot, S. (2022). Visual-processing deficits in children with neurofibromatosis type 1:
  A clinical marker of reading difficulties. *European Journal of Paediatric Neurology*, *38*,
  25–32. https://doi.org/10.1016/j.ejpn.2022.03.009
- 1254 Vidyasagar, T. R., & Pammer, K. (2010). Dyslexia: a deficit in visuo-spatial attention, not in
   phonological processing. *Trends in Cognitive Sciences*, 14(2), 57–63.
   https://doi.org/10.1016/j.tics.2009.12.003
- 1257 Vitu, F. (2003). The basic assumptions of E-Z Reader are not well-founded. *Behavioral and* 1258 *Brain Sciences*, 26(4), 506–507. https://doi.org/10.1017/S0140525X0351010X
- 1259 Vitu, F., O'Regan, J. K., Inhoff, A. W., & Topolski, R. (1995). Mindless reading: Eye1260 movement characteristics are similar in scanning letter strings and reading texts.
  1261 *Perception & Psychophysics*, 57(3), 352–364. https://doi.org/10.3758/BF03213060
- Vitu, F., O'Regan, J. K., & Mittau, M. (1990). Optimal landing position in reading isolated
  words and continuous text. *Perception & Psychophysics*, 47(6), 583–600.
  https://doi.org/10.3758/BF03203111
- Vorstius, C., Radach, R., & Lonigan, C. J. (2014). Eye movements in developing readers: A
  comparison of silent and oral sentence reading. *Visual Cognition*, 22(3–4), 458–485.
  https://doi.org/10.1080/13506285.2014.881445
- White, S. J., & Liversedge, S. P. (2006). Foveal processing difficulty does not modulate nonfoveal orthographic influences on fixation positions. *Vision Research*, 46(3), 426–437.
  https://doi.org/10.1016/j.visres.2005.07.006

- 1271 World Health Organisation. (2008). *59th general assembly* (A. WM (ed.)). WHO.
- 1272 Zegarra-Moran, O., & Geiger, G. (1993). Visual Recognition in the Peripheral Field: Letters
   1273 versus Symbols and Adults versus Children. *Perception*, 22(1), 77–90.
   1274 https://doi.org/10.1068/p220077
- 1275 Zorzi, M., Barbiero, C., Facoetti, A., Lonciari, I., Carrozzi, M., Montico, M., Bravar, L.,
- 1276 George, F., Pech-Georgel, C., & Ziegler, J. C. (2012). Extra-large letter spacing
- 1277 improves reading in dyslexia. *Proceedings of the National Academy of Sciences of the*
- 1278 United States of America, 109(28), 11455–11459.
- 1279 https://doi.org/10.1073/pnas.1205566109

Table 1. Mean of initial saccade size (in characters), saccade latency (in milliseconds) and refixation probability (in percent) as a function of stimulus type, visual field and group in Experiment 1.

	Initial saccade size Mean (SD)		Saccade latency Mean (SD)		Refixation probability Mean (SD)	
	TD	DD	TD	DD	TD	DD
Left Visual Field						
Words	4.33 (0.74)	3.95 (1.30)	170.55 (13.86)	194.94 (28.02)	74.32 (19.69)	55.75 (24.52)
Strings of hashes	4.34 (0.93)	3.72 (0.64)	173.59 (19.96)	193.64 (25.79)	69.24 (24.72)	49.20 (22.42)
Solid lines	3.59 (0.84)	3.64 (0.99)	217.92 (28.47)	220.19 (29.70)	69.02 (19.53)	48.93 (21.98)
Right Visual Field						
Words	3.40 (0.78)	3.27 (0.87)	173.85 (21.51)	191.36 (33.15)	53.64 (28.45)	72.34 (24.84)
Strings of hashes	3.49 (0.54)	3.63 (0.56)	173.91 (21.84)	192.18 (38.67)	70.88 (22.71)	59.44 (22.77)
Solid lines	3.54 (0.59)	3.60 (0.57)	221.18 (17.23)	220.63 (24.91)	71.43 (24.44)	60.17 (26.62)

*Notes.* Saccade size was measured according to the central fixation point. TD: typicallydeveloping children; DD: children with developmental dyslexia.

Table 2. Mean of initial saccade size (in characters), saccade latency (in milliseconds) and
refixation probability (in percent) as a function of stimulus type, visual field, and group in
Experiment 2.

	Initial saccade size Mean (SD)		Saccade latency Mean (SD)		Refixation probability Mean (SD)	
	NF1noRD	NF1RD	NF1noRD	NF1RD	NF1noRD	NF1RD
Left Visual Field						
Words	4.46 (0.71)	4.22 (0.52)	173.91 (8.81)	188.88 (12.88)	74.96 (25.10)	47.79 (26.49)
Strings of hashes	4.43 (0.26)	3.55 (0.55)	175.38 (11.29)	191.10 (19.02)	71.03 (23.59)	51.89 (23.82)
Solid lines	3.67 (0.96)	3.64 (1.22)	219.34 (32.24)	222.05 (35.39)	69.05 (24.85)	45.98 (23.06)
Right Visual Field						
Words	3.43 (0.57)	3.34 (0.63)	173.47 (15.93)	182.89 (12.20)	53.92 (33.42)	72.41 (21.02)
Strings of hashes	3.50 (0.90)	3.53 (0.70)	174.61 (14.54)	184.78 (17.63)	70.24 (25.97)	47.05 (29.55)
Solid lines	3.67 (1.18)	3.52 (0.65)	220.33 (22.14)	222.51 (25.85)	71.11 (30.09)	47.65 (23.47)

1290 *Notes.* Initial saccade size was measured according to the central fixation point. NF1RD: NF1

1291 children with reading disorders; NF1noRD: NF1 children without reading disorders.



Figure 1. Procedure used in the oculomotor lateralized bisection task. The different display conditions of the stimuli are represented in this figure with the 3 types of stimuli (words, hashes and lines) and the 2 visual fields of presentation (LVF and RVF). When the target item appeared, participants were instructed to move their eyes to look at what they considered to be the middle of the item (see arrows and dashed vertical lines in the figure).



Figure 2. Individual distribution of first fixation position specifically for words presented in
the RVF in NF1noRD (i.e., left part of the figure) and NF1RD (i.e., right part of the figure)
children.

*Notes.* This representation was made only for the words presented in the RVF to observe the
oculomotor behaviour in the most reading-like situation. Each line represents the individual
performance of a different participant. NF1RD: NF1 children with reading disorders;
NF1noRD: NF1 children without reading disorders.



Figure 3. Comparison of the initial landing position's frequency for words in the RVFbetween the 4 groups of children in Experiment 1 and 2.

*Notes.* This representation was made only for the words presented in the RVF to observe the
oculomotor behaviour in the most reading-like situation. TD: Typically developing children;
DD: children with developmental dyslexia; NF1RD: NF1 children with reading disorders;
NF1noRD: NF1 children without reading disorders.





Figure 4. Frequency of the initial landing of the first saccade (in percentage) at each positionof words according to reading age (in months) and groups.

Notes. These representations were made only for the words presented in the RVF to observe
the oculomotor behaviour in the most reading-like situation. TD: Typically developing
children; DD: children with developmental dyslexia; NF1RD: NF1 children with reading
disorders; NF1noRD: NF1 children without reading disorders.



Figure 5. Distributions of the initial landing position (in percentage) according to the 3clusters found for all groups combined.

1323 Notes. This figure was made only for the words presented in the RVF to observe the

1324 oculomotor behaviour in the most reading-like situation.