

# A historical record of benthic foraminifera in seasonally anoxic Lake Grevelingen, the Netherlands

Julien Richirt, Anaïs Guihéneuf, Aurélia Mouret, Magali Schweizer, Caroline Slomp, Frans Jorissen

## ▶ To cite this version:

Julien Richirt, Anaïs Guihéneuf, Aurélia Mouret, Magali Schweizer, Caroline Slomp, et al.. A historical record of benthic foraminifera in seasonally anoxic Lake Grevelingen, the Netherlands. Palaeo-geography, Palaeoclimatology, Palaeoecology, 2022, 599, pp.111057. 10.1016/j.palaeo.2022.111057. hal-03925917

## HAL Id: hal-03925917 https://hal.science/hal-03925917

Submitted on 8 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	A HISTORICAL RECORD OF BENTHIC FORAMINIFERA IN
2	SEASONALLY ANOXIC LAKE GREVELINGEN, THE NETHERLANDS
3	
4	Julien Richirt <sup>1</sup> , Anaïs Guihéneuf <sup>1</sup> , Aurélia Mouret <sup>1</sup> , Magali Schweizer <sup>1</sup> , Caroline P.
5	SLOMP <sup>2</sup> AND FRANS J. JORISSEN <sup>1</sup>
6	
7	
8 9 10 11 12 13	<sup>1</sup> UMR 6112 LPG, University of Angers, Nantes University, Le Mans University, CNRS, 2 Boulevard Lavoisier, 49045 Angers, France <sup>2</sup> Department of Earth Sciences (Geochemistry), Faculty of Geosciences, Utrecht University, Princetonlaan 8a, 3584 CB Utrecht, the Netherlands
14	*Corresponding author: richirt.julien@gmail.com
15	
16 17	
1/ 19	
10	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29 26	
30	

#### 31 Abstract

32 Lake Grevelingen is a former branch of the Rhine-Meuse-Scheldt estuary, which was 33 artificially transformed into a salt-water lake by a dam in 1971. This transformation 34 induced profound changes in the biological community of the basin, which have been 35 described as an ecodisaster, with seasonal hypoxia/anoxia occurring in the deepest parts 36 of the lake. Here, we investigate a sediment core sampled in 2012 in the Den Osse Basin 37 (34 m depth) of Lake Grevelingen, recording the last ~ 50 years and including the 38 transition from an estuary to a salt-water lake. Sediment molybdenum (Mo) concentrations were used to refine an existing age model based on <sup>210</sup>Pb, giving us an 39 40 estimated precision of  $\pm$  3 years. The benthic foraminiferal succession reflects the anthropogenic modifications in Lake Grevelingen and allows four successive stages to be 41 42 distinguished: 1) before 1971, when the system was estuarine; 2) from 1971 to 1978, when 43 the system rapidly changed into an enclosed brackish water lake; 3) from 1978 to 1999, 44 when a seaward sluice was opened during winter and 4) the period from 1999 to 2012, with a year-round opening of the sluice and doubling of water exchanges with the North 45 46 Sea. The foraminiferal record, showing communities mainly dominated by *Elphidium* selsevense, also highlights the appearance of the putatively exotic Ammonia confertitesta 47 48 in the mid-1980s, which thereafter progressively replaced the congeneric autochthonous 49 Ammonia aberdoveyensis. Finally, we hypothesise that the activity of cable bacteria, S-50 oxidising prokaryotes present in the surface sediments of Den Osse Basin in winter and 51 spring, causes dissolution of foraminiferal tests by decreasing the pore water carbonate saturation state in the sediments directly below the oxic zone. This explains the contrast 52 53 between the abundant living populations and the very low numbers of foraminiferal shells 54 preserved in the top ~ 15 cm of the sediment. 55 56 57 58 Keywords: foraminifera, historical record, Lake Grevelingen, anoxia, cable bacteria 59 60

61 62

## 76 **1. INTRODUCTION**

77 Foraminifera are among the most widespread groups of marine unicellular eukaryotes and 78 constitute one of the most diverse groups of shelled organisms. They have a very rich fossil 79 record and are widely used for palaeoenvironmental reconstructions and palaeoecological 80 interpretations (Murray, 2006; Katz et al., 2010). Foraminifera are known to be relatively 81 tolerant to low oxygen concentrations, anoxia and/or presence of sulphides in their habitat, on 82 various timescales. This has been documented for many species from different types of 83 environments in laboratory culture (e.g. Moodley and Hess, 1992; Bernhard, 1993; Alve and 84 Bernhard, 1995; Moodley et al., 1998; Duijnstee et al., 2003; Geslin et al., 2004; Pucci et al., 85 2009) as well as in field studies (e.g. Piña-Ochoa et al., 2010; Langlet et al., 2013, 2014). 86 Unfortunately, studies on the foraminiferal response in systems affected by seasonal hypoxia-87 anoxia with sulphidic conditions are still very sparse. To our knowledge, all earlier studies show 88 that the foraminiferal response to hypoxia-anoxia is species-specific (e.g. Bernhard and Alve, 89 1996; Ernst et al., 2005; Bouchet et al., 2007; Geslin et al., 2014; Langlet et al., 2014). However, 90 this species-specific response generally follows the same scheme (usually decrease in density, 91 reduction of growth and/or reproduction), with different response intensities. Recently, a study 92 investigated the temporal response of living foraminiferal communities to recent seasonal 93 anoxia in the Den Osse Basin of Lake Grevelingen (the Netherlands, Richirt et al. 2020). The 94 authors highlighted that the deepest part of this basin shows a severe decline in foraminiferal 95 populations in the months following late summer-early autumn seasonal anoxia. They 96 explained this by the presence of sulphides in the foraminiferal habitat, that would inhibit 97 foraminiferal reproduction and growth. In the shallower parts of the basin, where the duration 98 of anoxia is shorter, there was not such a major decline in the foraminiferal community.

99 The Den Osse Basin is located in Lake Grevelingen, a former estuary of the Scheldt-Meuse-100 Rhine delta, in the Netherlands. After a storm in 1953, which caused large scale flooding and 101 more than 1800 casualties, landward (1965) and seaward (1971) dams were constructed to avoid 102 such disasters in future (Delta Plan, Bannink et al., 1984; Figure 1). Thanks to the preservation 103 of shelled foraminifera in the sediment record, Lake Grevelingen provides a unique opportunity 104 to study the benthic foraminiferal response to the progressive transformation of the ecosystem from an estuary to a salt water lake, especially regarding the development of seasonal 105 106 hypoxia/anoxia.

107 The living foraminiferal community of the Den Osse Basin is largely dominated by 108 *Elphidium selseyense* (phylotype S5, Darling et al., 2016), *Elphidium magellanicum* 

109 (morphospecies not yet sequenced) and Ammonia confertitesta (Ammonia sp. T6 in Richirt et 110 al., 2020; phylotype T6 in Hayward et al., 2021). The latter species, A. confertitesta, is often 111 considered as an exotic and/or invasive species, originating from East Asia (e.g. Pawlowski & 112 Holzmann, 2008; Schweizer et al, 2011; Bird et al., 2020). Ammonia confertitesta, which is 113 almost the only representative of the genus Ammonia in the recent living assemblages (Richirt 114 et al., 2020), is characterised by very large pores (Richirt et al., 2019a) compared to Ammonia 115 aberdoveyensis and to a lesser extend Ammonia veneta (phylotypes T2 and T1, respectively, 116 Hayward et al., 2021). Interestingly, a preliminary study of a long core sampled in the Den Osse 117 Basin showed a clear shift from specimens with small pores to specimens with bigger pores in 118 recent times (Petersen et al., 2016). This indicatess that in a recent past, A. confertitesta has 119 progressively replaced A. aberdoveyensis (and/or A. veneta) in Lake Grevelingen. Richirt et al. 120 (2019b), suggesting that an increase in overall porosity could be an adaptation to low oxygen 121 concentrations. The putative replacement of A. aberdoveyensis (and/or A. veneta) by A. 122 *confertitesta* could therefore be due to the larger tolerance of the latter species to hypoxia. The 123 study of the sedimentary foraminiferal record should enable us to shed more light on this 124 important species take-over.

In this study, we investigate foraminiferal assemblages of a 90 cm long sediment core sampled in the deepest part of the Den Osse Basin, representing the last ~ 50 years. We will especially address the following questions:

How did the foraminiferal community change in response to the closure of Lake
 Grevelingen in the 1970s and the subsequent management changes of the basin?

2) Has there been a substitution of the autochthonous *Ammonia* species (*A. veneta* and *A. aberdoveyensis*) by the putative allochthonous species *A. confertitesta*, and, if so, is it the result
of an overall spreading of *A. confertitesta* across Europe, or rather of a better adaptation of this
phylotype to seasonal hypoxia/anoxia?

## **2. STUDY AREA**

In 1965, the former Grevelingen estuary was closed landward (Grevelingendam) and became a semi-enclosed embayment without river input. A second dam was constructed seaward in 1971 (Brouwersdam), closing the former Grevelingen estuary completely. This stopped the inflow of salt waters from the North Sea and increased the water residence time in the Grevelingen lake up to three to six years (instead of a few days, when the system was estuarine, Nienhuis, 1978; Bannink et al., 1984). The suppression of tidal currents and decrease

141 in salinity directly after the full closure strongly affected marine benthic animals (e.g. sponges, 142 sea anemones, tunicates). Mass mortality of these animals contributed to increased food 143 availability, accumulation of decomposition products and finally oxygen depletion. This 144 "ecodisaster" induced a simplification of the ecosystem concerning both its structure and 145 functioning (Nienhuis & Veld, 1984; Bannink et al., 1984). Within a few years after closure of 146 the estuary, seasonal bottom water hypoxia/anoxia started to occur, followed by mass mortality 147 of the remaining marine benthic animal populations (e.g. molluscs, worms), especially in the 148 deeper parts of the basin (Nienhuis & Veld, 1984).

149 To remedy eutrophication phenomena and seasonal hypoxia by limiting water column 150 stratification, the connection with the North Sea was partly reopened in 1978, by the construction of an underwater sluice in the Brouwersdam (capacity of about 100–140 m<sup>3</sup> s<sup>-1</sup>, 151 152 initially only opened in winter, Bannink et al. 1984; Muelen et al., 1984). This partial reopening 153 improved the bottom water oxygenation in the Den Osse Basin and decreased water residence 154 times to about half a year. Consequently, several species, which were common in the 155 Grevelingen estuary but had disappeared when the estuary was closed, were observed again 156 (e.g. some jellyfish and crab species; Nienhuis and Veld, 1984). A second, landward sluice 157 (siphon) was opened in 1983 (at the Grevelingendam) to ensure a weak (compared to the natural situation without a dam) freshwater inflow (about 100 m<sup>3</sup> s<sup>-1</sup>, Bannink et al., 1984; Muelen et 158 159 al., 1984).

In 1999, the water exchanges between the Lake and the North Sea were doubled (from 160 1245–1255.10<sup>6</sup> m<sup>3</sup> to 2688–2864.10<sup>6</sup> m<sup>3</sup> per year, Wetsteijn, 2011) by opening the 161 Brouwersdam sluice the whole year, except for a few weeks in autumn and during storm events. 162 163 This allowed a further decrease of the average water residence time in the Lake from 164 to 72 164 days and limited water stratification due to high temperatures in summer (Wetsteijn, 2011). 165 However, this situation was still very different compared to the estuarine situation before 1965, 166 when the water residence time was estimated to be a few days (Nienhuis, 1978; Bannink et al., 1984) and average flood and ebb discharges at the location of the present Brouwersdam were 167 about 126000.10<sup>6</sup> and 135000.10<sup>6</sup> m<sup>3</sup> per year, respectively (calculated for 1959, Louters et al., 168 169 1998), ~ 50 times higher than the current water exchanges.

Despite these management changes, seasonal bottom water oxygen depletion continued to occur every year, as is still the case today, especially in the deepest parts of Lake Grevelingen, such as in the Den Osse Basin that is investigated in this study (Wetsteyn, 2011; Sulu-Gambari et al. 2017). The transformation of Grevelingen from an estuary to a salt-water lake induced profound modifications at ecosystem (e.g. habitat losses, changes in trophic relationships), community (e.g. decrease of biodiversity, replacement of marine by estuarine species) as well
as species levels (e.g. disturbed or interrupted life cycles, change in densities, Nienhuis, 1978;
Nienhuis & Veld, 1984).

To summarise, Lake Grevelingen, a former estuarine system, was completely closed in 179 1971. The ensuing strong increase of water residence time, bottom water stagnation and lower 180 salinity caused a major biotope shift in Lake Grevelingen and led to strong modifications of its 181 biological community (Nienhuis, 1978).

Today, two antagonistic groups of sulphur-oxidizing bacteria showing contrasting seasonal 182 183 population dynamics are present in sediments of the Den Osse Basin: cable bacteria in January-184 May and Beggiatoaceae in September-December (Seitaj et al., 2015; Sulu-Gambari et al., 185 2016a). Cable bacteria activity is known to strongly impact biogeochemical cycles (e.g. of 186 sulphur, iron or calcium) in the sediment, by promoting, for example, precipitation of iron oxide 187 in the oxic layer and dissolution of iron sulphide and calcium carbonate in the anoxic layer 188 (Risgaard-Petersen et al., 2012; Meysman et al., 2015). In the deepest part of the Den Osse 189 Basin, where the maximum oxygen penetration depth in the sediment is about 2 mm (Seitaj et 190 al., 2015), this was shown to occur seasonally, as a result of cable bacteria activity (Sulu-191 Gambari et al., 2016a, b). This phenomenon could play a major role in foraminiferal calcium 192 carbonate test dissolution during taphonomic processes.



193

194 <u>Figure 1.</u> Timeline of the human-induced modifications following the Delta Plan at Lake
 195 Grevelingen. The sediment core used to investigate dead foraminiferal assemblages was
 196 sampled in May 2012 (red line).

## 197 **3. MATERIALS AND METHODS**

## 198 **3.1. SAMPLING AND LABORATORY TREATMENT**



199

Figure 2. A. Map of The Netherlands (on the left) showing the location of Lake Grevelingen.
B. Map of Lake Grevelingen (main panel) showing the location of the sediment sampling in the
Den Osse Basin (red star). C. The transversal section of the Den Osse basin (top right) shows
the depth at which the studied sediment core was sampled at Station 1 (S1, 34 m depth). Station
2(S2, 23 m depth) was investigated for living assemblages by Richirt et al. (2020), as noted in
the discussion. This figure was modified from Sulu-Gambari et al. (2016a).

The study site (station 1) is in the Den Osse Basin (51° 44.834'N, 3° 53.401'E), the main 206 207 former channel of the Grevelingen estuary (The Netherlands), at 34 m depth (Figure 2). Our 208 site is located at a distance of  $\sim 100$  m from a location where water column oxygen was 209 intensively monitored since the late 1970s by the Dutch Water Agency (DWA; e.g. Wetsteijn, 210 2011). Bottom water oxygen concentrations from 1978 to 2011 from the DWA database were summarised previously (Figure 3; Sulu-Gambari et al., 2017) and reveal recurring seasonal 211 212 hypoxia and anoxia in bottom water. These bottom water data were collected up to about 3 m 213 above the basin floor (and not at the sediment-water interface), which was at 37 m water depth 214 at this location. Consequently, bottom water anoxia measured by the water column monitoring 215 (Figure 3) is a reliable indicator of an anoxic sediment-water interface. However, oxic 216 conditions in the water column (as frequently recorded from late 1990s to 2010, Figure 3), does 217 not necessarily indicate that the basin floor was also oxic.





Figure 3. Periods of oxic (blue), hypoxic (grey) and anoxic (black) bottom water conditions
from 1978 to 2011 at the DWA site, which is located at a distance of about 100 m from station
1 in this study (figure after Sulu-Gambari et al., 2017).

222 A sediment core of 90 cm length and 6 cm diameter was collected at station 1 in May 2012 223 with a UWITEC gravity interface corer. In the laboratory, the core was cut into 0.5 cm thick slices with an individual sample volume of ~  $14.1 \text{ cm}^3$ . All samples were sieved at 63, 125, 150 224 225 and  $315 \,\mu\text{m}$ , before being dried and stored in the laboratory. For this study, only for a forminifera 226 of the fraction > 125  $\mu$ m (including all fractions above 125  $\mu$ m) were picked and stored in 227 micropalaeontological Plummer cells. Considering that picking foraminifera is very time-228 consuming, we decided to look at one sample every 2.5 cm (i.e., every fifth sample). Because 229 the 54.5–55, 84.5–85 and 89.5–90 cm samples had a very high foraminiferal density (Suppl. 230 Table 1), they were split into half (125–150, 150–315 and >315  $\mu$ m for the first two, 125–150 231 µm fraction for the 89.5–90 cm sample) and complete splits were counted. Conversely, in the 232 top 15 cm, many samples were very poor in foraminifera (Suppl. Table 1). Consequently, we 233 decided to increase the resolution around these samples by picking adjacent samples. In all, the 234 foraminiferal content of 40 samples was studied (Suppl. Table 1). When possible, determination 235 was made to the species level using a stereomicroscope.

A major change in sediment composition was observed, from sandy deposits below 82.5 cm depth, very probably representative of the strong hydrodynamics of the former estuarine system to muddy/silty sediments above 82.5 cm depth, indicative of the much lower hydrodynamics of the salt lake environment.

#### 240 **3.2.** FORAMINIFERAL ANALYSES

257

In cases where samples had very low foraminiferal abundances, two to three adjacent samples were merged (Table 1). This concerned samples in depth intervals 4.5–6, 6.5–8, 9– 10.5, 11.5–13, 32–35, 57–60, 69.5–72.5 and 74.5–77.5 cm. We compromised between (1) maximising the number of individuals (to obtain a target value of at least 50 individuals) and (2) maximising sampling resolution. The total thickness of merged samples never exceeded 3 cm (Table 1).

To document temporal changes in assemblage composition, a Principal Component Analysis (PCA, non-standardised) was performed on the relative abundances of the dominant species. We considered only dominant taxa (i.e. taxa representing at least 5% of the total assemblage in at least one sample) to decrease the signal/noise ratio.

Data on living assemblages (0–1 cm depth interval) for our station 1 and additional station 2(located at 23 m depth, impacted by a shorter seasonal oxygen depletion event; Figure 2) were taken from Richirt et al. (2020) and were computed as the sum of all seasonal samples of 2012. These samples were considered as supplementary individuals in the analysis, meaning that they did not contribute to the construction of the principal component axes. The data pre-treatment and multivariate analysis procedure are summarised graphically in Figure 4.



Figure 4. A: Data pre-treatment procedure performed from the initial data matrix, with data
 matrix lines representing samples (depth) and data matrix columns representing taxa. B:
 Principal Component Analysis (PCA) procedure.

## 261 **3.3. AGE MODEL**

For another 90 cm long core, sampled in May 2011 at station 1, an age model was developed using <sup>210</sup>Pb dating (Donders et al., 2012). This age model, which was not very well delimited, suggested an average sedimentation rate of 2.37 cm per year, meaning that the 90 cm long core covered about 37 years (1974 to 2011). This order of magnitude confirms our earlier hypothesis that the very abrupt shift in sediment composition at 82.5 cm core depth corresponds to the closure of the basin in 1971.

268 For a third core sampled at this site during the same cruise as the sediment core used in this 269 study (in May 2012, Sulu-Gambari et al., 2017), molybdenum (Mo) concentrations were 270 measured in the solid phase. Molybdenum is sequestered when sulphide is present in the 271 porewater near the sediment-water interface (Crusius et al., 1996). Therefore, the successive 272 Mo maxima in the sediment record should reflect the yearly seasonal hypoxia/anoxia coupled 273 to the presence of sulphide (Seitaj et al., 2015), and can therefore be used to refine the age 274 model based on <sup>210</sup>Pb levels. This principle was used earlier for a core from another site in Lake 275 Grevelingen with an exceptionally high sediment accumulation rate (~ 13 cm y<sup>-1</sup>), in the 276 Scharendijke Basin, slightly further west of the Den Osse Basin (Egger et al., 2016).





278Figure 5.Molybdenum concentrations (ppm, detection limit = 0.06 ppm) as a function of core279depth (cm), modified from Sulu-Gambari et al. (2017). All values were LOG10(x+1)280transformed to better identify peaks. Each peak is highlighted with a red arrow and is assumed281to represent a yearly summer hypoxia/anoxia event. Some additional yearly hypoxia/anoxia282events have been added tentatively in case of broad peaks with shoulders (e.g., between 31 and28333 cm depth), or broad intervals without clear peaks (e.g., between 50 and 52 cm depth). Dotted284lines represent the two constraints of the age model: the first observed Mo peak corresponding

to 1972 (first anoxic/sulphidic event assumed the year following the closure of the former
 estuary in 1971) and the surface layer corresponding to the sampling year in 2012.

287 In Figure 5, a very regular, apparently cyclic alternation of maximal and minimal Mo 288 concentrations can be observed, where Mo maxima should correspond to late summer/early 289 autumn anoxia. The first peak, at ~ 83 cm depth, which occurred after the closure of the estuary 290 seaward in 1971, is assumed to represent a first anoxic/sulphidic event in summer/autumn 1972. Between this first peak at ~ 83 cm and the last peak, representing summer/autumn 2012, we 291 292 tentatively placed 39 more annual peaks. In some years, seasonal anoxia may not have 293 developed, or may have been very short, so that no Mo peaks developed. Another possible 294 explanation for missing Mo peaks could be that in years with somewhat lower sedimentation 295 rates, two seasonal Mo peaks may have merged. Consequently, our age model may be two to 296 three years off.

297 A further potential issue is that Mo concentrations were not measured in the same core 298 as the one used for the foraminiferal analysis. If we assume that the first Mo peak observed at 299  $\sim 83$  cm is characteristic of the first anoxic/sulphidic event that occurred in 1972, the year 300 following the transition from an estuary to a lake, the closure of the estuary in 1971 should have 301 taken place directly before, corresponding to a core depth of ~ 84 cm. For the same core, based on a sedimentation rate of 2 cm  $y^{-1}$  proposed in Malkin et al. (2014), the transition from an 302 estuary to a lake was estimated to have occurred at 82 cm depth by Sulu-Gambari et al. (2017; 303 304 2018). The same transition was found at 82.5 cm in the core used for foraminiferal analysis. 305 Consequently, when applying our Mo-based age model on the foraminiferal record, there is an 306 additional uncertainty of at most 1-2 cm, which could be the result of small differences in 307 compaction between the cores. In conclusion, we consider that our age model is reliable with a 308 precision of  $\pm 3$  years, which is sufficient for the aims of our study.

The average sediment accumulation rate calculated using the putative age model based on the Mo peaks is 2.06 cm y<sup>-1</sup>. This is consistent with the average sediment accumulation rate of 2.37 cm y<sup>-1</sup> obtained with the <sup>210</sup>Pb method by Donders et al. (2012).

312

#### 2

3.4.

#### CALCIUM CARBONATE SATURATION STATE

We used the CO2SYS software (Lewis and Wallace, 1998) to determine the variation of the calcium carbonate saturation state in the sediment at our station throughout the year. Monthly temperature and salinity data of 2012 are from Hagens et al. (2015) and were measured in the water column at 32 m depth. Monthly values for alkalinity and PO4<sup>3-</sup> concentrations in 317 the pore waters are from Sulu-Gambari et al. (2016a) and were measured in 0.5 cm thick 318 sediment slices. The values of pH were obtained by profiling the first 2.5 cm of the sediment 319 with a microsensor in three replicate cores (data from Seitaj et al., 2015). To compute the 320 calcium carbonate saturation state, using alkalinity and PO<sub>4</sub><sup>3-</sup> concentrations, pH values had to be averaged for 0.5 cm layers. Following Orr et al. (2015), we used the set of constant  $K_1$  and 321 322 K<sub>2</sub> from Lueker et al. (2000), the formula of Dickson (1990) for KSO<sub>4</sub> and the total boron-to-323 salinity ratio from Uppström (1974) for computation. The results of the calcium carbonate 324 saturation state calculations will be presented in the discussion section because we will only 325 use it to corroborate a hypothesis about the role of cable bacteria in the putative dissolution of 326 foraminiferal shells.

## 327 **4. RESULTS**

### 328 **4.1. FORAMINIFERAL COMMUNITIES**

The black curve in Figure 6 represents the number of individuals per sample (Suppl. Table 1), whereas the red curve represents the number of individuals after merging of samples when we were below our target value of 50 individuals. However, even after this merging procedure, all samples in the top 17 cm of the core were still very poor in foraminifera, with less than 50 individuals (Figure 6 and Table 1).



334

Figure 6. Total number of individuals in non-merged samples (black curve) and after merging
 samples to a maximum thickness of 3 cm (red curve). The dashed line represents the target
 value of 50 individuals per analysed sample.

Eleven of the 23 observed taxa were considered as dominant ( $\geq 5$  % in at least one sample, Suppl. Table 1). Together, these 11 dominant taxa represent more than 90 % of the total assemblage in almost all samples (Table 1).

Total foraminiferal density was relatively high between 90 and 84.5 cm depth, ranging from 1033 to 1810 individuals per sample (of 14.1 cm<sup>3</sup>; Figure 6 and Table 1). At 82.5 cm depth, the density abruptly decreased by one order of magnitude. Total densities between 82.5 and 17 cm depth oscillated between 15 (57–57.5 cm depth) and 950 individuals (54.5–55 cm

depth) per sample, with an average of about 175 ind. per sample.

Finally, in the top 17.5 cm of the core, total densities were very low, between 1 and 27 individuals per sample (Figure 6 and Table 1).

348 **<u>Table 1.</u>** Census data for foraminifera >125  $\mu$ m in the final reconstituted samples. Relative 349 abundances of the dominant species of foraminifera ( $\geq 5$  % of the total assemblage in at least 350 one sample), total percentage of dominant taxa and total number of individuals in the 351 recomposed samples (consisting of a single sample of 14.1 cm<sup>3</sup>, or 2 or 3 merged samples). We 352 also included *A. veneta* (not considered as a dominant species), because we discuss this species 353 later in the manuscript. For raw data of all individual samples, see Suppl. Table 1.

Interval depth (cm)	Ammonia aberdoveyensis	Ammonia confertitesta	Ammonia veneta	Elphidium magellanicum	Elphidium margaritaceum	Elphidium oceanense	Elphidium selseyense	Elphidium williamsoni	Haynesina germanica	Haynesina depressula	Quinqueloculina spp.	Stainforthia sp.	Total dominant taxa (%)	Total number of individuals	Number of merged samples
2–2.5	7.4	0.0	3.7	11.1	14.8	0.0	18.5	11.1	25.9	3.7	0.0	0.0	96.3	27	1
4.5–6	6.3	0.0	0.0	18.8	6.3	0.0	18.8	12.5	0.0	0.0	12.5	0.0	75.0	16	3
6.5–8	0.0	15.0	0.0	15.0	10.0	0.0	30.0	20.0	0.0	0.0	5.0	0.0	95.0	20	3
9–10.5	0.0	0.0	3.3	26.7	10.0	0.0	16.7	6.7	10.0	6.7	6.7	0.0	86.7	30	3
11.5–13	0.0	12.5	0.0	9.4	25.0	0.0	46.9	3.1	0.0	3.1	0.0	0.0	100.0	32	3
17–17.5	0.0	0.0	0.0	10.9	21.8	0.0	40.0	0.0	3.6	1.8	5.5	5.5	83.6	55	1
19.5–20	2.8	8.3	0.0	5.6	29.2	1.4	33.3	0.0	0.0	8.3	1.4	4.2	90.3	72	1
22–22.5	1.1	1.1	2.1	0.0	11.7	0.0	74.5	2.1	3.2	1.1	0.0	0.0	96.8	94	1
27–27.5	4.1	6.8	2.7	8.1	12.2	2.7	47.3	5.4	5.4	0.0	0.0	0.0	94.6	74	1
32–35	11.0	15.1	0.0	0.0	9.6	2.7	41.1	6.8	8.2	1.4	0.0	0.0	95.9	73	2
37–37.5	0.0	0.0	0.6	0.0	6.3	0.0	87.3	3.2	0.6	1.3	0.0	0.0	99.4	158	1
42-42.5	7.1	0.9	0.9	3.5	21.2	0.9	40.7	4.4	8.8	7.1	0.0	0.0	95.6	113	1
44.5–45	6.1	2.3	3.4	6.8	19.8	0.0	38.0	3.4	3.8	7.2	1.1	0.0	92.0	263	1
47–47.5	1.4	3.8	0.5	2.4	6.8	0.0	77.0	2.4	1.4	1.6	1.6	0.0	98.9	370	1
49.5–50	2.8	3.3	0.6	3.3	20.0	1.7	43.3	2.2	11.1	5.6	0.0	0.0	93.9	180	1
52–52.5	7.0	0.8	2.3	1.6	20.3	1.6	35.9	5.5	7.0	10.9	0.8	0.0	93.8	128	1
54.5–55	6.1	1.1	0.6	4.0	14.7	0.6	43.6	5.1	5.9	10.5	1.7	0.0	93.9	950	1
57-60	1.4	0.0	2.8	14.1	9.9	0.0	50.7	1.4	2.8	11.3	4.2	0.0	98.6	71	2
62-62.5	8.5	0.0	1.2	0.6	12.1	2.4	46.7	6.1	9.7	4.8	1.8	0.0	93.9	165	1
64.5–65	10.2	0.0	0.9	4.6	14.8	2.8	36.1	4.6	7.4	7.4	0.9	0.0	89.8	108	1
67–67.5	11.2	0.0	0.5	4.3	23.0	0.5	32.1	7.0	2.7	8.0	4.8	0.0	94.1	187	1
69.5–72.5	11.0	0.0	1.2	7.3	31.7	1.2	18.3	6.1	4.9	6.1	2.4	0.0	90.2	82	2
74.5–77.5	4.4	0.0	1.5	35.3	19.1	0.0	14.7	4.4	4.4	5.9	5.9	0.0	95.6	68	2
79.5–80	5.7	0.0	0.6	3.8	15.3	4.5	42.0	5.1	11.5	6.4	1.9	0.0	96.8	157	1
82-82.5	3.9	0.0	0.0	9.4	7.8	6.3	47.7	5.5	6.3	6.3	0.8	0.0	93.8	128	1
84.5–85	10.2	0.0	1.7	3.4	0.8	1.5	52.3	6.4	10.2	4.5	1.0	0.1	91.9	1033	1
87-87.5	17.3	0.0	1.5	1.9	0.8	1.4	21.0	8.8	31.3	6.2	0.3	0.0	90.6	1810	1
89.5–90	13.6	0.0	1.3	0.9	0.2	0.9	46.9	7.1	18.8	3.5	0.5	0.0	93.7	1805	1

- *Elphidium selseyense* was the dominant species in most of the samples, accounting for 15 to 87 % of the total assemblage (Figure 7, Table 1). The levels 47–47.5, 37–37.5 and 22– 22.5 cm presented peak values for this taxon (>70 %), whereas the assemblages of the top 10.5 cm were rather poor in *E. selseyense* (between 17 and 30 %).
- The relative abundance of *E. magellanicum* oscillated between 0 and 35 %. A major peak was present at 74.5–77.5 cm (35 %). This species showed relatively high percentages (9– 27 %) in the low-density samples of the topmost 17.5 cm (Figure 7, Table 1).
- *Elphidium margaritaceum* was nearly absent between 89.5–90 and 84.5–85 cm. Above
  these levels, relative abundances varied between 6 and 32 % with maxima around 70 cm depth,
  from 52–52.5 to 42–42.5 cm depth, and from 19.5–20 to 11.5–13 cm depth (Figure 7, Table 1).

The relative proportions of *H. germanica* fluctuated generally between 1 and 10 %. This species was abundant (19 and 31 %) in the first two samples (89.5–90 cm and 87–87.5 cm depth), whereas it was present in low numbers, or even absent, in the 20–4.5 cm interval. *Haynesina germanica* showed again an elevated percentage (26 %) in the most recent sample (2–2.5 cm depth, Figure 7, Table 1).







Figure 7. Relative abundances (%) of the dominant species as a function of depth (cm). We
 also included *A. veneta* (not considered as a dominant species), because we discuss this species
 later in the manuscript.

373 All other taxa only rarely contributed more than 10 % in the total assemblage (Figure 7, 374 Table 1). *Elphidium williamsoni* showed maximum values (11–20 %) in the top 10 cm of the 375 core, whereas *Elphidium oceanense* always accounted for only low percentages (0-6 %), 376 slightly decreasing upward. This taxon was absent in the top 17.5 cm. Ammonia aberdovevensis 377 showed a general decrease from about 10-17 % at the bottom of the core to a near absence at 378 about 20 cm depth. However, it showed higher percentages again (6-7 %) in the topmost two 379 samples (4.5-6 and 2-2.5 cm depth). Ammonia confertitesta was totally absent until 55 cm core 380 depth and fluctuated between absence and 15 % thereafter. Ammonia veneta was always present 381 in low percentages, never exceeding 4 %, without a clear pattern. Quinqueloculina spp. showed 382 two main presence intervals, between 80 and 44.5 cm (up to 6 %) and from 20 to 4.5 cm depth 383 (up to 13 %). Haynesina depressula (up to 11 %) tended to be more abundant in the lower 40 384 cm of the core. Finally, *Stainforthia* sp. was absent in most samples, but was present at 19.5– 385 20 and 17–17.5 cm depth (with 4 and 6 %, respectively).

386

## 4.2. MULTIVARIATE ANALYSIS

The first two axes of the non-standardised PCA based on relative abundance data account for 61 % and 17 % of the total variability of the dataset, respectively (Suppl. Table 2).

389 Table 2 presents the contributions of the dominant taxa to the first two PCs. Elphidium 390 selseyense has a strong positive contribution to PC1, whereas E. magellanicum has a slight 391 negative contribution to this axis. Elphidium margaritaceum and E. magellanicum have a 392 positive contribution to PC2, whereas H. germanica and A. aberdoveyensis load negatively on 393 this axis. Figure 8 presents a scatter plot for PC1 and PC2, in which we added the living 394 assemblages of station 1 (same station as the long core) and of the nearby, shallower station 2 395 (water depth 23 m) which is less severely affected by seasonal hypoxia/anoxia (Richirt et al., 396 2020), as supplementary individuals.

397 <u>Table 2.</u> Taxa contributions to the first two PCs from the PCA. Values in bold are superior to
 398 the average contribution (|9.09|, if all species would contribute equally).

Species / PC contribution	PC 1	PC 2
Ammonia aberdoveyensis	-0.6	-10.9
Ammonia confertitesta	0.0	1.6
Elphidium selseyense	87.5	0.4
Elphidium magellanicum	-7.3	10.4
Elphidium margaritaceum	-1.8	32.0

Elphidium williamsoni	-0.8	-1.8
Elphidium oceanense	0.0	-0.2
Haynesina germanica	-1.3	-41.8
Haynesina depressula	-0.2	0.0
Quinqueloculina spp.	-0.5	0.8
Stainforthia sp.	0.0	0.2





Figure 8. Scatter plot showing the taxa loadings and sample scores on the first two PCA axis.
 Core depth (cm) is indicated in black. Dominant taxa are represented by arrows. Supplementary
 individuals (living assemblages observed in 2012 from stations 1 and 2; Richirt et al., 2020) are
 indicated by a cross and a circle for stations 1 and 2, respectively. Blue dotted lines show the
 different, arbitrarily defined, groups of samples.

405 In Figure 8, five groups of samples have been recognised arbitrarily based on their 406 position on the PCA scatter plot and their proximity in the core:

407 Group A is constituted of samples with high negative scores on PC2, reflecting high
408 proportions of *A. aberdoveyensis* and *H. germanica*. This concerns the samples from 90 to 84.5
409 cm core depth.

Group B contains samples with strongly negative scores on PC1 and positive scores on
PC2, characterised by high relative abundances of *E. magellanicum* and *E. margaritaceum* and
relatively low percentages of *E. selseyense*. This concerns the 74.5–77.5 and 69.5–72.5 cm

samples, which are both merged samples because the number of individuals in the separatesamples was very low.

Group C is composed of samples with low absolute scores on both PCs (i.e. centred on the scatter plot, Figure 8), indicative of average assemblages. These "typical" assemblages are dominated by *E. selseyense*, have a high percentage of *E. margaritaceum*, a lower percentage of *A. aberdoveyensis* and *H. germanica* (compared to group A), and contain *A. confertitesta* from 55 cm onward. This concerns most samples from 67.5 to 27 cm depth as well as the 82– 82.5 and 79.5–80 cm samples (subgroup C'). The living community of station 2 (Den Osse Basin, 23 m water depth) plots together with this group.

422 **Group D** is composed of three samples with very high scores on PC1, reflecting 423 uncommonly high percentages of *E. selseyense*. This concerns the 22–22.5 cm, 37–37.5 cm and 424 47–47.5 cm samples. The living assemblage of station 1 is very similar, as shown by its 425 proximity on the scatter plot (Figure 8).

426 Group E groups samples intermediate between groups B and C, mostly with negative 427 scores on PC1, and positive scores on PC2. Assemblages in these samples are characterised by 428 high relative abundances of E. magellanicum and E. margaritaceum, although lower than the 429 samples of group B. Group E contains all samples between 20 and 2 cm core depth. The 2–2.5 430 cm sample is rather far away from the other samples of this group due to its high percentage of 431 H. germanica (Table 1). However, most samples in group E show a low absolute total 432 abundance, so that minor changes in the counts may lead to substantial changes in relative 433 proportions.

## 434 **5. DISCUSSION**

### 435 **5.1. TEMPORAL EVOLUTION OF THE FORAMINIFERAL COMMUNITY**

436 Here we will focus on the temporal changes in foraminiferal densities and assemblage 437 composition through time, and we will investigate whether it is possible to link the main 438 assemblage changes to the anthropogenic modifications of Lake Grevelingen during the last 50 439 years (Figure 9). As we look at 0.5 cm thick layers in a context where the average sediment accumulation rate is about 2 cm  $y^{-1}$ , individual layers may represent a single season and not a 440 441 complete year. Consequently, samples with high foraminiferal densities could be deposited 442 during spring/early summer seasons, whereas samples with low foraminiferal densities could 443 be representative of the late summer/autumn/winter seasons.

444 Additionally, we judge that vertical displacements of foraminiferal shells, either caused by 445 the vertical migration of infaunal species (when alive) or by macrofaunal bioturbation (living 446 as well as dead individuals), are strongly limited in the sedimentary environment studied here. 447 The severe anoxic and sulphidic conditions at this site (sulphide was present in the pore water 448 below 2 cm depth during the whole year in 2012, Seitaj et al., 2015) largely restrict the presence 449 of macrofauna and probably limit their bioturbation to the top 2 cm at most (Seitaj et al., 2017; 450 Hermans et al., 2020). For the same reason, living foraminifera are not expected to dwell deeper 451 than the superficial oxic layer, being limited by the sulphidic conditions immediately below.

Finally, the high sedimentation rate (about 2 cm.y<sup>-1</sup>) further decreases the possibility of mixing of sediment layers with different ages, suggesting that most samples contain contemporaneous assemblages, representing short periods of time (a single season to a year at most).

As explained before, the Principal Component Analysis allows us to recognise five groups
of samples, each representing different vertical intervals from the sediment core, and therefore,
successive stages of the evolution of the Lake Grevelingen.

459 Group A includes samples from 90 to 84.5 cm depth (estimated period ~ 1968 to 1971). 460 They contain a sandy sediment, and are situated immediately below the shift to a much finer 461 sediment at 82.5 cm. These samples represent the estuarine phase of the basin, before its 462 seaward closure in 1971. The foraminiferal assemblages are characterised by high density (between 1000 and 2000 ind. per 14.1 cm<sup>3</sup> sample), and by high percentages of typical 463 464 estuarine/mudflat species such as H. germanica and A. aberdoveyensis (e.g. Alve & Murray, 465 1994; 2001; Thibault de Chanvalon et al., 2015; Saad & Wade, 2016; 2017). In view of the 466 location of station 1, in the middle of the Den Osse Basin at 34 m depth (the main former 467 channel of the Grevelingen estuary), the sandy sediment reflects the strong hydrodynamics of 468 this site, when it was still an estuary. Currents were too strong to allow deposits of clays and 469 silts, so that the former channel floor was covered by a sandy lag deposit, with a very high 470 foraminiferal density. The strong currents undoubtedly facilitated the transport of allochthonous 471 foraminifera, such as the taxa cited above, which are typical of estuarine mudflats. However, 472 H. depressula, a species typical of marine influence (Murray 1983; Darling et al., 2016) is 473 present with substantial proportions as well, showing that foraminifera were not only 474 transported from the inner to the outer estuary, but also from the North Sea into the outer 475 estuary.

476 Subgroup C' of group C includes the next two samples, from 82.5 to 79.5 cm depth
477 (estimated period ~ 1972 to 1975). The sediment is mainly composed of clay/silt, confirming

its deposition after the seaward closure of the basin, when hydrodynamics became much weaker. The strong drop of foraminiferal densities (from higher than 1000 to less than 100 ind. per sample) shows that these samples no longer constitute a lag deposit, but rather that the foraminiferal populations are diluted by supplies of fine-grained sediment (about 2 cm y<sup>-1</sup>). The weaker hydrodynamics, together with the suppression of tides explain why the foraminiferal assemblage is much less affected by transport, and probably represents an autochthonous community.

485 The assemblages of these two samples show a strongly increased percentage of the more 486 marine species E. margaritaceum and the euryhaline species E. oceanense (Darling et al., 487 2016), contrasting with a strong decrease of the percentages of the more estuarine species H. 488 germanica and A. aberdoveyensis. This suggests that in the first years after the closure of the 489 former estuary, the basin floor was still inhabited by species typical of marine influence (high 490 salinity), whereas inner estuarine species were no longer living, or transported to the study area. 491 Salinity, measured on surface waters, was consistently high (~ 30) in the years immediately 492 following the closure of the basin (1972–1974, Bannink et al., 1984), allowing marine taxa to 493 inhabit the former main channel.

Group B, which represents the 77.5 to 69.5 cm interval (estimated period ~ 1975–1980)
contains only two merged samples (i.e. 74.5–77.5 and 69.5–72.5 cm). Their assemblage
composition is somewhat peculiar, because of the high relative proportions of *E. magellanicum*and *E. margaritaceum*, contrasting with the relatively low, but still substantial contribution of *E. selseyense*.

499 In this period, surface water salinity showed a progressive decrease, from ~ 30 in 1974 500 to a minimum of ~ 23 in 1978, due to the input of rainwater (Bannink et al., 1984). In case these 501 salinity values partly reflect conditions on the basin floor, this could suggest that E. 502 magellanicum and E. margaritaceum are more tolerant to lower salinities than E. selseyense. 503 Unfortunately, the ecology of these three species is still imperfectly known. In the literature, the European representatives of E. magellanicum are generally considered as euryhaline (20-504 505 30 in Fjords, Gustafsson & Nordberg, 1999; 2000; 10-20 in the western Baltic Sea, Schönfeld 506 et al., 2018; shallow marine at Dunkirk beach, Lévy et al., 1969). Elphidium selseyense appears 507 to be opportunistic and tolerant to a large range of salinity (Darling et al., 2016). Conversely, 508 E. margaritaceum has been reported as a more marine species which tolerates only slight 509 salinity variations (Alve & Murray, 1999; Darling et al., 2016).

510 The transition to a lacustrine system also strongly increased the average water residence 511 in the lake, up to several years, compared to a few days when Grevelingen was an estuary. This 512 strongly modified the ecosystem structure and functioning of Lake Grevelingen (Nienhuis, 513 1978; Saeijs & Stortelder, 1982). The strengthened water column stratification and the much 514 longer residence times led to a progressive increase of nutrient concentrations between 1971 515 and 1978 (Bannink et al., 1984). The opening of the Brouwersdam in late autumn 1978, 516 resulting in limited salt water inflow, induced a further strengthening of the salinity 517 stratification of the water column, which led to hypoxia/anoxia and mass mortality in the 518 deepest parts of the basin in summer 1979 (Bannink et al., 1984).

The three *Elphidium* species that dominated the assemblages during this period were apparently more resistant to these rapidly changing conditions than other taxa. Their tolerance regarding low oxygen conditions is shown by their presence in the recent assemblages of this site (Richirt et al., 2020), but also by their presence in fjords affected by seasonal anoxia (Gustafsson & Nordberg, 1999; 2000).

524 The remaining samples of Group C represent the 67.5 to 27 cm interval (estimated 525 period ~ 1980 to 2000). To remedy the increase of eutrophication phenomena observed in the 526 1970s, the basin was partly reopened by an underground sluice in 1978. This sluice was initially 527 opened only in winter. Consequently, surface water salinity almost immediately rose to ~ 30 528 and nutrient concentrations showed a major decrease (Bannink et al., 1984). However, this did 529 not solve the problem of bottom water oxygen deficiency, and in the next 20 years, an 530 intermittent succession of major later summer/early autumn hypoxic/anoxic periods was 531 observed, especially in the deepest parts of Lake Grevelingen (Figure 3). This is confirmed by 532 the results of Lindeboom & Sandee (1984), who found anoxic conditions coupled with the 533 presence of sulphide at a few mm depth in the sediment in May and June 1982, at much 534 shallower stations, with depths < 10 m. In 1999, the sluice was opened year-round, to further 535 increase water exchanges.

536 During this period (~ 1980 to 2000), the total foraminiferal density showed strong 537 variations, between ~ 70 and 950 individuals per sample. This alternance of high and lower 538 foraminiferal density could be due to differences between successive years regarding bottom 539 water oxygen content (Figure 3; Seitaj et al., 2015; 2017), more specifically due to inter-annual variability in the intensity and duration of the summer hypoxia/anoxia (with or without sulphide 540 541 in the topmost sediment). Richirt et al. (2020) showed that in case of prolonged summer anoxia 542 combined with the presence of sulphides in the topmost sediment, the foraminiferal community 543 almost completely disappears, and the site is recolonised the following winter, leading to low 544 standing stocks the next spring. The strong Mo peak at ~ 32 cm (estimated age ~ 1998, Figure 545 9), could indicate particularly severe anoxic and sulphidic conditions in the sediment (anoxia with presence of sulphides). This could explain why we observe a very low foraminiferal density (16 ind. per sample) at ~ 35 cm, immediately after this period. This apparent depth shift of ~ 3 cm between these two events could be explained by the fact that Mo concentrations and foraminiferal densities were measured on two different cores (see Materials & Methods section). Conversely, layers with high foraminiferal densities (such as sample at 54.5–55 cm, estimated age ~ 1986) could correspond to periods following weaker and/or shorter anoxic events, eventually without sulphides in the foraminiferal habitat.

553 All samples of group C are strongly dominated by E. selseyense, with E. margaritaceum 554 being the second most dominant species. This seems to reflect increased salinity (more marine 555 influence) and development of seasonal hypoxia favouring opportunistic species such as E. 556 selsevense. The arrival of A. confertitesta, at about 55 cm core depth (~ 1986) corresponds to a 557 progressive decrease of H. depressula, Quinqueloculina spp. and A. aberdoveyensis. It is 558 interesting to note that the assemblage of group C is comparable to the composition of the living 559 assemblage at the shallower station 2 (depth 23 m), as shown by the PCA scatter plot (Figure 560 8), suggesting that the environmental conditions may have been comparable, with a rather short 561 anoxic event in summer (maximum of 1 month, Richirt et al. 2020).



562

563 **Figure 9.** Timeline of the human-induced modifications following the Delta Plan at Lake 564 Grevelingen. The sediment core used to investigate dead foraminiferal assemblage density 565 record (black curve, excluding the top layer 0.5 cm showing living assemblage in grey) and the 566 Mo record (red curve). Both long cores were sampled in May 2012. Blue letters emphasise the 567 different groups of layers highlighted in the PCA scatter plot.

568 Group D contains only three samples (47–47.5, 37–37.5 and 22–22.5 cm core depth, 569 deposited in ~ 1991, ~ 1996 and ~ 2003, respectively). The foraminiferal densities in these 570 samples are comparable to densities from group C and range from 94 to 370 ind. per sample. 571 The composition of these samples stands out by exceptionally high proportions of *E. selseyense* 572 (> ~75%). This is very similar to the composition of the living assemblage in 2012 at the same 573 station (Richirt et al., 2020), as shown by their proximity on the PCA scatter plot. This suggests 574 that the three samples of group D reflect similar conditions, which could be characterised by 575 rather severe anoxic conditions with the presence of sulphides in the topmost sediment, as 576 recorded in 2012 (Seitaj et al., 2015). It appears that E. selsevense could be particularly tolerant 577 to such low oxygen conditions; this is confirmed by its strong dominance in the living 578 community sampled at this site in 2012 (Richirt et al., 2020).

579 Finally, Group E comprises all samples in the top 20 cm of the core (estimated period 580  $\sim$  2003 to 2011). This group of samples has been deposited in the years after the decision had 581 been taken to open the seaward sluice year-round (in 1999), leading to a further decrease in 582 water residence times. The total foraminiferal densities are extremely low in this interval (rarely 583 exceeding 15 ind. per sample). Because of the low foraminiferal densities, several samples had 584 to be merged. In spite of the merging procedure, relative proportions are still based on low 585 numbers (from 16 to 72 specimens) and should be considered with caution. Even when taking 586 into account the high sedimentation accumulation rate at this location, the unexpected very low 587 densities strongly contrast with the elevated densities of living communities, which attained a 588 maximum of 634 individuals per 14.1 cm<sup>3</sup> sample in May 2012 when the long core was sampled 589 (recalculated from Richirt et al., 2020; Figure 9).

The foraminiferal assemblages of the samples from group E are relatively poor in *E. selseyense*, whereas the percentages of *E. margaritaceum*, *E. magellanicum* and *E. williamsoni* show a strong increase. *Quinqueloculina* spp. is also somewhat more frequent (Table 1). It is surprising that the taxonomic composition of the group E samples is very different from the year-averaged living assemblage from this site (Figure 8).

In view of the important density differences between the rich living populations and poor taphocoenoses, with a very different species composition, we suspect that the dead assemblages have suffered from important taphonomic losses during early diagenesis, specifically due to carbonate dissolution. In fact, in the deepest parts of Lake Grevelingen, two antagonistic bacterial populations occur: cable bacteria develop mainly between January and May, whereas *Beggiatoceae* mats occupy the sea floor from September to December (Seitaj et al., 2015). The reasons why these two filamentous S-oxidising bacteria show such an

- antagonistic seasonal succession is not well understood yet (Seitaj et al. 2015), but it is clear
  that they strongly affect the geochemistry of the topmost sediment layer.
- 604 605

# 5.2. CABLE BACTERIA ACTIVITY RESPONSIBLE FOR FORAMINIFERAL TEST DISSOLUTION

606 Cable bacteria activity promotes carbonate precipitation in the oxic zone (maximum oxygen 607 penetration depth about 2 mm at this site, even in winter, see Seitaj et al., 2015), and carbonate 608 dissolution immediately below, where oxygen is absent (Risgaard-Petersen et al., 2012; 609 Meysman et al., 2015). Consequently, the carbonate saturation state ( $\Omega$ Ca) is low in the first 610 centimetres of the sediment ( $\Omega Ca < 1$  at least down to 2 cm, see Table 3), which could result in 611 the partial or complete dissolution of calcareous foraminiferal tests. In fact, for 2012, calculated  $\Omega$ Ca are < 1 from January (excluding in the top 0.5 cm) to April, when cable bacteria were 612 active, indicating strongly increased calcium carbonate (CaCO<sub>3</sub>) dissolution (Table 3). This is 613 corroborated by a release of  $Ca^{2+}$  measured in the first centimetres in pore waters at the same 614 615 station in January and March (Sulu-Gambari et al. 2016a).

616 <u>**Table 3.**</u> Values of calcium carbonate saturation state at station 1 at different depths in the 617 sediment for each month of 2012. Red:  $\Omega$ Ca < 1, indicating increased carbonate dissolution. 618 Blue:  $\Omega$ Ca > 1, indicating no carbonate dissolution. NA: data not available.

Depth (cm)	January	February	March	April	May	June	July	August	September	October	November	December
0.25	2.30	NA	NA	0.91	NA	1.93	3.08	NA	1.86	NA	2.81	3.06
0.75	0.54	0.90	0.68	0.70	NA	2.39	2.67	NA	2.30	7.46	9.62	14.68
1.25	0.29	0.41	NA	0.55	NA	2.41	2.87	NA	2.90	NA	14.76	22.66
1.75	0.20	0.27	0.24	0.51	NA	2.13	2.87	NA	NA	4.60	NA	21.11
2.25	NA	0.22	0.23	0.67	NA	1.99	2.75	NA	3.95	5.03	5.90	13.49
2.75	NA	0.21	NA	0.61	NA	2.40	2.78	NA	NA	NA	NA	7.75
3.25	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	4.29	6.98
3.75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	4.68	NA

Living foraminifera have mechanisms to deal with a lower carbonate saturation state (e.g. active control of their internal and external pH during the calcification process, intracellular storage of  $Ca^{2+}$  ions, Toyofuku et al., 2008; de Nooijer et al., 2009; Toyofuku et al., 2017). When dwelling in environments with a lower carbonate saturation state, living specimens only rarely show serious signs of test dissolution (e.g. Charrieau et al., 2018). Hence, most dissolution is thought to occur after the death of the foraminifera. Furthermore, we note that the dense living populations, which are not much affected by carbonate dissolution, predominantly live in the thin oxic zone (only a few mm), which is not affected by the strongly decreased  $\Omega$ Ca (Seitaj et al. 2015). Only when the tests of dead foraminifera passively cross this low  $\Omega$ Ca zone (due to sediment deposition at the sediment water interface), assemblages are probably affected by important losses due to carbonate dissolution. The very low foraminiferal densities in the top 20 cm of the core suggest that this situation has existed since about 2003, some years after the intensification of water exchanges. In the cartoon presented in Figure 10, the seasonal succession of events is illustrated.



#### 633

**Figure 10.** Conceptual scheme showing two consecutive years to explain the low density of dead foraminifera (in grey) in the first 20 cm of the core. The seasonal succession of oxic (blue), suboxic (orange) and sulphidic (black) zones and presence of cable bacteria (red dotted curves) are from Seitaj et al. (2015). Presence and densities (size of the signs) of living foraminifera (in green) are from Richirt et al. (2020). Squares, circles and triangles represent the yearly successive foraminifera generations.

If our hypothesis that the low density of foraminifera in the top 20 cm is due to dissolution resulting from cable bacteria activity is correct, this implies that cable bacteria activity in the Den Osse Basin started, or substantially increased between 2000 and 2005. This fits remarkably well with the year-round opening of the Brouwersdam sluice in 1999.

The increased inflow of saline, warm and well oxygenated water induced by the opening of the Brouwersdam sluice in 1999 should have improved the oxygenation and led to a reduction of the duration of anoxic events in the bottom waters compared to the situation before 1999 (Figure 3; Wetsteijn, 2011; Sulu-Gambari et al., 2017). However, the increased concentration and large peaks in the Mo record in the last ~ 30 cm of the core suggest the presence of increased sulphide concentrations in the pore waters during the same period. Sulu-Gambari et al. (2017) explained this contradiction (i.e. the better oxygenation of bottom waters but development of more sulphidic conditions in sediment) by the introduction of large amounts of phytoplankton (mainly *Phaeocystis globosa*) into Lake Grevelingen from the North Sea (Hagens et al. 2015) since ~ 2000. The intensified remineralisation of organic matter induced by this extra input of dead algal material enhanced the sulphate reduction rate resulting in an increase of the porewater sulphide concentration. This process may have contributed to the exceptionally long hypoxia/anoxia in 2011 and 2012, compared to the period 1999–2010 (Sulu-Gambari et al. 2017).

The suggested increased cable bacteria activity resulting from the increased water flow in 1999 is corroborated by the fact that the presence of an oxygen-sulphide interface is a prerequisite for the development of S-oxidising bacteria, such as cable bacteria and *Beggiatocea* (Jørgensen, 1982).

## 662 **5.3.** TEMPORAL SUCCESSION OF AMMONIA SPECIES

663 The historical record for Lake Grevelingen presented here shows the arrival of A. 664 confertitesta at 54.5–55 cm core depth (estimated year ~ 1986), and its subsequent progressive 665 replacement of A. aberdovevensis (Figure 11). It has already been suggested that A. confertitesta 666 could be an exotic and/or invasive species originating from east Asia and introduced in the 667 eastern Atlantic Ocean by anthropogenic activities (Pawlowski & Holzmann, 2008; Schweizer 668 et al, 2011; Bird et al., 2020). Concerning the other two representatives of the Ammonia genus 669 observed in our sediment core, A. veneta is considered cosmopolitan (Holzmann & Pawlowski, 670 2000; Hayward et al., 2021) whereas A. aberdoveyensis seems to be restricted to the north 671 Atlantic (Hayward et al., 2004; Hayward et al., 2021). The latter two species are both assumed 672 autochthonous in Europe and could at present be progressively replaced by the seemingly exotic 673 A. confertitesta (Pawlowski & Holzmann, 2008; Schweizer et al. 2011).

674 This species replacement suggests that A. confertitesta is more resistant than A. 675 aberdoveyensis to the seasonal hypoxia/anoxia found in Lake Grevelingen after the closure in 1971. This hypothesis is corroborated by the much higher test porosity measured in A. 676 677 confertitesta compared to A. aberdoveyensis (22.45  $\pm$  3.85 % versus 12.18  $\pm$  3.8 %, 678 respectively, Richirt et al. 2019a), suggesting that this phylotype could resist oxygen deficiency 679 by increasing its gas exchanges with the surrounding environment. The fact that A. confertitesta 680 is rather poorly represented (about 5 % of the total assemblage) in the living assemblages in 681 2012 was explained by a phase offset between: 1) the time of repopulation of the site in winter

- 682 2011–2012 after its decimation due to anoxic and sulphidic conditions in summer 2011, and 2)
  683 the availability of its preferred food sources, in spring 2012 (Richirt et al. 2020).
- Alternatively, the rather poor density of *A. confertitesta* could also mean that this taxon tolerates hypoxia and short phases of anoxia but is not able to withstand prolonged periods of oxygen depletion, together with a presence of sulphides in its microhabitat. This could explain the much higher percentage of this taxon in the living assemblages of station 2 (shallower station in Den Osse Basin), where the duration of seasonal hypoxia/anoxia is shorter than at
- 689 station 1 (Richirt et al. 2020).



690

Figure 11. Cumulative relative abundances (in % of the total assemblage) of the three different
 *Ammonia* species *A. veneta* (red), *A. aberdoveyensis* (green) and *A. confertitesta* (blue) as a
 function of core depth (cm). Please keep in mind that total densities are relatively low in the top
 15 cm of the core (~15-30 ind. per sample), meaning that small variations in absolute
 abundances induce large variations in relative abundances.

696 Finally, A. confertitesta is virtually the only Ammonia species found in recent living 697 communities (~ 5 % of the 1044 individuals, total foraminiferal assemblage, Richirt et al., 698 2020). In view of the strong anoxic events with co-occurring sulphidic conditions in 2011 and 699 2012 at this site (Seitaj et al. 2015; Richirt et al. 2020), this underlines the higher resistance of 700 A. confertitesta, compared to A. veneta and A. aberdoveyensis, to such adverse conditions.

701 Today, A. confertitesta is found in all (closed or open) branches of the Rhine-Meuse-Scheldt 702 estuary (Jorissen, unpublished data). We propose two hypotheses for the arrival of A. 703 confertitesta in Lake Grevelingen (i.e. Den Osse Basin), around 1986:

- 704 705
- 1) Ammonia confertitesta is indeed an exotic species, and its arrival in Lake Grevelingen was synchronous with its arrival on the eastern Atlantic coast.
- 706 2) Ammonia confertitesta was already present in the North Sea, either as an autochthonous 707 species, or after a first anthropogenic introduction from eastern Asia, and its arrival in 708 Lake Grevelingen reflects a secondary, local migration event.

709 We note that A. confertitesta was not observed in the assemblages of the lower part of the 710 core (90–85 cm) when Grevelingen was open to the North Sea. This strongly suggests that this 711 species reached this part of the North Sea between the closure of the lake in 1971 and its arrival 712 in the basin circa 1986. This would rather argue in favour of its exotic nature, as suggested by 713 Pawlowski & Holzmann (2008), who hypothesised an introduction of A. confertitesta in the North Sea in the late 19<sup>th</sup> century by means of ship ballast water. Massive imports of Japanese 714 715 oysters, from 1964 to about 1980 could also represent a possible vector for A. confertitesta, as 716 this has been reported for a large number of different metazoa (e.g. Polychaeta, Amphipoda, 717 Tunicata, Cirripedia, etc., Wolff & Reise, 2002).

718 Unfortunately, this study does not allow us to definitely settle whether A. confertitesta is an 719 exotic and/or invasive species in Europe. Even today, A. confertitesta is not present in all 720 estuaries of the mid latitude East Atlantic coast. For instance, it is absent in the French Auray 721 estuary (Fouet et al., in prep.). Therefore, the absence of A. confertitesta in the Grevelingen 722 estuary before its closure in 1971 could also be due to the fact that this species was not present 723 in this particular estuary. Finally, the transformation of the Grevelingen estuary to a marine lake 724 with occurrence of seasonal anoxia may have facilitated the colonisation of the Den Osse Basin 725 by A. confertitesta, putatively more resistant to hypoxia/anoxia than its congeneric species.

## 726 **6. CONCLUSION**

727 Our investigation of the sediment record of one of the deepest basins in Lake Grevelingen 728 highlights the important changes undergone by the foraminiferal community over the last 50 729 years. These changes reflect the major anthropogenic modifications of Lake Grevelingen, 730 which was artificially turned from an estuary into a marine lake during this period. The Mo 731 concentrations in the sediment record mark the yearly late summer-early autumn anoxic events 732 with co-occurring presence of free sulphides in the sediment pore water. This record provides 733 a very useful tool to detail the more conventional age model based on <sup>210</sup>Pb, allowing us to 734 obtain a precision of about  $\pm 3$  years.

735 The seaward closure of the basin in 1971 induced an important change in foraminiferal 736 communities, from high proportions of *H. germanica* and *A. aberdoveyensis*, which are typical 737 estuarine mudflat species, to assemblages dominated by *E. margaritaceum*, typical for marine 738 influence, E. oceanense, considered as an opportunistic species, and E. magellanicum, which 739 seems to be tolerant to oxygen depletion. This taxonomic change could be explained by the 740 disappearance of tides, much weaker hydrodynamics, increased eutrophication and increased 741 salinity immediately after the closure of the basin. The opening of the seaward sluice (in the 742 Brouwersdam) in autumn 1978, to counterbalance the eutrophication, increased the water 743 renewal, but did not solve the problem of seasonally occurring hypoxia/anoxia. At first, the 744 sluice was opened only in winter and E. selseyense strongly dominated the foraminiferal 745 assemblages, suggesting that this species benefited of more marine influence and development 746 of seasonal hypoxia. Around 1986, the emergence of A. confertitesta was observed coupled 747 with a progressive diminution of the proportion of A. aberdoveyensis, H. depressula and 748 Quinqueloculina spp. in the assemblages. After the sluice was opened almost year-round in 749 1999 and a further reduction of the water residence time by doubling the exchange with the 750 North Sea, the foraminiferal assemblages became very poor, so that the assemblage 751 composition should be interpreted very carefully. The extreme scarcity of foraminiferal tests in 752 the top 15 cm of the core, which strongly contrasts with the high abundance of living 753 foraminiferal communities, is ascribed to post-mortem dissolution, resulting from the strongly 754 diminished carbonate saturation state due to cable bacteria activity in the anoxic part of the 755 sediment.

Our results indicate that *A. confertitesta* arrived in Lake Grevelingen around 1986 and has
progressively supplanted other *Ammonia* species (*A. veneta* and especially *A. aberdoveyensis*)
in the record. We hypothesise that this progressive takeover is the result of a greater tolerance

to seasonal hypoxia/anoxia. However, our results do not definitely confirm the exotic and/or invasive nature of *A. confertitesta*. The absence of this taxon in the lower part of the core could also indicate its absence along the Dutch coast in 1971 or that the human-induced transformation of Lake Grevelingen created more favourable environmental conditions for its settlement.

#### 764 ACKNOWLEDGMENTS

765 We acknowledge the support of P. van Rijswijk, M. Hagens, A. Tramper, and the crew

766 of the R/V Luctor (P. Coomans and M. Kristalijn) during the sampling campaigns. We want to

thank Filip Meysman for providing datasets for pH values for carbonate saturation state

768 calculation. This study benefited from funding of Rijkswaterstaat and of the CNRS-INSU

769 programme EC2CO-DRIL (project AMTEP).

#### 770 **References**

77Alve, E., Bernhard, J.M., 1995. Vertical migratory response of benthic foraminifera to controlled
oxygen concentrations in an experimental mesocosm. Marine Ecology Progress Series 116,
137–151.

77Alve, E., Murray, J.W., 2001. Temporal variability in vertical distributions of live (stained) intertidal
775 foraminifera, southern England. Journal of Foraminiferal Research 31, 12–24.
776 https://doi.org/10.2113/0310012

77% Ive, E., Murray, J.W., 1999. Marginal marine environments of the Skagerrak and Kattegat: a
baseline study of living (stained) benthic foraminiferal ecology. Palaeogeography,
Palaeoclimatology, Palaeoecology 146, 171–193. https://doi.org/10.1016/S00310182(98)00131-X

78Alve, E., Murray, J.W., 1994. Ecology and taphonomy of benthic foraminifera in a temperate
782 mesotidal inlet. Journal of Foraminiferal Research 24, 18–27.
783 https://doi.org/10.2113/gsjfr.24.1.18

78**B**annink, B.A., Van der Meulen, J.H.M., Nienhuis, P.H., 1984. Lake grevelingen: From an estuary
to a saline lake. An introduction. Netherlands Journal of Sea Research 18, 179–190.
https://doi.org/10.1016/0077-7579(84)90001-2

78Bernhard, J.M., 1993. Experimental and field evidence of Antarctic foraminiferal tolerance to anoxia
and hydrogen sulfide. Marine Micropaleontology 20, 203–213. https://doi.org/10.1016/03778398(93)90033-T

79Bernhard, J.M., Alve, E., 1996. Survival, ATP pool, and ultrastructural characterization of benthic
foraminifera from Drammensfjord (Norway): response to anoxia. Marine Micropaleontology
28, 5–17. https://doi.org/10.1016/0377-8398(95)00036-4

79Bird, C., Schweizer, M., Roberts, A., Austin, W.E.N., Knudsen, K.L., Evans, K.M., Filipsson, H.L.,
Sayer, M.D.J., Geslin, E., Darling, K.F., 2020. The genetic diversity, morphology,
biogeography, and taxonomic designations of Ammonia (Foraminifera) in the Northeast

796	Atlantic.	Marine	Micropaleontology	155,	101726.
797	https://doi.org	g/10.1016/j.marmicro	0.2019.02.001		

798 ouchet, V.M.P., Debenay, J.-P., Sauriau, P.-G., Radford-Knoery, J., Soletchnik, P., 2007. Effects

of short-term environmental disturbances on living benthic foraminifera during the Pacific solution over summer mortality in the Marennes-Oléron Bay (France). Marine Environmental

801 Research 64, 358–383. https://doi.org/10.1016/j.marenvres.2007.02.007

802 harrieau, L.M., Filipsson, H.L., Nagai, Y., Kawada, S., Ljung, K., Kritzberg, E., Toyofuku, T.,
2018. Decalcification and survival of benthic foraminifera under the combined impacts of
varying pH and salinity. Marine Environmental Research 138, 36–45.
https://doi.org/10.1016/j.marenvres.2018.03.015

806 rusius, J., Calvert, S., Pedersen, T., Sage, D., 1996. Rhenium and molybdenum enrichments in sediments as indicators of oxic, suboxic and sulfidic conditions of deposition. Earth and Planetary Science Letters 145, 65–78. https://doi.org/10.1016/S0012-821X(96)00204-X

80Darling, K.F., Schweizer, M., Knudsen, K.L., Evans, K.M., Bird, C., Roberts, A., Filipsson, H.L.,
Kim, J.-H., Gudmundsson, G., Wade, C.M., Sayer, M.D.J., Austin, W.E.N., 2016. The genetic
diversity, phylogeography and morphology of Elphidiidae (Foraminifera) in the Northeast
Atlantic. Marine Micropaleontology 129, 1–23.

813 https://doi.org/10.1016/j.marmicro.2016.09.001

81**4**e Nooijer, L.J., Toyofuku, T., Kitazato, H., 2009. Foraminifera promote calcification by elevating 815 their intracellular pH. PNAS, 106, 15374–15378. https://doi.org/10.1073/pnas.0904306106

81Dickson, A.G., 1990. Standard potential of the reaction:  $AgCl(s) + \frac{1}{2}H_2(g) = Ag(s) + HCl(aq)$ , and and the standard acidity constant of the ion  $HSO_4^-$  in synthetic sea water from 273.15 to 318.15 K. The Journal of Chemical Thermodynamics 22, 113–127. https://doi.org/10.1016/0021-9614(90)90074-Z

82Donders, T.H., Guasti, E., Bunnik, F.P.M., van Aken, H., 2012. Impact van de Brouwersdam op zuurstofcondities in de Grevelingen; reconstructies uit natuurlijke sediment archieven. TNOrapport, TNO-060-UT-2011-02116. Geological Survey of the Netherlands.

82Duijnstee, I. a. P., Ernst, S.R., Zwaan, G.J. van der, 2003. Effect of anoxia on the vertical migration
824 of benthic foraminifera. Marine Ecology Progress Series 246, 85–94.
825 https://doi.org/10.3354/meps246085

82Egger, M., Lenstra, W., Jong, D., Meysman, F.J.R., Sapart, C.J., Veen, C. van der, Röckmann, T.,
Gonzalez, S., Slomp, C.P., 2016. Rapid Sediment Accumulation Results in High Methane
Effluxes from Coastal Sediments. PLOS ONE 11, e0161609.
https://doi.org/10.1371/journal.pone.0161609

83Ernst, S., Bours, R., Duijnstee, I., Zwaan, B. van der, 2005. EXPERIMENTAL EFFECTS OF AN
ORGANIC MATTER PULSE AND OXYGEN DEPLETION ON A BENTHIC
FORAMINIFERAL SHELF COMMUNITY. Journal of Foraminiferal Research 35, 177–197.
https://doi.org/10.2113/35.3.177

83**#**ouet, M., Heliot, S., Singer, D., Barras, C., and Jorissen, F. J.: How to adapt foraminiferal index to 835 estuaries? Application of the MII index, in preparation, 2021.

83Geslin, E., Barras, C., Langlet, D., Nardelli, M.P., Kim, J.-H., Bonnin, J., Metzger, E., Jorissen, F.J.,
2014. Survival, Reproduction and Calcification of Three Benthic Foraminiferal Species in
Response to Experimentally Induced Hypoxia, in: Kitazato, H., M. Bernhard, J. (Eds.),
Approaches to Study Living Foraminifera: Collection, Maintenance and Experimentation,

840 Environmental Science and Engineering. Springer Japan, Tokyo, pp. 163–193.
841 https://doi.org/10.1007/978-4-431-54388-6\_10

84Øceslin, E., Heinz, P., Jorissen, F., Hemleben, Ch., 2004. Migratory responses of deep-sea benthic 843 foraminifera to variable oxygen conditions: laboratory investigations. Marine 844 Micropaleontology 53, 227–243. https://doi.org/10.1016/j.marmicro.2004.05.010

84Siustafsson, M., Nordberg, K., 2000. Living (Stained) Benthic Foraminifera and their Response to
the Seasonal Hydrographic Cycle, Periodic Hypoxia and to Primary Production in Havstens
Fjord on the Swedish West Coast. Estuarine, Coastal and Shelf Science 51, 743–761.
https://doi.org/10.1006/ecss.2000.0695

84@ustafsson, M., Nordberg, K., 1999. Benthic foraminifera and their response to hydrography, periodic hypoxic conditions and primary production in the Koljö fjord on the Swedish west coast. Journal of Sea Research 41, 163–178. https://doi.org/10.1016/S1385-1101(99)00002-7

85P Jagens, M., Slomp, C.P., Meysman, F.J.R., Seitaj, D., Harlay, J., Borges, A.V., Middelburg, J.J.,
2015. Biogeochemical processes and buffering capacity concurrently affect acidification in a
seasonally hypoxic coastal marine basin. Biogeosciences 12, 1561–1583.
https://doi.org/10.5194/bg-12-1561-2015

856 Jayward, B.W., Holzmann, M., Grenfell, H.R., Pawlowski, J., Triggs, C.M., 2004. Morphological
distinction of molecular types in *Ammonia* – towards a taxonomic revision of the world's most
commonly misidentified foraminifera. Marine Micropaleontology 50, 237–271.
https://doi.org/10.1016/S0377-8398(03)00074-4

86 Dayward, B.W., Holzmann, M., Pawlowski, J., Parker, J.H., Kaushik, T., Toyofuku, M.S., Tsuchiya,
M., 2021. Molecular and morphological taxonomy of living *Ammonia* and related taxa
(Foraminifera) and their biogeography. Micropaleontology 67, 109–274.

86Blermans, M., Risgaard-Petersen, N., Meysman, F.J.R., Slomp, C.P., 2020. Biogeochemical impact
of cable bacteria on coastal Black Sea sediment. Biogeosciences 17, 5919–5938.
https://doi.org/10.5194/bg-17-5919-2020

866 Bolzmann, M., Pawlowski, J., 2000. Taxonomic relationships in the genus Ammonia (Foraminifera)
867 based on ribosomal DNA sequences. Journal of Micropalaeontology 19, 85–95.
868 https://doi.org/10.1144/jm.19.1.85

869ørgensen, B.B., Postgate, J.R., Postgate, J.R., Kelly, D.P., 1982. Ecology of the bacteria of the
sulphur cycle with special reference to anoxic—oxic interface environments. Philosophical
Transactions of the Royal Society of London. B, Biological Sciences 298, 543–561.
https://doi.org/10.1098/rstb.1982.0096

87Katz, M.E., Cramer, B.S., Franzese, A., Hönisch, B., Miller, K.G., Rosenthal, Y., Wright, J.D., 2010.
874 Traditional and emerging geochemical proxies in foraminifera. Journal of Foraminiferal
875 Research 40, 165–192. https://doi.org/10.2113/gsjfr.40.2.165

87Langlet, D., Baal, C., Geslin, E., Metzger, E., Zuschin, M., Riedel, B., Risgaard-Petersen, N.,
877 Stachowitsch, M., Jorissen, F.J., 2014. Foraminiferal species responses to in situ,
878 experimentally induced anoxia in the Adriatic Sea. Biogeosciences 11, 1775–1797.
879 https://doi.org/10.5194/bg-11-1775-2014

880 anglet, D., Geslin, E., Baal, C., Metzger, E., Lejzerowicz, F., Riedel, B., Zuschin, M., Pawlowski,
J., Stachowitsch, M., Jorissen, F.J., 2013. Foraminiferal survival after long-term in situ
experimentally induced anoxia. Biogeosciences 10, 7463–7480. https://doi.org/10.5194/bg-107463-2013

88ª évy, A., Mathieu, R., Momeni, I., Poignant, A., Rosset-Moulinier, M., Rouvillois, A., Ubaldo, M.,
1969. Les représentants de la famille des Elphidiidae (foraminifères) dans les sables des plages
des environs de Dunkerque. Remarques sur les espèces de Polystomella signalées par O.
Terguam Barra de Micropaléontologie 2, 02, 08

887 Terquem, Revue de Micropaléontologie 2, 92–98.

88Bewis, E., Wallace, D.W., 1998. R: Program developed for CO2 system calculations ORNL/CDIAC105. Carbon Dioxide Information Analysis Centre, Oak Ridge National Laboratory.

89D indeboom, H.J., Sandee, A.J.J., 1984. The Effect of Coastal Engineering Projects on
Microgradients and Mineralization Reactions in Sediments. Water Science and Technology 16,
892 87–94. https://doi.org/10.2166/wst.1984.0046

89Bouters, T., Berg, J.H. van den, Mulder, J.P.M., 1998. Geomorphological Changes of the
894 Oosterschelde Tidal System During and After the Implementation of the Delta Project. Journal
895 of Coastal Research 14, 1134–1151.

89Eueker, T.J., Dickson, A.G., Keeling, C.D., 2000. Ocean  $pCO_2$  calculated from dissolved inorganic carbon, alkalinity, and equations for K<sub>1</sub> and K<sub>2</sub>: validation based on laboratory measurements of CO<sub>2</sub> in gas and seawater at equilibrium. Marine Chemistry 70, 105–119. https://doi.org/10.1016/S0304-4203(00)00022-0

900 Meysman, F.J.R., Risgaard-Petersen, N., Malkin, S.Y., Nielsen, L.P., 2015. The geochemical 901 fingerprint of microbial long-distance electron transport in the seafloor. Geochimica et 902 Cosmochimica Acta 152, 122–142. https://doi.org/10.1016/j.gca.2014.12.014

90Moodley, L., Hess, C., 1992. Tolerance of Infaunal Benthic Foraminifera for Low and High Oxygen
904 Concentrations. The Biological Bulletin 183, 94–98. https://doi.org/10.2307/1542410

90Moodley, L., van der Zwaan, G.J., Rutten, G.M.W., Boom, R.C.E., Kempers, A.J., 1998. Subsurface
activity of benthic foraminifera in relation to porewater oxygen content: laboratory
experiments. Marine Micropaleontology 34, 91–106. https://doi.org/10.1016/S03778398(97)00044-3

909Muelen, J.H.M. v d, Leentvaar, J., Rooy, N.M. de, 1984. Environmental impact statement on the 910 problem of a fresh or salt lake Grevelingen. Water science and technology 16, 107–120.

91Murray, J.W., 2006. Ecology and Applications of Benthic Foraminifera. Cambridge University912 Press, 426 p.

91Murray, J.W., 1983. Population dynamics of benthic foraminifera; results from the Exe Estuary,
914 England. Journal of Foraminiferal Research 13, 1–12. https://doi.org/10.2113/gsjfr.13.1.1

91Nienhuis, P.H., 1978. Lake grevelingen: A case study of ecosystem changes in a closed estuary.
916 Hydrobiological Bulletin 12, 246–259. https://doi.org/10.1007/BF02259186

91 Nienhuis, P.H., Veld, J.C.H. in 't, 1984. Grevelingen: From an Estuary to a Saline Lake. Water 918 Science and Technology 16, 27–50. https://doi.org/10.2166/wst.1984.0043

91Ørr, J.C., Epitalon, J.-M., Gattuso, J.-P., 2015. Comparison of ten packages that compute ocean carbonate chemistry. Biogeosciences 12, 1483–1510. https://doi.org/10.5194/bg-12-1483-2015

92Pawlowski, J., Holzmann, M., 2008. Diversity and geographic distribution of benthic foraminifera:
922 a molecular perspective. Biodiversity and Conservation 17, 317–328.
923 https://doi.org/10.1007/s10531-007-9253-8

92₽etersen, J., Riedel, B., Barras, C., Pays, O., Guihéneuf, A., Mabilleau, G., Schweizer, M., Meysman,
F.J.R., Jorissen, F.J., 2016. Improved methodology for measuring pore patterns in the benthic
foraminiferal genus Ammonia. Marine Micropaleontology 128, 1–13.
https://doi.org/10.1016/j.marmicro.2016.08.001

928iña-Ochoa, E., Høgslund, S., Geslin, E., Cedhagen, T., Revsbech, N.P., Nielsen, L.P., Schweizer,
M., Jorissen, F., Rysgaard, S., Risgaard-Petersen, N., 2010. Widespread occurrence of nitrate
storage and denitrification among Foraminifera and Gromiida. PNAS 107, 1148–1153.
https://doi.org/10.1073/pnas.0908440107

93Pucci, F., Geslin, E., Barras, C., Morigi, C., Sabbatini, A., Negri, A., Jorissen, F.J., 2009. Survival
of benthic foraminifera under hypoxic conditions: Results of an experimental study using the
CellTracker Green method. Marine Pollution Bulletin, Foraminifera and Marine Pollution 59,
336–351. https://doi.org/10.1016/j.marpolbul.2009.08.015Richirt, J., Schweizer, M., Bouchet,
V.M.P., Mouret, A., Quinchard, S., Jorissen, F.J., 2019a. Morphological distinction of three
Ammonia phylotypes occurring along european coasts. Journal of Foraminiferal Research 49,
77–94.

93Richirt, J., Champmartin, S., Schweizer, M., Mouret, A., Petersen, J., Ambari, A., Jorissen, F.J.,
2019b. Scaling laws explain foraminiferal pore patterns. Scientific Reports 9, 9149.
https://doi.org/10.1038/s41598-019-45617-x

94Richirt, J., Riedel, B., Mouret, A., Schweizer, M., Langlet, D., Seitaj, D., Meysman, F.J.R., Slomp,
943 C.P., Jorissen, F.J., 2020. Foraminiferal community response to seasonal anoxia in Lake
944 Grevelingen (the Netherlands). Biogeosciences 17, 1415–1435. https://doi.org/10.5194/bg-17945 1415-2020

94**R**isgaard-Petersen, N., Revil, A., Meister, P., Nielsen, L.P., 2012. Sulfur, iron-, and calcium cycling
associated with natural electric currents running through marine sediment. Geochimica et
Cosmochimica Acta 92, 1–13. https://doi.org/10.1016/j.gca.2012.05.036

94**S**aad, S.A., Wade, C.M., 2017. Seasonal and spatial variations of saltmarsh benthic foraminiferal
ommunities from north Norfolk, England. Microbial Ecology 73, 539–555.
https://doi.org/10.1007/s00248-016-0895-5

95**S**aad, S.A., Wade, C.M., 2016. Biogeographic distribution and habitat association of *Ammonia*genetic variants around the coastline of Great Britain. Marine Micropaleontology 124, 54–62.
https://doi.org/10.1016/j.marmicro.2016.01.004

95Saeijs, H.L.F., Stortelder, P.B.M., 1982. Converting an estuary to Lake Grevelingen: Environmental
956 review of a coastal engineering project. Environmental Management 6, 377–405.
957 https://doi.org/10.1007/BF01871888

958chönfeld, J., 2018. Monitoring benthic foraminiferal dynamics at Bottsand coastal lagoon (western
959 Baltic Sea). Journal of Micropalaeontology 37, 383–393. https://doi.org/10.5194/jm-37-383960 2018

96\$chweizer, M., Polovodova, I., Nikulina, A., Schönfeld, J., 2011. Molecular identification of *Ammonia* and *Elphidium* species (Foraminifera, Rotaliida) from the Kiel Fjord (SW Baltic Sea)
with rDNA sequences. Helgoland Marine Research 65, 1–10. https://doi.org/10.1007/s10152010-0194-3

96Seitaj, D., Schauer, R., Sulu-Gambari, F., Hidalgo-Martinez, S., Malkin, S.Y., Burdorf, L.D.W.,
Slomp, C.P., Meysman, F.J.R., 2015. Cable bacteria generate a firewall against euxinia in
seasonally hypoxic basins. PNAS 112, 13278–13283.
https://doi.org/10.1073/pnas.1510152112

96Seitaj, D., Sulu-Gambari, F., Burdorf, L.D.W., Romero-Ramirez, A., Maire, O., Malkin, S.Y.,
Slomp, C.P., Meysman, F.J.R., 2017. Sedimentary oxygen dynamics in a seasonally hypoxic
basin. Limnology and Oceanography 62, 452–473. https://doi.org/10.1002/lno.10434

97**S**ulu-Gambari, F., Hagens, M., Behrends, T., Seitaj, D., Meysman, F.J.R., Middelburg, J., Slomp,
C.P., 2018. Phosphorus Cycling and Burial in Sediments of a Seasonally Hypoxic Marine
Basin. Estuaries and Coasts 41, 921–939. https://doi.org/10.1007/s12237-017-0324-0

97**S**ulu-Gambari, F., Roepert, A., Jilbert, T., Hagens, M., Meysman, F.J.R., Slomp, C.P., 2017.
Molybdenum dynamics in sediments of a seasonally-hypoxic coastal marine basin. Chemical
Geology 466, 627–640. https://doi.org/10.1016/j.chemgeo.2017.07.015

978ulu-Gambari, F., Seitaj, D., Behrends, T., Banerjee, D., Meysman, F.J.R., Slomp, C.P., 2016a.
979 Impact of cable bacteria on sedimentary iron and manganese dynamics in a seasonally-hypoxic
980 marine basin. Geochimica et Cosmochimica Acta 192, 49–69.
981 https://doi.org/10.1016/j.gca.2016.07.028

98**S**ulu-Gambari, F., Seitaj, D., Meysman, F.J.R., Schauer, R., Polerecky, L., Slomp, C.P., 2016b.
Cable Bacteria Control Iron–Phosphorus Dynamics in Sediments of a Coastal Hypoxic Basin.
Environ. Sci. Technol. 50, 1227–1233. https://doi.org/10.1021/acs.est.5b04369Thibault de
Chanvalon, A., Metzger, E., Mouret, A., Cesbron, F., Knoery, J., Rozuel, E., Launeau, P.,
Nardelli, M.P., Jorissen, F.J., Geslin, E., 2015. Two-dimensional distribution of living benthic
foraminifera in anoxic sediment layers of an estuarine mudflat (Loire estuary, France).
Biogeosciences 12, 6219–6234. https://doi.org/10.5194/bg-12-6219-2015

989 yoyofuku, T., Matsuo, M.Y., Nooijer, L.J. de, Nagai, Y., Kawada, S., Fujita, K., Reichart, G.-J.,
Nomaki, H., Tsuchiya, M., Sakaguchi, H., Kitazato, H., 2017. Proton pumping accompanies
calcification in foraminifera. Nature Communications 8, 1–6.
https://doi.org/10.1038/ncomms14145

99**T**oyofuku, T., Nooijer, L.J. de, Yamamoto, H., Kitazato, H., 2008. Real-time visualization of
calcium ion activity in shallow benthic foraminiferal cells using the fluorescent indicator Fluo3 AM. Geochemistry, Geophysics, Geosystems 9. https://doi.org/10.1029/2007GC001772

996 ppström, L.R., 1974. The boron/chlorinity ratio of deep-sea water from the Pacific Ocean. Deep
997 Sea Research and Oceanographic Abstracts 21, 161–162. https://doi.org/10.1016/0011998 7471(74)90074-6

99Wetsteijn, L.P.M.J., 2011. Grevelingenmeer: meer kwetsbaar? Een beschrijving van de ecologische
ontwikkelingen voor de periode 1999 t/m 2008-2010 in vergelijking met de periode 1990 t/m
1001 1998., RWS Waterdienst. ed. Lelystad, Netherlands.

1002 Volff, W.J., Reise, K., 2002. Oyster Imports as a Vector for the Introduction of Alien Species into
Northern and Western European Coastal Waters, in: Leppäkoski, E., Gollasch, S., Olenin, S.
(Eds.), Invasive Aquatic Species of Europe. Distribution, Impacts and Management. Springer
Netherlands, Dordrecht, pp. 193–205. https://doi.org/10.1007/978-94-015-9956-6\_21