



HAL
open science

Scattering for non-radial 3D NLS with combined nonlinearities

Jacopo Bellazzini, van Duong Dinh, Luigi Forcella

► **To cite this version:**

Jacopo Bellazzini, van Duong Dinh, Luigi Forcella. Scattering for non-radial 3D NLS with combined nonlinearities. 2023. hal-03926533

HAL Id: hal-03926533

<https://hal.science/hal-03926533>

Preprint submitted on 6 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SCATTERING FOR NON-RADIAL 3D NLS WITH COMBINED NONLINEARITIES

JACOPO BELLAZZINI, VAN DUONG DINH, AND LUIGI FORCELLA

ABSTRACT. We show scattering below the ground state energy level for a class of nonlinear 3D Schrödinger equations (NLS) with mass-energy intracritical competing nonlinearities. Specifically, the NLS has a focusing leading order nonlinearity with a defocusing perturbation. Our strategy combines interaction Morawetz estimates and a new crucial bound for the Pohozaev functional of localized functions. Furthermore, we give the rate of blow-up for symmetric solutions.

1. INTRODUCTION

Aim of the present paper is to study long time dynamics of solutions to the following nonlinear Schrödinger equation with competing nonlinearities

$$i\partial_t u + \Delta u = \lambda_1 |u|^{q-1} u + \lambda_2 |u|^{p-1} u, \quad (1.1)$$

where $u(t, x) : I \times \mathbb{R}^d \rightarrow \mathbb{C}$, $I \subseteq \mathbb{R}$, and the parameters $\lambda_1, \lambda_2 \in \mathbb{R}$. Equation (1.1) is a nonlinear Schrödinger equation which arises in many physical contexts, and we refer the reader to [15, 17, 23, 31] for motivations and further discussions on the models. Since the early works by Zhang [35], Tao, Vişan, and Zhang [34], and Miao, Xu, and Zhao [28], equations of the type (1.1) have attracted a lot of attention leading to a wide literature concerning different problems: local and global theory, scattering, blow-up, stability of standing waves, and so on; see [2, 3, 5, 7, 8, 16, 22, 24–27, 29, 32, 33] and references therein.

In this article, we are interested in the 3D physical case, and we consider exponents satisfying $\frac{7}{3} < q < p < 5$, with a positive coefficient λ_1 and a negative coefficient λ_2 . Namely, the nonlinearities are defocusing and mass supercritical, and focusing and energy subcritical, respectively. We remark that we cover the physically relevant case of focusing cubic NLS with a mass supercritical defocusing perturbation. Without loss of generality, we can normalise the coefficients, by scaling and homogeneity, as $\lambda_1 = -\lambda_2 = 1$. Therefore, we are primarily interested in the Cauchy problem below:

$$\begin{cases} i\partial_t u + \Delta u = |u|^{q-1} u - |u|^{p-1} u \\ u(0, x) = u_0 \in H^1(\mathbb{R}^3), \end{cases} \quad (1.2)$$

where $u(t, x) : I \times \mathbb{R}^3 \rightarrow \mathbb{C}$, $I \subseteq \mathbb{R}$, with $\frac{7}{3} < q < p < 5$.

At the local level, it is well-known that (1.2) admits solutions, see [6], and by denoting $I_{\max} \ni 0$ the maximal interval of existence, solutions preserve (among other quantities) mass and energy, defined by

$$M(u(t)) := \int_{\mathbb{R}^3} |u(t)|^2 dx. \quad (1.3)$$

2010 *Mathematics Subject Classification.* 35Q55; 35B40; 35B44.

Key words and phrases. Nonlinear Schrödinger equation, Scattering, Blow-up, Interaction Morawetz estimates.

and

$$E(u(t)) := \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u(t)|^2 dx + \frac{1}{q+1} \int_{\mathbb{R}^3} |u(t)|^{q+1} dx - \frac{1}{p+1} \int_{\mathbb{R}^3} |u(t)|^{p+1} dx, \quad (1.4)$$

respectively. Hence (1.3) and (1.4) are independent of time.

We introduce the following Pohozaev functional

$$G(u) = \int_{\mathbb{R}^3} |\nabla u(t)|^2 dx + \frac{3}{2} \left(\frac{q-1}{q+1} \right) \int_{\mathbb{R}^3} |u(t)|^{q+1} dx - \frac{3}{2} \left(\frac{p-1}{p+1} \right) \int_{\mathbb{R}^3} |u(t)|^{p+1} dx,$$

which will be crucial for the characterization of the dynamics of solutions to (1.2). We note in particular that solutions to (1.2) of the form $u(t, x) = e^{i\omega t} \psi(x)$, with ω being a real parameter, i.e., standing wave solutions, fulfil $G(\psi) = 0$.

Furthermore, we introduce the action functional

$$\begin{aligned} S_\omega(\phi) &= E(\phi) + \frac{\omega}{2} M(\phi) \\ &= \frac{1}{2} \|\nabla \phi\|_{L^2(\mathbb{R}^3)}^2 - \frac{1}{p+1} \|\phi\|_{L^{p+1}(\mathbb{R}^3)}^{p+1} + \frac{1}{q+1} \|\phi\|_{L^{q+1}(\mathbb{R}^3)}^{q+1} + \frac{\omega}{2} \|\phi\|_{L^2(\mathbb{R}^3)}^2 \end{aligned}$$

and the following ground state energy

$$m_\omega := \inf \{ S_\omega(\phi) : \phi \in H^1(\mathbb{R}^3) \setminus \{0\}, G(\phi) = 0 \}.$$

Note that the action functional S_ω is a conserved quantity along the time evolution of a solution to (1.2), as sum of conserved quantities.

The existence of standing waves of the form $u(t, x) = e^{i\omega t} \psi(x)$ can be proved by showing that the aforementioned ground state energy is achieved, i.e., that the infimum of the action on the constraint $G = 0$ is indeed a minimum. The fact that ground state energy level m_ω is achieved for any $\omega > 0$ is proved in [8] while the instability properties of the corresponding standing waves is a recent result by [16]. The question concerning the existence of ground states with an assigned mass has been addressed in [4]. Eventually, for $\omega > 0$, we introduce

$$\mathcal{A}_\omega^+ := \{ u \in H^1(\mathbb{R}^3) : S_\omega(u) < m_\omega, G(u) \geq 0 \}$$

and

$$\mathcal{A}_\omega^- := \{ u \in H^1(\mathbb{R}^3) : S_\omega(u) < m_\omega, G(u) < 0 \},$$

and we recall the notion of scattering: we say that a solution $u(t, x)$ to (1.2) *scatters* provided that

$$\lim_{t \rightarrow \pm\infty} \|u(t) - e^{it\Delta} u_0^\pm\|_{H^1(\mathbb{R}^3)} = 0,$$

for suitable $u_0^\pm \in H^1(\mathbb{R}^3)$, where $e^{it\Delta}$ is the linear Schrödinger propagator.

Our purpose is to show that:

- (1) If $u_0 \in \mathcal{A}_\omega^+$, then the corresponding solution exists globally in time and scatters in $H^1(\mathbb{R}^3)$ in both directions.
- (2) If $u_0 \in \mathcal{A}_\omega^-$, then if we assume u_0 is radially symmetric or cylindrically symmetric (with some additional restrictions on p and some relaxations on the hypothesis on q), then the corresponding solution blows up in finite time with an explicit blow-up rate.

Our first result concerns the energy scattering for (1.2) with data in \mathcal{A}_ω^+ .

Theorem 1.1. *Let $\frac{7}{3} < q < p < 5$ and $\omega > 0$. Let $u_0 \in \mathcal{A}_\omega^+$. Then the corresponding solution to (1.2) exists globally in time and scatters.*

Our result covers the case when the defocusing term is mass supercritical without any symmetry assumption. We follow the strategy developed in Dodson-Murphy [11] and our new key ingredient is a coercivity property for solutions belonging to \mathcal{A}_ω^+ that we develop by a careful variational study of the functional G that we believe it is of independent interest. More precisely, we use the variational properties of the functional G , that appears in the virial-like estimates, to prove an interaction Morawetz estimate. We show that taken a suitable cut-off function $\chi_R(x) = \chi(\frac{x}{R})$ such that $\chi_R = 1$ when $|x| \leq (1 - \eta)R$ with some $\eta > 0$ small, the Pohozaev functional fulfils, for any R sufficiently large,

$$G(\chi_R(\cdot - z)u^\xi(t, \cdot)) \geq \delta \|\nabla(\chi_R(\cdot - z)u^\xi(t, \cdot))\|_{L^2(\mathbb{R}^3)}^2$$

for any space shift $z \in \mathbb{R}^3$, all time $t \in \mathbb{R}$, and some $\xi = \xi(t, z, R) \in \mathbb{R}^3$, where $u^\xi(t, x)$ stands for the modulated function $e^{ix \cdot \xi} u(t, x)$. Here $\delta > 0$ is a constant independent of time and the translation vector.

The localized coercivity result, i.e., that the coercivity is true if we localize the modulated and translated solution $u(t)$ in a sufficiently large ball centered anywhere, as far as we know is a new property and we think that it can be relevant in other contexts. In particular we are able to bypass the difficulties related to the lack of scaling due to the presence of two competing nonlinearities. We shall underline that in the paper by Dodson-Murphy, the coercivity property needed to prove interaction Morawetz estimates comes from the refined Gagliardo-Nirenberg inequality (see [11, Lemma 2.1])

$$\|f\|_{L^4(\mathbb{R}^3)}^4 \leq C_{GN} \|f\|_{L^2(\mathbb{R}^3)} \|\nabla f\|_{L^2(\mathbb{R}^3)} \|\nabla f^\xi\|_{L^2(\mathbb{R}^3)}^2 \quad (1.5)$$

which holds for any $f \in H^1(\mathbb{R}^3)$ and any $\xi \in \mathbb{R}^3$, and that we are prevented to use in our setting. Indeed, due to the non-homogeneity of nonlinearities, such an inequality (1.5) is not applicable in our framework.

Let us briefly recall what is known in the literature concerning scattering for NLS with a leading focusing term. In the homogeneous case when the nonlinearity is energy subcritical and mass supercritical, the scattering below ground state energy was proved in [18] and [13] (the former in the radial case, the latter in the non-radial case, both for the 3D cubic equation) and then in [1, 14] in the general dimension case. The argument in the these cited papers follows the Kenig-Merle road map [21] with a concentration-compactness-rigidity scheme. An alternative recent strategy to prove scattering below the ground state energy is given by Dodson-Murphy in [11] for the $\dot{H}^{1/2}$ -critical focusing equation, combining interaction Morawetz estimates together with a scattering criterion. See also [10] for a generalisation to intracritical powers.

In the case we consider here, namely a focusing leading term and a defocusing perturbation we quote the early work by Miao, Xu, and Zhao [28] that considered in the radial case an energy critical focusing power with a cubic defocusing perturbation. The key argument to overcome the difficulty to treat a non-scaling invariant equation in [28], deeply inspired by [19], consists in a new radial profile decomposition with the scaling parameter and then apply it to the scattering theory. More recently, Cheng, Miao, and Zhao [8], and Luo [26] treated, still in the radial case, the focusing energy critical or energy subcritical power with a mass critical defocusing term. The radial assumption in dimension $d \leq 4$ has been recently removed by Cheng [7].

Remark 1.2. To the best of our knowledge, this is the first result of scattering for non-radial defocusing-focusing solutions in the range of nonlinearities as in the statement of

the above Theorem (see also [34, Table 1]). Cheng in [7] removed the radially assumption when working with a defocusing mass-critical nonlinearity. Let us mention that we addressed the blow-up problem for (1.2) in our previous work [4]. See next paragraph.

Our second result concerns the blow-up for (1.2) with initial data in \mathcal{A}_ω^- . In [4], we proved that if $\frac{7}{3} < p < 5$, $1 < q < p$, and $\omega > 0$, for $u_0 \in \mathcal{A}_\omega^-$, if one of the following conditions is satisfied:

- $u_0 \in \Sigma(\mathbb{R}^3) := H^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^3, |x|^2 dx)$,
- $u_0 \in H^1(\mathbb{R}^3)$ is radial,
- $u_0 \in \Sigma_3(\mathbb{R}^3)$ and $p \leq 3$, where

$$\Sigma_3(\mathbb{R}^3) := \{f \in H^1(\mathbb{R}^3) : f(x) = f(\bar{x}, x_3) = f(|\bar{x}|, x_3), f \in L^2(\mathbb{R}^3, x_3^2 dx)\}, \quad \bar{x} = (x_1, x_2),$$

then the corresponding solution to (1.2) blows up in finite time.

Here we show that we have the following blow-up rate for (1.2).

Theorem 1.3. *Let $\frac{7}{3} < p < 5$, $1 < q < p$, $\omega > 0$, and $u_0 \in \mathcal{A}_\omega^-$.*

(i) *Assume that one of the following conditions is fulfilled:*

- (1) $u_0 \in H^1(\mathbb{R}^3)$ is radial,
- (2) $u_0 \in \Sigma_3(\mathbb{R}^3)$ and $p < 3$.

Then the corresponding solution to (1.2) blows up in finite time, i.e., $T^ < +\infty$, and for t close to T^* ,*

$$\int_t^{T^*} (T^* - \tau) \|\nabla u(\tau)\|_{L^2(\mathbb{R}^3)}^2 d\tau \leq \begin{cases} C(T^* - t)^{\frac{2(5-p)}{p+3}} & \text{if (1) holds,} \\ C(T^* - t)^{\frac{4(3-p)}{5-p}} & \text{if (2) holds.} \end{cases} \quad (1.6)$$

In addition, there exists a time sequence $t_n \nearrow T^$ such that*

$$\|\nabla u(t_n)\|_{L^2(\mathbb{R}^3)} \leq \begin{cases} C(T^* - t_n)^{-\frac{2(p-1)}{p+3}} & \text{if (1) holds,} \\ C(T^* - t_n)^{-\frac{p-1}{5-p}} & \text{if (2) holds.} \end{cases} \quad (1.7)$$

(ii) *If we do not assume any symmetry on the solution (nor any restriction on the exponents), then either $T^* < \infty$ or u grows up in infinite time, namely $T^* = \infty$ and $\limsup_{t \rightarrow \infty} \|\nabla u(t)\|_{L^2(\mathbb{R}^3)} = \infty$.*

The Theorem above is based on the results we obtained in [4], jointly with a Merle, Raphaël, and Szeftel argument [30]. Moreover, the grow-up result follows by means of the Du, Wu, and Zhang approach [12]. All of these results come from virial estimates for suitable localization functions.

Notations. In the rest of the paper, we will use the notations below.

Given two quantities A and B , we denote $A \lesssim B$ if there exists a positive constant C such that $A \leq CB$. For $1 \leq p \leq \infty$, the $L^p = L^p(\Omega; \mathbb{C})$ are the classical Lebesgue spaces endowed with norm $\|f\|_{L^p(\Omega)} = (\int_\Omega |f(x)|^p dx)^{1/p}$ if $p \neq \infty$ or $\|f\|_{L^\infty(\Omega)} = \text{ess sup}_{x \in \Omega} |f(x)|$ for $p = \infty$. We denote by $H^1 = H^1(\mathbb{R}^3; \mathbb{C})$ the usual Sobolev space of L^2 functions with gradient in L^2 . Here $\langle \cdot \rangle$ stands for the Japanese brackets $\langle \cdot \rangle = (1 + |\cdot|^2)^{1/2}$. Given an interval $I \subseteq \mathbb{R}$, bounded or unbounded, we define by $L_t^p X_x = L_t^p(I, X_x)$ the Bochner space of vector-valued functions $f : I \rightarrow X$ endowed with the norm $(\int_I \|f(s)\|_X^p ds)^{1/p}$ for $1 \leq p < \infty$, with similar modification as above for $p = \infty$. (In what follows, $f \in L_t^p X_x$ means that $f = f(t, x)$ is a function depending on the time variable $t \in I \subseteq \mathbb{R}$ and the space variable $x \in \mathbb{R}^3$, with finite $L_t^p X_x$ -norm). For any $p \in [1, \infty]$, p' denotes its

dual defined by $p' = \frac{p}{p-1}$. As we work in the 3D case, we omit the \mathbb{R}^3 notation when no confusion may arise.

2. VARIATIONAL ANALYSIS

In this section, we recall and prove some crucial variational tools used along the paper, used in particular to define the scattering/blow-up dichotomy regions of initial data for the Cauchy problem (1.2). Furthermore, some of the results illustrated below will be also essential in proving a new coercivity property that we need to prove scattering by means of suitable interaction Morawetz estimates.

Let $\omega > 0$. We consider the minimization problem

$$m_\omega := \inf \{ S_\omega(\phi) : \phi \in H^1 \setminus \{0\}, G(\phi) = 0 \},$$

where

$$S_\omega(\phi) = E(\phi) + \frac{\omega}{2}M(\phi) = \frac{1}{2}\|\nabla\phi\|_{L^2}^2 - \frac{1}{p+1}\|\phi\|_{L^{p+1}}^{p+1} + \frac{1}{q+1}\|\phi\|_{L^{q+1}}^{q+1} + \frac{\omega}{2}\|\phi\|_{L^2}^2$$

is the action functional and

$$G(\phi) = \|\nabla\phi\|_{L^2}^2 - \frac{3(p-1)}{2(p+1)}\|\phi\|_{L^{p+1}}^{p+1} + \frac{3(q-1)}{2(q+1)}\|\phi\|_{L^{q+1}}^{q+1}$$

is the Pohozaev functional.

Proposition 2.1. *Let $\frac{7}{3} < p < 5$, $1 < q < p$, and $\omega > 0$. Then $m_\omega > 0$ and there exists at least a minimizer for m_ω .*

Before giving the proof of Proposition 2.1, let us start with the following observation.

Lemma 2.2. *Let $\phi \in H^1 \setminus \{0\}$. Then there exists a unique $\lambda_0 > 0$ such that*

$$G(\phi_\lambda) \begin{cases} > 0 & \text{if } 0 < \lambda < \lambda_0, \\ = 0 & \text{if } \lambda = \lambda_0, \\ < 0 & \text{if } \lambda > \lambda_0, \end{cases}$$

where

$$\phi_\lambda(x) := \lambda^{\frac{3}{2}}\phi(\lambda x), \quad \lambda > 0. \quad (2.1)$$

Proof. We have

$$\begin{aligned} G(\phi_\lambda) &= \lambda^2\|\nabla\phi\|_{L^2}^2 - \frac{3(p-1)}{2(p+1)}\lambda^{\frac{3}{2}(p-1)}\|\phi\|_{L^{p+1}}^{p+1} + \frac{3(q-1)}{2(q+1)}\lambda^{\frac{3}{2}(q-1)}\|\phi\|_{L^{q+1}}^{q+1} \\ &= \lambda^2 \left(\|\nabla\phi\|_{L^2}^2 - \frac{3(p-1)}{2(p+1)}\lambda^{\frac{3}{2}(p-1)-2}\|\phi\|_{L^{p+1}}^{p+1} + \frac{3(q-1)}{2(q+1)}\lambda^{\frac{3}{2}(q-1)-2}\|\phi\|_{L^{q+1}}^{q+1} \right) \\ &=: \lambda^2 f(\lambda). \end{aligned}$$

Consider $f(\lambda) = a - b\lambda^\alpha + c\lambda^\beta$ with $a, b, c > 0$, $\alpha > 0$, and $\beta < \alpha$. We have

$$f'(\lambda) = \lambda^{\beta-1}(c\beta - b\alpha\lambda^{\alpha-\beta}).$$

In particular, there exists a unique $\lambda_1 > 0$ such that

$$f(\lambda_1) = \max_{\lambda > 0} f(\lambda).$$

Drawing the graph of f , we see that there exists a unique $\lambda_0 > 0$ such that $f(\lambda_0) = 0$. Moreover, $f(\lambda) > 0$ for $0 < \lambda < \lambda_0$ and $f(\lambda) < 0$ for $\lambda > \lambda_0$. This shows the lemma. \square

Proof of Proposition 2.1. We proceed in several steps, illuminated by the works of Ibrahim, Masmoudi, and Nakanishi [19], and Akahori, Ibrahim, Kikuchi, and Nawa [2].

Step 1. An auxiliary minimization problem. Denote

$$I_\omega(\phi) := S_\omega(\phi) - \frac{2}{3(q-1)}G(\phi) = \frac{3q-7}{6(q-1)}\|\nabla\phi\|_{L^2}^2 + \frac{p-q}{(p+1)(q-1)}\|\phi\|_{L^{p+1}}^{p+1} + \frac{\omega}{2}\|\phi\|_{L^2}^2$$

and consider

$$\tilde{m}_\omega := \inf \{ I_\omega(\phi) : \phi \in H^1 \setminus \{0\}, G(\phi) \leq 0 \}. \quad (2.2)$$

We claim that $m_\omega = \tilde{m}_\omega > 0$. In fact, it is clear that $\tilde{m}_\omega \leq m_\omega$. Now let $\phi \in H^1 \setminus \{0\}$ be such that $G(\phi) \leq 0$. By Lemma 2.2, there exists $\lambda_0 \in (0, 1]$ such that $G(\phi_{\lambda_0}) = 0$. Thus

$$m_\omega \leq S_\omega(\phi_{\lambda_0}) = I_\omega(\phi_{\lambda_0}) \leq I_\omega(\phi),$$

where we have used $\lambda_0 \leq 1$ in the last inequality. Taking the infimum, we get $m_\omega \leq \tilde{m}_\omega$, hence $m_\omega = \tilde{m}_\omega$.

To see that $\tilde{m}_\omega > 0$, we take a minimizing sequence $\{\phi_n\}_n$ for \tilde{m}_ω , i.e., $\phi_n \in H^1 \setminus \{0\}$, $G(\phi_n) \leq 0$, and $I_\omega(\phi_n) \rightarrow \tilde{m}_\omega$. As I_ω is non-negative, there exists $C = C(\omega) > 0$ such that

$$\|\phi_n\|_{L^2}^2 \leq C, \quad \forall n \geq 1.$$

On the other hand, since $G(\phi_n) \leq 0$, we have

$$\|\nabla\phi_n\|_{L^2}^2 \leq \frac{3(p-1)}{2(p+1)}\|\phi_n\|_{L^{p+1}}^{p+1}$$

which together with the standard Gagliardo-Nirenberg inequality yield

$$\|\nabla\phi_n\|_{L^2}^2 \leq C\|\nabla\phi_n\|_{L^2}^{\frac{3(p-1)}{2}}\|\phi_n\|_{L^2}^{\frac{5-p}{2}} \leq C\|\nabla\phi_n\|_{L^2}^{\frac{3(p-1)}{2}}.$$

This shows that

$$\|\nabla\phi_n\|_{L^2}^2 \geq C > 0, \quad \forall n \geq 1, \quad (2.3)$$

hence $\tilde{m}_\omega > 0$.

Step 2. Minimizers for \tilde{m}_ω . Let $\{\phi_n\}_n$ be a minimizing sequence for \tilde{m}_ω . Observe that if ϕ_n^* is the Schwarz symmetrization of ϕ_n , then $\{\phi_n^*\}_n$ is still a minimizing sequence for \tilde{m}_ω . Thus without loss of generality, we can assume that ϕ_n is radially symmetric. As I_ω is non-negative, we see that $\{\phi_n\}_n$ is a bounded sequence in H^1 . Passing to a subsequence if necessary, there exists $\phi \in H^1$ such that $\phi_n \rightharpoonup \phi$ weakly in H^1 and $\phi_n \rightarrow \phi$ strongly in L^r for all $2 < r < 6$. As $G(\phi_n) \leq 0$, using (2.3), we have

$$\|\phi_n\|_{L^{p+1}}^{p+1} \geq \frac{2(p+1)}{3(p-1)}\|\nabla\phi_n\|_{L^2}^2 \geq \frac{2(p+1)}{3(p-1)}C > 0, \quad \forall n \geq 1$$

which shows $\phi \neq 0$. We also have

$$G(\phi) \leq \liminf_{n \rightarrow \infty} G(\phi_n) = 0,$$

thus

$$\tilde{m}_\omega \leq I_\omega(\phi) \leq \liminf_{n \rightarrow \infty} I_\omega(\phi_n) = \tilde{m}_\omega.$$

This shows that ϕ is a minimizer for \tilde{m}_ω .

Step 3. Minimizers for m_ω . We now show that there exists at least a minimizer for m_ω . Let ϕ be a minimizer for \tilde{m}_ω . By Lemma 2.2, there exists $\lambda_0 \in (0, 1]$ such that $G(\phi_{\lambda_0}) = 0$. Thus

$$m_\omega \leq S_\omega(\phi_{\lambda_0}) = I_\omega(\phi_{\lambda_0}) \leq I_\omega(\phi) = \tilde{m}_\omega = m_\omega.$$

This shows that $\lambda_0 = 1$, $G(\phi) = 0$, and $S_\omega(\phi) = m_\omega$. In particular, ϕ is a minimizer for m_ω . \square

3. SCATTERING CRITERION

In this section, we state and prove the key dispersive/Strichartz estimates which will enable us to state a scattering criterion. The results contained in this section are inspired by the recent techniques developed in the paper by Dodson and Murphy [11].

We start by reporting the well-known 3D dispersive estimate, see [6] for a proof.

Lemma 3.1. *We have, for all $r \in [2, \infty]$ and for any $t \neq 0$,*

$$\|e^{it\Delta} f\|_{L_x^r} \lesssim |t|^{-\frac{3}{2}(1-\frac{2}{r})} \|f\|_{L_x^{r'}} \quad (3.1)$$

for any $f \in L^{r'}$.

The next ones are the well-known Strichartz estimates, arising from the dispersive estimate above. See [6, 20].

Proposition 3.2. *The following space-time bounds hold true.*

- (Homogeneous Strichartz estimates) For any $f \in L^2$ and any Schrödinger admissible pair (a, b) , i.e.,

$$\frac{2}{a} + \frac{3}{b} = \frac{3}{2}, \quad b \in [2, 6],$$

then

$$\|e^{it\Delta} f\|_{L_t^a(\mathbb{R}, L_x^b)} \lesssim \|f\|_{L_x^2}.$$

- (Inhomogeneous Strichartz estimates) Let $I \subset \mathbb{R}$ be an interval containing 0. Then

$$\left\| \int_0^t e^{i(t-s)\Delta} F(s) ds \right\|_{L_t^a(I, L_x^b)} \lesssim \|F\|_{L_t^{\rho'}(I, L_x^{\gamma'})}$$

for any $F \in L_t^{\rho'}(I, L_x^{\gamma'})$ and any Schrödinger admissible pairs (a, b) and (ρ, γ) .

- (Strichartz estimates for non-admissible pairs) Let $I \subset \mathbb{R}$ be an interval containing 0 and (a, b) be a Schrödinger admissible pair with $b > 2$. Fix $m > \frac{a}{2}$ and define n by

$$\frac{1}{m} + \frac{1}{n} = \frac{2}{a}.$$

Then

$$\left\| \int_0^t e^{i(t-s)\Delta} F(s) ds \right\|_{L_t^m(I, L_x^b)} \lesssim \|F\|_{L_t^{n'}(I, L_x^{b'})}$$

for any $F \in L_t^{n'}(I, L_x^{b'})$.

Let us introduce some notations.

$$\begin{aligned} a_1 &:= \frac{4(p+1)}{3(p-1)}, & m_1 &:= \frac{2(p-1)(p+1)}{5-p}, & n_1 &:= \frac{2(p-1)(p+1)}{3p^2-5p-2}, \\ a_2 &:= \frac{4(q+1)}{3(q-1)}, & m_2 &:= \frac{2(q-1)(q+1)}{5-q}, & n_2 &:= \frac{2(q-1)(q+1)}{3q^2-5q-2}, \end{aligned}$$

and

$$\begin{aligned} b_1 &:= p + 1, & r_1 &:= \frac{6(p-1)(p+1)}{3p^2 + 2p - 13}, \\ b_2 &:= q + 1, & r_2 &:= \frac{6(q-1)(q+1)}{3q^2 + 2q - 13}. \end{aligned}$$

We readily check that $(a_1, b_1), (a_2, b_2), (m_1, r_1), (m_2, r_2)$ are Schrödinger admissible pairs,

$$\frac{1}{m_1} + \frac{1}{n_1} = \frac{2}{a_1}, \quad \frac{1}{m_2} + \frac{1}{n_2} = \frac{2}{a_2},$$

and

$$\frac{1}{b_1} = \frac{1}{r_1} - \frac{\sigma_1}{3}, \quad \frac{1}{b_2} = \frac{1}{r_2} - \frac{\sigma_2}{3},$$

where

$$\sigma_1 := \frac{3p-7}{2(p-1)}, \quad \sigma_2 := \frac{3q-7}{2(q-1)}.$$

The following Lemma follows directly from the above choices, Hölder's inequality, and Sobolev embeddings.

Lemma 3.3. *Let $I \subset \mathbb{R}$ be an interval. We have*

$$\begin{aligned} \| |u|^{p-1} u \|_{L_t^{n'_1}(I, L_x^{b'_1})} &\lesssim \| u \|_{L_t^{m_1}(I, L_x^{b_1})}^p, \\ \| |u|^{q-1} u \|_{L_t^{n'_2}(I, L_x^{b'_2})} &\lesssim \| u \|_{L_t^{m_2}(I, L_x^{b_2})}^q, \\ \| \langle \nabla \rangle (|u|^{p-1} u) \|_{L_t^{a'_1}(I, L_x^{b'_1})} &\lesssim \| u \|_{L_t^{m_1}(I, L_x^{b_1})}^{p-1} \| \langle \nabla \rangle u \|_{L_t^{a_1}(I, L_x^{b_1})}, \\ \| \langle \nabla \rangle (|u|^{q-1} u) \|_{L_t^{a'_2}(I, L_x^{b'_2})} &\lesssim \| u \|_{L_t^{m_2}(I, L_x^{b_2})}^{q-1} \| \langle \nabla \rangle u \|_{L_t^{a_2}(I, L_x^{b_2})}, \\ \| u \|_{L_t^{m_1}(I, L_x^{b_1})} &\lesssim \| |\nabla|^{\sigma_1} u \|_{L_t^{m_1}(I, L_x^{r_1})} \lesssim \| \langle \nabla \rangle u \|_{L_t^{m_1}(I, L_x^{r_1})}, \\ \| u \|_{L_t^{m_2}(I, L_x^{b_2})} &\lesssim \| |\nabla|^{\sigma_2} u \|_{L_t^{m_2}(I, L_x^{r_2})} \lesssim \| \langle \nabla \rangle u \|_{L_t^{m_2}(I, L_x^{r_2})}. \end{aligned}$$

We next prove a global existence result for small data.

Lemma 3.4. *Let $\frac{7}{3} < q < p < 5$ and $T > 0$ be such that $u(T) \in H^1$. Then there exists $\delta > 0$ sufficiently small such that if*

$$\| e^{i(t-T)\Delta} u(T) \|_{L_t^{m_1}([T, \infty), L_x^{b_1}) \cap L_t^{m_2}([T, \infty), L_x^{b_2})} < \delta,$$

then there exists a unique solution to (1.2) with initial datum $u(T)$ satisfying

$$\| u \|_{L_t^{m_1}([T, \infty), L_x^{b_1}) \cap L_t^{m_2}([T, \infty), L_x^{b_2})} \leq 2 \| e^{i(t-T)\Delta} u(T) \|_{L_t^{m_1}([T, \infty), L_x^{b_1}) \cap L_t^{m_2}([T, \infty), L_x^{b_2})}$$

and

$$\| \langle \nabla \rangle u \|_{L_t^{a_1}([T, \infty), L_x^{b_1}) \cap L_t^{a_2}([T, \infty), L_x^{b_2})} \leq C \| u(T) \|_{H_x^1}$$

for some constant $C > 0$.

Proof. We consider

$$X_T := \left\{ u : \| u \|_{L_t^{m_1}(I, L_x^{b_1}) \cap L_t^{m_2}(I, L_x^{b_2})} \leq M, \quad \| \langle \nabla \rangle u \|_{L_t^{a_1}(I, L_x^{b_1}) \cap L_t^{a_2}(I, L_x^{b_2})} \leq L \right\}$$

equipped with the distance

$$d(u, v) := \| u - v \|_{L_t^{a_1}(I, L_x^{b_1}) \cap L_t^{a_2}(I, L_x^{b_2})},$$

where $I = [T, \infty)$ and $M, L > 0$ will be chosen later. By the persistence of regularity (see e.g., [6, Theorem 1.2.5]), we readily see that (X_T, d) is a complete metric space. Our purpose is to show that the Duhamel functional

$$\Phi_T(u(t)) := e^{i(t-T)\Delta}u(T) + i \int_T^t e^{i(t-s)\Delta} (|u(s)|^{p-1}u(s) - |u(s)|^{q-1}u(s)) ds \quad (3.2)$$

is a contraction on (X_T, d) . By Strichartz estimates and Lemma 3.3, we have

$$\begin{aligned} \|\Phi_T(u)\|_{L_t^{m_1}(I, L_x^{b_1})} &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}(I, L_x^{b_1})} + \left\| \int_T^t e^{i(t-s)\Delta} |u(s)|^{p-1}u(s) ds \right\|_{L_t^{m_1}(I, L_x^{b_1})} \\ &\quad + \left\| \int_T^t e^{i(t-s)\Delta} |u(s)|^{q-1}u(s) ds \right\|_{L_t^{m_1}(I, L_x^{b_1})} \\ &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}(I, L_x^{b_1})} + C \| |u|^{p-1}u \|_{L_t^{n'_1}(I, L_x^{b'_1})} \\ &\quad + \left\| \langle \nabla \rangle \int_T^t e^{i(t-s)\Delta} |u(s)|^{q-1}u(s) ds \right\|_{L_t^{m_1}(I, L_x^{r_1})} \\ &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}(I, L_x^{b_1})} + C \| |u|^{p-1}u \|_{L_t^{n'_1}(I, L_x^{b'_1})} \\ &\quad + C \| \langle \nabla \rangle (|u|^{q-1}u) \|_{L_t^{a'_2}(I, L_x^{b'_2})} \\ &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}(I, L_x^{b_1})} + C \|u\|_{L_t^{m_1}(I, L_x^{b_1})}^p \\ &\quad + \|u\|_{L_t^{m_2}(I, L_x^{b_2})}^{q-1} \| \langle \nabla \rangle u \|_{L_t^{a_2}(I, L_x^{b_2})}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \|\Phi_T(u)\|_{L_t^{m_2}(I, L_x^{b_2})} &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_2}(I, L_x^{b_2})} + C \|u\|_{L_t^{m_1}(I, L_x^{b_1})}^{p-1} \| \langle \nabla \rangle u \|_{L_t^{a_2}(I, L_x^{b_2})} \\ &\quad + C \|u\|_{L_t^{m_2}(I, L_x^{b_2})}^q. \end{aligned}$$

We next estimate

$$\begin{aligned} \| \langle \nabla \rangle \Phi_T(u) \|_{L_t^{a_1}(I, L_x^{b_1})} &\leq C \|u(T)\|_{H_x^1} + C \| \langle \nabla \rangle (|u|^{p-1}u) \|_{L_t^{a'_1}(I, L_x^{b'_1})} \\ &\quad + C \| \langle \nabla \rangle (|u|^{q-1}u) \|_{L_t^{a'_2}(I, L_x^{b'_2})} \\ &\leq C \|u(T)\|_{H_x^1} + C \|u\|_{L_t^{m_1}(I, L_x^{b_1})}^{p-1} \| \langle \nabla \rangle u \|_{L_t^{a_1}(I, L_x^{b_1})} \\ &\quad + C \|u\|_{L_t^{m_2}(I, L_x^{b_2})}^{q-1} \| \langle \nabla \rangle u \|_{L_t^{a_2}(I, L_x^{b_2})} \end{aligned}$$

and

$$\begin{aligned} \| \langle \nabla \rangle \Phi_T(u) \|_{L_t^{a_2}(I, L_x^{b_2})} &\leq C \|u(T)\|_{H_x^1} + C \|u\|_{L_t^{m_1}(I, L_x^{b_1})}^{p-1} \| \langle \nabla \rangle u \|_{L_t^{a_1}(I, L_x^{b_1})} \\ &\quad + C \|u\|_{L_t^{m_2}(I, L_x^{b_2})}^{q-1} \| \langle \nabla \rangle u \|_{L_t^{a_2}(I, L_x^{b_2})}. \end{aligned}$$

Next we have

$$\begin{aligned} \|\Phi_T(u) - \Phi_T(v)\|_{L_t^{a_1}(I, L_x^{b_1}) \cap L_t^{a_2}(I, L_x^{b_2})} &\leq C \| |u|^{p-1}u - |v|^{p-1}v \|_{L_t^{n'_1}(I, L_x^{b'_1})} \\ &\quad + C \| |u|^{q-1}u - |v|^{q-1}v \|_{L_t^{n'_2}(I, L_x^{b'_2})} \\ &\leq \left(\|u\|_{L_t^{m_1}(I, L_x^{b_1})}^{p-1} + \|v\|_{L_t^{m_1}(I, L_x^{b_1})}^{p-1} \right) \|u - v\|_{L_t^{a_1}(I, L_x^{b_1})} \\ &\quad + \left(\|u\|_{L_t^{m_2}(I, L_x^{b_2})}^{q-1} + \|v\|_{L_t^{m_2}(I, L_x^{b_2})}^{q-1} \right) \|u - v\|_{L_t^{a_2}(I, L_x^{b_2})}. \end{aligned}$$

Thus there exists $C > 0$ independent of T such that for any $u, v \in X_T$, we have

$$\begin{aligned} \|\Phi_T(u)\|_{L_t^{m_1}(I, L_x^{b_1}) \cap L_t^{m_2}(I, L_x^{b_2})} &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}(I, L_x^{b_1}) \cap L_t^{m_2}(I, L_x^{b_2})} + C(M^q + M^p) \\ &\quad + C(M^{q-1} + M^{p-1})L, \end{aligned}$$

$$\|\langle \nabla \rangle \Phi_T(u)\|_{L_t^{a_1}(I, L_x^{b_1}) \cap L_t^{a_2}(I, L_x^{b_2})} \leq C\|u(T)\|_{H_x^1} + C(M^{q-1} + M^{p-1})L,$$

and

$$d(\Phi_T(u), \Phi_T(v)) \leq C(M^{q-1} + M^{p-1})d(u, v).$$

By choosing $M = 2\|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}(I, L_x^{b_1}) \cap L_t^{m_2}(I, L_x^{b_2})}$ and $L = 2C\|u(T)\|_{H_x^1}$ and taking M sufficiently small, we see that Φ_T is a contraction on (X_T, d) . This completes the proof. \square

The next Lemma is a small data scattering result.

Lemma 3.5. *Let $\frac{7}{3} < q < p < 5$. Suppose that $u(t)$ is a global solution to (1.2) satisfying $\|u\|_{L_t^\infty(\mathbb{R}, H_x^1)} < \infty$. Then there exists $\delta > 0$ sufficiently small such that if*

$$\|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} < \delta$$

for some $T > 0$, then u scatters forward in time.

Proof. We first observe that for any interval $I \subset \mathbb{R}$,

$$\|u\|_{L_t^{m_2}(I, L_x^{b_2})} \leq \|u\|_{L_t^{m_1}(I, L_x^{b_1})}^\theta \|u\|_{L_t^\rho(I, L_x^\gamma)}^{1-\theta},$$

where

$$\theta = \frac{(3q-7)(p-1)}{(3p-7)(q-1)}, \quad \rho = \frac{(1-\theta)m_1m_2}{m_1 - \theta m_2}, \quad \gamma = \frac{(1-\theta)b_1b_2}{b_1 - \theta b_2}.$$

We readily check that for $\frac{7}{3} < q < p < 5$,

$$\theta \in (0, 1), \quad \frac{2}{\rho} + \frac{3}{\gamma} = \frac{3}{2}, \quad \gamma \in [2, 6],$$

namely (ρ, γ) is a Schrödinger admissible pair. It follows that

$$\begin{aligned} \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_2}([T, \infty), L_x^{b_2})} &\leq \|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}([T, \infty), L_x^{b_1})}^\theta \|e^{i(t-T)\Delta}u(T)\|_{L_t^\rho([T, \infty), L_x^\gamma)}^{1-\theta} \\ &\leq \delta^\theta \|u(T)\|_{H_x^1}^{1-\theta}. \end{aligned}$$

Thus

$$\|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}([T, \infty), L_x^{b_1}) \cap L_t^{m_2}([T, \infty), L_x^{b_2})} < \varepsilon(\delta)$$

for some $\varepsilon(\delta) > 0$ small depending on δ . By Lemma 3.4, we have

$$\|u\|_{L_t^{m_1}([T, \infty), L_x^{b_1}) \cap L_t^{m_2}([T, \infty), L_x^{b_2})} \leq 2\|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}([T, \infty), L_x^{b_1}) \cap L_t^{m_2}([T, \infty), L_x^{b_2})} \quad (3.3)$$

and

$$\|\langle \nabla \rangle u\|_{L_t^{a_1}([T, \infty), L_x^{b_1}) \cap L_t^{a_2}([T, \infty), L_x^{b_2})} \leq C\|u(T)\|_{H_x^1} \quad (3.4)$$

for some constant $C > 0$.

For $T < t_1 < t_2$, we use the Duhamel formula (3.2), (3.3), and (3.4) to have

$$\begin{aligned} \|e^{-it_2\Delta}u(t_2) - e^{-it_1\Delta}u(t_1)\|_{H_x^1} &\leq C\|\langle \nabla \rangle (|u|^{p-1}u)\|_{L_t^{a_1'}((t_1, t_2), L_x^{b_1})} \\ &\quad + C\|\langle \nabla \rangle (|u|^{q-1}u)\|_{L_t^{a_2'}((t_1, t_2), L_x^{b_2})} \\ &\leq C\|u\|_{L_t^{m_1}((t_1, t_2), L_x^{b_1})}^{p-1} \|\langle \nabla \rangle u\|_{L_t^{a_1}((t_1, t_2), L_x^{b_1})} \\ &\quad + C\|u\|_{L_t^{m_2}((t_1, t_2), L_x^{b_2})}^{q-1} \|\langle \nabla \rangle u\|_{L_t^{a_2}((t_1, t_2), L_x^{b_2})} \rightarrow 0 \end{aligned}$$

as $t_1, t_2 \rightarrow \infty$. This shows that $\{e^{-it\Delta}u(t)\}_{t \rightarrow \infty}$ is a Cauchy sequence in H_x^1 . Thus there exists

$$u_+ = e^{-iT\Delta}u(T) + i \int_T^\infty e^{-is\Delta} (|u(s)|^{p-1}u(s) - |u(s)|^{q-1}u(s)) ds \in H_x^1$$

such that $e^{-it\Delta}u(t) \rightarrow u_+$ strongly in H_x^1 as $t \rightarrow \infty$. By the unitary property of the propagator, we obtain

$$\|u(t) - e^{it\Delta}u_+\|_{H_x^1} \rightarrow 0 \text{ as } t \rightarrow \infty.$$

The proof is complete. \square

We are now ready to state the following scattering criterion. As explained in [11], the criterion infers that if on any time interval which is large enough, one can find a large interval where the scattering norm is small, then the global solution scatters.

Lemma 3.6. *Let $\frac{7}{3} < q < p < 5$. Suppose that $u(t)$ is a global solution to (1.2) satisfying $\|u\|_{L_t^\infty(\mathbb{R}, H_x^1)} < \infty$. Then there exist $\varepsilon > 0$ and $T_0 = T_0(\varepsilon) > 0$ such that if for any $a > 0$, there exists $t_0 \in (a, a + T_0)$ such that $[t_0 - \varepsilon^{-\sigma}, t_0] \subset (a, a + T_0)$ and*

$$\|u\|_{L_t^{m_1}([t_0 - \varepsilon^{-\sigma}, t_0], L_x^{b_1})} \lesssim \varepsilon \quad (3.5)$$

for some $\sigma > 0$, then the solution scatters forward in time.

Proof. By Lemma 3.5, it suffices to prove that there exists $T > 0$ such that

$$\|e^{i(t-T)\Delta}u(T)\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} \lesssim \varepsilon^\mu \quad (3.6)$$

for some $\mu > 0$. To show (3.6), we write

$$e^{i(t-T)\Delta}u(T) = e^{it\Delta}u_0 + i \int_0^T e^{i(t-s)\Delta} (|u(s)|^{p-1}u(s) - |u(s)|^{q-1}u(s)) ds.$$

By Sobolev embedding and Strichartz estimate, we have

$$\|e^{it\Delta}u_0\|_{L_t^{m_1}(\mathbb{R}, L_x^{b_1})} \lesssim \|u_0\|_{H_x^1}.$$

By the monotone convergence theorem, there exists $T_1 > 0$ sufficiently large such that for all $T > T_1$,

$$\|e^{it\Delta}u_0\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} \lesssim \varepsilon. \quad (3.7)$$

Taking $a = T_1$ and $T = t_0$ with a and t_0 as in (3.5), we write

$$\begin{aligned} & \int_0^T e^{i(t-s)\Delta} (|u(s)|^{p-1}u(s) - |u(s)|^{q-1}u(s)) ds = \\ & \int_{I_1 \cup I_2} e^{i(t-s)\Delta} (|u(s)|^{p-1}u(s) - |u(s)|^{q-1}u(s)) ds \end{aligned}$$

where $I_1 = [T - \varepsilon^{-\sigma}, T]$ and $I_2 = [0, T - \varepsilon^{-\sigma}]$. By using the linearity, we denote by $F_1(t)$ and $F_2(t)$ the integrals over I_1 and I_2 , respectively.

To estimate F_1 , we start with the following observation:

$$\|u\|_{L_t^{qn'_1}(I_1, L_x^{qb'_1})} \leq \|u\|_{L_t^{m_1}(I_1, L_x^{b_1})}^\theta \|u\|_{L_t^\rho(I_1, L_x^\gamma)}^{1-\theta}, \quad (3.8)$$

where

$$\theta = \frac{3pq - 3q - 4p}{q(3p - 7)}, \quad \rho = \frac{(1 - \theta)qm_1n'_1}{m_1 - \theta qn'_1}, \quad \gamma = \frac{(1 - \theta)qb_1b'_1}{b_1 - \theta qb'_1}.$$

Since $\frac{7}{3} < q < p < 5$, we see that

$$\theta \in (0, 1), \quad \frac{2}{\rho} + \frac{3}{\gamma} = \frac{3}{2}, \quad \gamma \in [2, 6].$$

In particular, (ρ, γ) is a Schrödinger admissible pair. Using (3.8), we have

$$\begin{aligned} \|F_1\|_{L_t^{m_1}([T, \infty) L_x^{b_1})} &\leq C \| |u|^{p-1} u \|_{L_t^{n'_1}(I_1, L_x^{b'_1})} + C \| |u|^{q-1} u \|_{L_t^{n'_1}(I_1, L_x^{b'_1})} \\ &\leq C \|u\|_{L_t^{m_1}(I_1, L_x^{b_1})}^p + C \|u\|_{L_t^{qn'_1}(I_1, L_x^{qb'_1})}^q \\ &\leq C \|u\|_{L_t^{m_1}(I_1, L_x^{b_1})}^p + C \|u\|_{L_t^{m_1}(I_1, L_x^{b_1})}^{q\theta} \|u\|_{L_t^{\rho}(I_1, L_x^{\gamma})}^{q(1-\theta)} \\ &\leq C \|u\|_{L_t^{m_1}(I_1, L_x^{b_1})}^p + C |I_1|^{\frac{q}{\rho}(1-\theta)} \|u\|_{L_t^{m_1}(I_1, L_x^{b_1})}^{q\theta} \|u\|_{L_t^{\infty}(I_1, H_x^1)}^{q(1-\theta)}. \end{aligned}$$

Since $\|u\|_{L_t^{\infty}(\mathbb{R}, H_x^1)} < \infty$, we infer from (3.5) that

$$\|F_1\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} \leq C\varepsilon^p + C\varepsilon^{q\theta - \frac{q}{\rho}\sigma(1-\theta)}.$$

Taking $\sigma > 0$ small, we have

$$\|F_1\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} \leq C\varepsilon^{\mu} \tag{3.9}$$

for some $\mu > 0$.

To estimate F_2 , we use Hölder's inequality to have

$$\|F_2\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} \leq \|F_2\|_{L_t^{\frac{r_1}{b_1}}([T, \infty), L_x^{r_1})} \|F_2\|_{L_t^{\frac{b_1-r_1}{b_1}}([T, \infty), L_x^{\infty})}.$$

As (m_1, r_1) is a Schrödinger admissible pair and

$$F_2(t) = e^{i(t-T+\varepsilon^{-\sigma})\Delta} u(T - \varepsilon^{-\sigma}) - e^{it\Delta} u_0,$$

we have

$$\|F_2\|_{L_t^{m_1}([T, \infty), L_x^{r_1})} \lesssim 1.$$

On the other hand, by the dispersive estimates (3.1) and Sobolev embedding, and noting that $m_1 > 2$, we have for all $t \geq T$,

$$\begin{aligned} \|F_2(t)\|_{L_x^{\infty}} &\lesssim \int_0^{T-\varepsilon^{-\sigma}} (t-s)^{-\frac{3}{2}} (\| |u(s)|^{p-1} u(s) \|_{L_x^1} + \| |u(s)|^{q-1} u(s) \|_{L_x^1}) ds \\ &= \int_0^{T-\varepsilon^{-\sigma}} (t-s)^{-\frac{3}{2}} (\|u(s)\|_{L_x^p}^p + \|u(s)\|_{L_x^q}^q) ds \\ &\lesssim \int_0^{T-\varepsilon^{-\sigma}} (t-s)^{-\frac{3}{2}} (\|u(s)\|_{H_x^1}^p + \|u(s)\|_{H_x^1}^q) ds \\ &\lesssim (t-T + \varepsilon^{-\sigma})^{-\frac{1}{2}}. \end{aligned}$$

It follows that

$$\|F_2\|_{L_t^{m_1}([T, \infty), L_x^{\infty})} \lesssim \left(\int_T^{\infty} (t-T + \varepsilon^{-\sigma})^{-\frac{m_1}{2}} dt \right)^{\frac{1}{m_1}} \lesssim \varepsilon^{\sigma(\frac{1}{2} - \frac{1}{m_1})}.$$

In particular, we get

$$\|F_2\|_{L_t^{m_1}([T, \infty), L_x^{b_1})} \lesssim \varepsilon^{\sigma(\frac{1}{2} - \frac{1}{m_1}) \frac{r_1 - b_1}{r_1}}. \tag{3.10}$$

Collecting (3.7), (3.9), (3.10), and choosing $\sigma > 0$ sufficiently small, we prove (3.6). The proof is complete. \square

4. ENERGY SCATTERING

This section is the bulk of the paper and contains the main novelty. Specifically, we will prove a coercivity property, see Lemma 4.2 below, and the interaction Morawetz estimates that will allow to prove the scattering for large data global solutions to (1.2).

4.1. A cutoff function. Let $\eta \in (0, 1)$ be a small constant. Let χ be a smooth decreasing radial function satisfying

$$\chi(x) = \chi(r) = \begin{cases} 1 & \text{if } r \leq 1 - \eta, \\ 0 & \text{if } r > 1, \end{cases} \quad |\chi'(r)| \lesssim \frac{1}{\eta}, \quad r = |x|. \quad (4.1)$$

For $R > 0$ large, we define the functions

$$\phi_R(x) := \frac{1}{\omega_3 R^3} \int \chi_R^2(x-z) \chi_R^2(z) dz$$

and

$$\phi_{p,R}(x) := \frac{1}{\omega_3 R^3} \int \chi_R^2(x-z) \chi_R^{p+1}(z) dz,$$

$$\phi_{q,R}(x) := \frac{1}{\omega_3 R^3} \int \chi_R^2(x-z) \chi_R^{q+1}(z) dz,$$

where $\chi_R(z) := \chi(z/R)$, and ω_3 is the volume of the unit ball in \mathbb{R}^3 . We see that ϕ_R , $\phi_{p,R}$, and $\phi_{q,R}$ are radial functions. We next define the radial function

$$\psi_R(x) = \psi_R(r) := \frac{1}{r} \int_0^r \phi_R(\tau) d\tau, \quad r = |x|. \quad (4.2)$$

We collect some basic properties of ϕ_R , $\phi_{p,R}$, $\phi_{q,R}$, and ψ_R as follows (see [9] for a proof.).

Lemma 4.1. *We have*

$$|\psi_R(x)| \lesssim \min \left\{ 1, \frac{R}{|x|} \right\}, \quad \partial_j \psi_R(x) = \frac{x_j}{|x|^2} (\phi_R(x) - \psi_R(x)), \quad j = 1, 2, 3$$

and

$$\psi_R(x) - \phi_R(x) \geq 0, \quad \phi_R(x) - \phi_{q,R}(x) \geq 0, \quad |\phi_R(x) - \phi_{p,R}(x)| \lesssim \eta \quad (4.3)$$

and

$$|\nabla \phi_R(x)| \lesssim \frac{1}{\eta R}, \quad |\psi_R(x) - \phi_R(x)| \lesssim \frac{1}{\eta} \min \left\{ \frac{|x|}{R}, \frac{R}{|x|} \right\}, \quad |\nabla \psi_R(x)| \lesssim \frac{1}{\eta} \min \left\{ \frac{1}{R}, \frac{R}{|x|^2} \right\} \quad (4.4)$$

for all $x \in \mathbb{R}^3$.

4.2. A coercivity property. The following is the essential new ingredient that will be exploited to prove suitable interaction Morawetz estimates. As already mentioned in the introduction, the scaling invariant equation with one nonlinearity is treated by taking advantage of the refined Gagliardo-Nirenberg inequality (1.5), that we cannot use in the present paper.

Lemma 4.2. *Let $\frac{7}{3} < q < p < 5$ and $\omega > 0$. Then \mathcal{A}_ω^+ is invariant under the flow of (1.2), i.e., if $u_0 \in \mathcal{A}_\omega^+$, then $u(t) \in \mathcal{A}_\omega^+$ for all $t \in I_{\max}$. In particular, the solution to (1.2) with initial data u_0 exists globally in time. In addition, there exists $R_0 > 0$ sufficiently large such that for all $R \geq R_0$, all $z \in \mathbb{R}^3$, and all $t \in \mathbb{R}$,*

$$G(\chi_R(\cdot - z)u^\xi(t)) \geq \delta \|\nabla(\chi_R(\cdot - z)u^\xi(t))\|_{L^2}^2, \quad (4.5)$$

where $u^\xi(t, x) := e^{ix \cdot \xi} u(t, x)$ with $\xi = \xi(t, z, R)$ and

$$\xi(t, z, R) := \begin{cases} -\frac{\int \operatorname{Im}(\chi_R^2(x-z)\bar{u}(t,x)\nabla u(t,x))dx}{\int \chi_R^2(x-z)|u(t,x)|^2 dx} & \text{if } \int \chi_R^2(x-z)|u(t,x)|^2 dx \neq 0, \\ 0 & \text{if } \int \chi_R^2(x-z)|u(t,x)|^2 dx = 0, \end{cases} \quad (4.6)$$

and $\chi_R(x) = \chi(x/R)$ with χ as in (4.1)

Proof. Let $u_0 \in \mathcal{A}_\omega^+$. We will show that $u(t) \in \mathcal{A}_\omega^+$ for all $t \in I_{\max}$. By the conservation of mass and energy, we have $S_\omega(u(t)) = S_\omega(u_0) < m_\omega$ for all $t \in I_{\max}$. Assume by contradiction that there exists t_0 such that $G(u(t_0)) < 0$. As $u : I_{\max} \rightarrow H^1$ is continuous, there exists t_1 such that $G(u(t_1)) = 0$. By the definition of m_ω , we have $S_\omega(u(t_1)) \geq m_\omega$ which is a contradiction. Thus $G(u(t)) \geq 0$ for all $t \in I_{\max}$ or \mathcal{A}_ω^+ is invariant under the flow of (1.2).

As $G(u(t)) \geq 0$, we have

$$\begin{aligned} & \frac{3q-7}{6(q-1)} \|\nabla u(t)\|_{L^2}^2 + \frac{p-q}{(p+1)(q-1)} \|u(t)\|_{L^{p+1}}^{p+1} + \frac{\omega}{2} \|u(t)\|_{L^2}^2 \\ & = I_\omega(u(t)) = S_\omega(u(t)) - \frac{2}{3(q-1)} G(u(t)) \leq S_\omega(u(t)) < m_\omega, \quad \forall t \in I_{\max}. \end{aligned}$$

This shows that

$$\|\nabla u(t)\|_{L^2}^2 \leq \frac{6(q-1)}{3q-7} m_\omega, \quad \forall t \in I_{\max}. \quad (4.7)$$

The blow-up alternative shows that the solution must exist globally in time.

Next we prove (4.5). To this end, we observe that

$$I_\omega(u(t)) = S_\omega(u(t)) - \frac{2}{3(q-1)} G(u(t)) \leq S_\omega(u(t)) = S_\omega(u_0) = m_\omega - \nu, \quad \forall t \in I_{\max}, \quad (4.8)$$

for some $\nu > 0$ as $S_\omega(u_0) < m_\omega$. Using

$$\int |\nabla(\chi u)|^2 dx = \int \chi^2 |\nabla u|^2 dx - \int \chi \Delta \chi |u|^2 dx, \quad (4.9)$$

we have

$$\int |\nabla(\chi u^\xi)|^2 dx = |\xi|^2 \int \chi^2 |u|^2 dx + \int \chi^2 |\nabla u|^2 dx - \int \chi \Delta \chi |u|^2 dx + 2\xi \cdot \int \operatorname{Im}(\chi^2 \bar{u} \nabla u) dx.$$

By the definition of ξ , we have

$$\begin{aligned} I_\omega(\chi_R(\cdot - z)u^\xi(t)) &= \frac{3q-7}{6(q-1)} \left(\int \chi_R^2(\cdot - z) |\nabla u(t)|^2 dx \right. \\ &\quad \left. - \int \chi_R(\cdot - z) \Delta \chi_R(\cdot - z) |u(t)|^2 dx \right) \\ &\quad - \frac{3q-7}{6(q-1)} \frac{\left| \int \operatorname{Im}(\chi_R^2(\cdot - z) \bar{u}(t, x) \nabla u(t, x)) dx \right|^2}{\int \chi_R^2(\cdot - z) |u(t, x)|^2 dx} \\ &\quad + \frac{p-q}{(p+1)(p-1)} \int \chi_R^{p+1}(\cdot - z) |u(t)|^{p+1} dx + \frac{\omega}{2} \int \chi_R^2(\cdot - z) |u(t)|^2 dx \\ &\leq I_\omega(u(t)) + O(R^{-2}). \end{aligned}$$

Thus, by (4.8), there exists $R_0 > 0$ sufficiently large such that

$$I_\omega(\chi_R(\cdot - z)u^\xi(t)) \leq m_\omega - \frac{\nu}{2}, \quad \forall R \geq R_0, \forall z \in \mathbb{R}^3, \forall t \in \mathbb{R}. \quad (4.10)$$

We claim that

$$G(\chi_R(\cdot - z)u^\xi(t)) > 0, \quad \forall R \geq R_0, \forall z \in \mathbb{R}^3, \forall t \in \mathbb{R}. \quad (4.11)$$

Suppose now that there exists $R_1 \geq R_0$, $z_1 \in \mathbb{R}^3$, $t_1 \in \mathbb{R}$, and $\xi_1 = \xi(t_1, z_1, R_1) \in \mathbb{R}^3$ such that $G(\chi_{R_1}(\cdot - z_1)u^{\xi_1}(t_1)) \leq 0$. Using the definition of \tilde{m}_ω (see (2.2)) and the fact that $m_\omega = \tilde{m}_\omega$, we have

$$I_\omega(\chi_{R_1}(\cdot - z_1)u^{\xi_1}(t_1)) \geq m_\omega$$

which contradicts (4.10).

For $R \geq R_0$, $z \in \mathbb{R}^3$, $t \in \mathbb{R}$, and $\xi = \xi(t, z, R) \in \mathbb{R}^3$, we denote $\Theta := \chi_R(x - z)u^\xi(t)$. We consider two cases.

Case 1. Assume that

$$4\|\nabla\Theta\|_{L^2}^2 - \frac{3(p-1)(3p+1)}{4(p+1)}\|\Theta\|_{L^{p+1}}^{p+1} + \frac{3(q-1)(3q+1)}{4(q+1)}\|\Theta\|_{L^{q+1}}^{q+1} \geq 0.$$

Then we have

$$\begin{aligned} G(\Theta) &= \|\nabla\Theta\|_{L^2}^2 - \frac{3(p-1)}{2(p+1)}\|\Theta\|_{L^{p+1}}^{p+1} + \frac{3(q-1)}{2(q+1)}\|\Theta\|_{L^{q+1}}^{q+1} \\ &\geq \frac{3p-7}{3p+1}\|\nabla\Theta\|_{L^2}^2 + \frac{9(q-1)(p-q)}{2(q+1)(3p+1)}\|\Theta\|_{L^{q+1}}^{q+1} \\ &\geq \frac{3p-7}{3p+1}\|\nabla\Theta\|_{L^2}^2. \end{aligned}$$

This proves (4.5).

Case 2. We now assume that

$$4\|\nabla\Theta\|_{L^2}^2 - \frac{3(p-1)(3p+1)}{4(p+1)}\|\Theta\|_{L^{p+1}}^{p+1} + \frac{3(q-1)(3q+1)}{4(q+1)}\|\Theta\|_{L^{q+1}}^{q+1} < 0. \quad (4.12)$$

We first observe that as $I_\omega(\Theta) < m_\omega$ (see (4.10)), an argument leading to (4.7) yields

$$\|\nabla\Theta\|_{L^2}^2 \leq \frac{6(q-1)}{3q-7}m_\omega. \quad (4.13)$$

Now set $f(\lambda) := S_\omega(\Theta_\lambda)$, with Θ_λ as in the rescaling (2.1). We have

$$f'(\lambda) = \lambda \|\nabla \Theta\|_{L^2}^2 - \frac{3(p-1)}{2(p+1)} \lambda^{\frac{3(p-1)}{2}-1} \|\Theta\|_{L^{p+1}}^{p+1} + \frac{3(q-1)}{2(q+1)} \lambda^{\frac{3(q-1)}{2}-1} \|\Theta\|_{L^{q+1}}^{q+1} = \frac{G(\Theta_\lambda)}{\lambda}$$

and

$$(\lambda f'(\lambda))' = 2\lambda \|\nabla \Theta\|_{L^2}^2 - \frac{9(p-1)^2}{4(p+1)} \lambda^{\frac{3(p-1)}{2}-1} \|\Theta\|_{L^{p+1}}^{p+1} + \frac{9(q-1)^2}{4(q+1)} \lambda^{\frac{3(q-1)}{2}-1} \|\Theta\|_{L^{q+1}}^{q+1}.$$

We write

$$\begin{aligned} (\lambda f'(\lambda))' &= -2f'(\lambda) + \lambda \left(4\|\nabla \Theta\|_{L^2}^2 - \frac{3(p-1)(3p+1)}{4(p+1)} \lambda^{\frac{3p-7}{2}} \|\Theta\|_{L^{p+1}}^{p+1} \right. \\ &\quad \left. + \frac{3(q-1)(3q+1)}{4(q+1)} \lambda^{\frac{3q-7}{2}} \|\Theta\|_{L^{q+1}}^{q+1} \right) \\ &=: -2f'(\lambda) + \lambda h(\lambda). \end{aligned}$$

Thanks to (4.12), we have

$$\begin{aligned} h'(\lambda) &= \lambda^{\frac{3q-7}{2}} \left(-\frac{3(p-1)(3p+1)(3p-7)}{8(p+1)} \lambda^{\frac{3(p-q)}{2}} \|\Theta\|_{L^{p+1}}^{p+1} \right. \\ &\quad \left. + \frac{3(q-1)(3q+1)(3p-7)}{8(q+1)} \|\Theta\|_{L^{q+1}}^{q+1} \right) \\ &\leq \lambda^{\frac{3q-7}{2}} \left(-\frac{3(p-1)(3p+1)(3p-7)}{2(p+1)} \lambda^{\frac{3(p-q)}{2}} \|\nabla \Theta\|_{L^2}^2 \right. \\ &\quad \left. + \frac{3(q-1)(3q+1)}{4(q+1)} \left(-\frac{3p-7}{2} \lambda^{\frac{3(p-q)}{2}} + \frac{3q-7}{2} \right) \|\Theta\|_{L^{q+1}}^{q+1} \right) \\ &\leq \lambda^{\frac{3q-7}{2}} \left(-\frac{3(p-1)(3p+1)(3p-7)}{2(p+1)} \lambda^{\frac{3(p-q)}{2}} \|\nabla \Theta\|_{L^2}^2 \right. \\ &\quad \left. - \frac{9(q-1)(3q+1)(p-q)}{8(q+1)} \|\Theta\|_{L^{q+1}}^{q+1} \right) < 0 \end{aligned}$$

for all $\lambda \geq 1$. This shows that $h(\lambda) \leq h(1) < 0$ for all $\lambda \geq 1$. In particular, we have

$$(\lambda f'(\lambda))' \leq -2f'(\lambda), \quad \forall \lambda \geq 1. \quad (4.14)$$

As $G(\Theta) > 0$ due to (4.11), Lemma 2.2 shows $G(\Theta_{\lambda_0}) = 0$ for some $\lambda_0 > 1$. Integrating (4.14) over $(1, \lambda_0)$, we get

$$G(\Theta) \geq 2(S_\omega(\Theta_{\lambda_0}) - S_\omega(\Theta)) = 2 \left(I_\omega(\Theta_{\lambda_0}) - I_\omega(\Theta) - \frac{2}{3(q-1)} G(\Theta) \right).$$

It follows that

$$\begin{aligned} G(\Theta) &\geq \frac{6(q-1)}{3q+1} (I_\omega(\Theta_{\lambda_0}) - I_\omega(\Theta)) \\ &\geq \frac{6(q-1)}{3q+1} (m_\omega - I_\omega(\Theta)) \\ &\geq \frac{6(q-1)}{3q+1} \left(1 - \frac{I_\omega(\Theta)}{m_\omega} \right) m_\omega \\ &\geq \frac{(3q-7)\nu}{2(3q+1)m_\omega} \|\nabla \Theta\|_{L^2}^2, \end{aligned}$$

where we have used (4.10) and (4.13) to get the last inequality. This also proves (4.5). \square

4.3. An interaction Morawetz estimate. We next define the interaction Morawetz action

$$\mathcal{M}_R^{\otimes 2}(t) := 2 \iint |u(t, y)|^2 \psi_R(x - y)(x - y) \cdot \text{Im}(\bar{u}(t, x) \nabla u(t, x)) dx dy, \quad (4.15)$$

where ψ_R is as in (4.2). We start by the following interaction Morawetz identity.

Lemma 4.3. *Let u be a solution to (1.2) satisfying*

$$\sup_{t \in [0, T^*)} \|u(t)\|_{H_x^1} \leq A$$

for some constant $A > 0$. Let $\mathcal{M}_R(t)$ be as in (4.15). Then we have

$$\sup_{t \in [0, T^*)} |\mathcal{M}_R^{\otimes 2}(t)| \lesssim_A R. \quad (4.16)$$

Moreover, we have

$$\begin{aligned} \frac{d}{dt} \mathcal{M}_R^{\otimes 2}(t) &= -4 \sum_{j,k} \iint \partial_j (\text{Im}(\bar{u}(t, y) \partial_j u(t, y))) \\ &\quad \times \psi_R(x - y)(x_k - y_k) \text{Im}(\bar{u}(t, x) \partial_k u(t, x)) dx dy \end{aligned} \quad (4.17)$$

$$\begin{aligned} &- 4 \sum_{j,k} \iint |u(t, y)|^2 \psi_R(x - y)(x_j - y_j) \\ &\quad \times \partial_k (\text{Re}(\partial_j u(t, x) \partial_k \bar{u}(t, x))) dx dy \end{aligned} \quad (4.18)$$

$$+ \iint |u(t, y)|^2 \psi_R(x - y)(x - y) \cdot \nabla \Delta(|u(t, x)|^2) dx dy \quad (4.19)$$

$$+ \frac{2(p-1)}{p+1} \iint |u(t, y)|^2 \psi_R(x - y)(x - y) \cdot \nabla(|u(t, x)|^{p+1}) dx dy \quad (4.20)$$

$$- \frac{2(q-1)}{q+1} \iint |u(t, y)|^2 \psi_R(x - y)(x - y) \cdot \nabla(|u(t, x)|^{q+1}) dx dy \quad (4.21)$$

for all $t \in [0, T^*)$.

Proof. By the support property of χ , we have $\phi_R(\tau) = 0$ for all $|\tau| \geq 2R$. Thus we get

$$\psi_R(x)|x| \leq \int_0^{2R} \phi_R(\tau) d\tau = \frac{R}{\omega_3} \int_0^2 \int \chi^2(x-z) \chi^2(z) dz d\tau = CR \quad (4.22)$$

for some constant $C > 0$ independent of R . The estimate (4.16) follows directly from Hölder's inequality and (4.22). The identities (4.17)–(4.21) follow from a direct computation using

$$\partial_t(|u|^2) = -2 \sum_j \partial_j (\text{Im}(\bar{u} \partial_j u))$$

and

$$\begin{aligned} \partial_t (\text{Im}(\bar{u} \partial_j u)) &= - \sum_k \partial_k \left(2 \text{Re}(\partial_j u \partial_k \bar{u}) - \frac{1}{2} \delta_{jk} \Delta(|u|^2) \right) \\ &\quad + \frac{p-1}{p+1} \partial_j(|u|^{p+1}) - \frac{q-1}{q+1} \partial_j(|u|^{q+1}), \end{aligned}$$

for $j = 1, 2, 3$, where δ_{jk} is the Kronecker symbol. \square

We are now in position to prove our interaction Morawetz estimates, that jointly to the scattering criterion of the previous section will yield to the main result of the paper. The coercivity result in Lemma 4.2 is essential for the proof of the estimates below.

Proposition 4.4. *Let $\frac{7}{3} < q < p < 5$ and $\omega > 0$. Let $u_0 \in \mathcal{A}_\omega^+$ and $u(t)$ be the corresponding solution to (1.2). Define $\mathcal{M}_R^{\otimes 2}(t)$ as in (4.15). Then for $\varepsilon > 0$ sufficiently small, there exist $T_0 = T_0(\varepsilon)$, $J = J(\varepsilon)$, $R_0 = R_0(\varepsilon, u_0)$ sufficiently large, and $\eta = \eta(\varepsilon) > 0$ sufficiently small such that for any $a \in \mathbb{R}$,*

$$\frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \frac{1}{R^3} \iiint |\chi_R(y-z)u(t,y)|^2 |\nabla(\chi_R(x-z)u^\xi(t,x))|^2 dx dy dz \frac{dR}{R} dt \lesssim \varepsilon, \quad (4.23)$$

where $\chi_R(x) = \chi(x/R)$ with χ as in (4.1) and $u^\xi(t,x) = e^{ix \cdot \xi} u(t,x)$ with some $\xi = \xi(t, z, R) \in \mathbb{R}^3$.

Proof. Denote

$$P_{jk}(x) := \delta_{jk} - \frac{x_j x_k}{|x|^2}.$$

As

$$\partial_j(\psi_R x_k) = \delta_{jk} \psi_R + \frac{x_j x_k}{|x|^2} (\phi_R - \psi_R),$$

the integration by parts yields

$$\begin{aligned} (4.17) &= 4 \sum_{j,k} \iint \operatorname{Im}(\bar{u}(t,y) \partial_j u(t,y)) \partial_j^y(\psi_R(x-y)(x_k - y_k)) \operatorname{Im}(\bar{u}(t,x) \partial_k u(t,x)) dx dy \\ &= -4 \sum_{j,k} \iint \operatorname{Im}(\bar{u}(t,y) \partial_j u(t,y)) \delta_{jk} \phi_R(x-y) \operatorname{Im}(\bar{u}(t,x) \partial_k u(t,x)) dx dy \end{aligned} \quad (4.24)$$

$$\begin{aligned} &- 4 \sum_{j,k} \iint \operatorname{Im}(\bar{u}(t,y) \partial_j u(t,y)) P_{jk}(x-y) \\ &\quad \times (\psi_R - \phi_R)(x-y) \operatorname{Im}(\bar{u}(t,x) \partial_k u(t,x)) dx dy, \end{aligned} \quad (4.25)$$

where ∂_j^y is ∂_j with respect to the y -variable. Similarly, we have

$$\begin{aligned} (4.18) &= 4 \sum_{j,k} \iint |u(t,y)|^2 \partial_k^x(\psi_R(x-y)(x_j - y_j)) \operatorname{Re}(\partial_j u(t,x) \partial_k \bar{u}(t,x)) dx dy \\ &= 4 \sum_{j,k} \iint |u(t,y)|^2 \delta_{jk} \phi_R(x-y) \operatorname{Re}(\partial_j u(t,x) \partial_k \bar{u}(t,x)) dx dy \end{aligned} \quad (4.26)$$

$$\begin{aligned} &- 4 \sum_{j,k} \iint |u(t,y)|^2 P_{jk}(x-y) (\psi_R - \phi_R)(x-y) \operatorname{Re}(\partial_j u(t,x) \partial_k \bar{u}(t,x)) dx dy, \end{aligned} \quad (4.27)$$

where ∂_k^x is ∂_k with respect to the x -variable. We have

$$\begin{aligned} (4.25) + (4.27) &= 4 \iint |u(t,y)|^2 |\nabla_y u(t,x)|^2 (\psi_R - \phi_R)(x-y) dx dy \\ &\quad - 4 \iint \operatorname{Im}(\bar{u}(t,y) \nabla_x u(t,y)) \cdot \operatorname{Im}(\bar{u}(t,x) \nabla_y u(t,x)) (\psi_R - \phi_R)(x-y) dx dy, \end{aligned}$$

where

$$\nabla_y u(t,x) := \nabla u(t,x) - \frac{x-y}{|x-y|} \left(\frac{x-y}{|x-y|} \nabla u(t,x) \right)$$

is the angular derivative centered at y , and similarly for $\nabla_x u(t, y)$. As $\psi_R - \phi_R \geq 0$, the Cauchy-Schwarz inequality yields

$$(4.25) + (4.27) \geq 0.$$

Next, using the fact that

$$\phi_R(x - y) = \frac{1}{\omega_3 R^3} \int \chi_R^2(x - y - z) \chi_R^2(z) dz = \frac{1}{\omega_3 R^3} \int \chi_R^2(x - z) \chi_R^2(y - z) dz,$$

we have

$$\begin{aligned} (4.24) + (4.26) &= 4 \iint \phi_R(x - y) (|u(t, y)|^2 |\nabla u(t, x)|^2 \\ &\quad - \operatorname{Im}(\bar{u}(t, y) \nabla u(t, y)) \cdot \operatorname{Im}(\bar{u}(t, x) \nabla u(t, x))) dx dy \\ &= \frac{4}{\omega_3 R^3} \iiint \chi_R^2(x - z) \chi_R^2(y - z) (|u(t, y)|^2 |\nabla u(t, x)|^2 \\ &\quad - \operatorname{Im}(\bar{u}(t, y) \nabla u(t, y)) \cdot \operatorname{Im}(\bar{u}(t, x) \nabla u(t, x))) dx dy dz. \end{aligned}$$

For $z \in \mathbb{R}^3$ fixed, we consider the quantity defined by

$$\begin{aligned} &\iint \chi_R^2(x - z) \chi_R^2(y - z) (|u(t, y)|^2 |\nabla u(t, x)|^2 \\ &\quad - \operatorname{Im}(\bar{u}(t, y) \nabla u(t, y)) \cdot \operatorname{Im}(\bar{u}(t, x) \nabla u(t, x))) dx dy. \end{aligned}$$

It is not hard to see that the above quantity is invariant under the Galilean transformation $u(t, x) \mapsto u^\xi(t, x)$ for all $\xi \in \mathbb{R}^3$ due to the symmetry of χ . We will choose a suitable $\xi \in \mathbb{R}^3$ such that

$$\int \operatorname{Im}(\chi_R^2(x - z) \bar{u}^\xi(t, x) \nabla(u^\xi(t, x))) dx = 0.$$

Specifically, we select ξ as follows (compare with (4.6)):

$$\xi = \xi(t, z, R) = - \frac{\int \operatorname{Im}(\chi_R^2(x - z) \bar{u}(t, x) \nabla u(t, x)) dx}{\int \chi_R^2(x - z) |u(t, x)|^2 dx},$$

provided that the denominator is non-zero (otherwise $\xi = 0$ suffices). With this ξ , we get

$$(4.24) + (4.26) = \frac{4}{\omega_3 R^3} \iiint |\chi_R(y - z) u(t, y)|^2 |\chi_R(x - z) \nabla(u^\xi(t, x))|^2 dx dy dz.$$

By integration by parts twice and using

$$\sum_j \partial_j(\psi_R x_j) = 3\psi_R + \sum_j x_j \partial_j \psi_R = 3\phi_R + 2(\psi_R - \phi_R),$$

we have

$$\begin{aligned} (4.19) &= \sum_{j,k} \iint |u(t, y)|^2 \psi_R(x - y) (x_j - y_j) \partial_j \partial_k^2 (|u(t, x)|^2) dx dy \\ &= - \sum_{j,k} \iint |u(t, y)|^2 \partial_j^x (\psi_R(x - y) (x_j - y_j)) \partial_k^2 (|u(t, x)|^2) dx dy \\ &= \sum_k \iint |u(t, y)|^2 \partial_k^x (3\phi_R(x - y) + 2(\psi_R - \phi_R)(x - y)) \partial_k (|u(t, x)|^2) dx dy. \end{aligned}$$

We also have

$$\begin{aligned}
(4.20) &= -\frac{2(p-1)}{p+1} \sum_j \iint |u(t, y)|^2 \partial_j^x (\psi_R(x-y)(x_j - y_j)) |u(t, x)|^{p+1} dx dy \\
&= -\frac{6(p-1)}{p+1} \iint |u(t, y)|^2 \phi_{p,R}(x-y) |u(t, x)|^{p+1} dx dy \\
&\quad - \frac{6(p-1)}{p+1} \iint |u(t, y)|^2 (\phi_R - \phi_{p,R})(x-y) |u(t, x)|^{p+1} dx dy \\
&\quad - \frac{4(p-1)}{p+1} \iint |u(t, y)|^2 (\psi_R - \phi_R)(x-y) |u(t, x)|^{p+1} dx dy.
\end{aligned} \tag{4.28}$$

We can rewrite (4.28) as

$$(4.28) = -\frac{6(p-1)}{(p+1)\omega_3 R^3} \iiint |\chi_R(y-z)u(t, y)|^2 |\chi_R(x-z)u(t, x)|^{p+1} dx dy dz.$$

Similarly, we have

$$\begin{aligned}
(4.21) &= \frac{6(q-1)}{(q+1)\omega_3 R^3} \iiint |\chi_R(y-z)u(t, y)|^2 |\chi_R(x-z)u(t, x)|^{q+1} dx dy dz \\
&\quad + \frac{6(q-1)}{q+1} \iint |u(t, y)|^2 (\phi_R - \phi_{q,R})(x-y) |u(t, x)|^{q+1} dx dy \\
&\quad + \frac{4(q-1)}{q+1} \iint |u(t, y)|^2 (\psi_R - \phi_R)(x-y) |u(t, x)|^{q+1} dx dy.
\end{aligned}$$

Collecting the above identities, we obtain

$$\begin{aligned}
\frac{d}{dt} \mathcal{M}_R^{\otimes 2}(t) &\geq \frac{4}{\omega_3 R^3} \iiint |\chi_R(y-z)u(t, y)|^2 |\chi_R(x-z)\nabla(u^\xi(t, x))|^2 dx dy dz \\
&\quad + \iint |u(t, y)|^2 \nabla(3\phi_R(x-y) + 2(\psi_R - \phi_R)(x-y)) \cdot \nabla(|u(t, x)|^2) dx dy \\
&\quad - \frac{6(p-1)}{(p+1)\omega_3 R^3} \iiint |\chi_R(y-z)u(t, y)|^2 |\chi_R(x-z)u(t, x)|^{p+1} dx dy dz \\
&\quad - \frac{6(p-1)}{p+1} \iint |u(t, y)|^2 (\phi_R - \phi_{p,R})(x-y) |u(t, x)|^{p+1} dx dy \\
&\quad - \frac{4(p-1)}{p+1} \iint |u(t, y)|^2 (\psi_R - \phi_R)(x-y) |u(t, x)|^{p+1} dx dy \\
&\quad + \frac{6(q-1)}{(q+1)\omega_3 R^3} \iiint |\chi_R(y-z)u(t, y)|^2 |\chi_R(x-z)u(t, x)|^{q+1} dx dy dz \\
&\quad + \frac{6(q-1)}{q+1} \iint |u(t, y)|^2 (\phi_R - \phi_{q,R})(x-y) |u(t, x)|^{q+1} dx dy \\
&\quad + \frac{4(q-1)}{q+1} \iint |u(t, y)|^2 (\psi_R - \phi_R)(x-y) |u(t, x)|^{q+1} dx dy.
\end{aligned}$$

As $\psi_R - \phi_R \geq 0$ and $\phi_R - \phi_{q,R} \geq 0$, we get

$$\begin{aligned}
& \frac{4}{\omega_3 R^3} \iiint |\chi_R(y-z)u(t,y)|^2 \left(|\chi_R(x-z)\nabla(u^\xi(t,x))|^2 \right. \\
& \quad \left. - \frac{3(p-1)}{2(p+1)} |\chi_R(x-z)u(t,x)|^{p+1} + \frac{3(q-1)}{2(q+1)} |\chi_R(x-z)u(t,x)|^{q+1} \right) dx dy dz \\
& \leq \frac{d}{dt} \mathcal{M}_R^{\otimes 2}(t) - \iint |u(t,y)|^2 \nabla (3\phi_R(x-y) + 2(\psi_R - \phi_R)(x-y)) \cdot \nabla (|u(t,x)|^2) dx dy \\
& \quad + \frac{6(p-1)}{p+1} \iint |u(t,y)|^2 (\phi_R - \phi_{p,R})(x-y) |u(t,x)|^{p+1} dx dy \\
& \quad + \frac{4(p-1)}{p+1} \iint |u(t,y)|^2 (\psi_R - \phi_R)(x-y) |u(t,x)|^{p+1} dx dy.
\end{aligned}$$

By (4.16), we see that

$$\begin{aligned}
\left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \frac{d}{dt} \mathcal{M}_R^{\otimes 2}(t) \frac{dR}{R} dt \right| & \leq \frac{1}{JT_0} \int_{R_0}^{R_0 e^J} \sup_{t \in [a, a+T_0]} |\mathcal{M}_R^{\otimes 2}(t)| \frac{dR}{R} \\
& \lesssim \frac{1}{JT_0} \int_{R_0}^{R_0 e^J} dR \lesssim \frac{R_0 e^J}{JT_0}.
\end{aligned} \tag{4.29}$$

As $|\nabla \phi_R(x)| \lesssim \frac{1}{\eta R}$ and $\sup_{t \in \mathbb{R}} \|u(t)\|_{H^1} < \infty$, we see that

$$\begin{aligned}
& \left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \iint |u(t,y)|^2 \nabla \phi_R(x-y) \cdot \nabla (|u(t,x)|^2) dx dy \frac{dR}{R} dt \right| \\
& \lesssim \frac{1}{\eta JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \|u(t)\|_{L^2}^3 \|\nabla u(t)\|_{L^2} \frac{dR}{R^2} dt \\
& \lesssim \frac{1}{\eta JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \frac{dR}{R^2} dt \\
& \lesssim \frac{1}{\eta J R_0}.
\end{aligned} \tag{4.30}$$

Similarly, as $|\nabla(\psi_R - \phi_R)(x)| \lesssim \frac{1}{\eta} \min \left\{ \frac{1}{R}, \frac{R}{|x|^2} \right\} < \frac{1}{\eta R}$, we have

$$\left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \iint |u(t,y)|^2 \nabla(\psi_R - \phi_R)(x-y) \cdot \nabla (|u(t,x)|^2) dx dy \frac{dR}{R} dt \right| \lesssim \frac{1}{\eta J R_0}. \tag{4.31}$$

Using (4.4), we see that

$$\begin{aligned}
& \left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \iint |u(t,y)|^2 (\psi_R - \phi_R)(x-y) |u(t,x)|^{p+1} dx dy \frac{dR}{R} dt \right| \\
& \lesssim \frac{1}{\eta JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \iint |u(t,y)|^2 \min \left\{ \frac{|x-y|}{R}, \frac{R}{|x-y|} \right\} |u(t,x)|^{p+1} dx dy \frac{dR}{R} dt \\
& \lesssim \frac{1}{\eta JT_0} \int_a^{a+T_0} \iint |u(t,y)|^2 |u(t,x)|^{p+1} \left(\int_{R_0}^{R_0 e^J} \min \left\{ \frac{|x-y|}{R}, \frac{R}{|x-y|} \right\} \frac{dR}{R} \right) dx dy dt \\
& \lesssim \frac{1}{\eta J}.
\end{aligned} \tag{4.32}$$

Here we have used the fact that $\sup_{t \in \mathbb{R}} \|u(t)\|_{H^1} < \infty$ and

$$\int_0^\infty \min \left\{ \frac{|x-y|}{R}, \frac{R}{|x-y|} \right\} \frac{dR}{R} \lesssim 1.$$

Using (4.3), we have

$$\begin{aligned} & \left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \iint |u(t, y)|^2 (\phi_R - \phi_{p,R})(x-y) |u(t, x)|^{p+1} dx dy \frac{dR}{R} dt \right| \\ & \lesssim \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \eta \frac{dR}{R} dt \lesssim \eta. \end{aligned} \quad (4.33)$$

By glueing up together (4.29), (4.30), (4.31), (4.32), and (4.33), we obtain

$$\begin{aligned} & \left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \frac{1}{R^3} \iiint |\chi_R(y-z)u(t, y)|^2 (|\chi_R(x-z)\nabla(u^\xi(t, x))|^2 \right. \\ & \quad \left. - \frac{3(p-1)}{2(p+1)} |\chi_R(x-z)u(t, x)|^{p+1} + \frac{3(q-1)}{2(q+1)} |\chi_R(x-z)u(t, x)|^{q+1}) dx dy dz \frac{dR}{R} dt \right| \\ & \lesssim \eta + \frac{R_0 e^J}{\eta J T_0} + \frac{1}{\eta J} + \frac{1}{\eta J R_0}. \end{aligned} \quad (4.34)$$

Now, for fixed $z, \xi \in \mathbb{R}^3$, we have from (4.9) that

$$\int |\chi_R(x-z)\nabla(u^\xi(t, x))|^2 dx = \int |\nabla[\chi_R(x-z)u^\xi(t, x)]|^2 dx + O(R^{-2}\|u(t)\|_{L^2}^2).$$

From the conservation of mass and (4.5) that for $R \geq R_0$ with R_0 sufficiently large,

$$\begin{aligned} & \int |\chi_R(x-z)\nabla(u^\xi(t, x))|^2 - \frac{3(p-1)}{2(p+1)} |\chi_R(x-z)u(t, x)|^{p+1} \\ & \quad + \frac{3(q-1)}{2(q+1)} |\chi_R(x-z)u(t, x)|^{q+1} dx \\ & = G(\chi_R(\cdot-z)u^\xi(t)) + O(R^{-2}) \\ & \geq \delta \|\nabla(\chi_R(\cdot-z)u^\xi(t))\|_{L^2}^2 + O(R^{-2}). \end{aligned}$$

The term $O(R^{-2})$ can be treated analogously to (4.30). We thus infer from (4.34) that

$$\begin{aligned} & \left| \frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \frac{1}{R^3} \iiint |\chi_R(y-z)u(t, y)|^2 |\nabla(\chi_R(x-z)u^\xi(t, x))|^2 dx dy dz \frac{dR}{R} dt \right| \\ & \lesssim \eta + \frac{R_0 e^J}{\eta J T_0} + \frac{1}{\eta J} + \frac{1}{\eta J R_0}. \end{aligned} \quad (4.35)$$

This proves (4.23) by taking $\eta = \varepsilon, J = \varepsilon^{-3}, R_0 = \varepsilon^{-1}$ and $T_0 = e^{\varepsilon^{-3}}$. The proof is complete. \square

4.4. Proof of the main result. We can now proceed with the proof of the main result.

Proof of Theorem 1.1. The global existence is proved in Lemma 4.2. It remains to prove the scattering. We only consider the positive times since the one for negative times is similar. Our purpose is to check the scattering criteria given in Lemma 3.6. To this end, we fix $a \in \mathbb{R}$ and let $\varepsilon > 0$ sufficiently small and $T_0 > 0$ sufficiently large to be determined later. We will show that there exists $t_0 \in (a, a + T_0)$ such that $[t_0 - \varepsilon^{-\sigma}, t_0] \subset (a, a + T_0)$ and

$$\|u\|_{L_t^{m_1}([t_0 - \varepsilon^{-\sigma}, t_0], L_x^{b_1})} \lesssim \varepsilon^\mu \quad (4.36)$$

for some $\sigma, \mu > 0$ to be determined later. By (4.23), there exist $T_0 = T_0(\varepsilon)$, $J = J(\varepsilon)$, $R_0 = R_0(\varepsilon, u_0)$ and $\eta = \eta(\varepsilon)$ such that

$$\frac{1}{JT_0} \int_a^{a+T_0} \int_{R_0}^{R_0 e^J} \frac{1}{R^3} \iiint |\chi_R(y-z)u(t,y)|^2 |\nabla(\chi_R(x-z)u^\xi(t,x))|^2 dx dy dz \frac{dR}{R} dt \lesssim \varepsilon.$$

It follows that there exists $R_1 \in [R_0, R_0 e^J]$ such that

$$\frac{1}{T_0} \int_a^{a+T_0} \frac{1}{R_1^3} \iiint |\chi_{R_1}(y-z)u(t,y)|^2 |\nabla(\chi_{R_1}(x-z)u^\xi(t,x))|^2 dx dy dz dt \lesssim \varepsilon$$

hence

$$\frac{1}{T_0} \int_a^{a+T_0} \frac{1}{R_1^3} \int \|\chi_{R_1}(\cdot-z)u(t)\|_{L_x^2}^2 \|\nabla(\chi_{R_1}(\cdot-z)u^\xi(t))\|_{L_x^2}^2 dz dt \lesssim \varepsilon.$$

By the change of variable $z = \frac{R_1}{4}(w + \theta)$ with $w \in \mathbb{Z}^3$ and $\theta \in [0, 1]^3$, we deduce that there exists $\theta_1 \in [0, 1]^3$ such that

$$\frac{1}{T_0} \int_a^{a+T_0} \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^2}^2 \left\| \nabla \left(\chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u^\xi(t) \right) \right\|_{L_x^2}^2 dt \lesssim \varepsilon.$$

Let $\sigma > 0$ to be chosen later. By dividing the interval $[a + \frac{T_0}{2}, a + \frac{3T_0}{4}]$ into $T_0 \varepsilon^\sigma$ intervals of length $\varepsilon^{-\sigma}$, we infer that there exists $t_0 \in [a, \frac{T_0}{2}, a + \frac{3T_0}{4}]$ such that $[t_0 - \varepsilon^{-\sigma}, t_0] \subset (a, a + T_0)$ and

$$\int_{t_0 - \varepsilon^{-\sigma}}^{t_0} \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^2}^2 \left\| \nabla \left(\chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u^\xi(t) \right) \right\|_{L_x^2}^2 dt \lesssim \varepsilon^{1-\sigma}.$$

This, together with the Gagliardo-Nirenberg inequality

$$\|u\|_{L_x^3}^4 \lesssim \|u\|_{L_x^2}^2 \|\nabla u^\xi\|_{L_x^2}^2,$$

implies that

$$\int_{t_0 - \varepsilon^{-\sigma}}^{t_0} \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^4 dt \lesssim \varepsilon^{1-\sigma}. \quad (4.37)$$

On the other hand, by Hölder's inequality, Cauchy-Schwarz inequality, and Sobolev embedding, we have

$$\begin{aligned} & \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^2 \\ & \leq \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^2} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^6} \\ & \leq \left(\sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^2}^2 \right)^{1/2} \left(\sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^6}^2 \right)^{1/2} \\ & \lesssim \|u(t)\|_{L_x^2} \|u(t)\|_{H_x^1} \lesssim 1. \end{aligned} \quad (4.38)$$

Here we have used the following estimate to get the last line:

$$\begin{aligned}
& \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^6}^2 \\
& \lesssim \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) \nabla u(t) \right\|_{L_x^2}^2 + \frac{1}{R_1^2} \left\| (\nabla \chi)_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^2}^2 \\
& \lesssim \|\nabla u(t)\|_{L_x^2}^2 + \frac{1}{R_1^2 \eta^2} \|u(t)\|_{L_x^2}^2 \lesssim \|u(t)\|_{H_x^1}^2
\end{aligned}$$

as $R_1 > R_0 = \varepsilon^{-1} = \eta^{-1}$ (see after (4.35)). Note that $|\nabla \chi| \lesssim \eta^{-1}$ by the choice of χ . Combining (4.37) and (4.38), we get from the property of χ_{R_1} , in conjunction with the Hölder and the Cauchy-Schwarz inequalities, that

$$\begin{aligned}
\|u\|_{L_{t,x}^3([t_0 - \varepsilon^{-\sigma}, t_0] \times \mathbb{R}^3)}^3 & \lesssim \int_{t_0 - \varepsilon^{-\sigma}}^{t_0} \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^3 dt \\
& \lesssim \int_{t_0 - \varepsilon^{-\sigma}}^{t_0} \left(\sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^4 \right)^{\frac{1}{2}} \\
& \quad \times \left(\sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^2 \right)^{\frac{1}{2}} dt \\
& \lesssim \left(\int_{t_0 - \varepsilon^{-\sigma}}^{t_0} \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^4 dt \right)^{\frac{1}{2}} \\
& \quad \times \left(\int_{t_0 - \varepsilon^{-\sigma}}^{t_0} \sum_{w \in \mathbb{Z}^3} \left\| \chi_{R_1} \left(\cdot - \frac{R_1}{4}(w + \theta_1) \right) u(t) \right\|_{L_x^3}^2 dt \right)^{\frac{1}{2}} \\
& \lesssim \varepsilon^{\frac{1-\sigma}{2}} \varepsilon^{-\frac{\sigma}{2}} = \varepsilon^{\frac{1}{2}-\sigma},
\end{aligned}$$

which implies that

$$\|u\|_{L_{t,x}^3([t_0 - \varepsilon^{-\sigma}, t_0] \times \mathbb{R}^3)} \lesssim \varepsilon^{\frac{1}{3}(\frac{1}{2}-\sigma)}. \quad (4.39)$$

Let $\theta \in (0, 1)$ to be chosen shortly. We define (γ, ρ) by

$$\frac{1}{m_1} = \frac{\theta}{3} + \frac{1-\theta}{\gamma}, \quad \frac{1}{b_1} = \frac{\theta}{3} + \frac{1-\theta}{\rho}.$$

Pick $\beta, s > 0$ such that

$$\frac{1}{\rho} = \frac{1}{\beta} - \frac{s}{3}, \quad \frac{2}{\gamma} + \frac{3}{\beta} = \frac{3}{2}.$$

We readily check that

$$\frac{2}{m_1} + \frac{3}{b_1} = \frac{2}{p-1} = \frac{5\theta}{3} + (1-\theta) \left(\frac{3}{2} - s \right).$$

In particular,

$$s = \frac{3}{2} - \frac{1}{1-\theta} \left(\frac{2}{p-1} - \frac{5\theta}{3} \right).$$

As $7/3 < p < 5$, we can take $\theta > 0$ sufficiently small so that $0 < s < 1$. In particular, (γ, β) is a Schrödinger admissible pair. By Hölder's inequality, Sobolev embedding, and

(4.39), we have

$$\begin{aligned}
\|u\|_{L_t^{m_1}([t_0-\varepsilon^{-\sigma}, t_0], L_x^{b_1})} &\leq \|u\|_{L_{t,x}^3([t_0-\varepsilon^{-\sigma}, t_0] \times \mathbb{R}^3)}^\theta \|u\|_{L_t^\gamma([t_0-\varepsilon^{-\sigma}, t_0], L_x^\rho)}^{1-\theta} \\
&\lesssim \|u\|_{L_{t,x}^3([t_0-\varepsilon^{-\sigma}, t_0] \times \mathbb{R}^3)}^\theta \|\langle \nabla \rangle^s u\|_{L_t^\gamma([t_0-\varepsilon^{-\sigma}, t_0], L_x^\beta)}^{1-\theta} \\
&\lesssim \|u\|_{L_{t,x}^3([t_0-\varepsilon^{-\sigma}, t_0] \times \mathbb{R}^3)}^\theta \|\langle \nabla \rangle^s u\|_{L_t^\gamma([t_0-\varepsilon^{-\sigma}, t_0], L_x^\beta)}^{1-\theta} \\
&\lesssim \varepsilon^{\frac{\theta}{3}(\frac{1}{2}-\sigma)} \varepsilon^{-\frac{\sigma}{\gamma}(1-\theta)} \\
&\lesssim \varepsilon^{\frac{\theta}{6} - (\frac{\theta}{3} + \frac{1-\theta}{\gamma})\sigma}.
\end{aligned}$$

Here we have used the fact that

$$\|\langle \nabla \rangle u\|_{L_t^\gamma(I, L_x^\beta)} \lesssim \langle I \rangle^{\frac{1}{\gamma}}$$

which follows from the local theory. This proves (4.36) by choosing $\sigma > 0$ small enough. The proof is complete. \square

5. BLOW-UP

In this last section, we prove the blow-up rate results as stated in Theorem 1.3. We start with the following upper bound for the Pohozaev functional for solutions arising from initial data in \mathcal{A}_ω^- . The result of the next Lemma is also contained in [4, 16], but we report the short proof for sake of completeness.

Lemma 5.1. *Let $\frac{7}{3} < p < 5$, $1 < q < p$, and $\omega > 0$. Then \mathcal{A}_ω^- is invariant under the flow of (1.2), i.e., if $u_0 \in \mathcal{A}_\omega^-$, then $u(t) \in \mathcal{A}_\omega^-$ for all $t \in I_{\max}$. In addition, we have*

$$G(u(t)) \leq -\frac{3(p-1)}{2}(m_\omega - S_\omega(u_0)), \quad \forall t \in I_{\max}. \quad (5.1)$$

Furthermore, there exists $\delta > 0$ small such that

$$G(u(t)) + \delta \|\nabla u(t)\|_{L^2}^2 \leq -\frac{3(p-1)}{4}(m_\omega - S_\omega(u_0)), \quad \forall t \in I_{\max}. \quad (5.2)$$

Proof. Let $u_0 \in \mathcal{A}_\omega^-$. We will show that $u(t) \in \mathcal{A}_\omega^-$ for all $t \in I_{\max}$. By the conservation of mass and energy, we have

$$S_\omega(u(t)) = S_\omega(u_0) < m_\omega, \quad \forall t \in I_{\max}. \quad (5.3)$$

Assume by contradiction that there exists t_0 such that $G(u(t_0)) \geq 0$. As $u : I_{\max} \rightarrow H^1$ is continuous, there exists t_1 such that $G(u(t_1)) = 0$. By the definition of m_ω , we have $S_\omega(u(t_1)) \geq m_\omega$ which contradicts (5.3). Thus $G(u(t)) < 0$ for all $t \in I_{\max}$ or \mathcal{A}_ω^- is invariant under the flow of (1.2).

For simplicity, we denote $u := u(t)$ and set $f(\lambda) := S_\omega(u_\lambda)$, where u_λ is the scaling (2.1). We have

$$f'(\lambda) = \lambda \|\nabla u\|_{L^2}^2 - \frac{3(p-1)}{2(p+1)} \lambda^{\frac{3(p-1)}{2}-1} \|u\|_{L^{p+1}}^{p+1} + \frac{3(q-1)}{2(q+1)} \lambda^{\frac{3(q-1)}{2}-1} \|u\|_{L^{q+1}}^{q+1} = \frac{G(u_\lambda)}{\lambda}$$

and

$$(\lambda f'(\lambda))' = 2\lambda \|\nabla u\|_{L^2}^2 - \frac{9(p-1)^2}{4(p+1)} \lambda^{\frac{3(p-1)}{2}-1} \|u\|_{L^{p+1}}^{p+1} + \frac{9(q-1)^2}{4(q+1)} \lambda^{\frac{3(q-1)}{2}-1} \|u\|_{L^{q+1}}^{q+1}.$$

We then write

$$(\lambda f'(\lambda))' = \frac{3(p-1)}{2} f'(\lambda) - \frac{3p-7}{2} \lambda \|\nabla u\|_{L^2}^2 - \frac{9(q-1)(p-q)}{2(q+1)} \|u\|_{L^{q+1}}^{q+1}.$$

As $\frac{7}{3} < p < 5$ and $1 < q < p$, we get

$$(\lambda f'(\lambda))' \leq \frac{3(p-1)}{2} f'(\lambda), \quad \forall \lambda > 0. \quad (5.4)$$

On the other hand, as $G(u) < 0$, by Lemma 2.2, there exists $\lambda_0 \in (0, 1)$ such that $G(u_{\lambda_0}) = 0$. In particular, we have $\lambda_0 f'(\lambda_0) = 0$ and $f(\lambda_0) = S_\omega(u_{\lambda_0}) \geq m_\omega$. Integrating (5.4) over $(\lambda_0, 1)$, we obtain

$$G(u) \leq \frac{3(p-1)}{2} (S_\omega(u) - S_\omega(u_{\lambda_0})) \leq \frac{3(p-1)}{2} (S_\omega(u) - m_\omega)$$

which is (5.1).

Finally, we prove (5.2). Observe that

$$\begin{aligned} \|\nabla u\|_{L^2}^2 &= \frac{6(q-1)}{3q-7} \left(I_\omega(u) - \frac{p-q}{(p+1)(q-1)} \|u\|_{L^{p+1}}^{p+1} - \frac{\omega}{2} \|u\|_{L^2}^2 \right) \\ &= \frac{6(q-1)}{3q-7} \left(S_\omega(u) - \frac{2}{3(q-1)} G(u) - \frac{p-q}{(p+1)(q-1)} \|u\|_{L^{p+1}}^{p+1} - \frac{\omega}{2} \|u\|_{L^2}^2 \right). \end{aligned}$$

It follows that

$$\begin{aligned} G(u) + \delta \|\nabla u\|_{L^2}^2 &= \left(1 - \frac{4\delta}{3q-7} \right) G(u) + \frac{6\delta(q-1)}{3q-7} S_\omega(u) \\ &\quad - \frac{6\delta(p-q)}{(3q-7)(p+1)} \|u\|_{L^{p+1}}^{p+1} - \frac{3\delta\omega(q-1)}{3q-7} \|u\|_{L^2}^2 \\ &\leq \left(1 - \frac{4\delta}{3q-7} \right) G(u) + \frac{6\delta(q-1)}{3q-7} S_\omega(u) \end{aligned}$$

as $\omega > 0$ and $q < p$. By the energy and mass conservation laws, and (5.1), we get

$$G(u(t)) + \delta \|\nabla u(t)\|_{L^2}^2 \leq - \left(1 - \frac{4\delta}{3q-7} \right) \frac{3(p-1)}{2} (m_\omega - S_\omega(u_0)) + \frac{6\delta(q-1)}{3q-7} S_\omega(u_0),$$

$\forall t \in I_{\max}$. By taking $\delta > 0$ sufficiently small, we obtain (5.2). \square

We are now ready to give the proof of the blow-up results in Theorem 1.3. They are based on virial identities/estimates. We classically introduce a sufficiently smooth and decaying function $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}$. We denote the virial quantity

$$V_\phi(t) := \int \phi |u(t)|^2 dx.$$

The following identities are nowadays standard (see e.g., [6]):

$$V'_\phi(t) = 2 \operatorname{Im} \int \nabla \phi \cdot \nabla u(t) \bar{u}(t) dx$$

and

$$\begin{aligned} V''_\phi(t) &= - \int \Delta^2 \phi |u(t)|^2 dx + 4 \sum_{j,k} \operatorname{Re} \int \partial_{jk}^2 \phi \partial_j u(t) \partial_k \bar{u}(t) dx \\ &\quad + \frac{2(q-1)}{q+1} \int \Delta \phi |u(t)|^{q+1} dx - \frac{2(p-1)}{p+1} \int \Delta \phi |u(t)|^{p+1} dx. \end{aligned}$$

A first application of the above virial identities is the following virial identity for finite variance solutions. More precisely, if $u_0 \in \Sigma$, then the corresponding solution to (1.2) satisfies

$$V''_{|x|^2}(t) = 8G(u(t)), \quad \forall t \in I_{\max}. \quad (5.5)$$

Provided $u_0 \in \mathcal{A}_\omega^-$, by Lemma 5.1 $G(u(t)) \lesssim -1$ for any $t \in I_{\max}$. Then (5.5) implies finite time blow-up for solutions in Σ via a convexity argument.

Another applications are virial estimates for radial and cylindrical solutions [4]. To state these estimates, we let $\theta : [0, \infty) \rightarrow [0, 2]$ a smooth function satisfying

$$\theta(r) = \begin{cases} 2 & \text{if } 0 \leq r \leq 1, \\ 0 & \text{if } r \geq 2. \end{cases} \quad (5.6)$$

We define the function $\vartheta : [0, \infty) \rightarrow [0, \infty)$ by

$$\vartheta(r) := \int_0^r \int_0^s \theta(\tau) d\tau ds.$$

For $\varrho > 0$, we define the radial function $\phi_\varrho : \mathbb{R}^3 \rightarrow \mathbb{R}$ by

$$\phi_\varrho(x) = \phi_\varrho(r) := \varrho^2 \vartheta(r/\varrho), \quad r = |x|. \quad (5.7)$$

Then the following virial estimate for radial solutions: If $u_0 \in H^1$ is radial, then the corresponding solution to (1.2) satisfies

$$V''_{\phi_\varrho}(t) \leq 8G(u(t)) + C\varrho^{-2} + C\varrho^{-(p-1)} \|\nabla u(t)\|_{L^2}^{\frac{p-1}{2}}, \quad \forall t \in I_{\max}. \quad (5.8)$$

To state virial estimates for cylindrical solutions, we define, for $\varrho > 0$, the function

$$\phi_\varrho(x) := \varrho^2 \vartheta(r/\varrho) + x_3^2, \quad r := |\bar{x}|, \quad \bar{x} = (x_1, x_2). \quad (5.9)$$

If $u_0 \in \Sigma_3$, then the corresponding solution to (1.2) satisfies

$$V''_{\phi_\varrho}(t) \leq 8G(u(t)) + C\varrho^{-2} + C\varrho^{-\frac{p-1}{2}} \|\nabla u(t)\|_{L^2}^{p-1}, \quad \forall t \in I_{\max}. \quad (5.10)$$

We refer the readers to [4] for a proof of (5.8) and (5.10). We are now able to prove our blow-up result for (1.2).

Proof of Theorem 1.3. Let us start with the point (i). The proof is based on an idea of Merle, Raphaël, and Szeftel [30].

(1) Let us consider the radial case. Using

$$G(u) = \frac{3(p-1)}{2} E(u) - \frac{3p-7}{4} \|\nabla u\|_{L^2}^2 - \frac{3(p-q)}{2(q+1)} \|u\|_{L^{q+1}}^{q+1},$$

we infer from (5.8) that

$$\begin{aligned} V''_{\phi_\varrho}(t) &\leq 12(p-1)E(u(t)) - 2(3p-7)\|\nabla u(t)\|_{L^2}^2 - \frac{12(p-q)}{q+1} \|u(t)\|_{L^{q+1}}^{q+1} \\ &\quad + C\varrho^{-2} + C\varrho^{-(p-1)} \|\nabla u(t)\|_{L^2}^{\frac{p-1}{2}}, \quad \forall t \in I_{\max}, \end{aligned}$$

where ϕ_ϱ is as in (5.7). As $p < 5$, by Young's inequality, we have for any $\varepsilon > 0$,

$$V''_{\phi_\varrho}(t) \leq 12(p-1)E(u(t)) - 2(3p-7)\|\nabla u(t)\|_{L^2}^2 + C\varrho^{-2} + \varepsilon \|\nabla u(t)\|_{L^2}^2 + C\varepsilon^{-\frac{p-1}{5-p}} \varrho^{-\frac{4(p-1)}{5-p}},$$

$\forall t \in I_{\max}$. Taking $\varepsilon = 3p-7$, we get

$$V''_{\phi_\varrho}(t) \leq 12(p-1)E(u(t)) - (3p-7)\|\nabla u(t)\|_{L^2}^2 + C\varrho^{-2} + C\varrho^{-\frac{4(p-1)}{5-p}},$$

$\forall t \in I_{\max}$. By the conservation of energy and $2 < \frac{4(p-1)}{5-p}$, we obtain

$$(3p-7)\|\nabla u(t)\|_{L^2}^2 + V_{\phi_\varrho}''(t) \leq C\varrho^{-\frac{4(p-1)}{5-p}}, \quad (5.11)$$

$\forall t \in I_{\max}$ provided that $\varrho > 0$ is taken sufficiently small. Let $0 < t_0 < t < T^*$. We integrate (5.11) twice on (t_0, t) and get

$$\begin{aligned} (3p-7) \int_{t_0}^t \int_{t_0}^s \|\nabla u(\tau)\|_{L^2}^2 d\tau ds + V_{\phi_\varrho}(t) &\leq C\varrho^{-\frac{4(p-1)}{5-p}}(t-t_0)^2 + (t-t_0)V_{\phi_\varrho}'(t_0) + V_{\phi_\varrho}(t_0) \\ &\leq C\varrho^{-\frac{4(p-1)}{5-p}}(t-t_0)^2 + C\varrho(t-t_0)\|\nabla u(t_0)\|_{L^2} \\ &\quad + C\varrho^2, \end{aligned}$$

where we have used the conservation of mass and

$$\begin{aligned} V_{\phi_\varrho}(t_0) &\leq C\varrho^2\|u(t_0)\|_{L^2}^2 \leq C\varrho^2, \\ V_{\phi_\varrho}'(t_0) &\leq C\varrho\|\nabla u(t_0)\|_{L^2}\|u(t_0)\|_{L^2} \leq C\varrho\|\nabla u(t_0)\|_{L^2}. \end{aligned}$$

Note that the constant $C > 0$ may vary from line to line. By Fubini's Theorem, we have

$$\int_{t_0}^t \int_{t_0}^s \|\nabla u(\tau)\|_{L^2}^2 d\tau ds = \int_{t_0}^t \left(\int_{\tau}^t ds \right) \|\nabla u(\tau)\|_{L^2}^2 d\tau = \int_{t_0}^t (t-\tau)\|\nabla u(\tau)\|_{L^2}^2 d\tau.$$

As V_{ϕ_ϱ} is non-negative, we get

$$\int_{t_0}^t (t-\tau)\|\nabla u(\tau)\|_{L^2}^2 d\tau \leq C\varrho^{-\frac{4(p-1)}{5-p}}(t-t_0)^2 + C\varrho(t-t_0)\|\nabla u(t_0)\|_{L^2} + C\varrho^2.$$

Letting $t \nearrow T^*$, we obtain

$$\int_{t_0}^{T^*} (T^*-\tau)\|\nabla u(\tau)\|_{L^2}^2 d\tau \leq C\varrho^{-\frac{4(p-1)}{5-p}}(T^*-t_0)^2 + C\varrho(T^*-t_0)\|\nabla u(t_0)\|_{L^2} + C\varrho^2.$$

Optimizing in ϱ by choosing $\varrho^{-\frac{4(p-1)}{5-p}}(T^*-t_0)^2 = \varrho^2$ or equivalently $\varrho = (T^*-t_0)^{\frac{5-p}{p+3}}$, we have

$$\begin{aligned} \int_{t_0}^{T^*} (T^*-\tau)\|\nabla u(\tau)\|_{L^2}^2 d\tau &\leq C(T^*-t_0)^{\frac{2(5-p)}{p+3}} + C(T^*-t_0)^{\frac{5-p}{p+3}+1}\|\nabla u(t_0)\|_{L^2} \\ &\leq C(T^*-t_0)^{\frac{2(5-p)}{p+3}} + C(T^*-t_0)^2\|\nabla u(t_0)\|_{L^2}^2, \end{aligned}$$

for any $0 < t_0 < T^*$. Now set

$$g(t) := \int_t^{T^*} (T^*-\tau)\|\nabla u(\tau)\|_{L^2}^2 d\tau. \quad (5.12)$$

We have

$$g(t) \leq C(T^*-t)^{\frac{2(5-p)}{p+3}} - (T^*-t)g'(t), \quad \forall 0 < t < T^*$$

which is rewritten as

$$\left(\frac{g(t)}{T^*-t} \right)' = \frac{1}{(T^*-t)^2}(g(t) + (T^*-t)g'(t)) \leq \frac{C}{(T^*-t)^{\frac{4(p-1)}{p+3}}}.$$

Integrating it over $(0, t)$, we obtain

$$\frac{g(t)}{T^*-t} \leq \frac{g(0)}{T^*} + \frac{C}{(T^*-t)^{\frac{3p-7}{p+3}}} - \frac{C}{(T^*)^{\frac{3p-7}{p+3}}}$$

which yields

$$\frac{g(t)}{T^* - t} \leq \frac{C}{(T^* - t)^{\frac{3p-7}{p+3}}} \quad \text{for } t \text{ close to } T^*.$$

In particular, we have $g(t) \leq C(T^* - t)^{\frac{2(5-p)}{p+3}}$ which is (1.6).

To see (1.7), we rewrite (1.6) as

$$\frac{1}{T^* - t} \int_t^{T^*} (T^* - \tau) \|\nabla u(\tau)\|_{L^2}^2 d\tau \leq \frac{C}{(T^* - t)^{\frac{3p-7}{p+3}}}. \quad (5.13)$$

Take $T_n \nearrow T^*$. For a fixed n , g defined in (5.12) is a continuous function on $[T_n, T^*]$ and differentiable on (T_n, T^*) . By the mean value theorem, there exists $t_n \in (T_n, T^*)$ such that

$$-(T^* - t_n) \|\nabla u(t_n)\|_{L^2}^2 = g'(t_n) = \frac{g(T^*) - g(T_n)}{T^* - T_n} = -\frac{\int_{T_n}^{T^*} (T^* - \tau) \|\nabla u(\tau)\|_{L^2}^2 d\tau}{T^* - T_n}.$$

Using (5.13), we have

$$(T^* - t_n) \|\nabla u(t_n)\|_{L^2}^2 \leq \frac{C}{(T^* - T_n)^{\frac{3p-7}{p+3}}} \leq \frac{C}{(T^* - t_n)^{\frac{3p-7}{p+3}}}$$

hence

$$\|\nabla u(t_n)\|_{L^2}^2 \leq \frac{C}{(T^* - t_n)^{\frac{4(p-1)}{p+3}}}.$$

This proves (1.7).

(2) We now consider the cylindrical case. By (5.10), we have

$$\begin{aligned} V_{\phi_\varrho}''(t) &\leq 8G(u(t)) + C\varrho^{-2} + C\varrho^{-\frac{p-1}{2}} \|\nabla u(t)\|_{L^2}^{p-1} \\ &\leq 12(p-1)E(u(t)) - 2(3p-7) \|\nabla u(t)\|_{L^2}^2 - \frac{12(p-q)}{q+1} \|u(t)\|_{L^{q+1}}^{q+1} \\ &\quad + C\varrho^{-2} + C\varrho^{-\frac{p-1}{2}} \|\nabla u(t)\|_{L^2}^{p-1}, \quad \forall t \in I_{\max}, \end{aligned}$$

where ϕ_ϱ is as in (5.9). By Young's inequality with $p < 3$ and $q < p$, we have

$$\begin{aligned} V_{\phi_\varrho}''(t) &\leq 12(p-1)E(u(t)) - 2(3p-7) \|\nabla u(t)\|_{L^2}^2 + C\varrho^{-2} + \varepsilon \|\nabla u(t)\|_{L^2}^2 + C\varepsilon^{-\frac{p-1}{3-p}} \varrho^{-\frac{p-1}{3-p}} \\ &\leq 12(p-1)E(u(t)) - (3p-7) \|\nabla u(t)\|_{L^2}^2 + C\varrho^{-2} + C\varrho^{-\frac{p-1}{3-p}}, \quad \forall t \in I_{\max}, \end{aligned}$$

where we have chosen $\varepsilon = 3p - 7$ to get the second line. The energy conservation and $2 < \frac{p-1}{3-p}$ yield

$$(3p-7) \|\nabla u(t)\|_{L^2}^2 + V_{\phi_\varrho}''(t) \leq C\varrho^{-\frac{p-1}{3-p}}, \quad \forall t \in I_{\max}$$

provided that $\varrho > 0$ is taken sufficiently small. By the same reasoning as above, we prove (1.6) and (1.7).

Point (ii). The result is a straightforward application of the estimate in Lemma 5.1 yielding $G(u(t)) \lesssim -1$ uniformly in time in the maximal time of existence, and the Du, Wu, and Zhang scheme [12]. Indeed, with respect to the NLS equation with one focusing nonlinearity, the extra defocusing term accounts for negative contributions in the virial estimates. Hence by repeating the argument in [12] jointly with the uniform negative upper bound for G , the proof is complete. \square

REFERENCES

- [1] T. Akahori and H. Nawa, Blowup and scattering problems for the nonlinear Schrödinger equations, *Kyoto J. Math.*, Vol. 53, No. 3 (2013), 629–672. [3](#)
- [2] T. Akahori, S. Ibrahim, H. Kikuchi, and H. Nawa, Existence of a ground state and blow-up problem for a nonlinear Schrödinger equation with critical growth, *Differential Integral Equations*, 25:(3-4) 383–402, 2012. [1](#), [6](#)
- [3] T. Akahori, S. Ibrahim, H. Kikuchi, and H. Nawa, Existence of a ground state and and scattering for a nonlinear Schrödinger equation with critical growth, *Selecta Math. (N.S.)* 19(2): 545–609, 2013. [1](#)
- [4] J. Bellazzini, L. Forcella, and V. Georgiev, Ground state energy threshold and blow-up for NLS with competing nonlinearities, *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze (in press)*. arXiv:2012.10977 [math.AP] (2021), DOI: 10.2422/2036-2145.202005_044. [2](#), [4](#), [25](#), [27](#)
- [5] R. Carles and C. Sparber, Orbital stability vs. scattering in the cubic-quintic Schrödinger equation, *Rev. Math. Phys.* 33 (2021), no. 3, Paper No. 2150004, 27 pp. [1](#)
- [6] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics 10, New York University, AMS, 2003. [1](#), [7](#), [9](#), [26](#)
- [7] X. Cheng, Scattering for the mass supercritical perturbations of the mass critical nonlinear Schrödinger equations, *Illinois J. Math.* 64 (2020), 21–48. [1](#), [3](#), [4](#)
- [8] X. Cheng, C. Miao, and L. Zhao, Global well-posedness and scattering for nonlinear Scdinger equations with combined nonlinearities in the radial case, *J. Differential Equations* 261 (2016), 2881–2934. [1](#), [2](#), [3](#)
- [9] V. D. Dinh, Non-radial scattering theory for nonlinear Schrödinger equations with potential, *Non-linear Differ. Equ. Appl.* 28 (2021), art. no. 61. [13](#)
- [10] V. D. Dinh, A unified approach for energy scattering for focusing nonlinear Schrödinger equations, *Discrete Contin. Dyn. Syst.* 40 (2020), no. 11, 6441–6471. [3](#)
- [11] B. Dodson and J. Murphy, A new proof of scattering below the ground state for the non-radial focusing NLS, *Math. Res. Lett.* 25 (2018), no. 6, 1805–1825. [3](#), [7](#), [11](#)
- [12] D. Du, Y. Wu, and K. Zhang, On blow-up criterion for the nonlinear Schrödinger equation, *Discrete Contin. Dyn. Syst.* 36 (2016), no. 7, 3639–3650. [4](#), [29](#)
- [13] T. Duyckaerts, J. Holmer, and S. Roudenko, Scattering for the non-radial 3D cubic nonlinear Schrödinger equation, *Math. Res. Lett.* 15 (2008), 1233–1250. [3](#)
- [14] D. Fang, J. Xie, and T. Cazenave, Scattering for the focusing energy-subcritical nonlinear Schrödinger equation, *Sci. China Math.*, 54 (2011), 2037–2062. [3](#)
- [15] G. Fibich, *The nonlinear Schrödinger equation. Singular solutions and optical collapse*. Applied Mathematical Sciences, Springer, Cham, 2015. [1](#)
- [16] N. Fukaya and M. Hayashi, Instability of algebraic standing waves for nonlinear Schrödinger, *Trans. Amer. Math. Soc* 374 (2021) 1421–1447. [1](#), [2](#), [25](#)
- [17] A. Gammal, T. Federico. L. Tomio, and P. Chomaz, Atomic Bose-Einstein condensation with three-body interactions and collective excitations, *J. Phys. B* 33 (2000) 4053–4067. [1](#)
- [18] J. Holmer and S. Roudenko, A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation, *Comm. Math. Phys.* 282 (2008), 435–467. [3](#)
- [19] S. Ibrahim, N. Masmoudi, and K. Nakanishi, Scattering threshold for the focusing nonlinear Klein-Gordon equation. *Analysis PDE*. 4(3), 405–460 (2011). [3](#), [6](#)
- [20] M. Keel and T. Tao, Endpoint Strichartz estimates, *Amer. J. Math.* 120 (1998), 955–980. [7](#)
- [21] C. E. Kenig and F. Merle, Global well-posedness, scattering and blow-up for the energy critical, focusing, nonlinear Schrödinger equation in the radial case, *Invent. Math.* 166 (2006), 645–675. [3](#)
- [22] R. Killip, T. Oh, O. Pocovnicu, and M. Viřan. Solitons and scattering for the cubic-quintic nonlinear Schrödinger equation on \mathbb{R}^3 , *Arch. Ration. Mech. Anal.*, 225(1):469–548, 2017. [1](#)
- [23] Y. S. Kivshar and G. P. Agrawal, *Optical Solitons: From Fibers to Photonic Crystals*, Academic Press (2003). [1](#)
- [24] L. Jeanjean, J. Jendrej, T. T. Le, and N. Visciglia, Orbital stability of ground states for a Sobolev critical Schrödinger equation, *J. Math. Pures Appl.* (9) 164 (2022), 158–179. [1](#)
- [25] M. Lewin and S. Rota-Nodari, The double-power nonlinear Schrödinger equation and its generalizations: uniqueness, non-degeneracy and applications, *Calc. Var. Partial Differential Equations* 59 (2020), no. 6, Paper No. 197, 49 pp. [1](#)
- [26] Y. Luo, Scattering threshold for radial defocusing-focusing mass-energy double critical nonlinear Schrödinger equation in $d \geq 5$, arXiv:2106.06993 [math.AP] (2021). [1](#), [3](#)

- [27] Y. Luo, Sharp scattering for the cubic-quintic nonlinear Schrödinger equation in the focusing-focusing regime, *J. Funct. Anal.*, Volume 283, (2022) Issue 1, Paper No. 109489, 34. [1](#)
- [28] C. Miao, G. Xu, and L. Zhao, The dynamics of the 3D radial NLS with the combined terms, *Comm. Math. Phys.* 318 (3) (2013) 767–808. [1](#), [3](#)
- [29] C. Miao, T. Zhao, and J. Zheng, On the 4D nonlinear Schrödinger equation with combined terms under the energy threshold, *Calc. Var. and Partial Differential Equations*, 56(6):Paper No. 179, 39, 2017. [1](#)
- [30] F. Merle, P. Raphaël, and J. Szeftel, On collapsing ring blow-up solutions to the mass supercritical nonlinear Schrödinger equation, *Duke Mathematical Journal* 163 (2014), no. 2, 369–431. [4](#), [27](#)
- [31] D. E. Pelinovsky, V. V. Afanasjev, and Y. S. Kivshar, Nonlinear theory of oscillating, decaying, and collapsing solitons in the generalized nonlinear Schrödinger equation, *Phys. Rev. E* **2** (1996), no. 53, 1940–1953. [1](#)
- [32] N. Soave, Normalized ground states for the NLS equation with combined nonlinearities. *J. Differential Equations*, 269(9):6941–6987, 2020. [1](#)
- [33] N. Soave, Normalized ground states for the NLS equation with combined nonlinearities: the Sobolev critical case. *J. Funct. Anal.*, 279(6):108610, 43, 2020. [1](#)
- [34] T. Tao, M. Viřan, and X. Zhang, The nonlinear Schrödinger equation with combined power-type nonlinearities, *Comm. Partial Differential Equations* 32:1281–1343, 2007. [1](#), [4](#)
- [35] X. Zhang, On the Cauchy problem of 3D energy-critical Schrödinger equations with subcritical perturbations, *J. Differential Equations* 230:422–445, 2006. [1](#)

JACOPO BELLAZZINI, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI PISA, LARGO BRUNO PONTECORVO, 5, 56127, PISA, ITALY

Email address: jacopo.bellazzini@unipi.it

VAN DUONG DINH, ECOLE NORMALE SUPÉRIEURE DE LYON & CNRS, UMPA (UMR 5669), FRANCE

Email address: contact@duongdinh.com

LUIGI FORCELLA, DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E MATEMATICA, UNIVERSITÀ DEGLI STUDI DELL'AQUILA, VIA VETOIO, 67100, L'AQUILA, ITALY

Email address: luigi.forcella@univaq.it