



HAL
open science

Discrete Morse Functions and Watersheds

Gilles Bertrand, Nicolas Boutry, Laurent Najman

► **To cite this version:**

Gilles Bertrand, Nicolas Boutry, Laurent Najman. Discrete Morse Functions and Watersheds. 2023. hal-03928064v1

HAL Id: hal-03928064

<https://hal.science/hal-03928064v1>

Preprint submitted on 9 Jan 2023 (v1), last revised 4 Jul 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Discrete Morse Functions and Watersheds

Gilles Bertrand¹, Nicolas Boutry² and Laurent Najman¹

¹Univ Gustave Eiffel, CNRS, LIGM, F-77454 Marne-la-Vallée, France.

²EPITA, Research and Development Laboratory (LRDE), France.

Contributing authors: gilles.bertrand@esiee.fr; nicolas.boutry@lrde.epita.fr;
laurent.najman@esiee.fr;

Abstract

Any watershed, when defined on a stack on a normal pseudomanifold of dimension d , is a pure $(d - 1)$ -subcomplex that satisfies a drop-of-water principle. In this paper, we introduce Morse stacks, a class of functions that are equivalent to discrete Morse functions. We show that the watershed of a Morse stack on a normal pseudomanifold is uniquely defined, and can be obtained with a linear-time algorithm relying on a sequence of collapses. Last, but not the least, we prove that such a watershed is the cut of the unique minimum spanning forest, rooted in the minima of the Morse stack, of the facet graph of the pseudomanifold.

Keywords: Topological Data Analysis, Mathematical Morphology, Discrete Morse Theory, Simplicial Stacks, Minimum Spanning Forest.

1 Introduction

Watershed is a fundamental tool in computer vision, since its inception as an algorithm by the school of mathematical morphology [1, 2]. It is still true in this era of deep learning, where it is used as a post-processing tool [3]. From a discrete, theoretical point of view, the first topologically-sound approach was proposed in [4, 5]. Building on those results, in [6–8], it is demonstrated that watersheds are included in skeletons on pseudomanifolds of arbitrary dimension.

In this paper, we continue exploring the link between watershed and topology, in the framework of discrete Morse theory [9, 10]. Indeed, mathematical morphology [11] and discrete Morse theory, although they pursue different objectives, share many similar ideas. In particular, as demonstrated in [12–14], filtering minima using *morphological dynamics* [15] in watershed-based

image-segmentation, is equivalent to filtering the minima by *persistence*, a fundamental tool from Persistent Homology [16] used for topological data analysis [17, 18].

Although the main ideas of the present paper originate in [19], we have worked towards a simpler, unifying framework for exposing these ideas. This leads us to introduce *Morse stacks*: these functions correspond to the inverse of flat Witten-Morse functions that, according to R. Forman [20], *seem to have shown themselves to be the appropriate combinatorial analogue of smooth non-degenerate Morse functions*. We also propose a new definition for normal pseudomanifolds, a class of manifolds on which path-connectivities are all equivalent. Relying on these notions, we prove that a watershed, a pure $(d - 1)$ -subcomplex of a normal pseudomanifold, has several interesting properties when defined on a Morse stack F . In particular, in this setting, a watershed is uniquely

defined, and can be obtained thanks to a linear-time algorithm, relying on a sequence of collapses. Furthermore, a watershed is the cut of the unique minimum spanning forest of the facet graph of the normal pseudomanifold weighted by F , rooted in the minima of F . Relations between watersheds and Morse theory have long been informally known [21], but this is the first time that a link is presented in the discrete setting, relying on a precise definition of the watershed. Furthermore, as far as we know, this is the first time that a concept from Discrete Morse Theory is linked to a classical combinatorial optimization problem.

The plan of this paper is the following. Section 2 provides some basic definitions of simplicial complexes. We introduce here the notion of a covering pair, that is fundamental for the definition of Morse stacks. Section 3 recalls some definitions of simplicial stacks, which are a class of weighted simplicial complexes whose upper threshold sets are also complexes. Section 4 proposes a new definition for normal pseudomanifolds. Section 5 provides the necessary definitions for watersheds on stacks defined on normal pseudomanifolds. We propose here an algorithm for computing watershed relying on the collapse operation. In section 6, we introduce Morse stacks. Section 7 studies the properties of watersheds on Morse stacks. Section 8 links watershed and the minimum spanning forest. We conclude the paper with a discussion in section 9, in which we highlight the importance of our results, both from a theoretical and a practical point of views, and we propose some perspective for future work.

Finally, appendix A shows that our definition of normal pseudomanifold is equivalent to the classical one, and appendix B demonstrates that Morse stacks are equivalent to classical discrete Morse functions.

2 Simplicial complexes

A *simplex* x is a non-empty finite set; the dimension of x , written $\dim(x)$, is the number of its elements minus one. We also say that x is a p -*simplex* if $\dim(x) = p$.

Let S be a finite set of simplexes. A p -simplex in S is a (p -)face of S . A (p -)facet of S is a p -face of S that is maximal for inclusion. If x and y are two distinct faces of S such that $x \subseteq y$, we say that x is a face of y (in S). The *simplicial closure*

of S is the set $S^- = \{y \subseteq x \mid y \neq \emptyset \text{ and } x \in S\}$. The set S is a (*simplicial*) *complex* if $S = S^-$.

Let X be a complex. The *dimension of X* , written $\dim(X)$, is the largest dimension of its simplices, the *dimension of \emptyset* being defined to be -1 .

A 0-face of X is a *vertex of X* and a 1-face of X is an *edge of X* .

A complex X is a *graph* if the dimension of X is at most 1.

Let X be a complex and let $S \subseteq X$. If S is a complex, we say that S is *closed for X* or that S is a *subcomplex of X* . We say that S is *open for X* or that S is an *open subset of X* if, for any $x \in S$, we have $y \in S$ whenever $x \subseteq y$ and $y \in X$. If X is a complex and $S \subseteq X$, we note that S is closed for X if and only if $X \setminus S$ is open for X . In particular, \emptyset and X are both closed and open for X .

Let S be a finite set of simplexes. Let $\pi = \langle x_0, \dots, x_k \rangle$ be a sequence of elements of S . The sequence π is a *path in S* (from x_0 to x_k) if, for any $i \in [0, k-1]$, either $x_i \subseteq x_{i+1}$ or $x_{i+1} \subseteq x_i$. We say that S is *connected* if, for any $x, y \in S$, there exists a path from x to y in S . We say that $T \subseteq S$ is a *connected component of S* if T is connected and maximal, with respect to set inclusion, for this property.

Remark 1 We observe that:

- If X is a complex, then X is connected if and only if, for any vertices $x, y \in X$, there exists a sequence $\langle x = x_0, \dots, x_k = y \rangle$ of vertices of X such that, for any $i \in [0, k-1]$, $x_i \neq x_{i+1}$ and $x_i \cup x_{i+1}$ is an edge of X .
- If S is an open subset of a complex X , then S is connected if and only if, for any facets x, y of S , there exists a sequence $\langle x = x_0, \dots, x_k = y \rangle$ of facets of S such that, for any $i \in [0, k-1]$, $x_i \cap x_{i+1}$ is in S .

The following simple definition of a covering pair (or a p -pair) will play an important role in the sequel of the paper:

- It will first allow us to define a free pair (Definition 3). This corresponds to the operation of collapse of a simplicial complex introduced by J.H.C. Whitehead [22], which is a discrete analogue of a retraction, that is, a continuous (homotopic) deformation of an object onto one

of its subsets. In Section 3, free pairs for a simplicial complex will be extended to free pairs on stacks, which are maps on simplicial complexes (Definition 4).

- In Section 6, we introduce the notion of a flat pair, which is a special case of a covering pair. This permits us to have a very simple and concise presentation of a Morse stack (Definition 16). Indeed, a basic link between Morse stacks and the collapse operation is straightforward (Proposition 17). Furthermore, the notions of a gradient vector field and a gradient path follow immediately from covering and flat pairs.

Definition 2 (Covering pair) *Let X be a complex and $x, y \in X$, with $\dim(y) = p$. We say that (x, y) is a covering pair of X or a p -pair of X if x is a face of y and $\dim(x) = p - 1$.*

Definition 3 (Free pair) *Let X be a complex and let (x, y) be a p -pair of X . We say that (x, y) is a free (p -)pair of X if y is the only face of X that contains x .*

Thus, if (x, y) is a free pair of X , we have necessarily $\dim(x) = \dim(y) - 1$. Furthermore, we observe that y is necessarily a facet of X .

If (x, y) is a free p -pair of a complex X , then $Y = X \setminus \{x, y\}$ is an elementary (p -)collapse of X . We say that X collapses (resp. p -collapses) onto Y , if there exists a sequence $\langle X = X_0, \dots, X_k = Y \rangle$ such that X_i is an elementary collapse (resp. elementary p -collapse) of X_{i-1} , $i \in [1, k]$. If, furthermore, Y has no free pair (resp. free p -pair), then Y is an ultimate collapse (resp. ultimate p -collapse) of X . A complex X is collapsible if X collapses onto a single vertex.

3 Simplicial stacks

Let X be a simplicial complex, and let F be a map from X to \mathbb{Z} . If x is a face of X , the value $F(x)$ is called the altitude of F at x . For any $\lambda \in \mathbb{Z}$, we write $F[\lambda] = \{x \in X \mid F(x) \geq \lambda\}$, $F[\lambda]$ is the λ -section of F . We say that F is a (simplicial) stack on X if any λ -section of F is a simplicial complex. In other words, any λ -section of F is a closed set for X .

Let F be a map from a complex X to \mathbb{Z} . It may be easily seen that F is a simplicial stack if and only if, for any $x, y \in X$ such that $x \subseteq y$, we have

$F(x) \geq F(y)$. Also, a map F is a simplicial stack if and only if, for any covering pair (x, y) in X , we have $F(x) \geq F(y)$.

Now, we extend the notion of free pairs of simplicial complexes to simplicial stacks. This extension allows us to introduce some fundamental discrete homotopic transforms of these maps.

Definition 4 *Let F be a simplicial stack on X . We set $\lambda_m = \min\{F(x) \mid x \in X\}$. Let (x, y) be a p -pair of X . We say that (x, y) is a free (p -)pair of F if (x, y) is a free (p -)pair of $F[\lambda]$, with $\lambda = F(x)$ and $\lambda > \lambda_m$.*

If (x, y) is a free pair of F , then both x and y are in $F[\lambda]$, with $\lambda = F(x)$. Thus $F(y) \geq F(x)$. Also, we have $x \subseteq y$. Then, since F is a stack, we have $F(y) \leq F(x)$. Thus, we have $F(x) = F(y)$ whenever (x, y) is a free pair of F .

Let F be a simplicial stack on a complex X .

1. Let (x, y) be a free (p -)pair of F . Let G be the map such that $G(z) = F(z) - 1$ if $z = x$ or $z = y$, and $G(z) = F(z)$ if $z \in X \setminus \{x, y\}$. We can see that G is a simplicial stack on X , the map G is called an elementary (p -)collapse of F through (x, y) or simply an elementary (p -)collapse of F .
2. If G is the result of a sequence of elementary collapses (resp. p -collapses) of F , then we say that F collapses (resp. p -collapses) onto G .
3. If F collapses (resp. p -collapses) onto a stack G that has no free pair (resp. no free p -pair), then G is an ultimate collapse (resp. ultimate p -collapse) of F .

We conclude this section by giving a definition of a (regional) minimum of a stack, which plays a crucial role for the notion of a watershed.

Let F be a simplicial stack on a complex X and let $\lambda \in \mathbb{Z}$. A subset A of X is a minimum of F (at altitude λ) if A is a connected component of $X \setminus F[\lambda + 1]$ and $A \cap (X \setminus F[\lambda]) = \emptyset$. The divide of F is the set composed of all faces of X that are not in a minimum of F . Note that any minimum of F is an open set for X , and the divide of F is a simplicial complex.

4 Normal pseudomanifolds

The results of this paper hold true in a large family of n -dimensional discrete spaces, namely the

normal pseudomanifolds. This section provides a presentation of these spaces.

Let S be a finite set of simplexes. A *strong p -path in S (from x_0 to x_k)* is a path $\langle x_0, \dots, x_k \rangle$ such that, for each $i \in [0, k-1]$, either (x_i, x_{i+1}) is a p -pair, or (x_{i+1}, x_i) is a p -pair. The set S is *(d -)pure* if all facets of S have the same dimension d . If S is d -pure, we say that a strong d -path in S is a *strong path in S* . Also, we say that S is *strongly connected* if, for any two facets x, y in S , there exists a strong path in S from x to y . A subset T of S is a *strong connected component of S* if T is strongly connected and maximal, with respect to set inclusion, for this property.

Definition 5 (Normal pseudomanifold) *A connected and d -pure complex X , with $d \geq 1$, is a normal pseudomanifold (or a normal d -pseudomanifold) if:*

1. *The complex X is non-branching, that is, each $(d-1)$ -face of X is included in exactly two d -faces of X .*
2. *The complex X is strictly connected, that is, each connected open subset of X is strongly connected.*

Recall that a pure complex X is a *pseudomanifold* if it is non-branching and strongly connected [23]. Since the very set X is open for a complex X , we see that any normal pseudomanifold is a pseudomanifold.

In fact, the above definition is a new definition for a normal pseudomanifold. In Appendix A, we show that it is equivalent to the classical definition [24], [25], [26], which consists of a local condition together with the conditions that must be satisfied by a pseudomanifold.

Let us consider Fig. 1. The triangulated torus (a) is a normal pseudomanifold. The triangulated pinched torus (b) is a pseudomanifold that is not normal: the set of all faces containing the pinch vertex is a connected open subset of the complex, but this set is not strongly connected. The triangulated pinched torus (c) is not a pseudomanifold: the pinch segment does not satisfy the non-branching condition.

Let X be a proper subcomplex of a d -pseudomanifold M . We can see that, if $\dim(X) = d$, the complex X has necessarily a boundary, that is, there exists a free d -pair for X . By induction,

it means that the dimension of an ultimate d -collapse of X is necessarily $d-1$. See [7] for a formal proof.

Important notations. In the sequel of the paper:

- We denote by \mathbb{S} the collection of all simplicial complexes.
- If $X \in \mathbb{S}$, we write $Y \preceq X$ whenever $Y \subseteq X$ and $Y \in \mathbb{S}$, that is, whenever Y is a subcomplex of X .
- If $X \in \mathbb{S}$, and $S \subseteq X$, we write $S \sqsubseteq X$ whenever S is an open subset of X .
- We denote by \mathbb{M} (resp. \mathbb{M}_d) the collection of all normal pseudomanifolds (resp. all normal d -pseudomanifolds).
- If F is a stack on $M \in \mathbb{M}$, the notation $\min(F)$ stands for the union of all minima of F , and we write $\mathfrak{div}(F)$ for the divide of F . Thus, we have $\mathfrak{div}(F) = M \setminus \min(F)$, $\mathfrak{div}(F) \preceq M$, and $\min(F) \sqsubseteq M$.

We are now ready to introduce the notion of a watershed in the context of simplicial complexes. We will consider a normal pseudomanifold $M \in \mathbb{M}$ and a simplicial stack F on M , the map F may be seen as a “topographical relief” on the space M . A simplicial complex $W \preceq M$ may be a watershed of F if W “separates the minima of F ”. It means that if $A \sqsubseteq M$ is a connected component of $M \setminus W$, then A contains one and only one set $B \sqsubseteq M$ that is a minimum of F . Furthermore, the complex W must satisfy a “drop of water principle”: from each face of W , we may reach two distinct minima of F by following a descending path. Each connected component A of $M \setminus W$ will correspond to a “catchment basin” of the map F .

5 Watersheds

Let $X \in \mathbb{S}$, and let $A \sqsubseteq X$, with $A \neq \emptyset$. We say that $B \sqsubseteq X$ is an *extension of A* if $A \subseteq B$, and if each connected component of B includes exactly one connected component of A . We also say that B is an extension of A if $A = B = \emptyset$.

Proposition 6 *Let $M \in \mathbb{M}$ and let $A \sqsubseteq M$.*

1. *A subset S of A is a connected component of A if and only if S is a strong connected component of A .*




Fig. 1 (a): A normal pseudomanifold, which is a torus, (b): A pseudomanifold, which is a pinched torus, and where the pinch face is a vertex, (c): A pinched torus where the pinch face is a segment. This is not a pseudomanifold.

2. Let $B \sqsubseteq M$, with $A \subseteq B$. The set B is an extension of A if and only if each strong connected component of B includes exactly one strong connected component of A .

Proof 1) It may be seen that, if S is a connected component of the open set A , then S is necessarily an open set for M . Since M is a normal pseudomanifold, we deduce that S is strongly connected. Furthermore, S is a strong connected component of A , otherwise S would not be a maximal connected subset of A . Now, if S is a strong connected component of A , then S is a connected subset of A . Again, we see that S is a connected component of A . Otherwise, S would be a proper subset of a connected open subset T of A . Since M is a normal pseudomanifold, this subset T would be strongly connected, and S would not be a maximal strongly connected subset of A .

2) is a direct consequence of 1). \square

Let $X \in \mathbb{S}$ and $Y \preceq X$. Let $A \sqsubseteq X$, with $A \neq \emptyset$. We say that Y is a cut for A , if $X \setminus Y$ is an extension of A , and if Y is minimal for this property. That is, if $Z \preceq Y$, and if $X \setminus Z$ is an extension of A , then we have necessarily $Z = Y$.

Proposition 7 (from [7]) *Let $M \in \mathbb{M}$, $A \sqsubseteq M$ and $X \preceq M$, with $A \neq \emptyset$. If X is a cut for A , then the complex X is either empty or a pure $(n-1)$ -complex.*

Remark 8 *It could be seen that the previous result no longer holds if we consider arbitrary pseudomanifolds instead of normal pseudomanifolds. For example, the pinched vertex of the pinched torus of Figure 1.(b) could be in a cut.*

In fact, it is possible to bypass this situation by considering only strong paths between faces, as it is done in [7]. In this paper, in order to handle general connectedness and arbitrary paths, we have made the choice to settle our results in normal pseudomanifolds.

Let F be a stack on $M \in \mathbb{M}$. If $\pi = \langle x_0, \dots, x_k \rangle$ is a path in M , we say that π is *ascending* for F

(resp. *descending* for F) if, for any $i \in [0, k]$, we have $F(x_i) \leq F(x_{i+1})$ (resp. $F(x_i) \geq F(x_{i+1})$).

Definition 9 (Cut) *Let F be a stack on $M \in \mathbb{M}$ and let $X \preceq M$ be a cut for $\min(F)$. We say that X is a watershed of F if, for each $x \in X$, there exist two strong paths $\pi_1 = \langle x_0, \dots, x_k \rangle$ and $\pi_2 = \langle y_0, \dots, y_l \rangle$ in $M \setminus X$, such that:*

- $x \subseteq x_0$ and $x \subseteq y_0$;
- π_1 and π_2 are descending paths for F ; and
- x_k and y_l are simplices of two distinct minima of F .

Let $M \in \mathbb{M}$, F be a stack on M , and let W be a watershed for F . We say that $B \sqsubseteq M$ is a (*catchment*) *basin* of W if B is a connected component of $\overline{W} = M \setminus W$. Since W is a cut for $\min(F)$,

- any catchment basin B of W contains a unique minimum A of F , we say that A is the *minimum* of B ;
- any minimum A of F is included in a unique basin B of W , we say that B is the *catchment basin* of A .

Proposition 10 (from [7]) *Let $M \in \mathbb{M}_d$, F be a stack on M , and W be a watershed of F . Then, for any d -face x in M , there exists a strong path in $M \setminus W$ from x to a d -face of a minimum of F , that is descending for F .*

From the previous result, we easily derive the following proposition.

Proposition 11 *Let $M \in \mathbb{M}$, F be a stack on M , and W be a watershed of F . Let B be the catchment basin of a minimum A of F . Then, for any $x \in B$, there exists a descending path in B from x to a face of A .*

The two following results are crucial for linking a watershed of a stack F and the homotopy of F .

Proposition 12 (from [7]) *Let $M \in \mathbb{M}_d$. If F is a stack on M and H is a collapse of F , then:*

1. $\min(H)$ is an extension of $\min(F)$.
2. $\text{div}(H)$ is a collapse of $\text{div}(F)$.

It should be noted that the previous property is no longer true if we consider a stack on an arbitrary complex $X \in \mathbb{S}$ rather than a complex $M \in \mathbb{M}$. See Fig 2 which provides a simple counter-example.

Proposition 13 (from [7]) *Let $M \in \mathbb{M}_d$ and F be a stack on M .*

1. F contains a free d -pair if and only if $\text{div}(F)$ contains a free d -pair.
2. If $\dim(\text{div}(F)) = d$, then there exists a free d -pair for F .

Let F be a stack on $M \in \mathbb{M}_d$ and x be a $(d-1)$ -face of M . Let y, z be the two d -faces containing x . We say that x is (locally) separating for F if $F(y) < F(x)$ and $F(z) < F(x)$. We say that x is biconnected for F if y and z belong to distinct minima of F .

Definition 14 *Let F be a stack on $M \in \mathbb{M}_d$. Let $X \preceq M$. We say that X is a cut by collapse of F , or a \mathcal{C} -watershed of F , if there exists an ultimate d -collapse H of F such that X is the simplicial closure of the set of all faces of M that are biconnected for H .*

Theorem 15 (from [7]) *Let $M \in \mathbb{M}$ and let F be a stack on M . A complex $X \preceq M$ is a watershed of F if and only if X is a \mathcal{C} -watershed of F .*

Theorem 15 is illustrated in Fig. 3.

From Theorem 15 we may derive the procedure **WatershedCollapse**(F, M) (Fig. 4) for obtaining a watershed of a stack F on $M \in \mathbb{M}_d$.

The result W depends on the choices of the free pairs that are made at step 1. In any case, any watershed of F may be obtained by this procedure.

As explained hereafter, a direct implementation of the algorithm **WatershedCollapse**(F, M) can be slow.

- Step 1 is the more complex one. A naive implementation of this step is in the order of $n^2 * h$, where n is the number of d -faces, and h is the number of different altitudes of F . However, this step can be done in quasi-linear time, relying on a straightforward adaptation to simplicial complexes of the algorithm presented in [27]. This algorithm relies on a tree structure, where the nodes of the tree are the connected components of all the level sets of F , and where the edges of the tree correspond to the parenthood relationships between those connected components.
- Step 2 is a simple labelling, and may be done in linear time with respect to the number of d -faces. By using such a labelling, checking if a d -face is biconnected can be done in constant time.
- Finally, step 3 may be implemented in linear time, with respect to the number of incidence relations of M , that is the cardinal of the set $\{(x, y) \mid x, y \in M \text{ and } x \subsetneq y\}$.

6 Morse stacks

In this section, we transpose some basic notions of discrete Morse theory to stacks. We proceed by defining a Morse stack, which is the counterpart of a classical discrete Morse function. Morse stacks simply correspond to the inverse of flat discrete Morse functions. See Appendix B, which provides the few facts linking these two notions.

Let F be a map from a complex X to \mathbb{Z} . We say that a covering pair (x, y) of X is a flat pair of F whenever we have $F(x) = F(y)$.

Definition 16 (Morse stack) *Let F be a simplicial stack on a complex X . We say that F is a Morse stack (on X) if any face of X is in at most one flat pair of F .*

Let F be an arbitrary simplicial stack, and let (x, y) be a covering pair of F . We have seen that, if (x, y) is a free pair of F , then necessarily (x, y) is a flat pair of F . Suppose now that (x, y) is a flat pair of F . Then, there may exist another covering pair (x, z) that is also a flat pair of F . In this case, we see that (x, y) is not a free pair of F . By the very definition of a Morse stack, this situation cannot occur. In fact, we have the following result.




Fig. 2 Two stacks F and H with two levels: altitude 0 (faces in light grey) and altitude 1 (faces in black). The stack H is an elementary collapse of F (at altitude 1). But F has three minima whereas H has only two. Thus, $\min(H)$ is not an extension of $\min(F)$.




Fig. 3 (a) A simplicial stack F on a subset of a normal 2-pseudomanifold. (b) An ultimate 2-collapse of F . (c) A watershed of F . The pair (y, x) in (a) is a free-pair for F .

Proposition 17 *Let F be a Morse stack on a complex X . A covering pair (x, y) of X is a free pair of F if and only if (x, y) is a flat pair of F .*

Definition 18 (Regular and critical simplex) *Let F be a Morse stack on a complex X and let $x \in X$ with $\dim(x) = p$.*

- We say that x is regular or p -regular for F if x is in a flat pair of F .
- We say that x is critical or p -critical for F if x is not regular for F .

A Morse stack defined on a normal pseudo-manifold, together with its critical and regular simplices, can be seen on Fig. 5.

Let F be a Morse stack on a complex X . The gradient vector field of F , written $\overrightarrow{\text{grad}}(F)$, is the set of all flat pairs of F . If (x, y) is a covering pair of F such that $F(x) > F(y)$, we say that (y, x) is a differential pair of F . We write $\overrightarrow{\text{diff}}F$ for the set of all differential pairs for F . We also set:

WatershedCollapse(F, M)

1. Set $H = F$. Until H has no free d -pair, select arbitrarily a free d -pair (x, y) of H and replace H by the elementary collapse of H through (x, y) ;
2. Label all d -faces of distinct minima of H with distinct labels;
3. Extract from H the complex W that is the simplicial closure of the set of all d -faces of M which are biconnected for H .

Fig. 4 The procedure **WatershedCollapse**(F, M) computes a watershed W of a stack F defined on a normal pseudomanifold M .

- $\overrightarrow{\text{grad}}_p(F) = \{(x, y) \in \overrightarrow{\text{grad}}(F) \mid \dim(y) = p\}$,
- and
- $\overleftarrow{\text{diff}}_p(F) = \{(y, x) \in \overleftarrow{\text{diff}}(F) \mid \dim(y) = p\}$.

A Λ_p -path in F (from x_0 to x_k) is a sequence $\pi = \langle x_0, x_1, \dots, x_k \rangle$ composed of faces of X such that, for all $i \in [0, k - 1]$, the pair (x_i, x_{i+1}) is either in $\overrightarrow{\text{grad}}_p(F)$ or in $\overleftarrow{\text{diff}}_p(F)$. A sequence π is a *gradient path* for F if π is a Λ_p -path for F for some p .

Let $\pi = \langle x_0, x_1, \dots, x_k \rangle$ be a Λ_p -path in F . We observe that:

- For any $i \in [1, k - 1]$, the pair (x_i, x_{i+1}) is in $\overrightarrow{\text{grad}}_p(F)$ (resp. $\overleftarrow{\text{diff}}_p(F)$) whenever (x_{i-1}, x_i) is in $\overleftarrow{\text{diff}}_p(F)$ (resp. $\overrightarrow{\text{grad}}_p(F)$).
- Each face of π is either a p -face or a $(p - 1)$ -face. For any $i \in [0, k - 1]$, if x_i is a p -face, then x_{i+1} is a $(p - 1)$ -face, and if x_i is a $(p - 1)$ -face, then x_{i+1} is a p -face.
- The path π is a strong p -path.
- If π is not trivial, then π cannot be closed, that is, we have necessarily $k = 0$ whenever $x_k = x_0$.
- The path π is an ascending path, that is, we have $F(x_i) \leq F(x_{i+1})$ for any $i \in [0, k - 1]$. Furthermore, we have $F(x_i) < F(x_{i+2})$ for any $i \in [0, k - 2]$.

The following result is a basic fact about Morse functions.

Proposition 19 *Let F be a Morse stack on a complex $X \in \mathbb{S}$, and let S be a subset of X . If S is a minimum of F , then S is composed of a single facet of X .*

In the sequel, we will say that a face $x \in X$ is a *minimum* (of F) whenever the set $\{x\}$ is a minimum of F .

7 Morse stacks and watersheds

Let F be a Morse stack on a complex $X \in \mathbb{S}$. Let x, y be two faces of X . We say that x is Λ_p -linked to y if there is a Λ_p -path in F from x to y . Let $\pi = \langle x = x_0, \dots, x_k = y \rangle$ be a Λ_p -path in F from x to y . We write $\tilde{\pi} = \langle y = x_k, \dots, x_0 = x \rangle$ and we say that $\tilde{\pi}$ is a $\tilde{\Lambda}_p$ -path in F from y to x . We say that a face $z \in X$ is an *extension* of π if $\langle x = x_0, \dots, x_k = y, z \rangle$ is a Λ_p -path in F from x to z . We say that z is an *extension* of $\tilde{\pi}$ if $\langle y = x_k, \dots, x_0 = x, z \rangle$ is a $\tilde{\Lambda}_p$ -path in F from y to z .

Proposition 20 *Let F be a Morse stack on $M \in \mathbb{M}_d$, and let x be a facet of X . Let $\tilde{\pi}$ be a $\tilde{\Lambda}_d$ -path in F from the facet x to a face $y \in X$. Then one and only one of the following statements is true:*

1. The face y is a minimum.
2. There exists a unique face that is an extension of $\tilde{\pi}$.

Proof We set $\tilde{\pi} = \langle x = x_0, \dots, x_k = y \rangle$.

i) Suppose y is a $(d - 1)$ -face. In this case, y cannot be a minimum. Furthermore, we have $k \geq 1$ (since x is a d -face), and the face $t = x_{k-1}$ is necessarily a d -face. By the definition of a $\tilde{\Lambda}_p$ -path, the pair (y, t) is a flat pair. Since X is a pseudomanifold, there exists a unique d -face $z \in X$ such that $t \cap z = y$. We must have $F(z) < F(y)$, otherwise y would belong to more than one flat pair. Therefore, the pair (z, y) is a differential pair and z is an extension of $\tilde{\pi}$. Since (y, t) and (y, z) are the only covering pairs that contain y , the face z is the unique extension of $\tilde{\pi}$.

ii) Suppose y is a d -face. By the definition of a $\tilde{\Lambda}_p$ -path, a face z is an extension of $\tilde{\pi}$ if and only if (z, y) is a flat pair. Now we observe that y is not a minimum if and only if there is a face z such that (z, y) is a flat pair. But y belongs to at most one flat pair. Thus, $\tilde{\pi}$ has a unique extension whenever y is not a minimum. \square

Let F be a Morse stack on $M \in \mathbb{M}_d$. It should be noted that, if π is a Λ_d -path in F from a facet x to a face y , then π may have more than one extension. Nevertheless, by induction, we obtain the following result from Prop. 20.

Proposition 21 *Let F be a Morse stack on $M \in \mathbb{M}_d$, and let x be a facet of X . There exists a unique minimum m of F such that m is Λ_d -linked to x . Furthermore, there exists a unique Λ_d -path in F from m to x .*

We now consider the case where a Λ_d -path has no extension. Recall that a $(d-1)$ -face x is separating for F if the two d -faces y, z which contain x , are such that $F(y) < F(x)$ and $F(z) < F(x)$.

Proposition 22 *Let F be a Morse stack on $M \in \mathbb{M}_d$. Let x be a facet of X , and let π be a Λ_d -path in F from x to a face $y \in M$. If π has no extension, then the face y is necessarily separating for F .*

Proof Let $\pi = \langle x = x_0, \dots, x_k = y \rangle$ be a Λ_d -path in F from x to $y \in M$.

If $\dim(y) = d$, then π has necessarily an extension. Now suppose $\dim(y) = d-1$, thus $k \geq 1$. The face y is a face of two d -faces, the face $z = x_{k-1}$ and another face t . Since π has no extension, we have $F(t) < F(y)$. Furthermore, by the very definition of a Λ_d -path, the pair (z, y) is a differential pair, thus we have $F(z) < F(y)$. Therefore, y is separating for F . \square

Let F be a Morse stack on a complex $X \in \mathbb{S}$. Let π be a Λ_p -path in F . We say that π is *maximal* if neither π nor $\tilde{\pi}$ has an extension. The following result is a direct consequence of Prop. 20 and 22.

Corollary 23 *Let F be a Morse stack on $M \in \mathbb{M}_d$. Let π be a Λ_d -path in F from x to y . If π is maximal, then x is a minimum of F and y is a separating face for F .*

Let F be a Morse stack on $M \in \mathbb{M}_d$. Let x be a $(d-1)$ -face of X , and let y, z be the two distinct d -faces containing x . According to Prop. 21, each of these faces is Λ_d -linked to a single minimum. We say that the face x is Λ -biconnected (for F) if these two minima are distinct. Observe that a face is necessarily separating whenever it is Λ -biconnected.

Definition 24 (Morse watershed) *Let F be a Morse stack on $M \in \mathbb{M}$. The Morse watershed of F is the complex that is the simplicial closure of the set composed of all faces that are Λ -biconnected for F .*

Theorem 25 *Let F be a Morse stack on $M \in \mathbb{M}$. The Morse watershed of F is a watershed of F . Furthermore, the Morse watershed of F is the unique watershed of F .*

Th. 25 is illustrated in Fig. 5.

Proof Let W be the Morse watershed of F .

1. Let A be a connected component of $M \setminus W$. By Prop. 6, the set A is a strong connected component of $M \setminus W$. Let f be a facet of A . By Prop. 21, there exists a Λ_d -path π from a minimum m of F to f . By the very definition of a Λ_d -path, it may be seen that π does not contain any separating face. Thus π is included in $M \setminus W$, and m is in A . Therefore, A contains a minimum m .

Now let x be an arbitrary facet of A . Since A is strongly connected, there exists a strong path $\pi = \langle m = x_0, \dots, x_k = x \rangle$ in A from m to x . By Prop. 21, for each facet of π , there is a unique minimum of F that is Λ_d -linked to this facet. Let x_i be a facet of π , with $i \leq k-2$. Thus x_{i+1} is a $(d-1)$ -face and x_{i+2} is a facet. Let m_i (resp. m_{i+2}) be the unique minimum of F that is Λ_d -linked to x_i (resp. to x_{i+2}). Since x_{i+1} is not Λ -biconnected for F , it may be checked that we have necessarily $m_i = m_{i+2}$. By induction, it follows that m is Λ_d -linked to the facet x . Since this result holds for any facet of A , this clearly implies that m is the unique minimum of F which is in A . Thus, any connected component of $M \setminus W$ contains exactly one minimum of F . But any minimum of F is included in $M \setminus W$ (since a minimum is a d -face). It follows that $M \setminus W$ is an extension of $\min(F)$. Furthermore, by the definition of a Λ -biconnected face, W is minimal for this last property. Therefore, W is a cut for $\min(F)$. Since any Λ_d -path is a descending strong path, it may be checked that W fulfills all the conditions of Definition 9: W is a watershed.

2. Let W' be a watershed of F . Let x be a $(d-1)$ -face of M that is Λ -biconnected for F , and let




Fig. 5 The Morse watershed (in red) on a Morse stack defined on a normal pseudomanifold, a 2d-torus. In black, critical simplices: those of dimension 2 are minima, those of dimension 1 are saddle, and those of dimension 0 are maxima. Arrows represent gradient/collapse of dimension 2. We observe that the critical simplex 20 do not belong to the watershed.

y, z be the two distinct d -faces containing x . By Prop. 10 there exist a descending strong path in $M \setminus W'$ from y to a minimum m and a descending strong path in $M \setminus W'$ from z to a minimum m' . We observe that any descending strong path in M is also a $\tilde{\Lambda}_d$ -path in M . Thus, by the very definition of a Λ -biconnected face, we must have $m \neq m'$. Therefore, the face x must be in W' , otherwise $y, z, m,$ and m' , would belong to the same connected component of $M \setminus W'$. Thus $W \subseteq W'$. Since $M \setminus W' \subseteq M \setminus W$, each connected component of $M \setminus W'$ is included in one connected component of $M \setminus W$. But $M \setminus W'$ must be maximal for this last property, otherwise W' would not be a cut for $\min(F)$. It follows that we have $W' = W$.

□

By Theorem 15, the Morse watershed may be obtained by the algorithm **WatershedCollapse**(F, M). In this case we have a greedy procedure since the result W does not depend on the choice of the free pair of H that is made at each iteration.

In fact, since F is a Morse stack, we can simplify this procedure. The algorithm **MorseWatershed**(F, M) (Fig. 6) extracts the Morse watershed W of a Morse stack F on $M \in \mathbb{M}_d$. Also, it gives the catchment basin B of each minimum of F .

The soundness of this algorithm is a direct consequence of the above results. It may be implemented in linear time, with respect to the number of incidence relations of M , that is the cardinal of the set $\{(x, y) \mid x, y \in M \text{ and } x \subsetneq y\}$.

MorseWatershed(F, M)

- 1) Label all d -faces $x \in M$ with the label $B(x) = 0$;
 Label all faces $x \in M$ with the label $W(x) = \text{False}$;
 Label all d -faces $x \in M$ of distinct minima of F with distinct labels $B(x) > 0$;
- 2) Insert all d -faces x such that $B(x) > 0$ in a list L ;
- 3) Until L is empty, do:
 - 3.1) Extract a face x from L ;
 - 3.2) For all y such that $z = x \cap y$ is a $(d - 1)$ -face do:
 - 3.2.1) If $F(y) = F(z)$ insert y in L and do $B(y) := B(x)$;
 - 3.2.2) If $B(y) > 0$ and $B(y) \neq B(x)$ do $W(z) := \text{True}$;
- 4) For all $(d - 1)$ -faces $x \in M$ with $W(x) = \text{True}$ and all $y \subsetneq x$, do $W(y) := \text{True}$;
- 5) For all d -faces $x \in M$ and all $y \subsetneq x$ with $W(y) = \text{False}$, do $B(y) = B(x)$.

Fig. 6 The **MorseWatershed**(F, M) algorithm computes the Morse watershed W of a Morse stack F defined on a normal pseudomanifold M .

8 Morse watersheds and minimum spanning forests

In [7], an equivalence result which links the notion of a watershed in an arbitrary stack with the one of a minimum spanning forest is given. In this section, we refine this result in the case of Morse stacks.

Recall that we have defined a graph as a complex $X \in \mathbb{S}$ such that the dimension of X is at most 1.

Let $X \in \mathbb{S}$ with $\dim(X) = 0$, that is X is a non-empty set of vertices. Let Y be a graph such that $X \preceq Y$. We say that Y is a *forest rooted by* X if:

- we have $X = Y$, or
- there exists a free pair $\{x, y\}$ of Y such that $Y \setminus \{x, y\}$ is a forest rooted by X . If $\{x, y\}$ is a free pair for Y , we say that x is a *leaf* for Y .

If X is made of a single vertex, then it may be seen that the previous definition is an inductive definition, which is equivalent to the notion of a rooted tree in the sense of graph theory. If X is made of k vertices, then Y has k connected components. Each of these connected components is a rooted tree for some vertex of X .

Let $M \in \mathbb{M}_d$. The *facet graph* of M is the graph, denoted by Υ_M , such that:

- A vertex $\{x\}$ is in Υ_M if and only if x is a d -face of M ;
- An edge $\{x, y\}$ is in Υ_M if and only if $x \cap y$ is a $(d - 1)$ -face of M .

Let F be a Morse stack on $M \in \mathbb{M}$, and let $X = \{\{x\} \mid x \in \min(F)\}$. By Prop. 19, each $\{x\} \in X$ is a vertex of Υ_M . Let $Y \preceq \Upsilon_M$ be a forest rooted by X . We say that Y is a *spanning forest for* $\min(F)$ if all vertices of Υ_M are in Y . We define the *weight* of Y as the sum of all numbers $F(x \cap y)$, where $\{x, y\}$ is an edge of Y . We say that Y is a *minimum spanning forest for* $\min(F)$, if Y is a spanning forest for $\min(F)$ whose weight is minimum.

Let F be a Morse stack on $M \in \mathbb{M}_d$. We denote by S the set of all couples of d -faces (x, y) in \mathbb{M} such that $(x, x \cap y)$ is a differential pair of F , and $(x \cap y, y)$ is a flat pair of F . Thus, $\pi = \langle x, x \cap y, y \rangle$ is a Λ_d -path in F from x to y .

The *watershed forest* of F is the graph $G \preceq \Upsilon_M$ such that:

- All vertices of Υ_M are in G ;
- An edge $\{x, y\}$ is in G if and only if (x, y) or (y, x) is a couple in S .

From Proposition 21, we can check that the watershed forest is indeed a forest. More precisely, we can derive the following result.

Proposition 26 *If F is a Morse stack on $M \in \mathbb{M}_d$, then the watershed forest of F is a spanning forest for $\min(F)$.*

Theorem 27 *Let F be a Morse stack on $M \in \mathbb{M}$. The watershed forest of F is a minimum spanning forest for $\min(F)$. Furthermore, the watershed forest of F is the unique minimum spanning forest for $\min(F)$.*




Fig. 7 The Morse watershed (in red) on a Morse stack F defined on a 2d-torus. In blue, the watershed forest, which is the minimum spanning forest for $\min(F)$.

This theorem is illustrated in Fig. 7.

Proof Let G be a minimum spanning forest for $\min(F)$. By a minimum spanning tree lemma [28], [29], if $\{x\}$ is a vertex of G , then G must contain an edge $\{x, y\}$ that is a minimum weighted edge containing $\{x\}$. Now let W be the watershed forest of F and let $\{x, y\}$ be an edge of W . By the very definition of W , either $(x \cap y, x)$ or $(x \cap y, y)$ is a flat pair. By the definition of a Morse stack, if $(x \cap y, x)$ is a flat pair, then $\{x, y\}$ is the only minimum weighted edge containing $\{x\}$. Similarly, if $(x \cap y, y)$ is a flat pair, then $\{x, y\}$ is the only minimum weighted edge containing $\{y\}$. It follows that, if e is an edge of W , then e is the only minimum weighted edge containing some vertex v of W . Since v is necessarily in G , we deduce by the above lemma that $e \in G$. Therefore, we have $W \subseteq G$. Since both G and W are spanning forests, we must have $G = W$. This shows that W is the unique minimum spanning forest for $\min(F)$. \square

9 Discussion, future work and conclusion

In this paper, we propose, for the first time, a definition of watersheds for discrete Morse functions, and we study its properties. We are working in this paper on normal pseudomanifolds. This allows us in particular to exhibit a link between watershed and minimum spanning tree, that relies on gradient paths.

While a watershed definition have been proposed in the continuous Morse setting [30], the watershed notion was only a source of inspiration in discrete Morse theory. We mention in particular the following.

- A watershed algorithm was used as a preprocessing in [31] for computing a gradient vector

field, but without proof that watershed basins were related to any topological notion. Our approach directly provides watershed basins that are defined with gradient paths.

- In [32], watershed ideas were used as a motivation for obtaining Morse cells similar to catchment basins, with application to image segmentation. Our framework allows clarifying the difference between Morse cells and watershed basins. For example, in Fig. 5, the critical 1-simplex σ_0 is not part of the watershed cut, while it is part of the boundary of the Morse cell. We intend to explore in more details those differences in future work. We also envision studying the Morse-Smale decomposition.

One important difference between Morse stacks and discrete Morse functions is that minima are d -dimensional simplices in our framework, while they are 0-dimensional ones in Morse theory. Although this might appear minor, such a difference has important consequences. In particular, the watershed is a pure $(d - 1)$ -subcomplex, while a similar property is not possible with the boundary of Morse cells as classically defined (see [32] for example). Our approach allows for easily extracting topological features linking two regions, following the seminal paper [33]: indeed, we can for instance weight any simplex of the watershed cut with the persistence/dynamics [14] at which it disappears in a filtering. Such a representation, illustrated in Fig. 8, is called a *geodesic saliency map* in mathematical morphology, and is widely used (under the name *ultrametric contour map* [3]) as a post-processing behind deep-learning approaches. See [34, 35] for theoretical studies of this notion, and [36] for a toolbox implementing many variations around it.

Data analysis heavily relies on data simplification and data visualization. We advocate that the watershed, together with filtering operators such as morphological dynamics, is a cornerstone for data analysis [37]. We aim at controlling the topological simplification, and understanding what is discarded in the simplification. The results of this paper is a first step in this direction. We envision using skeleton algorithms such as [38], and tools from cross-section topology [39], that, up to now, have been used mainly for image analysis. Indeed, these tools can be applied to general data. In this regard, an important perspective of the

current paper is to bring together the persistence homology framework with the morphological one, reaching an audience as large as possible.

Acknowledgements

The authors would like to thank both Julien Tierny and Thierry Géraud, for many insightful discussions.

Appendix A Normal pseudomanifolds

A normal pseudomanifold is usually defined as a pseudomanifold that satisfies a certain link condition, which corresponds to a local property [24], [25], [26]. In this section, we show that this definition is equivalent to the one given in Definition 5.

Let S be a finite set of simplexes. If x and y are facets of S , a *p -chain (in S) from x to y* is a sequence $\langle x = x_0, \dots, x_k = y \rangle$ of facets of S such that, for each $i \in [0, k - 1]$, $x_i \cap x_{i+1}$ is a q -face of S , with $q \geq p$. The set S is *p -connected* if, for any two facets x, y in S , there is a p -chain in S from x to y .

We observe that:

- A complex is connected if and only if it is 0-connected.
- A d -pure complex is strongly connected if and only if it is $(d - 1)$ -connected.

Let X be a complex. Two faces $x, y \in X$ are *adjacent* if $x \cup y \in X$. The *link of $x \in X$ in X* is the complex $lk(x, X) = \{y \in X \mid x \cap y = \emptyset \text{ and } x \cup y \in X\}$.

The *star of $x \in X$ in X* is the set $st(x, X) = \{y \in X \mid x \subseteq y\}$.

Let X be a d -pseudomanifold. We say that X *satisfies the link condition* if $lk(x, X)$ is connected whenever x is a p -face of X and $p \leq d - 2$.

Let X be a complex and x be a p -face of X . Let $st^*(x, X) = st(x, X) \setminus \{x\}$. We have $lk(x, X) = \{y \setminus x \mid y \in st^*(x, X)\}$ and $st^*(x, X) = \{z \cup x \mid z \in lk(x, X)\}$.

We note that there is a set isomorphism between $lk(x, X)$ and $st^*(x, X)$, which preserves set inclusion. If $y \in st^*(x, X)$, the corresponding face $y \setminus x$ of $lk(x, X)$ is such that $dim(y \setminus x) = dim(y) -$




Fig. 8 Top: an image (left), together with a geodesic saliency map (right) where the darker a contour is, the more persistent it is. Bottom: two different views of a triangular mesh, superimposed with a geodesic saliency map where the whiter a contour is, the more persistent it is.

$(p + 1)$. Thus, $\dim(y \setminus x) = \dim(y) - p^+$, where $p^+ = p + 1$ is the number of elements in x .

Let X be a d -pseudomanifold and x be a p -face of X . Let $p^+ = p + 1$ and $d' = d - p^+$. The following facts are a direct consequence of the above isomorphism:

- The complex $lk(x, X)$ is d' -pure.
- The complex $lk(x, X)$ is non-branching.
- The set $st^*(x, X)$ is q -connected if and only if $lk(x, X)$ is q' -connected, with $q' = q - p^+$.

Proposition 28 *A pseudomanifold is normal if and only if it satisfies the link condition.*

Proof Let X be a d -pseudomanifold.

1. Suppose X satisfies the link condition and let S be a connected open subset of X .

Let x and y be two d -faces of S . By Remark 1, there exists a p -chain π in S from x to y . Thus $\pi = \langle x = x_0, \dots, x_k = y \rangle$ is a sequence of facets of S such that, for each $i \in [0, k - 1]$, $x_i \cap x_{i+1}$ is a q -face of S , with $q \geq p$. We choose π such that p is maximal and, if p is maximal, such that the number $K(\pi)$ of p -faces $x_i \cap x_{i+1}$, with $i \in [0, k - 1]$, is minimal. If $p = d - 1$, it means that S is strongly connected; then we are done. Suppose $p < d - 1$ and let x_i, x_{i+1} such that $z = x_i \cap x_{i+1}$ is a p -face.

Since X satisfies the link condition, $lk(z, X)$ is connected. By the isomorphism between $lk(z, X)$ and $st^*(z, X)$, it follows there is a q -chain $\langle x_i = w_0, \dots, w_l = x_{i+1} \rangle$ in $st^*(z, X)$ with $q > p$. Therefore, $\pi' = \langle x = x_0, \dots, x_i = w_0, \dots, w_l = x_{i+1}, \dots, x_k = y \rangle$ is a p -chain in S from x to y . But we have $K(\pi') < K(\pi)$, a contradiction. Thus, each connected open subset of X is strongly connected.

2. Suppose X is strictly connected. That is, any connected open subset of X is $(d - 1)$ -connected. Let x be a p -face of X with $p \leq d - 2$. The set $st(x, X)$ is a connected open subset of X , thus it is $(d - 1)$ -connected. Since $p < d - 1$, it means that $st^*(x, X)$ is $(d - 1)$ -connected. By the isomorphism between $lk(x, X)$ and $st^*(x, X)$, it follows that $lk(x, X)$ is strongly connected. Thus $lk(x, X)$ is connected. \square

In the second part of the proof of Prop. 28, we showed that $lk(x, X)$ is strongly connected. Consequently, we have the following characterization of a normal pseudomanifold.

Proposition 29 *A pseudomanifold X is normal if and only if, for each p -face x of X , with $p \leq d - 2$, the complex $lk(x, X)$ is a pseudomanifold.*

Appendix B discrete Morse functions

Let us consider the following definition of a discrete Morse function:

Definition 30 (Morse function) *Let X be a complex and let F be a map from X to \mathbb{Z} . We say that F is a discrete Morse function on X if any face of X is in at most one covering pair (x, y) in X such that $F(x) \geq F(y)$. If F is a discrete Morse function, we say that such a pair is a regular pair of F .*

It may be checked that this definition is equivalent to the classical one given by Forman (See Def. 2.1 and Lemma 2.5 of [40]).

In this way, the *gradient vector field* of a discrete Morse function F , written $\overrightarrow{\text{grad}}(F)$, is the set composed of all regular pairs of F .

The following restriction of a discrete Morse function will lead us to Morse stacks.

We say that a discrete Morse function F on X is *flat* if we have $F(x) = F(y)$ whenever (x, y) is a regular pair of F , that is, if each regular pair of F is a flat pair of F .

We can check that a map F from X to \mathbb{Z} is a flat discrete Morse function if and only if:

1. Each covering pair (x, y) in X is such that $F(x) \leq F(y)$;
2. Each face of X is in at most one flat pair of F .

Therefore, if we consider the function $-F$, we obtain the following:

Proposition 31 *Let X be a complex and let F be a map from X to \mathbb{Z} . The map F is a Morse stack on X if and only if the map $-F$ is a flat discrete Morse function on X .*

The following proposition claims that, up to an equivalence, we may assume that any discrete Morse function is flat (see Def. 2.27 and Prop. 4.16 of [10]).

Proposition 32 (from [10]) *If F is a discrete Morse function on X , then there exists a flat discrete Morse function G on X such that, for every covering pair (x, y) in X , we have $F(x) \geq F(y)$ if and only if*

$G(x) \geq G(y)$. In other words, the function G is such that $\overrightarrow{\text{grad}}(G) = \overrightarrow{\text{grad}}(F)$.

References

- [1] Digabel, H., Lantuéjoul, C.: Iterative algorithms. In: Proc. 2nd European Symp. Quantitative Analysis of Microstructures in Material Science, Biology and Medicine, vol. 19, p. 8 (1978). Riederer Verlag
- [2] Vincent, L., Soille, P.: Watersheds in digital spaces: an efficient algorithm based on immersion simulations. *IEEE Transactions on Pattern Analysis & Machine Intelligence* **13**(6), 583–598 (1991)
- [3] Arbelaez, P., Maire, M., Fowlkes, C., Malik, J.: Contour detection and hierarchical image segmentation. *IEEE transactions on pattern analysis and machine intelligence* **33**(5), 898–916 (2010)
- [4] Couprie, M., Bertrand, G.: Topological gray-scale watershed transformation. In: *Vision Geometry VI*, vol. 3168, pp. 136–146 (1997). SPIE
- [5] Bertrand, G.: On topological watersheds. *Journal of Mathematical Imaging and Vision* **22**(2), 217–230 (2005)
- [6] Cousty, J., Bertrand, G., Najman, L., Couprie, M.: Watershed cuts: Minimum spanning forests and the drop of water principle. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **31**(8), 1362–1374 (2009)
- [7] Cousty, J., Bertrand, G., Couprie, M., Najman, L.: Collapses and watersheds in pseudomanifolds. In: *International Workshop on Combinatorial Image Analysis. Lecture Notes in Computer Science*, vol. 5852, pp. 397–410 (2009). Springer
- [8] Cousty, J., Bertrand, G., Couprie, M., Najman, L.: Collapses and watersheds in pseudomanifolds of arbitrary dimension. *Journal of mathematical imaging and vision* **50**(3), 261–285 (2014)

- [9] Forman, R.: A Discrete Morse Theory for cell complexes. In: Yau, S.-T. (ed.) *Geometry, Topology for Raoul Bott*. International Press, Somerville, MA, USA (1995)
- [10] Scoville, N.A.: *Discrete Morse Theory* vol. 90. American Mathematical Soc., Providence, RI, USA (2019)
- [11] Najman, L., Talbot, H.: *Mathematical Morphology: from Theory to Applications*. John Wiley & Sons, Hoboken, NJ, USA (2013)
- [12] Boutry, N., Géraud, T., Najman, L.: An equivalence relation between Morphological Dynamics and Persistent Homology in 1D. In: *International Symposium on Mathematical Morphology. Lecture Notes in Computer Science Series*, vol. 11564, pp. 57–68 (2019). Springer
- [13] Boutry, N., Géraud, T., Najman, L.: An equivalence relation between morphological dynamics and persistent homology in n -D. In: *International Conference on Discrete Geometry and Mathematical Morphology*, pp. 525–537 (2021). Springer
- [14] Boutry, N., Najman, L., Géraud, T.: Some equivalence relation between persistent homology and morphological dynamics. *Journal of Mathematical Imaging and Vision* (2022). <https://doi.org/10.1007/s10851-022-01104-z>
- [15] Grimaud, M.: New measure of contrast: the dynamics. In: *Image Algebra and Morphological Image Processing III*, vol. 1769, pp. 292–306 (1992). International Society for Optics and Photonics
- [16] Edelsbrunner, H., Harer, J.: *Persistent Homology - A survey*. *Contemporary mathematics* **453**, 257–282 (2008)
- [17] Tierny, J.: *Introduction to Topological Data Analysis*. Technical report, Sorbonne University, LIP6, APR team, France (May 2017). <https://hal.archives-ouvertes.fr/cel-01581941>
- [18] Munch, E.: A user’s guide to topological data analysis. *Journal of Learning Analytics* **4**(2), 47–61 (2017)
- [19] Boutry, N., Bertrand, G., Najman, L.: Gradient vector fields of discrete morse functions and watershed-cuts. In: Baudrier, É., Naegel, B., Krähenbühl, A., Tajine, M. (eds.) *Discrete Geometry and Mathematical Morphology*, pp. 35–47. Springer, Cham (2022)
- [20] Forman, R.: Witten-morse theory for cell complexes. *Topology* **37**(5), 945–980 (1998)
- [21] De Floriani, L., Fugacci, U., Iuricich, F., Magillo, P.: Morse complexes for shape segmentation and homological analysis: discrete models and algorithms. *Computer Graphics Forum* **34**(2), 761–785 (2015). Wiley Online Library
- [22] Whitehead, J.H.C.: Simplicial spaces, nuclei and m -groups. *Proceedings of the London mathematical society* **2**(1), 243–327 (1939)
- [23] Massey, W.S.: *A Basic Course in Algebraic Topology*. Springer, Berlin, Germany (1991)
- [24] Bagchi, B., Datta, B.: Lower bound theorem for normal pseudomanifolds. *Expositiones Mathematicae* **26**(4), 327–351 (2008)
- [25] Basak, B., Swartz, E.: Three-dimensional normal pseudomanifolds with relatively few edges. *Advances in Mathematics* **365**, 107035 (2020)
- [26] Datta, B., Nilakantan, N.: Three-dimensional pseudomanifolds on eight vertices. *Int. J. Math. Mathematical Sciences* (2008)
- [27] Couprie, M., Najman, L., Bertrand, G.: Algorithms for the topological watershed. In: Andres, E., Damiand, G., Lienhardt, P. (eds.) *Discrete Geometry for Computer Imagery*, pp. 172–182. Springer, Berlin, Heidelberg (2005)
- [28] Cormen, T.H., Leiserson, C.E., Rivest, R.L.: *Introduction to Algorithms*. 23rd printing. The MIT Press and McGraw-Hill (1999)
- [29] Motwani, R., Raghavan, P.: *Randomized*

- Algorithms. Cambridge university press, Cambridge, UK (1995)
- [30] Najman, L., Schmitt, M.: Watershed of a continuous function. *Signal Processing* **38**(1), 99–112 (1994)
- [31] Čomić, L., De Floriani, L., Iuricich, F., Magillo, P.: Computing a discrete Morse gradient from a watershed decomposition. *Computers & Graphics* **58**, 43–52 (2016)
- [32] Delgado-Friedrichs, O., Robins, V., Sheppard, A.: Skeletonization and partitioning of digital images using discrete morse theory. *IEEE transactions on pattern analysis and machine intelligence* **37**(3), 654–666 (2014)
- [33] Najman, L., Schmitt, M.: Geodesic saliency of watershed contours and hierarchical segmentation. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **18**(12), 1163–1173 (1996)
- [34] Najman, L.: On the equivalence between hierarchical segmentations and ultrametric watersheds. *Journal of Mathematical Imaging and Vision* **40**(3), 231–247 (2011)
- [35] Cousty, J., Najman, L., Kenmochi, Y., Guimarães, S.: Hierarchical segmentations with graphs: quasi-flat zones, minimum spanning trees, and saliency maps. *Journal of Mathematical Imaging and Vision* **60**(4), 479–502 (2018)
- [36] Perret, B., Chierchia, G., Cousty, J., Guimarães, S.J.F., Kenmochi, Y., Najman, L.: Higr: Hierarchical graph analysis. *SoftwareX* **10**, 100335 (2019)
- [37] Challa, A., Danda, S., Sagar, B.D., Najman, L.: Watersheds for semi-supervised classification. *IEEE Signal Processing Letters* **26**(5), 720–724 (2019)
- [38] Bertrand, G., Couprie, M.: Powerful parallel and symmetric 3d thinning schemes based on critical kernels. *Journal of Mathematical Imaging and Vision* **48**(1), 134–148 (2014)
- [39] Bertrand, G., Everat, J.-C., Couprie, M.: Image segmentation through operators based on topology. *Journal of Electronic Imaging* **6**(4), 395–405 (1997)
- [40] Forman, R.: Morse Theory for cell complexes. *Advances in Mathematics* **134**, 90–145 (1998)