



**HAL**  
open science

# A lithospheric magnetic field model to spherical harmonic degree 1300

Erwan Thébault, Gauthier Hulot

► **To cite this version:**

Erwan Thébault, Gauthier Hulot. A lithospheric magnetic field model to spherical harmonic degree 1300. 12th Swarm Data Quality Workshop, Oct 2022, Uppsala, Sweden. hal-03928768

**HAL Id: hal-03928768**

**<https://hal.science/hal-03928768>**

Submitted on 7 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A lithospheric magnetic field model to spherical harmonic degree 1300

Thébault E.<sup>1</sup> and G. Hulot<sup>2</sup>

<sup>1</sup>Laboratoire Magmas et Volcans, Observatoire de Physique du Globe de Clermont-Ferrand, Université Clermont-Auvergne, France.

<sup>2</sup>Institut de Physique du Globe de Paris (IPGP) - CNRS - Université Paris Cité, France.


## SUMMARY

Detailed mapping of the Earth's magnetic field brings key constraints on the composition, dynamics, and history of the crust. Satellite and near-surface measurements detect different length scales and are complementary. Here, we build a model, selecting and processing magnetic field measurements from the German CHAMP and ESA Swarm satellites, which we merge with near-surface scalar anomaly data. We follow a regional approach for modeling the magnetic field measurements and next transform the series of regional models into a unique set of spherical harmonic (SH) Gauss coefficients. This produces the first global model to SH degree 1300 derived by inversion of all available measurements. The new model agrees with previous satellite-based models at large wavelengths and fits the CHAMP and Swarm satellite data down to expected noise levels. Further assessment in the geographical and spectral domains are in progress but preliminary results show that the model is stable when downward continued to the Earth's surface.

## ALGORITHM

We apply a procedure developed during the Swarm preparation phase (Thébault et al., 2013). We consider the Swarm A and C (A/C) measurements from March 2014 to August 2022 (baseline 0602) and the last 2 years of the CHAMP data (2008-2010). We select night time (23LT-5LT) vector/scalar measurements corresponding to Dst value lower than  $\pm 5$  nT, Kp index smaller than 2 $\sigma$  and  $|IMFBy| < 2$  nT and  $IMFBz > 0$ . The measurements are corrected for the main and external magnetic fields. Along-track scalar and vector gradients for Swarm and CHAMP and across track Swarm A/C gradients are built. We extend the satellite dataset with the World Digital Magnetic Anomaly grid data (WDMAM 2.0: Lesur et al., 2015) at 5 km altitude above the Earth's surface. The full dataset ( $> 110 \cdot 10^6$  data) is processed by a series of 700 equally distributed regional models derived to 30 km resolution, which are then transformed into Spherical Harmonics to SH degree 1300 (Thébault et al., 2021).

## THE LITHOSPHERIC MAGNETIC FIELD MODEL



**Figure :** (top left): Down vertical component; (top right) North component ; (bottom left) East component ;(bottom right) residuals between the anomaly field predicted by the SH model and the WDMAM scalar anomaly gridded data. Projections are Mollweid. The field components in the unit of nT are sketched at the Earth's mean radius to SH degree 1300.

## ACKNOWLEDGMENTS AND REFERENCES

The European Spatial Agency (ESA) is gratefully acknowledged for providing access to the L1b Swarm data. The authors would like to thank J. Dyment and V. Lesur for making the World Digital Magnetic Anomaly map grid available online. This study was partly funded by CNES in the framework of the project "Exploitation de la mission spatiale Swarm" and by the ESA DISC through the project "Extending the dedicated lithospheric field inversion".

- Lesur, V., Hamoudi, M., Choi, Y., Dyment, J., & Thébault, E. (2016). Building the second version of the world digital magnetic anomaly map (WDMAM). *Earth, Planets and Space*, 68(1), 1-13.
- Thébault E., P. Vigneron, S. Maus, A. Chulliat, O. Sirol, G. Hulot, Swarm SCARF Dedicated Lithospheric Field Inversion chain, *Earth Planets and Space*, Vol. 65 (No. 11), pp. 1257-1270, 2013
- Thébault, E., Hulot, G., Langlais, B., & Vigneron, P. (2021). A Spherical Harmonic Model of Earth's Lithospheric Magnetic Field up to Degree 1050. *Geophysical Research Letters*, 48(21), e2021GL095147.