



HAL
open science

Arylcyclohexylamine Derivatives: Pharmacokinetic, Pharmacodynamic, Clinical and Forensic Aspects

Romain Pelletier, Brendan Le Daré, Diane Le Bouedec, Angéline Kernalléguen, Pierre-Jean Ferron, Isabelle Morel, Thomas Gicquel

► **To cite this version:**

Romain Pelletier, Brendan Le Daré, Diane Le Bouedec, Angéline Kernalléguen, Pierre-Jean Ferron, et al.. Arylcyclohexylamine Derivatives: Pharmacokinetic, Pharmacodynamic, Clinical and Forensic Aspects. International Journal of Molecular Sciences, 2022, 23 (24), pp.15574. 10.3390/ijms232415574 . hal-03930104

HAL Id: hal-03930104

<https://hal.science/hal-03930104>

Submitted on 9 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Review

Arylcyclohexylamine Derivatives: Pharmacokinetic, Pharmacodynamic, Clinical and Forensic Aspects

Romain Pelletier ^{1,2,*}, Brendan Le Daré ^{1,3}, Diane Le Bouëdec ², Angéline Kernalléguen ¹, Pierre-Jean Ferron ¹, Isabelle Morel ^{1,2} and Thomas Gicquel ^{1,2}

¹ Univ Rennes, INSERM, INRAE, CHU Rennes, Institut NuMeCan (Nutrition, Métabolisme et Cancer), F-35000 Rennes, France

² Centre Hospitalier Universitaire de Rennes, Laboratoire de Toxicologie Biologique et Medico-Légale, CHU Pontchaillou, 2 Rue Henri Le Guilloux, F-35000 Rennes, France

³ Centre Hospitalier Universitaire de Rennes, Service Pharmacie, F-35000 Rennes, France

* Correspondence: romain.pelletier@chu-rennes.fr

Abstract: Since the 2000s, an increasing number of new psychoactive substances (NPS) have appeared on the drug market. Arylcyclohexylamine (ACH) compounds such as ketamine, phencyclidine and eticyclidine derivatives are of particular concern, given their rapidly increasing use and the absence of detailed toxicity data. First used mainly for their pharmacological properties in anesthesia, their recreational use is increasing. ACH derivatives have an antagonistic activity against the N-methyl-D-aspartate receptor, which leads to dissociative effects (dissociation of body and mind). Synthetic ketamine derivatives produced in Asia are now arriving in Europe, where most are not listed as narcotics and are, thus, legal. These structural derivatives have pharmacokinetic and pharmacodynamic properties that are sometimes very different from ketamine. Here, we describe the pharmacology, epidemiology, chemistry and metabolism of ACH derivatives, and we review the case reports on intoxication.

Keywords: arylcyclohexylamines; ketamine; phencyclidine; NPS; review



Citation: Pelletier, R.; Le Daré, B.; Le Bouëdec, D.; Kernalléguen, A.; Ferron, P.-J.; Morel, I.; Gicquel, T. Arylcyclohexylamine Derivatives: Pharmacokinetic, Pharmacodynamic, Clinical and Forensic Aspects. *Int. J. Mol. Sci.* **2022**, *23*, 15574. <https://doi.org/10.3390/ijms232415574>

Academic Editor: Ylenia Zambito

Received: 14 October 2022

Accepted: 24 November 2022

Published: 8 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Since the 2000s, an increasing number of new psychoactive substances (NPS) have appeared on the drug market. In most cases, a single functional or chemical group constitutes the only difference between the NPS and the parent recreational drug. This change not only alters the molecule's pharmacological properties, but also makes the use of a molecule legal, simply because the derivative is not registered on the list of narcotic products—making it a “legal high” [1–3]. Due to their very low cost (5 to 20 euros per gram) and ready availability (facilitated by deliveries via the Internet), NPSs are an emerging public health problem. Arylcyclohexylamine (ACH) derivatives, such as ketamine, are of particular concern, given their rapidly increasing use and the absence of detailed toxicity data.

Although ketamine is currently the best-known member of the ACH family, the latter's history began with the lead compound phencyclidine (PCP). Clinical trials of PCP as an analgesic (Sernyl[®]) were initiated in 1958, but abandoned in 1965 due to sometimes uncontrollable “psychotic” adverse drug reactions in ~15% of the study participants [4,5]. PCP's psychedelic properties led to a new chapter in its history as a street drug (“angel dust”) and made it the first of many synthetic drugs to appear on the market as an illicit recreational substance [6]. PCP had its heyday in 1976, with many newspaper and television reports on its recreational use [7]. Despite stricter law enforcement controls, a move to Schedule II of the United States' Controlled Substances Act in January 1978 and a fall in media coverage, consumption of PCP has been increasing steadily after a slight decline in the late 1980s.

First-generation PCP derivatives appeared at street markets between the 1960s and 1990s. The first cases of non-medical PCP use were described in 1967 and 1968 in the USA (mainly in San Francisco and New York) [8]. Most of the PCP derivatives correspond to basic chemical changes. Morris and Wallach estimated that there are 14 PCP derivatives of recreational interest [9]. Over the last 10 years, only a few ACH derivatives (e.g., 3-MeO-PCP, 2-fluorodeschloroketamine (2F-DCK) and 3-OH-PCP) have been commonly used for recreational purposes. Like ketamine, these molecules can be extremely toxic and sometimes fatal (see Section 3). Ketamine was synthesized from PCP in 1962 by the pharmaceutical company Parke-Davis (Detroit, MI, USA), which was seeking to develop new ACH anesthetics with analgesic properties [10]. The first-in-man trial took place in 1964 at Michigan State Prison (Jackson, MI, USA). Despite the presence of dissociative effects (corresponding to a dissociation between body and mind) associated with short-acting anesthesia, a patent was filed in 1966, and ketamine was approved for sale by the United States Food and Drug Administration (FDA) in 1970 [11,12]. During the Vietnam War (1955–1975), ketamine was used as an anesthetic and an analgesic [13]. In 1989, the FDA's approval of propofol led to a decrease in the use of ketamine as an anesthetic [14]. The use of ketamine was revived a few years later, following the discovery of the drug's value in the treatment of opioid-induced hyperalgesia as well as the treatment of depression [15]. At present, ketamine is widely used in veterinary medicine, pediatrics, anesthesia and emergency medicine. Along with ketamine's medical uses, its powerful dissociative effect and low price make it a sought-after recreational product. In Europe, ketamine has been classified as a narcotic product since the 1990s [16].

Recreational ketamine use has been mainly described in electro-alternative culture, rave parties and, more particularly, in Asian countries. The synthetic ketamine derivatives produced in Asia are now arriving in Europe, where most are not listed as narcotics and are, thus, legal. The first synthetic derivatives (eticyclidine (PCE), 2-oxo-eticyclidine, etc.) were observed in the 1960s (at the same time as ketamine), although their effects were considered to be too powerful. Furthermore, supply problems prevented the drugs from being used widely. However, the advent of the Internet (and especially the Dark Web in the 2010s) and better logistics led to a huge increase in the consumption of ketamine and its derivatives. Cases of intoxication were first described in Asia, and are becoming more frequent in Europe [17–20]. Herein, we describe the pharmacology and metabolism of ACH derivatives and review the case reports on intoxication.

2. Pharmacology of Ketamine and Its Synthetic Derivatives

2.1. Physical-Chemical Properties

The first-generation PCP derivatives keep a cyclohexane ring, in order to retain the antagonistic activity against the N-methyl-D-aspartate (NMDA) receptor and, thus, the dissociative effects. The ACH family encompasses three main subfamilies: ketamine-like molecules, phencyclidine-like (PCP-like) molecules and eticyclidine-like (PCE-like) molecules. These compounds are mainly derived through modification of the aryl ring, i.e., through the addition of an alkyl chain or substitutions of the amine group (Figure 1).

Ketamine is characterized by three rings: an aryl ring, a cyclohexyl ring, and an amine ring (Figure 1). The cyclohexane ring is usually intact because it is required for antagonism of the NMDA receptor. ACHs are chiral; the asymmetric carbon in the C2 position has two optical isomers. Ketamine is marketed as a racemate (i.e., a mixture of S(+)-ketamine, also known as esketamine and R(-)-ketamine, also known as arketamine) or as the pure S enantiomer. Ketamine is the only ACH to have a halogen (chlorine) on the aryl ring. In 2-F-DCK, the chlorine is replaced by fluorine. In other ACH derivatives, hydroxy (OH) or methoxy (MeO) groups are added to the aryl ring [21] (Figure 1).




Figure 1. The ACH derivatives most frequently involved in cases of intoxication. PCP: phencyclidine (1-(1-phenylcyclohexyl)piperidine); DCK: deschloroketamine (2-(methylamino)-2-phenylcyclohexan-1-one); PCE: eticyclidine (N-ethyl-1-phenylcyclohexan-1-amine); MXE: methoxetamine (2-(ethylamino)-2-(3-methoxyphenyl)cyclohexan-1-one); MXPr: methoxpropamine (2-(3-methoxyphenyl)-2-(propylamino)cyclohexan-1-one).

2.2. Pharmacodynamics

2.2.1. Mechanism of Action

ACH derivatives act mainly by antagonizing the glutamate NMDA receptor in the brain and spinal cord [22]. The NMDA receptor is an ion channel receptor consisting of a combination of four subunits encoded by seven genes: GluN1, GluN2_{A→D} and GluN3_{A→B} [23]. The channel is permeable to calcium ions, and triggers many intracellular pathways [24]. ACH derivatives bind to an allosteric site (the PCP-binding site) in the channel and act as non-competitive antagonists [25]. This antagonism is responsible for the dissociative anesthetic and amnesic effects, and (perhaps) the antidepressant and analgesic effects (Figure 2).

An ACH's three-dimensional structure is important for its antagonistic action. Compared with R(−)-ketamine, S(+)-ketamine has four times the activity and four times the affinity for the PCP site of the NMDA receptor. Hence, S(+)-ketamine is twice as active as the racemate [26].

Although data on the pharmacology of the new synthetic ACHs are scarce, all the derivatives (including PCP and ketamine) bind to the PCP binding site [9,27] (Figure 2).

Although it is thought that ACH derivatives bind to other receptors (e.g., the muscarinic and opioid receptors, for ketamine), there are no data on relevant clinical effects [28–30].




Figure 2. Mechanism of action of arylcyclohexylamine derivatives.

2.2.2. Clinical Effects

Ketamine has a powerful, rapid, anesthetic and dissociative effect, but (unlike opioids) does not cause hypotension or a respiratory depressant effect [6]. Domino et al. (1965) defined dissociative syndrome as the failure of cortical sensory information to reach associative areas, i.e., dissociation between the thalamocortical and limbic systems.

As is the case for the ACH derivatives' anesthetic effects, the analgesic uses are limited by the compound's psychodysleptic effects. Ketamine has several mechanisms of action, resulting in (i) a dissociative effect via the decoupling of nociceptive information from pain, (ii) inhibition of descending inhibitory systems and astrocytic pronociceptive systems, (iii) a local anesthetic effect and (iv) anti-inflammatory properties [31].

Interestingly, the mechanism of neuropathic pain involves activation of the NMDA receptor (leading to the loss of downward inhibition) and inflammation-associated changes in the spinal cord [32–36]. Ketamine's antagonism of the NMDA receptor meant that it was originally used as a painkiller.

S(+)-ketamine can also be used as antidepressant, although the NMDA receptor's role in this depression has not been characterized [37]. S(+)-ketamine's antidepressant effect is thought to come from activation of the mammalian target of rapamycin pathway and inactivation of glycogen synthase kinase-3 beta—especially because inhibitors of the latter enzyme potentiate and prolong the effects of ketamine, even at low doses [38,39].

2.3. Pharmacokinetics

Little is known about the pharmacokinetics of ACH derivatives. However, it is suspected that these derivatives are pharmacokinetically similar to ketamine, the only molecule used therapeutically, and for which detailed studies have been undertaken. However, sometimes very small structural modifications can greatly modify a drug's pharmacokinetics. Given this context, we primarily describe the pharmacokinetics of ketamine below.

2.3.1. Absorption

The bioavailability of orally administered ketamine is very low (between 8% and 24%), due to a major hepatic first pass effect. The bioavailability of orally administered S(+)-ketamine is even lower (~10%) [40–42]. The peak blood concentration is usually obtained after 40 to 55 min, and the first metabolites are detected after 10 to 30 min [6,42,43]. Similarly, the bioavailability of sublingual preparations (used for analgesic purposes) is low (~30%) [44].

In contrast, snorting or inhaling (the most frequent techniques in recreational use) leads to almost complete absorption [45]. The intramuscular route also allows rapid, high absorption, with a bioavailability of about 93%, and the injected ketamine can be detected

in plasma after just 4 min [45,46]. After intravenous injection, ketamine can be observed after 5 min, and the plasma peak is seen at 5 to 30 min [45,47].

2.3.2. Distribution

The ACH derivatives' high lipid solubility (estimated by LogP) is an important factor for bioavailability and enables them to cross the various lipid membranes in tissues. The plasma protein-bound fraction of ketamine is low (around 10–30%), which allows the drug to cross the blood–brain barrier easily and, thus, to exert its best-known effects (anesthesia, analgesia and anti-inflammatory properties) in the brain [48].

After systemic absorption, ketamine (LogP = 2.2) distributes rapidly through the tissues, and especially the brain. Ketamine's very short absorption half-life (2 to 4 min) gives it very rapid effects [6,48]. The drug's volume of distribution is high (3.5 L/kg) due to its liposolubility [41,48,49] (Table 1).

2.3.3. Metabolism

The metabolism of ketamine has been extensively studied, and is summarized in Figure 3 [26,50–57]. The primary metabolic pathway gives rise to norketamine (an active metabolite) and hydroxylated derivatives [45,46]. Most of these compounds are then glucuronoconjugated and excreted via the renal or biliary route. Ketamine is mainly metabolized in the liver by cytochromes P450 (CYP450); more particularly, CYP3A4 and CYP2B6 give rise to norketamine and dehydronorketamine (the other main metabolite), respectively [26,57] (Figure 3).




Figure 3. Overview of ketamine's metabolism [26,45,46,51–53]; bold arrows indicate the major pathway and dotted arrows indicate minor metabolic pathways; FMO: flavin-containing monooxygenase.

Table 1. The metabolic fate of ACH derivatives.

Molecule	Main Metabolites	Biotransformation	CYP Involved	Predicted LogP	Reference
Ketamine derivatives					
DCK	Dihydro-DCK	Hydrogenation	Unknown	2.7	[58]
	Dihydro-nor-DCK	Hydrogenation + demethylation			
2F-DCK	Nor-2F-DCK	Demethylation	Unknown	2.9	[21,58,59]
	Dihydro-2F-DCK	Hydrogenation			
	Dihydro-nor-2F-DCK	Hydrogenation + demethylation			
Phencyclidine derivatives					
PCP	c-PPC	Hydroxylation	CYP1A,3A	3.6	[60,61]
	t-PPC				
	PCHP				
3-OH-PCP	M1	Hydroxylation	Unknown	3.3	[62]
	M2	N-dealkylation + carboxylation			
	M3	O-glucuronidation			
3-MeO-PCP	Hydroxy-3-MeO-PCP	Hydroxylation	CYP2B6	3.6	[63–65]
	Demethyl-dihydroxy-3-MeO-PCP	Demethylation + hydroxylation	CYP2C19/2D6		
	Piperidine-dihydroxy-3-MeO-PCP	Hydroxylation	CYP2B6		
Eticyclidine derivatives					
Methoxpropamine	N-despropyl(nor)MXPr	Depropylation	Unknown	2.8	[58,66]
	O-desmethylMXPr	Demethylation			
	DihydroMXPr	Hydrogenation			
2-oxo-PCE	2-en-PCA-N-Glu	Dehydration + glucuronidation	Unknown	2.5	[67]
	M3	Oxidative deamination + dehydration			
	O-PCA-N-Glu	Glucuronidation			
2-FDCNEK	2-fluorodeschloro-norketamine	N-dealkylation	Unknown		[68]

LogP was calculated with XLogP3 software. DCK: Deschloroketamine, PCP: Phencyclidine, MXPr: Methoxpropamine, PCE: Eticyclidine; c-PPC: cis-1-(1-phenyl-4-hydroxycyclohexyl)piperidine; t-PPC: trans-1-(1-phenyl-4-hydroxycyclohexyl)piperidine; 1-(1-phenylcyclohexyl)-4-hydroxypiperidine; 2-FDCNEK: 2-fluoro-deschloro-N-ethyl-ketamine aM: metabolite.

2.3.4. Excretion

Ketamine has a relatively short duration of action. The excretion half-life is approximately 2 to 4 h, and the drug is excreted mainly by the kidneys [40]. Only 2% of ketamine and norketamine are excreted in unchanged form; the corresponding percentage for dehydronorketamine is about 16% [35,50,65–67]. Approximately 80% of hydroxyketamine and hydroxynorketamine are glucuroconjugated, which facilitates biliary and urinary excretion [69]. The excretion half-life of S(+)-ketamine appears to be longer (about 4 to 7 h) than that of the racemic mixture, due to CYP450's greater stereoselectivity for R(–)-ketamine [40,42,43,70,71].

The limited data on the excretion of other ACH derivatives come from studies of human metabolism [21,59,63,72]. 2F-DCK and 3-MeO-PCE were detected in the urine and bile of an intoxicated patient, suggesting the presence of both urinary and biliary excretion. Similarly, 2-F-DCK metabolites (including phase II metabolites) were found in the urine and bile of an intoxicated patient who subsequently died [19].

3. Arylcyclohexylamines: Clinical Toxicity and Forensic Cases

3.1. Clinical Toxicity

Ketamine has long been neglected by recreational users—partially because its effects were considered to be too strong. Despite its low addictive power and mild withdrawal syndromes, ketamine remains a highly toxic drug when taken under poorly controlled conditions. Psychic dependence and craving phenomena have often been reported. Ketamine is also often taken in combination with other molecules, which makes the drug's effects even harder to predict. Most of the fatal cases of intoxications involve the use of multiple substances.

Ketamine was originally used as an anesthetic, but the absence of a respiratory depressant effect has caused the drug's role to change gradually over time [73,74]. Ketamine has mainly been used recreationally by experienced users who wanted to discover new sensations or new emotions. The consumption of ketamine can induce euphoric effects, such as “cottony” drunkenness, at low doses, and hallucinations and powerful dissociation effects at high doses.

The ketamine trafficking and dealing network is very small, and is currently limited to free parties and (in some large cities) clubs. Ketamine has a reputation for not being cut, which has improved its image among consumers. For recreational use, the ingested or snorted doses are typically between 125 and 500 mg [75].

3.1.1. Acute Toxicity

Initially, ketamine was only used by a very small proportion of people in the “rave” culture, i.e., a population of nomads who used the drug for its powerful dissociative and hallucinogenic effects. Ketamine had a rather negative reputation, due to its radical effects and its use in veterinary medicine.

This poor reputation faded in the 2000s, when ketamine experienced a revival among recreational users who were younger, less extreme, more regular consumers of lower doses. These low doses produce a drowsy sensation but do not go as far as the so-called “K-hole”—a “black hole” effect characteristic of ketamine that can include cognitive and amnesic disorders, mood and behavioral problems, hallucinatory delirium, nightmares and loss of identity and contact with reality. This new form of consumption is more conventional, and ketamine is now used more frequently in nightclubs and among friends (“ketamine aperitifs”). A group of inexperienced users consumes ketamine in a more unreasonable way (often combined with high doses of alcohol), and actively seeks the “K-hole” and loss of consciousness.

The main mode of use is snorting, although some people inject ketamine intravenously or intramuscularly.

Few cases of death by overdose have been described, notably because chronic users and poly-users tend to be aware of the doses that should not be exceeded [76–79]. The main injuries reported (bruising, fractures, drowning, etc.) are related to the drug's anesthetic effect.

In France, ketamine is relatively inexpensive, at between 40 and 50 euros per g; this quantity can be divided into several doses [80].

Users also like the fact that in contrast to alcohol, police forces cannot perform roadside tests for ACHs. Furthermore, ketamine's effects wear off quickly (after 20 to 40 min) and so users can (for example) drive a motor vehicle soon after taking the drug.

In addition to the marked neurological effects, digestive disorders (such as diarrhea, nausea and vomiting), tachycardia and hypertension have also been described [81–83].

The medical management of these patients is sometimes complicated by the fact that several other substances have been ingested concomitantly. Management is essentially based on symptomatic treatment, with hyperhydration (to promote substance excretion) and tracheal intubation (to counter the effects of sedation). Clinical biochemistry variables (especially cytolysis markers, creatinine and creatine phosphokinase) and vital signs (ar-

terial oxygen saturation, heart rate and blood pressure) must be monitored closely. An antipsychotic may also need to be administered, as the patients are often agitated [84].

3.1.2. Chronic Toxicity

The toxicity profile of chronic intake is still unclear, but appears to include severe, irreversible damage to the bladder, such as thickening of the bladder wall, leading to secondary kidney damage [85]. Although there are no data on the bladder toxicity of other ACH derivatives, specific clinical monitoring for this entire chemical family makes sense. Digestive disorders and abdominal pain (known as “K-pain”, ranging from simple abdominal pain to intense colic) have also been observed [73,86]. Liver damage has also been described: in 2017, 10 cases of choledochal cysts and dilatation of the bile duct were linked to repeated and/or prolonged use of high-dose ketamine (>100 mg/d). Four of these cases required liver transplantation [73,86–89]. Cognitive problems include addictive behavior, impaired color perception, loss of memory and attention, longer reaction times, impaired perception of time and dissociative effects [90,91].

Ketamine tolerance in humans has been described, although there have been few cases of withdrawal syndrome. Cases of dependence have also been described [88,92–95]. The arrival of new synthetic derivatives on the market will doubtless change this safety profile, with greater variability in doses and product quality.

3.2. Epidemiology

The epidemiology of users of ketamine and its derivatives has changed markedly since the 2000s, with the appearance of a new wave of younger people using lower doses, but in a chronic manner [80]. Ketamine is usually supplied as a liquid formulation for injection, but can be prepared as small white crystals, or even (to facilitate use) as a powder. Some sellers report that ketamine is bought in 1- to 10-liter drums in India, sometimes colored, and then transported in shampoo bottles or other smaller containers. The drug can then be “cooked” into a solid form for easier use. A few cases of theft of hospital or veterinary supplies of ketamine (often of higher quality) have been reported [96]. The drug’s nicknames include “keta”, “ké”, “kéké”, “special K”, “kate”, “pony drug”, and “horse drug”. With the advent of the Internet and the Dark Web, many novel ACH derivatives have attracted interest. Polyconsumption is very frequently described, because ketamine is known to be easily combined with other molecules (e.g., cocaine) and helps to moderate the withdrawal symptoms of other drugs. Asia, and especially Hong Kong, have been greatly affected by the arrival of novel ACH derivatives since the 2000s [18,97]. Even though the raw materials are not easy to obtain, novel ACH derivatives now seem to be spreading across over the world, and consumption is growing exponentially. Ketamine consumption is lower in Europe than in Asia, although the United Kingdom and Spain are markets hubs for drugs produced in Asia. A European Monitoring Center for Drugs and Drug Addictions (EMCDDA) report estimated that only 16 ACHs were reported for the first time between 2005 and 2017—far fewer than the novel synthetic cathinones and cannabinoids [98]. The French TREND/SINTES network has reported significant increases in ketamine dealing and use. Supplies in France are, nevertheless, very limited for the moment, with most of the drug coming from India, the United Kingdom and Spain [99]. Table 2 summarizes the reported cases of ACH-derivative intoxication worldwide, and Figure 4 shows the cases’ geographical distribution.

Table 2. Literature reports of intoxications involving ACH derivatives other than ketamine.

Co-Intoxication		Matrix	Concentration	Mortality	Country	Source
3-MeO-PCP	No	Peripheral blood, urine	Case 1: Peripheral blood = 71.1 ng/mL and urine = 706.9 ng/mL/Case 2 none	2 non-fatal	France	[100]
	Ethanol, diazepam, cocaine	Femoral blood, urine, bile, hair	Femoral blood = 63 ng/mL, bile = 64 ng/mL, urine = 94 ng/mL	1 fatal	France	[101]
	Methadone, THC	Femoral blood, urine	Femoral blood = 3 525 ng/mL and urine = 7 384 ng/mL	1 fatal	France	[64]
	Alcohol	Peripheral blood, urine	Case 1: Peripheral blood = 350.0 ng/mL and urine = 6109.2 ng/mL/Case 2: peripheral blood = 180.1 ng/mL and urine = 3003.5 ng/mL	2 non-fatal	Italy	[102]
	4 cases with 4-MeO-PCP, but the majority included other novel derivatives	Serum, urine	Serum = 1 to 242 ng/mL, urine = 2 to 52,759 ng/mL	56 cases	Sweden	[17]
	Several substances	Femoral blood	Serum = 0.05–0.38 µg/g	1 fatal and 7 non-fatal	Sweden	[103]
	Diphenhydramine, THC, amphetamine	Postmortem blood	Blood = 139 ± 41 µg/L	1 fatal	USA	[104]
	Amphetamine, alcohol	Postmortem blood	Blood = 152 µg/L	1 fatal	The Netherlands	[105]
	Methamphetamine (Case 1), Ethanol/bupropion/paroxetine (Case 2)	Postmortem blood	Blood = Case 1: 0.63 and Case 2: 3.2 mg/L	2 fatal	USA	[106]
	3-OH-PCP, 3-MeO-PCP, 2F-DCK, N-ethylhexedrone, CMC	Urine	Urine = 110 mg/L	1 non-fatal	France	[72]
	Alcohol, amphetamine (Case 1)	Peripheral blood, urine	Blood = 49 ng/mL (Case 1) and 66 ng/mL (Case 2)	2 non-fatal	Czech Republic	[107]
	Several substances	Peripheral and femoral blood, urine, bile, hair	Urine = 498 ng/mL to 16,700 ng/mL Blood = 63 ng/mL	3 non-fatal and 2 fatal	France	[108]
	No data	Femoral blood, urine	Case 1: Femoral blood = 63 ng/mL, bile = 94 ng/mL/Case 2: femoral blood = 498 ng/mL and urine = 16 700 ng/mL	2 fatal	France	[20]
	No data	Urine	Qualitative test	1 non-fatal	USA	[109]

Table 2. Cont.

Co-Intoxication	Matrix	Concentration	Mortality	Country	Source	
2F-DCK	3-MeO-PCE, 5-MeO-DMT, amphetamine and cocaine	Peripheral blood, urine, bile, vitreous humor	Peripheral blood = 1780 µg/L, urine = 6106 µg/L, bile = 12,200 µg/L, vitreous humor = 1500 µg/L	1 fatal	France	[19]
	Majority of ketamine and derivatives, cocaine, methamphetamine	Urine	Qualitative test	20 cases	Hong Kong	[59]
	3-MeO-PCP, 3-OH-PCP, CMC, N-ethylhexedrone	Urine	Urine = 147 mg/L	1 non-fatal	France	[72]
3-OH-PCP	3-MeO-PCP, CMC, 2F-DCK, N-ethylhexedrone	Urine	Urine = 12,085 mg/L	1 non-fatal	France	[72]
3-MeO-PCE	2F-DCK, 5-MeO-DMT, amphetamine and cocaine	Peripheral blood, urine, bile, vitreous humor	Peripheral blood = 90 µg/L, urine = 6.3 µg/L, bile = 3.5 µg/L, vitreous humor = 66 µg/L	1 fatal	France	[19]
MXE	Several substances	Peripheral blood, urine	Qualitative test	8 fatal	UK	[110]
	Alcohol/benzofuran (Case 1)	Serum	Serum = 0.09 to 0,2 mg/L	3 non-fatal	UK	[83]
	Methamphetamine, dextromethorphan	Serum	Serum = 160 ng/mL	1 non-fatal	USA	[111]
	Three synthetic cannabinoids	Femoral blood	Femoral blood = 8.6 µg/g	1 fatal	Sweden	[112]
	Clonazepam, THC, diphenhydramine, MDMA	Peripheral blood	Blood = 10 ng/mL	1 fatal	USA	[113]
	Amphetamine	Peripheral blood	Peripheral blood = 0.32 µg/ml	1 fatal	Poland	[114]
	AH-7921, benzodiazepines	Serum	Unknown	1 non-fatal case	Norway	[115]
	5- or 6-APB	Serum	Serum = 0.16 to 0.45 mg/L	3 non-fatal case	United Kingdom	[116]
	Ketamine, psychotics	Serum, urine, hair	Serum = 30 and urine = 408 µg/L	1 non-fatal case	France	[117]

Table 2. Cont.

	Co-Intoxication	Matrix	Concentration	Mortality	Country	Source
	No data	Serum, urine	Serum = 5.8 µg/mL, urine = 85 µg/mL	1 fatal	Poland	[118]
	No data	Serum, urine	Serum = 270 ng/ml and urine = 660 ng/mL	1 non-fatal	Poland	[119]
2-oxo-PCE	Several substances	Serum, urine	No data	56 non-fatal cases	Hong Kong	[97]
	Venlafaxine	Liver, urine, bile, gastric content, heart blood, femoral blood	Liver = 6137 ng/g, urine = 3468 µg/L, bile fluid = 3290 µg/L, gastric contents = 3086 µg/L, heart blood = 2159 µg/L liquor = 1564 µg/L, femoral blood = 375 µg/L	1 fatal case	Germany	[120]
2-FDCNEK	No data	Urine	Urine (metabolite only: 2-FDCNEK) = 133 ng/mL	1 non-fatal case	China	[68]

THC: tetrahydrocannabinol; CMC: chloromethcathinone; MDMA: 3,4-methyl-enedioxy-methamphetamine; ABP: 6-(2-aminopropyl)benzofuran); AH-7921: 3,4-dichloro-N-[[1-(dimethylamino)cyclohexyl]methyl]benzamide; 5-MeO-DMT: O-methyl-bufotenin.



Figure 4. Geographical distribution of published cases of ACH-derivative intoxication.

3.2.1. Phencyclidine Derivatives

Phencyclidine

In 2013, the US Center for Behavioral Health Statistics and Quality reported that PCP use and related emergency department admissions had increased substantially since the 2000s. Consumption increased by 400% between 2005 and 2011, and then by 200% between 2009 and 2011. This report, as well as Domici et al.'s (2015) observations in a Philadelphia hospital, show that consumption fluctuates greatly over time [121,122].

4-MeO-PCP

Although 4-MeO-PCP was synthesized in the 1960s, it appeared on the narcotics market in 2008 as one of the first “dissociative research chemicals” [4]. Many isomers or other structurally similar molecules have since appeared, which circumvent the legislation on drugs of abuse.

The first notification of 4-MeO-PCP from the European authorities dates back to 2012 [123]. Users described oral intakes between 50 and 100 mg, and 4-MeO-PCP appeared to be less potent than PCP and 3-MeO-PCP. Few cases of intoxication have been described, other than a case in Sweden in 2015 and a case in Korea in 2019 [17,124] (Table 2).

3-MeO-PCP

3-MeO-PCP is an isomer of 4-MeO-PCP. Few metabolic and kinetic data are available, although several cases of intoxication have been published [20,63,107]. 3-MeO-PCP was first noted by the EMCDDA in 2012 [123]. This molecule has been reported in two studies performed in the toxicology laboratory of the Rennes University Medical Center (Rennes, France). Berar et al. (2019) first described 3-MeO-PCP intoxication in a patient, with initial toxic concentrations in blood (71.1 ng/mL) and urine (706.9 ng/mL) [100]. Allard et al. (2019) then used molecular networking to describe the hepatic metabolism of 3-MeO-PCP [63]. In another French study, Grossenbacher et al. (2018) described five cases of 3-MeO-PCP intoxication (two of which were fatal) [108] (Table 2).

3-OH-PCP

3-OH-PCP was first reported in 2009. The drug is reportedly active at low doses (between 1 and 10 mg). However, no cases of intoxication have yet been described in the literature.

3.2.2. Ketamine Derivatives

The “DCKs” (deschloroketamine (DCK) and the fluorinated derivative 2F-deschloroketamine (2F-DCK)) are the least well-described ACH derivatives. The first report on these molecules dates to 2015, with the analysis of powders found in the USA, China and Europe [125]. DCK and 2F-DCK are mainly available on the Dark Web [126]. These derivatives are particularly consumed in Asia in general, and in Hong Kong in particular. Although DCK was synthesized by Stevens in 1962, its misuse was not reported until 2015, and the first analytical data were published in 2016 (187,188).

2F-DCK was first synthesized in 1987, and consumption was first described in Internet forums in 2015 [59]. Between January and July 2019, Tang et al. described 20 cases of patients in whom 2F-DCK was detected [59]. In most cases, 2F-DCK was combined with ketamine. Tang et al. reported that 2F-DCK had the same toxic effects as ketamine, although poly-use prevented the identification of specific clinical effects [127]. Similarly, Weng et al. (2020) described 11 cases of 2F-DCK or DCK poisoning [128]. To broaden the detection window, Davidsen et al. (2020) developed a hair analysis technique, which provided the first pharmacokinetic data from samples collected from roadside offenders in Denmark [129]. Snorted doses of 2F-DCK are generally between 87.5 and 330 mg, whereas those for ketamine are between 60 and 250 mg [75]. There are only two case reports on 2F-DCK consumption: a fatal case of DCK and 2-oxo-PCE intoxication in Germany [113] and our report on a patient at Lille University Medical Center (Lille, France) [19] (Table 2).

3.2.3. Eticyclidine Derivatives

Like the other molecules described here, eticyclidine derivatives are particularly present in Asian countries, and especially in Hong Kong. Although these derivatives are not very common, a few cases of intoxication have yielded clinical and toxicological data (Table 2).

Methoxetamine

Methoxetamine (3-MeO-2-oxo-PCE) appears to be the most widespread ACH in Europe, as dealers sometimes supply it instead of ketamine when supplies of the latter run out [9,17,100,103,130]. Its effects are reportedly similar to those of ketamine, but last longer and are more intense [81,131,132]. The doses are also very different, meaning that the switch from ketamine is particularly dangerous. The first consumer discussions on Internet forums date back to 2010 [133]. Methoxetamine’s nicknames include “Mexxy”, “M-ket”, “MEX”, “Kmax”, and “legal ketamine” [134]. It is the most widely used ketamine derivative in Europe, prompting the European authorities to classify it as a narcotic in 2013 [135]. This is also the case in Japan, the United Kingdom, China, Russia, Turkey and Korea [136]. Nasal and oral administration routes are the most common, although cases of intravenous, anal and sublingual administration have been described [132,134,137]. The most commonly reported doses are 20 to 100 mg for the nasal route, 40 to 100 mg for the oral route and 10 to 80 mg for the intramuscular route [131]. The first effects are felt after 10 to 90 min, with a peak between 1 and 7 h (depending on the administration route) [132,137]. In 2015, the World Health Organization reported 120 cases of non-fatal poisoning and 22 deaths worldwide [136]. The adverse effects are similar to those of ketamine, with nausea/diarrhea/vomiting, tachycardia, hypertension and loss of consciousness [81–83] (Table 2).

2-oxo-PCE

The first report to EMCDDA was made in 2016 [98]. Tang et al. (2018) described a cluster of 56 cases of intoxication in the Hong Kong area in October 2017. The toxidrome was similar to that of ketamine, with predominant tachycardia and hypertension. Users report

dissociative and hallucinogenic effects similar to those of ketamine, but more powerful (five times more than ketamine and three times more than methoxetamine). Oral doses are reportedly between 6 and 12 mg [138,139]. Nasal swab samples from three users contained only 2-oxo-PCE and no ketamine [97] (Table 2).

3-MeO-PCE

Although an assay for 3-MeO-PCE in biological fluids has been developed and validated, there are no published in vivo data on concentrations or metabolism [140].

Chemists have recently developed other structurally similar PCP derivatives (e.g., diarylethylamines), but these are not yet widely used [141] (Table 2).

As reported in Table 2, ACHs are frequently co-consumed with other drugs of abuse, including alcohol, THC, amphetamines or cocaine. These cocktails are likely to modify the pharmacokinetics, pharmacodynamics and toxicity profile of these drugs.

4. Conclusions

ACH consumption is growing exponentially, and is associated with an increase in the number of cases of fatal intoxication. The rapid detection of ACHs is a challenge for toxicology laboratories. Given the small number of reports on intoxication, few data on the metabolism and clinical effects of ACHs have been published. The number of cases has certainly been underestimated because the molecules are not detected or not easily detected. However, constant improvement in analytical techniques and greater access to databases should enable the epidemiology of this phenomenon to be better defined in the future. Assessment of the ACHs' metabolism and pharmacologic properties might make it possible to provide more suitable medical management, rather than symptomatic treatment alone. Metabolic studies will provide insights into the fate of these drugs in the body, the presence of active metabolites, the metabolic pathways involved and, potentially, valuable biomarkers for consolidating diagnostic evidence and broadening detection windows when the parent molecule is no longer detectable.

Author Contributions: Conceptualization: R.P., B.L.D. and T.G.; investigation: R.P., B.L.D., D.L.B., A.K., P.-J.F., I.M. and T.G.; writing—original draft preparation: R.P., B.L.D. and T.G.; writing—review and editing: T.G. and I.M.; visualization: R.P. and D.L.B.; supervision: B.L.D., I.M. and T.G.; funding acquisition: R.P. All authors have read and agreed to the published version of the manuscript.

Funding: This project was supported by financial allowance from “Défi Scientifique” Université Rennes 1. Direction de la Recherche et de l’Innovation; CHU de Rennes-Hôpital Pontchaillou-2, rue Henri Le Guilloux, 35033 Rennes 352 Cedex 9.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors thank David Fraser (Biotech Communication SARL, Ploudalmezeau, France) for copy-editing assistance.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Shafi, A.; Berry, A.J.; Sumnall, H.; Wood, D.M.; Tracy, D.K. New psychoactive substances: A review and updates. *Ther. Adv. Psychopharmacol.* **2020**, *10*, 2045125320967197. [[CrossRef](#)] [[PubMed](#)]
2. Néfau, T.; Martinez, M. Nouveaux Produits de Synthèse Identifiés en France Depuis 2000. L’Observatoire Français des Drogues et Toxicomanies: Paris, France, 2000; Volume 3.
3. Ameline, A.; Alvarez, J.-C.; Dumestre, V.; Eysseric, H.; Gaulier, J.-M.; Kintz, P.; Labat, L.; Lelievre, B.; Muckensturm, A.; Péliissier, A.-L.; et al. Recommandations de La SFTA Pour La Réalisation Des Analyses Toxicologiques Dans Les Cas de Décès Impliquant Des NPS – Version 2019 [SFTA Guidelines for the Achievement of Toxicological Analyzes for Deaths Involving NPS – 2019 Version]. *Toxicol. Anal. Clin.* **2019**, *31*, 337–339. [[CrossRef](#)]
4. Johnstone, M.; Evans, V.; Baigel, S. SERNYL (C1-395) IN CLINICAL ANAESTHESIA. *Br. J. Anaesth.* **1959**, *31*, 433–439. [[CrossRef](#)] [[PubMed](#)]

5. Mion, G. Histoire de la kétamine et du psychédélisme. *Ann. Méd.-Psychol. Rev. Psychiatr.* **2017**, *175*, 661–664. [CrossRef]
6. Bertron, J.L.; Seto, M.; Lindsley, C.W. DARK Classics in Chemical Neuroscience: Phencyclidine (PCP). *ACS. Chem. Neurosci.* **2018**, *9*, 2459–2474. [CrossRef]
7. Jenkins, P. *Synthetic Panics: The Symbolic Politics of Designer Drugs*; NYU Press: New York, NY, USA, 1999; 260p.
8. Harvey, W.; Feldman. Angel Dust in Four American Cities: An Ethnographic Study of PCP Users. Available online: <https://onlinebooks.library.upenn.edu/webbin/book/lookupid?key=ha011389822> (accessed on 24 January 2022).
9. Morris, H.; Wallach, J. From PCP to MXE: A comprehensive review of the non-medical use of dissociative drugs. *Drug Test. Anal.* **2014**, *6*, 614–632. [CrossRef]
10. Li, L.; Vlisides, P.E. Ketamine: 50 Years of Modulating the Mind. *Front. Hum. Neurosci.* **2016**, *10*, 612. [CrossRef]
11. Dundee, J.W. Twenty-five years of ketamine A report of an international meeting. *Anaesthesia* **1990**, *45*, 159–160. [CrossRef]
12. Ketalar. U.S. Food and Drug Administration, Center for Drug Evaluation and Research, in Label and Approval History. 1970. Available online: http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Label_ApprovalHistory#aphist (accessed on 24 January 2022).
13. Orhurhu, V.J.; Vashisht, R.; Claus, L.E.; Cohen, S.P. Ketamine Toxicity. StatPearls Publishing. 2022. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK541087/> (accessed on 15 September 2022).
14. Chidambaran, V.; Costandi, A.; D’Mello, A. Propofol: A review of its role in pediatric anesthesia and sedation. *CNS Drugs* **2015**, *29*, 543–563. [CrossRef]
15. Nowacka, A.; Borczyk, M. Ketamine applications beyond anesthesia—A literature review. *Eur. J. Pharmacol.* **2019**, *860*, 172547. [CrossRef]
16. Inscription sur la Liste des Stupéfiants des Préparations Injectables à base de Kétamine à Compter du 24 Avril 2017-Point D’information-ANSM: Agence Nationale de Sécurité du Médicament et des Produits de Santé. Available online: <https://ansm.sante.fr/S-informer/Points-d-information-Points-d-information/Inscription-sur-la-liste-des-stupefiants-des-preparations-injectables-a-base-de-ketamine-a-compter-du-24-avril-2017-Point-d-Information> (accessed on 19 December 2020).
17. Bäckberg, M.; Beck, O.; Helander, A. Phencyclidine analog use in Sweden—intoxication cases involving 3-MeO-PCP and 4-MeO-PCP from the STRIDA project. *Clin. Toxicol. Phila. Pa.* **2015**, *53*, 856–864. [CrossRef]
18. Chong, Y.K.; Tang, M.H.; Chan, C.L.; Li, Y.K.; Ching, C.K.; Mak, T.W. 2-oxo-PCE: Ketamine analogue on the streets. *Hong Kong Med. J. Xianggang Yi Xue Za Zhi* **2017**, *23*, 665–666. [CrossRef]
19. Gicquel, T.; Richeval, C.; Mesli, V.; Gish, A.; Hakim, F.; Pelletier, R.; Raphael, C.; Axelle, B.; Delphine, A.; Jean-Michel, G. Fatal intoxication related to two new arylcyclohexylamine derivatives (2F-DCK and 3-MeO-PCE). *Forensic. Sci. Int.* **2021**, *324*, 110852. [CrossRef]
20. Ameline, A.; Greney, H.; Monassier, L.; Raul, J.S.; Kintz, P. Metabolites to parent 3-MeO-PCP ratio in human urine collected in two fatal cases. *J. Anal. Toxicol.* **2019**, *43*, 321–324. [CrossRef]
21. Gicquel, T.; Pelletier, R.; Richeval, C.; Gish, A.; Hakim, F.; Ferron, P.-J.; Mesli, V.; Allorge, D.; Morel, I.; Gaulier, J.-M. Metabolite Elucidation of 2-Fluoro-Deschloroketamine (2F-DCK) Using Molecular Networking across Three Complementary in Vitro and in Vivo Models. *Drug Test. Anal.* **2021**, *14*, 144–153. [CrossRef]
22. Ihmsen, H.; Geisslinger, G.; Schüttler, J. Stereoselective pharmacokinetics of ketamine: R(-)-ketamine inhibits the elimination of S(+)-ketamine. *Clin. Pharmacol. Ther.* **2001**, *70*, 431–438. [CrossRef]
23. Zorumski, C.F.; Izumi, Y.; Mennerick, S. Ketamine: NMDA Receptors and Beyond. *J. Neurosci. Off. J. Soc. Neurosci.* **2016**, *36*, 11158–11164. [CrossRef]
24. Vyklícky, V.; Korinek, M.; Smejkalova, T.; Balik, A.; Krausova, B.; Kaniakova, M. Structure, function, and pharmacology of NMDA receptor channels. *Physiol. Res.* **2014**, *63* (Suppl. S1), S191–S203. [CrossRef]
25. Rudgley, R. *The Encyclopedia of Psychoactive Substances*; Macmillan: New York, NY, USA, 2014; 321p.
26. Dinis-Oliveira, R.J. Metabolism and metabolomics of ketamine: A toxicological approach. *Forensic. Sci. Res.* **2017**, *2*, 2–10. [CrossRef]
27. Karinen, R.; Høiseith, G. A literature review of blood concentrations of new psychoactive substances classified as phenethylamines, aminoindanes, arylalkylamines, arylcyclohexylamines, and indolalkylamines. *Forensic. Sci. Int.* **2017**, *276*, 120–125. [CrossRef]
28. Morris, P.J.; Moaddel, R.; Zanos, P.; Moore, C.E.; Gould, T.D.; Zarate, C.A.; Thomas, C.J. Synthesis and N-Methyl-d-Aspartate (NMDA) Receptor Activity of Ketamine Metabolites. *Org. Lett.* **2017**, *19*, 4572–4575. [CrossRef] [PubMed]
29. Pham, T.H.; Mendez-David, I.; Defaix, C.; Guiard, B.P.; Tritschler, L.; David, D.J.; Gardier, A.M. Ketamine Treatment Involves Medial Prefrontal Cortex Serotonin to Induce a Rapid Antidepressant-like Activity in BALB/CJ Mice. *Neuropharmacology* **2017**, *112*, 198–209. [CrossRef] [PubMed]
30. Stein, C. Opioid Receptors. *Annu. Rev. Med.* **2016**, *67*, 433–451. [CrossRef] [PubMed]
31. Lois, F.; De Kock, M. Something new about ketamine for pediatric anesthesia? *Curr. Opin. Anaesthesiol.* **2008**, *21*, 340–344. [CrossRef] [PubMed]
32. Petrenko, A.B.; Yamakura, T.; Baba, H.; Shimoji, K. The role of N-methyl-D-aspartate (NMDA) receptors in pain: A review. *Anesth. Analg.* **2003**, *97*, 1108–1116. [CrossRef]
33. Marchand, F.; Perretti, M.; McMahon, S.B. Role of the immune system in chronic pain. *Nat. Rev. Neurosci.* **2005**, *6*, 521–532. [CrossRef]

34. Watkins, L.R.; Maier, S.F. Immune regulation of central nervous system functions: From sickness responses to pathological pain. *J. Intern. Med.* **2005**, *257*, 139–155. [[CrossRef](#)]
35. Ossipov, M.H.; Dussor, G.O.; Porreca, F. Central modulation of pain. *J. Clin. Invest.* **2010**, *120*, 3779–3787. [[CrossRef](#)]
36. Woolf, C.J. Central sensitization: Implications for the diagnosis and treatment of pain. *Pain* **2011**, *152* (Suppl. S3), S2–S15. [[CrossRef](#)]
37. Segmiller, F.; Rütther, T.; Linhardt, A.; Padberg, F.; Berger, M.; Pogarell, O.; Möller, H.-J.; Kohler, C.; Schüle, C. Repeated S-Ketamine Infusions in Therapy Resistant Depression: A Case Series. *J. Clin. Pharmacol.* **2013**, *53*, 996–998. [[CrossRef](#)]
38. Scheuing, L.; Chiu, C.T.; Liao, H.M.; Chuang, D.M. Antidepressant mechanism of ketamine: Perspective from preclinical studies. *Front. Neurosci.* **2015**, *9*, 249. [[CrossRef](#)]
39. Liu, R.J.; Fuchikami, M.; Dwyer, J.M.; Lepack, A.E.; Duman, R.S.; Aghajanian, G.K. GSK-3 inhibition potentiates the synaptogenic and antidepressant-like effects of subthreshold doses of ketamine. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* **2013**, *38*, 2268–2277. [[CrossRef](#)]
40. Schüttler, J.; Schwilden, H.; Modern Anesthetics. Springer-Verlag: Berlin Heidelberg, Germany. 2008. (Handbook of Experimental Pharmacology). Available online: <https://www.springer.com/gp/book/9783540728139> (accessed on 22 March 2021).
41. Carr, D.B.; Goudas, L.C.; Denman, W.T.; Brookoff, D.; Staats, P.S.; Brennen, L.; Green, G.; Albin, R.; Hamilton, D.; Rogers, M.C.; et al. Safety and Efficacy of Intranasal Ketamine for the Treatment of Breakthrough Pain in Patients with Chronic Pain: A Randomized, Double-Blind, Placebo-Controlled, Crossover Study. *Pain* **2004**, *108*, 17–27. [[CrossRef](#)]
42. Fanta, S.; Kinnunen, M.; Backman, J.T.; Kalso, E. Population pharmacokinetics of S-ketamine and norketamine in healthy volunteers after intravenous and oral dosing. *Eur. J. Clin. Pharmacol.* **2015**, *71*, 441–447. [[CrossRef](#)]
43. Dayton, P.G.; Stiller, R.L.; Cook, D.R.; Perel, J.M. The binding of ketamine to plasma proteins: Emphasis on human plasma. *Eur. J. Clin. Pharmacol.* **1983**, *24*, 825–831. [[CrossRef](#)]
44. Sinner, B.; Graf, B.M. Ketamine. *Handb. Exp. Pharmacol.* **2008**, *182*, 313–333.
45. Peltoniemi, M.A.; Hagelberg, N.M.; Olkkola, K.T.; Saari, T.I. Ketamine: A Review of Clinical Pharmacokinetics and Pharmacodynamics in Anesthesia and Pain Therapy. *Clin. Pharmacokinet.* **2016**, *55*, 1059–1077. [[CrossRef](#)]
46. Clements, J.A.; Nimmo, W.S.; Grant, I.S. Bioavailability, pharmacokinetics, and analgesic activity of ketamine in humans. *J. Pharm. Sci.* **1982**, *71*, 539–542. [[CrossRef](#)]
47. Grant, I.S.; Nimmo, W.S.; Clements, J.A. Pharmacokinetics and analgesic effects of i.m. and oral ketamine. *Br. J. Anaesth.* **1981**, *53*, 805–810. [[CrossRef](#)]
48. Hagelberg, N.M.; Peltoniemi, M.A.; Saari, T.I.; Kurkinen, K.J.; Laine, K.; Neuvonen, P.J.; Olkkola, K.T. Clarithromycin, a potent inhibitor of CYP3A, greatly increases exposure to oral S-ketamine. *Eur. J. Pain. Lond. Engl.* **2010**, *14*, 625–629. [[CrossRef](#)]
49. Persson, J.; Hasselström, J.; Maurset, A.; Öye, I.; Svensson, J.O.; Almqvist, O.; Gustafsson, L. Pharmacokinetics and non-analgesic effects of S- and R-ketamines in healthy volunteers with normal and reduced metabolic capacity. *Eur. J. Clin. Pharmacol.* **2002**, *57*, 869–875. [[CrossRef](#)] [[PubMed](#)]
50. Zanos, P.; Gould, T.D. Mechanisms of ketamine action as an antidepressant. *Mol. Psychiatry* **2018**, *23*, 801–811. [[CrossRef](#)] [[PubMed](#)]
51. Rao, L.K.; Flaker, A.M.; Friedel, C.C.; Kharasch, E.D. Role of Cytochrome P4502B6 Polymorphisms in Ketamine Metabolism and Clearance. *Anesthesiology* **2016**, *125*, 1103–1112. [[CrossRef](#)] [[PubMed](#)]
52. Desta, Z.; Moaddel, R.; Ogburn, E.T.; Xu, C.; Ramamoorthy, A.; Venkata SL, V.; Wainer, I.W. Stereoselective and regiospecific hydroxylation of ketamine and norketamine. *Xenobiotica Fate Foreign Compd. Biol. Syst.* **2012**, *42*, 1076–1087. [[CrossRef](#)] [[PubMed](#)]
53. Portmann, S.; Kwan, H.Y.; Theurillat, R.; Schmitz, A.; Mevissen, M.; Thormann, W. Enantioselective capillary electrophoresis for identification and characterization of human cytochrome P450 enzymes which metabolize ketamine and norketamine in vitro. *J. Chromatogr. A* **2010**, *1217*, 7942–7948. [[CrossRef](#)]
54. Hijazi, Y.; Bolon, M.; Bouliou, R. Stability of Ketamine and Its Metabolites Norketamine and Dehydronorketamine in Human Biological Samples. *Clin. Chem.* **2001**, *47*, 1713–1715. [[CrossRef](#)]
55. Moaddel, R.; Venkata SL, V.; Tanga, M.J.; Bupp, J.E.; Green, C.E.; Iyer, L.; Wainer, I.W. A parallel chiral-achiral liquid chromatographic method for the determination of the stereoisomers of ketamine and ketamine metabolites in the plasma and urine of patients with complex regional pain syndrome. *Talanta* **2010**, *82*, 1892–1904. [[CrossRef](#)]
56. Adams, J.D.; Baillie, T.A.; Trevor, A.J.; Castagnoli, N. Studies on the biotransformation of ketamine. 1-Identification of metabolites produced in vitro from rat liver microsomal preparations. *Biomed. Mass. Spectrom* **1981**, *8*, 527–538. [[CrossRef](#)]
57. Karch's Pathology of Drug Abuse. CRC Press. 2015. Available online: <https://www.taylorfrancis.com/books/karch-pathology-drug-abuse-hank-ellison-olaf-drummer/10.1201/b18962> (accessed on 10 February 2021).
58. Mestria, S.; Odoardi, S.; Biosia, G.; Valentini, V.; Di Masi, G.; Cittadini, F.; Strano-Rossi, S. Method development for the identification of methoxpropamine, 2-fluoro-deschloroketamine and deschloroketamine and their main metabolites in blood and hair and forensic application. *Forensic. Sci. Int.* **2021**, *323*, 110817. [[CrossRef](#)]
59. Tang, M.H.Y.; Li, T.C.; Lai, C.K.; Chong, Y.K.; Ching, C.K.; Mak, T.W.L. Emergence of new psychoactive substance 2-fluorodeschloroketamine: Toxicology and urinary analysis in a cluster of patients exposed to ketamine and multiple analogues. *Forensic. Sci. Int.* **2020**, *312*, 110327. [[CrossRef](#)]
60. Laurenzana, E.M.; Owens, S.M. Metabolism of Phencyclidine by Human Liver Microsomes. *Drug Metab. Dispos.* **1997**, *25*, 557–563.

61. Meyer, M.R.; Maurer, H.H. Absorption, distribution, metabolism and excretion pharmacogenomics of drugs of abuse. *Pharmacogenomics* **2011**, *12*, 215–233. [CrossRef]
62. Davidsen, A.B.; Mardal, M.; Johansen, S.S.; Dalsgaard, P.W.; Linnet, K. In vitro and in vivo metabolism and detection of 3-HO-PCP, a synthetic phencyclidine, in human samples and pooled human hepatocytes using high resolution mass spectrometry. *Drug Test. Anal.* **2020**, *12*, 987–993. [CrossRef]
63. Allard, S.; Allard, P.M.; Morel, I.; Gicquel, T. Application of a molecular networking approach for clinical and forensic toxicology exemplified in three cases involving 3-MeO-PCP, doxylamine, and chlormequat. *Drug Test. Anal.* **2019**, *11*, 669–677. [CrossRef]
64. Arbouche, N.; Kintz, P.; Zagdoun, C.; Gheddar, L.; Raul, J.S.; Ameline, A. Determination of 3-MeO-PCP in human blood and urine in a fatal intoxication case, with a specific focus on metabolites identification. *Forensic. Sci. Res.* **2021**, *6*, 208–214. [CrossRef]
65. Michely, J.; Manier, S.; Caspar, A.; Brandt, S.; Wallach, J.; Maurer, H. New Psychoactive Substances 3-Methoxyphencyclidine (3-MeO-PCP) and 3-Methoxyrolyclidine (3-MeO-PCPy): Metabolic Fate Elucidated with Rat Urine and Human Liver Preparations and their Detectability in Urine by GC-MS, “LC-(High Resolution)-MSn” and “LC-(High Resolution)-MS/MS”. *Curr. Neuropharmacol.* **2017**, *15*, 692–712.
66. Goncalves, R.; Castaing, N.; Richeval, C.; Ducint, D.; Titier, K.; Morvan, E.; Molimard, M. Methoxpropamine (MXPr) in powder, urine and hair samples: Analytical characterization and metabolite identification of a new threat. *Forensic. Sci. Int.* **2022**, *333*, 111215. [CrossRef]
67. Larabi, I.A.; Zerizer, F.; Ameline, A.; Etting, I.; Joseph, D.; Kintz, P.; Alvarez, J.C. Metabolic Profiling of Deschloro-N-Ethyl-Ketamine and Identification of New Target Metabolites in Urine and Hair Using Human Liver Microsomes and High-Resolution Accurate Mass Spectrometry. *Drug Test. Anal.* **2021**, *13*, 1108–1117. [CrossRef]
68. Yen, Y.T.; Tsai, Y.S.; Su, W.L.; Huang, D.Y.; Wu, H.H.; Tseng, S.H.; Chyueh, S.C. New ketamine analogue: 2-fluorodeschloro-N-ethyl-ketamine and its suggested metabolites. *Forensic. Sci. Int.* **2022**, *341*, 111501. [CrossRef]
69. Chang, T.; Glazko, A.J. Biotransformation and disposition of ketamine. *Int. Anesthesiol. Clin.* **1974**, *12*, 157–177. [CrossRef]
70. Adamowicz, P.; Kala, M. Urinary excretion rates of ketamine and norketamine following therapeutic ketamine administration: Method and detection window considerations. *J. Anal. Toxicol.* **2005**, *29*, 376–382. [CrossRef]
71. Peltoniemi, M.A.; Saari, T.I.; Hagelberg, N.M.; Laine, K.; Neuvonen, P.J.; Olkkola, K.T. St John’s wort greatly decreases the plasma concentrations of oral S-ketamine. *Fundam. Clin. Pharmacol.* **2012**, *26*, 743–750. [CrossRef] [PubMed]
72. Pelletier, R.; Le Dare, B.; Grandin, L.; Couette, A.; Ferron, P.J.; Morel, I.; Gicquel, T. New psychoactive substance cocktail in an intensive care intoxication case elucidated by molecular networking. *Clin. Toxicol. Phila. Pa.* **2021**, *60*, 1–4. [CrossRef] [PubMed]
73. Jansen, K.L. A review of the nonmedical use of ketamine: Use, users and consequences. *J. Psychoact. Drugs.* **2000**, *32*, 419–433. [CrossRef]
74. Curran, H.V.; Morgan, C. Cognitive, dissociative and psychotogenic effects of ketamine in recreational users on the night of drug use and 3 days later. *Addict. Abingdon. Engl.* **2000**, *95*, 575–590. [CrossRef] [PubMed]
75. Rao, S.S.; Wood, D.M.; Dargan, P.I. Ketamine—Epidemiology of Misuse and Patterns of Acute and Chronic Toxicity. In *Ketamine*; CRC Press: Boca Raton, FL, USA, 2015.
76. Tao, Y.; Chen, X.P.; Qin, Z.H. A fatal chronic ketamine poisoning. *J. Forensic. Sci.* **2005**, *50*, 173–176. [CrossRef]
77. Licata, M.; Pierini, G.; Popoli, G. A fatal ketamine poisoning. *J. Forensic. Sci.* **1994**, *39*, 1314–1320. [CrossRef]
78. Moore, K.A.; Kilbane, E.M.; Jones, R.; Kunsman, G.W.; Levine, B.; Smith, M. Tissue distribution of ketamine in a mixed drug fatality. *J. Forensic. Sci.* **1997**, *42*, 1183–1185. [CrossRef]
79. Lalonde, B.R.; Wallage, H.R. Postmortem blood ketamine distribution in two fatalities. *J. Anal. Toxicol.* **2004**, *28*, 71–74. [CrossRef]
80. Gandilhon, M.; Cadet-Tairou, A.; Martinez, M. *L’usage de Kétamine En France: Tendances Récentes (2012-2013)*; OFDT: Saint-Denis, 2014 Gandilhon M, Cadet-Tairou A, Martinez M. *L’usage de kétamine en France: Tendances récentes (2012-2013)*. OFDT, 2014, Note 2014-7. Available online: <https://www.ofdt.fr/BDD/publications/docs/eisxmg6.pdf> (accessed on 25 January 2022).
81. Hofer, K.E.; Grager, B.; Müller, D.M.; Rauber-Lüthy, C.; Kupferschmidt, H.; Rentsch, K.M.; Ceschi, A. Ketamine-like effects after recreational use of methoxetamine. *Ann. Emerg. Med.* **2012**, *60*, 97–99. [CrossRef]
82. Advisory Council on the Misuse of Drugs (ACMD) Methoxetamine report. 2012; GOV.UK. Available online: <https://www.gov.uk/government/publications/advisory-council-on-the-misuse-of-drugs-acmd-methoxetamine-report-2012> (accessed on 9 March 2021).
83. Wood, D.M.; Davies, S.; Puchnarewicz, M.; Johnston, A.; Dargan, P.I. Acute toxicity associated with the recreational use of the ketamine derivative methoxetamine. *Eur. J. Clin. Pharmacol.* **2012**, *68*, 853–856. [CrossRef]
84. MacNeal, J.J.; Cone, D.C.; Sinha, V.; Tomassoni, A.J. Use of haloperidol in PCP-intoxicated individuals. *Clin. Toxicol. Phila. Pa.* **2012**, *50*, 851–853. [CrossRef]
85. Winstock, A.R.; Mitcheson, L.; Gillatt, D.A.; Cottrell, A.M. The prevalence and natural history of urinary symptoms among recreational ketamine users. *BJU Int.* **2012**, *110*, 1762–1766. [CrossRef]
86. Morgan, C.J.A.; Curran, H.V. Independent Scientific Committee on Drugs. Ketamine use: A review. *Addict. Abingdon. Engl.* **2012**, *107*, 27–38. [CrossRef]
87. Wong, S.W.; Lee, K.F.; Wong, J.; Ng, W.W.C.; Cheung, Y.S.; Lai, P.B.S. Dilated common bile ducts mimicking choledochal cysts in ketamine abusers. *Hong Kong Med. J. Xianggang Yi Xue Za Zhi* **2009**, *15*, 53–56.
88. Sassano-Higgins, S.; Baron, D.; Juarez, G.; Esmaili, N.; Gold, M. A REVIEW OF KETAMINE ABUSE AND DIVERSION. *Depress. Anxiety* **2016**, *33*, 718–727. [CrossRef]

89. Kétamine: Risque D'atteintes Hépatiques Graves Lors D'utilisations Prolongées et/ou à Doses élevées-Point D'information-ANSM: Agence Nationale de Sécurité du Médicament et des Produits de Santé. Available online: <https://ansm.sante.fr/S-informer/Points-d-information-Points-d-information/Ketamine-risque-d-atteintes-hepatiques-graves-lors-d-utilisations-prolongees-et-ou-a-doses-elevees-Point-d-Information> (accessed on 18 February 2021).
90. Bokor, G.; Anderson, P.D. Ketamine: An update on its abuse. *J. Pharm. Pract.* **2014**, *27*, 582–586. [CrossRef]
91. Liao, Y.; Tang, J.; Ma, M.; Wu, Z.; Yang, M.; Wang, X.; Hao, W. Frontal white matter abnormalities following chronic ketamine use: A diffusion tensor imaging study. *Brain J. Neurol.* **2010**, *133*, 2115–2122. [CrossRef]
92. Byer, D.E.; Gould, A.B. Development of tolerance to ketamine in an infant undergoing repeated anesthesia. *Anesthesiology* **1981**, *54*, 255–256. [CrossRef]
93. Cheng, W.J.; Chen, C.H.; Chen, C.K.; Huang, M.C.; Pietrzak, R.H.; Krystal, J.H.; Xu, K. Similar psychotic and cognitive profile between ketamine dependence with persistent psychosis and schizophrenia. *Schizophr. Res.* **2018**, *199*, 313–318. [CrossRef]
94. Garg, A.; Sinha, P.; Kumar, P.; Prakash, O. Use of naltrexone in ketamine dependence. *Addict Behav.* **2014**, *39*, 1215–1216. [CrossRef] [PubMed]
95. Pal, H.R.; Berry, N.; Kumar, R.; Ray, R. Ketamine dependence. *Anaesth. Intensive Care* **2002**, *30*, 382–384. [CrossRef] [PubMed]
96. Shbair, M.K.S.; Eljabour, S.; Lhermitte, M. Drugs involved in drug-facilitated crimes: Part I: Alcohol, sedative-hypnotic drugs, gamma-hydroxybutyrate and ketamine. A review. *Ann. Pharm. Fr.* **2010**, *68*, 275–285. [CrossRef] [PubMed]
97. Tang, M.H.; Chong, Y.K.; Chan, C.Y.; Ching, C.K.; Lai, C.K.; Li, Y.K.; Mak, T.W. Cluster of acute poisonings associated with an emerging ketamine analogue, 2-oxo-PCE. *Forensic. Sci. Int.* **2018**, *290*, 238–243. [CrossRef] [PubMed]
98. EMCDDA–Europol 2017 Annual Report on the Implementation of Council Decision 2005/387/JHA. Available online: https://www.emcdda.europa.eu/publications/implementation-reports/2017_en (accessed on 8 March 2021).
99. 2000 Annual Report on the State of the Drugs Problem in the European Union. Available online: https://www.emcdda.europa.eu/html.cfm/index37279EN.html_en (accessed on 17 February 2021).
100. Berar, A.; Allain, J.S.; Allard, S.; Lefevre, C.; Baert, A.; Morel, I.; Gicquel, T. Intoxication with 3-MeO-PCP alone: A case report and literature review. *Medicine* **2019**, *98*, e18295. [CrossRef]
101. Kintz, P.; Ameline, A.; Walch, A.; Farrugia, A.; Raul, J.S. Murdered while under the influence of 3-MeO-PCP. *Int. J. Legal Med.* **2019**, *133*, 475–478. [CrossRef]
102. Bertol, E.; Pascali, J.; Palumbo, D.; Catalani, V.; Di Milia, M.G.; Fioravanti, A.; Vaiano, F. 3-MeO-PCP intoxication in two young men: First in vivo detection in Italy. *Forensic. Sci. Int.* **2017**, *274*, 7–12. [CrossRef]
103. Johansson, A.; Lindstedt, D.; Roman, M.; Thelander, G.; Nielsen, E.I.; Lennborn, U.; Kugelberg, F.C. A non-fatal intoxication and seven deaths involving the dissociative drug 3-MeO-PCP. *Forensic. Sci. Int.* **2017**, *275*, 76–82. [CrossRef]
104. Bakota, E.; Arndt, C.; Romoser, A.A.; Wilson, S.K. Fatal Intoxication Involving 3-MeO-PCP: A Case Report and Validated Method. *J. Anal. Toxicol.* **2016**, *40*, 504–510. [CrossRef]
105. de Jong, L.A.A.; Olyslager, E.J.H.; Duijst, W.L.J.M. The risk of emerging new psychoactive substances: The first fatal 3-MeO-PCP intoxication in The Netherlands. *J. Forensic. Leg. Med.* **2019**, *65*, 101–104. [CrossRef]
106. Mitchell-Mata, C.; Thomas, B.; Peterson, B.; Couper, F. Two Fatal Intoxications Involving 3-Methoxyphencyclidine. *J. Anal. Toxicol.* **2017**, *41*, 503–507. [CrossRef]
107. Zidkova, M.; Hlozek, T.; Balik, M.; Kopecky, O.; Tesinsky, P.; Svanda, J.; Balikova, M.A. Two Cases of Non-fatal Intoxication with a Novel Street Hallucinogen: 3-Methoxy-Phencyclidine. *J. Anal. Toxicol.* **2017**, *41*, 350–354. [CrossRef]
108. Grossebacher, F.; Cazaubon, Y.; Feliu, C.; Ameline, A.; Kintz, P.; Passouant, O.; Mourvillier, B.; Djerada, Z. About 5 Cases with 3 Meo-PCP Including 2 Deaths and 3 Non-Fatal Cases Seen in France in 2018. *Toxicol. Anal. Clin.* **2019**, *31*, 332–336. [CrossRef]
109. Chang, B.N.; Smith, M.P. A Case of Unusual Drug Screening Results. *Clin. Chem.* **2017**, *63*, 958–961. [CrossRef]
110. Chiappini, S.; Claridge, H.; Corkery, J.M.; Goodair, C.; Loi, B.; Schifano, F. Methoxetamine-related deaths in the UK: An overview. *Hum. Psychopharmacol.* **2015**, *30*, 244–248. [CrossRef]
111. Fassette, T.; Martinez, A. An Impaired Driver Found to be Under the Influence of Methoxetamine. *J. Anal. Toxicol.* **2016**, *40*, 700–702. [CrossRef]
112. Wikström, M.; Thelander, G.; Dahlgren, M.; Kronstrand, R. An accidental fatal intoxication with methoxetamine. *J. Anal. Toxicol.* **2013**, *37*, 43–46. [CrossRef]
113. Elian, A.A.; Hackett, J. A polydrug intoxication involving methoxetamine in a drugs and driving case. *J. Forensic. Sci.* **2014**, *59*, 854–858. [CrossRef]
114. Wiergowski, M.; Anand, J.; Krzyzanowski, M.; Jankowski, Z. Acute methoxetamine and amphetamine poisoning with fatal outcome: A case report. *Int. J. Occup. Med. Environ. Health* **2014**, *27*, 683–690. [CrossRef]
115. Karinen, R.; Tuv, S.S.; Rogde, S.; Peres, M.D.; Johansen, U.; Frost, J.; Øiestad, Å.M.L. Lethal poisonings with AH-7921 in combination with other substances. *Forensic. Sci. Int.* **2014**, *244*, e21–e24. [CrossRef]
116. Shields, J.E.; Dargan, P.I.; Wood, D.M.; Puchnarewicz, M.; Davies, S.; Waring, W.S. Methoxetamine associated reversible cerebellar toxicity: Three cases with analytical confirmation. *Clin. Toxicol. Phila. Pa.* **2012**, *50*, 438–440. [CrossRef] [PubMed]
117. Imbert, L.; Boucher, A.; Delhome, G.; Cueto, T.; Boudinaud, M.; Maublanc, J.; Gaulier, J.M. Analytical findings of an acute intoxication after inhalation of methoxetamine. *J. Anal. Toxicol.* **2014**, *38*, 410–415. [CrossRef] [PubMed]
118. Adamowicz, P.; Zuba, D. Fatal intoxication with methoxetamine. *J. Forensic. Sci.* **2015**, *60* (Suppl. S1), S264–S268. [CrossRef] [PubMed]

119. Łukasik-Glebocka, M.; Sommerfeld, K.; Tezyk, A.; Zielińska-Psujka, B.; Druzd, A. Acute methoxetamine intoxication—a case report with serum and urine concentrations. *Przegl. Lek.* **2013**, *70*, 671–673. [PubMed]
120. Theofel, N.; Möller, P.; Vejmelka, E.; Kastner, K.; Roscher, S.; Scholtis, S.; Tsokos, M. A Fatal Case Involving N-Ethyl-deschloroketamine (2-Oxo-PCE) and Venlafaxine. *J. Anal. Toxicol.* **2019**, *43*, e2–e6. [CrossRef]
121. Bush, D.M. Emergency Department Visits Involving Phencyclidine (PCP) In The CBHSQ Report. Rockville (MD): Substance Abuse and Mental Health Services Administration (US). 2013. Available online: <http://www.ncbi.nlm.nih.gov/books/NBK385063/> (accessed on 8 March 2021).
122. Dominici, P.; Kopec, K.; Manur, R.; Khalid, A.; Damiron, K.; Rowden, A. Phencyclidine Intoxication Case Series Study. *J. Med. Toxicol. Off. J. Am. Coll. Med. Toxicol.* **2015**, *11*, 321–325. [CrossRef]
123. EMCDDA–Europol 2012 Annual Report on the Implementation of Council Decision 2005/387/JHA (New Drugs in Europe, 2012). Available online: https://www.emcdda.europa.eu/publications/implementation-reports/2012_en (accessed on 9 March 2021).
124. Abiero, A.; Botanas, C.J.; Custodio, R.J.; Sayson, L.V.; Kim, M.; Lee, H.J.; Cheong, J.H. 4-MeO-PCP and 3-MeO-PCMo, new dissociative drugs, produce rewarding and reinforcing effects through activation of mesolimbic dopamine pathway and alteration of accumbal CREB, deltaFosB, and BDNF levels. *Psychopharmacology* **2020**, *237*, 757–772. [CrossRef]
125. Wallach, J.; Brandt, S.D. 1,2-Diarylethylamine- and Ketamine-Based New Psychoactive Substances. *Handb. Exp. Pharmacol.* **2018**, *252*, 305–352.
126. Van Hout, M.C.; Hearne, E. New psychoactive substances (NPS) on cryptomarket fora: An exploratory study of characteristics of forum activity between NPS buyers and vendors. *Int. J. Drug Policy.* **2017**, *40*, 102–110. [CrossRef]
127. Li, C.; Lai, C.K.; Tang, M.H.Y.; Chan, C.C.K.; Chong, Y.K.; Mak, T.W.L. Ketamine analogues multiplying in Hong Kong. *Hong Kong Med J Xianggang Yi Xue Za Zhi* **2019**, *25*, 169. [CrossRef]
128. Weng, T.I.; Chin, L.W.; Chen, L.Y.; Chen, J.Y.; Chen, G.Y.; Fang, C.C. Clinical characteristics of patients admitted to emergency department for the use of ketamine analogues with or without other new psychoactive substances. *Clin. Toxicol. Phila. Pa.* **2020**, *59*, 528–531. [CrossRef] [PubMed]
129. Davidsen, A.B.; Mardal, M.; Holm, N.B.; Andreassen, A.K.; Johansen, S.S.; Noble, C.; Linnet, K. Ketamine analogues: Comparative toxicokinetic in vitro-in vivo extrapolation and quantification of 2-fluorodeschloroketamine in forensic blood and hair samples. *J. Pharm. Biomed. Anal.* **2020**, *180*, 113049. [CrossRef]
130. Zarantonello, P.; Bettini, E.; Paio, A.; Simoncelli, C.; Terreni, S.; Cardullo, F. Novel analogues of ketamine and phencyclidine as NMDA receptor antagonists. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 2059–2063. [CrossRef] [PubMed]
131. Zawilska, J.B. Methoxetamine—a novel recreational drug with potent hallucinogenic properties. *Toxicol. Lett.* **2014**, *230*, 402–407. [CrossRef] [PubMed]
132. Kjellgren, A.; Jonsson, K. Methoxetamine (MXE)—A Phenomenological Study of Experiences Induced by a “Legal High” from the Internet. *J. Psychoact. Drugs* **2013**, *45*, 276–286. [CrossRef]
133. EMCDDA home page. Available online: <https://www.emcdda.europa.eu/> (accessed on 23 November 2020).
134. Roth, B.L.; Gibbons, S.; Arunotayanun, W.; Huang, X.P.; Setola, V.; Treble, R.; Iversen, L. The ketamine analogue methoxetamine and 3- and 4-methoxy analogues of phencyclidine are high affinity and selective ligands for the glutamate NMDA receptor. *PLoS ONE* **2013**, *8*, e59334. [CrossRef]
135. Report on the Risk Assessment of 2-(3-Methoxyphenyl)-2-(Ethylamino)Cyclohexanone (Methoxetamine) in the Framework of the Council Decision on New Psychoactive Substances. Available online: https://www.emcdda.europa.eu/publications/risk-assessment/methoxetamine_en (accessed on 26 January 2022).
136. Methoxetamine-Critical Review Report (WHO). Available online: https://legal-high-inhaltsstoffe.de/sites/default/files/uploads/methoxetamine_-_mxe.pdf (accessed on 15 September 2022).
137. Corazza, O.; Schifano, F.; Simonato, P.; Fergus, S.; Assi, S.; Stair, J.; Scherbaum, N. Phenomenon of new drugs on the Internet: The case of ketamine derivative methoxetamine. *Hum. Psychopharmacol. Clin. Exp.* **2012**, *27*, 145–149. [CrossRef]
138. Erowid Ketamine Vault: Dosage. Available online: https://erowid.org/chemicals/ketamine/ketamine_dose.shtml (accessed on 10 March 2021).
139. Erowid Methoxetamine Vault: Dose/Dosage. Available online: https://erowid.org/chemicals/methoxetamine/methoxetamine_dose.shtml (accessed on 10 March 2021).
140. De Paoli, G.; Brandt, S.D.; Wallach, J.; Archer, R.P.; Pounder, D.J. From the street to the laboratory: Analytical profiles of methoxetamine, 3-methoxyeticyclidine and 3-methoxyphencyclidine and their determination in three biological matrices. *J. Anal. Toxicol.* **2013**, *37*, 277–283. [CrossRef]
141. Katselou, M.; Papoutsis, I.; Nikolaou, P.; Misailidi, N.; Spiliopoulou, C.; Athanaselis, S. Diphenidine: A dissociative NPS makes an entrance on the drug scene. *Forensic. Toxicol.* **2018**, *36*, 233–242. [CrossRef]