Liquid chromatography-mass spectrometry based metabolomics investigation of five different tissues of *Mytilus galloprovincialis*

Supplementary material


Comparative Biochemistry and Physiology

Part D: Genomics and Proteomics


Lea James, Elena Gomez, Gaelle Ramirez, Thibaut Dumas and Frédérique Courant * HydroSciences Montpellier, University of Montpellier, IRD, CNRS, Montpellier, France

HydroSciences Montpellier, 15 avenue Charles Flahault, 34090 Montpellier

* Corresponding author: email: frederique.courant@umontpellier.fr; tel: +0033411759414



Supplementary Fig. 1 PCA score plot of ESI - metabolic fingerprints from digestive gland (yellow circle), foot (red square), gill (green triangle), gonad (inverted blue triangle) and soft (orange diamond) mussel tissues showing a clear separation between these organs. QC_{organ} are shown in black (circles). (Note that interpretation of the references to colour in this figure legend are provided in the web version of this article)



Supplementary Fig. 2 Molecular network highlighted by GNPS analysis based on MS/MS fragmentation spectra obtained from different Mediterranean mussel organs. The color code is as follows: digestive gland (yellow), foot (red), gill (green), gonad (blue) and soft (orange) tissues. The relative intensity of the metabolite in each organ is represented as a pie chart on each node