



HAL
open science

A Lyapunov approach to stability of positive semigroups: An overview with illustrations

Marc Arnaudon, Pierre Del Moral, El Maati Ouhabaz

► To cite this version:

Marc Arnaudon, Pierre Del Moral, El Maati Ouhabaz. A Lyapunov approach to stability of positive semigroups: An overview with illustrations. 2023. hal-03931007

HAL Id: hal-03931007

<https://hal.science/hal-03931007>

Preprint submitted on 9 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Lyapunov approach to stability of positive semigroups: An overview with illustrations

Marc Arnaudon¹, Pierre Del Moral² & El Maati Ouhabaz¹

¹University of Bordeaux, Institut de Mathématiques de Bordeaux, France. E-Mail: marc.arnaudon@math.u-bordeaux.fr, Elmaati.Ouhabaz@math.u-bordeaux.fr

²Centre de Recherche Inria Bordeaux Sud-Ouest, Talence, 33405, FR. E-Mail: pierre.del-moral@inria.fr

January 9, 2023

Abstract

The stability analysis of possibly time varying positive semigroups on non necessarily compact state spaces, including Neumann and Dirichlet boundary conditions is a notoriously difficult subject. These crucial questions arise in a variety of areas of applied mathematics, including nonlinear filtering, rare event analysis, branching processes, physics and molecular chemistry. This article presents an overview of some recent Lyapunov-based approaches, focusing principally on practical and powerful tools for designing Lyapunov functions. These techniques include semigroup comparisons as well as conjugacy principles on non necessarily bounded manifolds with locally Lipschitz boundaries. All the Lyapunov methodologies discussed in the article are illustrated in a variety of situations, ranging from conventional Markov semigroups on general state spaces to more sophisticated conditional stochastic processes possibly restricted to some non necessarily bounded domains, including locally Lipschitz and smooth hypersurface boundaries, Langevin diffusions as well as coupled harmonic oscillators.

Keywords: Integral operators, semigroups, Markov and Sub-Markov semigroups, harmonic oscillators, Langevin diffusions, Lyapunov function, hypersurfaces, shape matrices, boundary problems.

Mathematics Subject Classification: Primary 47D08, 60J25, 47D06, 47D07, 47H07; secondary 47B65, 37A30, 37M25.

Contents

1	Introduction	3
1.1	Description of the models	3
1.2	Some basic notation	5
1.3	Regularity conditions	7
2	A brief review on semigroups	8
2.1	A V -norm contraction theorem	8
2.2	Normalized semigroups	10
2.3	Time homogenous models	13
2.4	Markov diffusion semigroups	15
2.5	Sub-Markov semigroups	20
3	Lyapunov design principles	25
3.1	Foster-Lyapunov conditions	25
3.2	Semigroup domination	27
3.3	Some conjugacy principles	29
4	Boundary problems	30
4.1	Bounded domains	31
4.2	Unbounded domains	32
4.3	Smooth boundaries	36
5	Riccati type processes	39
5.1	Positive diffusions	39
5.2	Matrix valued diffusions	40
5.3	Logistic birth and death process	41
5.4	Multivariate birth and death processes	42
6	Some conditional diffusions	43
6.1	Coupled harmonic oscillators	43
6.2	Half-harmonic linear diffusions	45
6.3	Linear diffusions in some domains	47
6.4	Langevin diffusions in some domains	48
6.5	Coupled oscillators in some domains	50
7	Some hypersurface boundaries	51
7.1	Defining functions and charts	51
7.2	The shape matrix	52
7.3	Surface and volume forms	55
7.4	Boundary decompositions	57

1 Introduction

1.1 Description of the models

Let $\mathcal{B}(E)$ be the algebra of locally bounded measurable functions on a locally compact Polish space E . We denote by $\mathcal{B}_b(E) \subset \mathcal{B}(E)$ the sub-algebra of bounded measurable functions endowed with the supremum norm $\|\cdot\|$. Let $Q_{s,t}$, be a semigroup of positive integral operators on $\mathcal{B}_b(E)$ indexed by a continuous time indices $s, t \in \mathcal{T} = \mathbb{R}_+ := [0, \infty[$ or by a discrete time index set $\mathcal{T} = \mathbb{N}$, with $s \leq t$.

For a given uniformly positive function $V \in \mathcal{B}(E)$, we let $\mathcal{B}_V(E) \subset \mathcal{B}(E)$ be the sub-space of functions $f \in \mathcal{B}(E)$ equipped with the norm $\|f\|_V := \|f/V\|$.

We also let $\mathcal{B}_\infty(E) \subset \mathcal{B}(E)$ be the subalgebra of locally bounded and uniformly positive functions V that grow at infinity; that is, $\sup_K V < \infty$ for any compact set $K \subset E$, and for any $r \geq V_\star := \inf_E V > 0$ the r -sub-level set $\mathcal{V}(r) := \{V \leq r\} \subset E$ is a non empty compact subset. We denote by $\mathcal{B}_0(E) := \{1/V : V \in \mathcal{B}_\infty\} \subset \mathcal{B}_b(E)$ the sub-algebra of positive functions, locally lower bounded and that vanish at infinity. For a given $V \in \mathcal{B}_\infty(E)$, consider the subspace

$$\mathcal{B}_{0,V}(E) := \{f \in \mathcal{B}(E) : |f|/V \in \mathcal{B}_0(E)\}.$$

We say that $Q_{s,t}$ is a V -positive semigroup on $\mathcal{B}_V(E)$ for some Lyapunov function $V \in \mathcal{B}_\infty(E)$ as soon as there exists some $\tau > 0$ and some function $\Theta_\tau \in \mathcal{B}_0(E)$ such that for any $0 < f \in \mathcal{B}_V(E)$ and $s < t$ we have $0 < Q_{s,t}(f) \in \mathcal{B}_{0,V}(E)$ as well as

$$Q_{s,s+\tau}(V)/V \leq \Theta_\tau \quad \text{and} \quad \sup_{|t-s| \leq \tau} (\|Q_{s,t}\| \vee \|Q_{s,t}\|_V) < \infty. \quad (1)$$

The growth conditions stated above are discussed in some details in Section 1.3. As shown in Section 3.1, the l.h.s. criterion in (1) can be seen as a uniform Foster-Lyapunov condition (a.k.a. drift condition).

Foster-Lyapunov criterion dates back to the 1950s with the seminal articles [32, 38]. These criteria are nowadays an essential tool to analyze the stability properties of Markov semigroups on general state spaces [7, 26, 37, 39, 50, 51]. Their use in the context of positive semigroup arising in discrete time nonlinear filtering goes back to the pioneering articles [27, 64], based on coupling techniques developed in [41, 42]. The extension of Foster-Lyapunov criterion to discrete or continuous time varying positive semigroups and their normalized versions on general state spaces were further developed in [21], extending Dobrushin's ergodic coefficient techniques introduced in [11, 12] and further developed in [15, 16, 17, 14, 22] to unbounded state space models.

Recall that the Dobrushin's ergodic coefficient of a Markov semigroup is the operator norm of the Markov integral operator acting on probability measures equipped with the total variation norm (see for instance [16] and references therein). In the same vein, the V -Dobrushin's ergodic coefficient of a Markov transition is defined as the operator norm of the Markov integral operator acting on probability measures

equipped with the V -norm. In this operator theoretical framework, the contraction w.r.t. V -norms is deduced by coupling the Foster-Lyapunov criterion with a local contraction on a sufficiently large compact sub-level set of the Lyapunov function. A brief overview on this subject is provided in Section 2.

The local contraction on the compact sub-level sets of the Lyapunov function is generally an easily verifiable condition. This property is often deduced from a Doeblin type local minorization property of integral operators on the compact sub-level sets of the Lyapunov function. For instance, this local minorization condition is satisfied as soon as the semigroup is lower bounded by an absolutely continuous integral operator (a.k.a. transition kernel operator). This class of models includes hypo-elliptic diffusion semigroups as well as some regular jump processes on non necessarily bounded domains.

Even for diffusion semigroups with smooth densities on bounded manifolds with entrance boundaries (i.e. boundary states that cannot be reached from the inside), the existence of a sufficiently strong Lyapunov function is essential to ensure the stability of the semigroup. In this context, the transition densities are null on entrance boundary states so that the local minorization condition alone applied to some exhausting sequence of compact subsets is not sufficient to ensure the stability of the process. The exhausting sequence of compact subsets needs to be equivalent to the sub-level sets of some sufficiently strong Lyapunov function near entrance boundaries. For a more thorough discussion on this subject we refer to Section 2 and the article [21], see also the series of Riccati-type diffusions discussed in Section 5.

The general problem of constructing Lyapunov functions for positive semigroups, including for Markov semigroups often requires to have some good intuition about a candidate for a Lyapunov function on some particular class of model. As for deterministic dynamical systems, the design of Lyapunov functions for sub-Markov semigroups associated with a non-absorbed stochastic process requires to use some physical insight on the stability and the behavior of the free evolution stochastic process near possible absorbing boundaries.

Constructing Lyapunov functions for general classes of positive semigroups is well known as a very hard problem in system theory as well as in applied probability literature. The main subject of this article is to find *practical ways* to design these Lyapunov functions for various classes of positive semigroups that have been discussed in the literature, including conditional diffusions on manifolds with Neumann and Dirichlet boundaries. We did our best to cover the subject as broadly as possible, we also refer to the article [21] for additional historical and reference pointers. Due to the vast literature on this subject we apologize for possible omissions of some important contributions due to the lack of knowledge.

The remainder of this article is structured as follows:

In Section 2, we begin with a brief review of V -norm contraction theorems and semigroup stability properties stemming from an assumed Lyapunov structure. Section 2.1 is dedicated to time varying Markov semigroups. The extension of these results to time varying positive semigroups are discussed in Section 2.2. In Section 2.3,

we present some consequences of these results in the context of time homogenous models, including existence of ground states and quasi-invariant measures. Section 2.4 and Section 2.5 present different tools to design Lyapunov functions for continuous time Markov semigroups and sub-Markov semigroups. We also illustrate these results through different examples of semigroups arising in physics and applied probability, including overdamped Langevin diffusions, Langevin and hypo-elliptic diffusions, as well as typical examples of solvable one-dimensional sub-Markov semigroups such as the harmonic oscillator, the half-harmonic oscillator and the Dirichlet heat kernel. General comparison and conjugacy principles to construct Lyapunov functions for positive semigroups are provided in Section 3. Boundary problems are discussed in some details in Section 4. We then turn in Section 5 to the design of Lyapunov functions for Riccati type processes, including positive definite matrix valued diffusions, logistic and multivariate birth and death processes arising respectively in Ensemble Kalman-Bucy filter theory and population dynamic analysis.

In Section 6 we illustrate the power of the Lyapunov approach in the context of multivariate conditional diffusions. Section 7 is dedicated to illustrations with explicit computations of geometrical objects for the Lyapunov functions discussed in Section 4.3 in the context of hypersurface Dirichlet boundaries.

1.2 Some basic notation

We denote by $\mathcal{C}(E) \subset \mathcal{B}(E)$ the sub-algebra of continuous functions and by $\mathcal{C}_b(E) \subset \mathcal{C}(E)$ the sub-algebra of bounded continuous functions.

We also set $\mathcal{C}_V(E) := \mathcal{B}_V(E) \cap \mathcal{C}(E)$, $\mathcal{C}_0(E) := \mathcal{B}_0(E) \cap \mathcal{C}(E)$ and $\mathcal{C}_\infty(E) := \mathcal{B}_\infty(E) \cap \mathcal{C}(E)$ and $\mathcal{C}_{0,V}(E) := \mathcal{B}_{0,V}(E) \cap \mathcal{C}(E)$. Note that none of the sub-algebras $\mathcal{B}_0(E)$ and $\mathcal{B}_\infty(E)$ have an unit unless E is compact, the null function $0 \notin \mathcal{B}_0(E)$ but the unit function $1 \in \mathcal{C}_{0,V}(E)$ as soon as $V \in \mathcal{B}_\infty(E)$.

Let $\mathcal{M}_b(E)$ be the set of bounded signed measures μ on E equipped with the total variation norm $\|\mu\|_{tv} := |\mu|(E)/2$, where $|\mu| := \mu_+ + \mu_-$. It stands for the total variation measure associated with a Hahn-Jordan decomposition $\mu = \mu_+ - \mu_-$ of the measure. Also let $\mathcal{P}(E) \subset \mathcal{M}_b(E)$ be the subset of probability measures on E .

With a slight abuse of notation, we denote by 0 and 1 the null and unit scalars as well as the null and unit function on E .

The action of $Q_{s,t}$ on $\mathcal{B}_b(E)$ is given for any $f \in \mathcal{B}_b(E)$ by the formulae

$$Q_{s,t}(f)(x) := \int Q_{s,t}(x, dy) f(y). \quad (2)$$

The left action of $Q_{s,t}$ on $\mathcal{M}_b(E)$ is given for any $\eta \in \mathcal{M}_b(E)$ by the formulae

$$(\eta Q_{s,t})(dy) := \int \eta(dx) Q_{s,t}(x, dy). \quad (3)$$

In this notation, the semigroup property takes the following form

$$Q_{s,u}Q_{u,t} = Q_{s,t} \quad \text{with} \quad Q_{s,s} = I, \quad \text{the identity operator.} \quad (4)$$

In the above display, $Q_{s,u}Q_{u,t}$ is a shorthand notation for the composition $Q_{s,u} \circ Q_{u,t}$ of the left or right-action operators. Unless otherwise stated, all the semigroups discussed in this article are indexed by conformal indices $s \leq t$ in the set \mathcal{T} . To avoid repetition, we often write $Q_{s,t}$ without specifying the order $s \leq t$ of the indices $s, t \in \mathcal{T}$.

We denote by $\mathcal{M}_V(E)$ be the space of measures $\mu \in \mathcal{M}_b(E)$ equipped with the operator V -norm $\| \mu \|_V := |\mu|(V)$, and by $\mathcal{P}_V(E) \subset \mathcal{M}_V(E)$ be the convex set of probability measures. We associate with a function $h \in \mathcal{B}_{0,V}(E)$ the Boltzmann-Gibbs transformation

$$\Psi_h : \mu \in \mathcal{P}_V(E) \mapsto \Psi_h(\mu) \in \mathcal{P}_{Vh}(E) \quad (5)$$

with the probability measure

$$\Psi_h(\mu)(dx) := \frac{h(x)}{\mu(h)} \mu(dx) \quad \text{and} \quad V^h := V/h \in \mathcal{B}_\infty(E).$$

We also denote by $\|Q\|_V$ the operator norm of a bounded linear operator $Q : f \in \mathcal{B}_V(E) \mapsto Q(f) \in \mathcal{B}_V(E)$; that is

$$\|Q\|_V := \sup\{\|Q(f)\|_V : f \in \mathcal{B}_V(E) \text{ such that } \|f\|_V \leq 1\}. \quad (6)$$

In terms of the V -conjugate semigroup

$$f \in \mathcal{B}_b(E) \mapsto Q^V(f) := Q(Vf)/V \in \mathcal{B}_b(E)$$

we have

$$\|Q\|_V = \|Q^V(1)\| = \|Q^V\| := \sup\{\|Q^V(f)\| : f \in \mathcal{B}_b(E) \text{ such that } \|f\| \leq 1\}.$$

For a given measurable function f and a given measurable subset, we use the shorthand notation

$$-\infty \leq \inf_A f := \inf_{x \in A} f(x) \leq \sup_A f := \sup_{x \in A} f(x) \leq +\infty.$$

For a given $s \in \mathcal{T}$ and $\tau \in \mathcal{T}$ with $\tau > 0$, we consider the time mesh

$$[s, \infty[_\tau := \{s + n\tau \in [s, \infty[: n \in \mathbb{N}\}.$$

Throughout, unless otherwise is stated we write c for some positive constants whose values may vary from line to line, and we write c_α , as well as $c(\beta)$ and $c_\alpha(\beta)$ when their values may depend on some parameters α, β defined on some parameter sets. We also set $a \wedge b = \min(a, b)$, $a \vee b = \max(a, b)$, and $a_+ = a \vee 0$ for $a, b \in \mathbb{R}$.

1.3 Regularity conditions

The irreducibility condition $f > 0 \implies Q_{s,t}(f) > 0$ is satisfied if and only if we have $Q_{s,t}(1) > 0$. We check this claim by contradiction. Assume that $Q_{s,t}(1) > 0$ and consider a function $f > 0$ and some $x \in E$ such that $Q_{s,t}(f)(x) = 0$. In this case, for any $\epsilon > 0$ we would have

$$0 = \epsilon Q_{s,t}(1_{f \geq \epsilon})(x) \leq Q_{s,t}(f)(x)$$

by Fatou's lemma we would find the contraction

$$\liminf_{\epsilon \rightarrow 0} Q_{s,t}(1_{f \geq \epsilon})(x) = 0 \geq Q_{s,t}(1)(x) \implies Q_{s,t}(1)(x) = 0.$$

Without further mention, all semigroups $Q_{s,t}$ considered in this article are assumed to be semigroups of positive integral operators $Q_{s,t}$ on $\mathcal{B}_b(E)$ satisfying the irreducibility condition $Q_{s,t}(1) > 0$ for any $s \leq t$. Notice that the condition

$$0 < f \in \mathcal{B}_V(E) \implies \forall s < t \quad 0 < Q_{s,t}(f) \in \mathcal{B}_{0,V}(E)$$

is met as soon as $Q_{s,t}$ is a strong V -Feller semigroup, in the sense that for any $s < t$ we have $Q_{s,t}(\mathcal{B}_V(E)) \subset \mathcal{C}_V(E)$ and when we have $Q_{s,t}(V)/V \in \mathcal{B}_0(E)$. To check this claim, observe that for any positive function $f \in \mathcal{B}_V(E)$ and $s < t$ the function $Q_{s,t}(f)$ is positive and continuous; and thus locally lower bounded. In this situation, whenever $\|f\|_V \leq 1$, for any $s < t$ we have the comparison property

$$Q_{s,t}(f)/V \leq Q_{s,t}(V)/V \in \mathcal{B}_0(E) \implies Q_{s,t}(f)/V \in \mathcal{B}_0(E) \iff Q_{s,t}(f) \in \mathcal{C}_{0,V}(E).$$

In summary, a strong V -Feller semigroup $Q_{s,t}$ is V -positive on $\mathcal{B}_V(E)$ as soon as there exists some $\tau > 0$ and some function $\Theta_\tau \in \mathcal{B}_0(E)$ such that the l.h.s. condition in (1) is met and for any $s < t$ we have

$$Q_{s,t}(V)/V \in \mathcal{B}_0(E) \quad \text{and} \quad Q_{s,s+\tau}(V)/V \leq \Theta_\tau \in \mathcal{B}_0(E).$$

When $V \in \mathcal{C}_\infty(E)$, we say that $Q_{s,t}$ is a V -positive semigroup on $\mathcal{C}_V(E)$ as soon as $Q_{s,t}(\mathcal{C}_V(E)) \subset \mathcal{C}_{0,V}(E)$ for any $s < t$ and condition (1) is met.

A V -Feller semigroup $Q_{s,t}$ for some $V \in \mathcal{C}_\infty(E)$, in the sense that for any $s < t$ we have $Q_{s,t}(\mathcal{C}_V(E)) \subset \mathcal{C}_V(E)$, is also said to be V -positive on $\mathcal{C}_V(E)$ as soon as there exists some $\tau > 0$ and some function $\Theta_\tau \in \mathcal{B}_0(E)$ such that the l.h.s. condition in (1) is met and for any $s < t$ we have

$$Q_{s,t}(V)/V \in \mathcal{C}_0(E) \quad \text{and} \quad Q_{s,s+\tau}(V)/V \leq \Theta_\tau \in \mathcal{B}_0(E).$$

Last but not least, observe that positive semigroups $Q_{s,t}$ with continuous time indices $s \leq t \in \mathbb{R}_+$ can be turned into discrete time models by setting $Q_{p,n} = Q_{p\tau,n\tau}$ for any $p \leq n \in \mathbb{N}$ and some parameter $\tau > 0$. Up to a time rescaling, the parameter $\tau > 0$ arising in the definition of a discrete time V -positive semigroups $Q_{p,n}$ can be chosen as the unit time parameter. In this context, the r.h.s. condition in (1) is automatically satisfied.

2 A brief review on semigroups

2.1 A V -norm contraction theorem

The aim of this section is to present some stability theorems for uniform V -positive semigroups. We first examine the situation where $Q_{s,t} = P_{s,t}$ is a semigroup of Markov integral operators $P_{s,t}$ on $\mathcal{B}_b(E)$. Note that the Lyapunov condition stated in the l.h.s. of (1) ensures the following geometric drift condition

$$P_{s,s+\tau}(V) \leq \epsilon_\tau V + c_\tau \quad (7)$$

some parameter $\epsilon_\tau \in [0, 1[$ and some finite constant $c_\tau < \infty$. The geometric drift condition (7) ensures that the sequence $\|P_{s,s+n\tau}\|_V$ indexed by $s \geq 0$ and $n \geq 1$ is uniformly bounded. In this context, the r.h.s. condition in (1) applied to $Q_{s,t} = P_{s,t}$ ensures that the operator norms of $P_{s,t}$ are uniformly bounded w.r.t. any time horizon. More precisely, whenever (7) is met we have the equivalence

$$\sup_{s \geq 0} \sup_{t \geq s} \|P_{s,t}\|_V < \infty \iff \sup_{|t-s| \leq \tau} \|P_{s,t}\|_V < \infty. \quad (8)$$

Note that (8) is automatically satisfied whenever (7) is met for any $\tau > 0$ with $\sup_{\tau \in [0,1]} c_\tau < \infty$. For instance, consider the Markov transition semigroup $P_{s,t}$ of a continuous time stochastic flow $X_{s,t}(x)$ on some locally compact normed vector space $(E, \|\cdot\|)$ with generator L_t defined on some common domain $\mathcal{D}(L) \subset \mathcal{B}(E)$. In this context, for any non negative function $V \in \mathcal{D}(L)$ and any parameters $a > 0$, $c < \infty$ and $\tau > 0$ we have

$$L_\tau(V) \leq -aV + c \implies (7) \text{ and } (8) \quad \text{with} \quad \epsilon_\tau = (1 + a\tau)^{-1} < 1 \quad \text{and} \quad c_\tau = c\tau. \quad (9)$$

The above estimate is rather well known, a detailed proof is provided in the appendix on page 63. Further examples of Markov diffusion semigroups on \mathbb{R}^n satisfying (7) are discussed in Section 2.4. We further assume there exists some $r_0 \geq 1$ and some function $\alpha_\tau : r \in [r_0, \infty[\mapsto \alpha_\tau(r) \in]0, 1]$, such that for any $r \geq r_0$ we have

$$\sup_{(x,y) \in \mathcal{V}(r)^2} \|\delta_x P_{s,s+\tau} - \delta_y P_{s,s+\tau}\|_{tv} \leq 1 - \alpha_\tau(r) \quad \text{with} \quad \mathcal{V}(r) := \{V \leq r\}. \quad (10)$$

Consider the V -norm operator $\beta_V(P_{s,t})$ (a.k.a. the V -Dobrushin coefficient) of $P_{s,t}$ defined by

$$\beta_V(P_{s,t}) := \sup_{\mu, \eta \in \mathcal{P}_V(E)} \|(\mu - \eta)P_{s,t}\|_V / \|\mu - \eta\|_V. \quad (11)$$

In this notation, conditions (7), (8) and (10) ensure the existence of some parameter $\bar{\tau} > 0$ such that

$$\sup_{|t-s| \leq \bar{\tau}} \beta_V(P_{s,t}) < \infty \quad \text{and} \quad \sup_{s \geq 0} \beta_V(P_{s,s+\bar{\tau}}) < 1. \quad (12)$$

The proof of the above assertion can be found in [21] (see also [22] in the context of time homogeneous models). The next exponential contraction theorem is a direct consequence of the operator norm estimates (12) and it is valid on abstract measurable spaces as well as for any function $V \geq 1$.

Theorem 2.1. *Let $P_{s,t}$ be a semigroup of Markov integral operators $P_{s,t}$ on some measurable state space E satisfying condition (12) for some function $V \geq 1$ and some parameter $\bar{\tau} > 0$. In this situation, there exists a parameter $b > 0$ and some finite constant $c < \infty$ such that for any $s \leq t$ and $\mu, \eta \in \mathcal{P}_V(E)$ we have the exponential estimate*

$$\|(\mu - \eta)P_{s,t}\|_V \leq c e^{-b(t-s)} \|\mu - \eta\|_V. \quad (13)$$

In particular, the above exponential Lipschitz estimates are met as soon as conditions (7), (8) and (10) are satisfied. The estimates (13) also hold for any $s \geq 0$ and $t \in [s, \infty[_{\tau}$ as soon as (7) and (10) are satisfied for some $\tau > 0$.

The proof of Theorem 2.1 is based on discrete time type V -norm operator contraction techniques combining the geometric drift condition (7) with the total variation estimates (10). The r.h.s. condition in (8) is a technical condition only made for continuous time semigroups to ensure that (13) also holds for continuous time indices.

For time homogeneous semigroups $P_t := P_{s,s+t}$ the contraction estimate (13) ensures the existence of a single invariant probability measure $\mu_\infty = \mu_\infty P_t \in \mathcal{P}_V(E)$. In this context, similar approaches are presented in the article [37], simplifying the Foster-Lyapunov methodologies and the small-sets return times estimation techniques developed in [50]. Theorem 2.1 can be seen as an extension of Harris' theorem to time varying Markov semigroup. The operator-theoretic framework discussed above provides a very direct proof based on the V -Dobrushin coefficient (11). For a more thorough discussion on this subject we refer to [21, 22].

Note that the strength of conditions (7) and (10) depends on the strength of the function V : when the function V is bounded, the geometric drift condition (7) and the uniform norm condition (8) are trivially met but in this case condition (10) is a uniform contraction condition on the state E . In the reverse angle, when $V \in \mathcal{B}_\infty(E)$ is a function with compact sub-level sets, the geometric drift condition (7) combined with (8) ensures that $\mu P_{s,t}$ is a tight collection of probability measures indexed by $s \leq t$. In this context, the local contraction condition (10) is met if and only if for any $s \geq 0$ and any $(x, y) \in \mathcal{V}(r)^2$ there exists some probability measure μ on E (that may depends on the parameters (τ, r, s, x, y)) such that

$$\forall z \in \{x, y\} \quad \delta_z P_{s,s+\tau}(dy) \geq \alpha_\tau(r) \mu(dy).$$

For instance, the above condition is met as soon as

$$P_{s,s+\tau}(x, dy) \geq p_{s,s+\tau}(x, y) \nu_\tau(dy) \quad (14)$$

for some Radon positive measure ν_τ on E and some density function $p_{s,s+\tau}$, satisfying for any $r \geq r_0$ the local minorization condition

$$0 < \inf_{s \in \mathcal{T}} \inf_{\mathcal{V}(r)^2} p_{s,s+\tau} \quad \text{and} \quad 0 < \nu_\tau(\mathcal{V}(r)) < \infty. \quad (15)$$

For locally compact Polish spaces condition $0 < \nu_\tau(\mathcal{V}(r)) < \infty$ is met as soon as V has compact sub-levels sets $\mathcal{V}(r)$ with non empty interior and ν_τ is a Radon measure of full support; that is ν_τ is finite on compact sets and strictly positive on non-empty open sets. For time homogeneous models, also note that the l.h.s. minorization condition (15) is satisfied as soon as $(x, y) \in (E^\circ)^2 \mapsto p_\tau(x, y)$ is a continuous positive function on the interior E° of the set E .

Several illustrations of Theorem 2.1 are discussed in Section 2.4 in the context of diffusion processes on Euclidean spaces as well as in Section 5 in the context of Riccati-type diffusion on positive definite matrix spaces and multivariate birth and death jump type processes on countable state spaces. The stability of Markov semigroups on manifolds with entrance boundaries can also be analyzed using the Lyapunov techniques developed in Section 4. For instance, as shown in Section 4.1, any absolutely continuous Markov semigroup $P_{s,t}$ on a bounded connected subset $E \subset \mathbb{R}^n$ with locally Lipschitz boundary ∂E satisfies the conditions of Theorem 2.1 with the (non unique) Lyapunov function $V(x) = 1/\sqrt{d(x, \partial E)}$ and the distance to the boundary defined for any $x \in E$ by

$$d(x, \partial E) := \inf \{ \|x - y\| : y \in \partial E \}.$$

We illustrate the above discussion with some elementary one dimensional examples.

Example 2.2. *Consider a one dimensional Brownian on the compact interval $E = [0, 1]$ with reflected boundaries. In this situation, $P_t := P_{0,t}$ coincides with the Neumann heat semigroup on $[0, 1]$. In this context, recalling that the Neumann heat kernel is smooth and strictly positive on the compact interval $[0, 1]$, the conditions of Theorem 2.1 are satisfied with the unit Lyapunov function $V(x) = 1$, as well as for any of the Lyapunov functions $V(x) = 1/\sqrt{x}$, $V(x) = 1/\sqrt{1-x}$ or $V(x) = 1/\sqrt{x} + 1/\sqrt{1-x}$.*

The same reasoning applies to the one dimensional positive Riccati-type diffusions with an entrance boundary at the origin discussed in Section 5. Reflecting this class of positive diffusions at $x = 1$, the conditions of Theorem 2.1 are satisfied on $E =]0, 1]$ with the Lyapunov functions $V(x) = 1/\sqrt{x}$ as well as for $V(x) = 1/\sqrt{x} + 1/\sqrt{1-x}$.

2.2 Normalized semigroups

For non necessarily Markov V -positive semigroups $Q_{s,t}$ one natural idea is to normalise the semigroups. For any probability measure $\eta \in \mathcal{P}_V(E)$ we let $\Phi_{s,t}(\eta) \in \mathcal{P}_V(E)$ be the normalised distribution defined for any $f \in \mathcal{B}_V(E)$ by the formula

$$\Phi_{s,t}(\eta)(f) := \frac{\eta Q_{s,t}(f)}{\eta Q_{s,t}(1)} \quad \text{and we set} \quad \bar{Q}_{s,t}(f)(x) := \frac{Q_{s,t}(f)(x)}{Q_{s,t}(1)(x)} = \Phi_{s,t}(\delta_x)(f). \quad (16)$$

The mapping $\Phi_{s,t}$ is a well defined semigroup on $\mathcal{P}_V(E)$. The denormalisation formula connecting these semigroups is given for any $t \in [s, +\infty[_\tau$ by

$$\mu Q_{s,t}(f) = \Phi_{s,t}(\mu)(f) \prod_{u \in [s,t[_\tau} \Phi_{s,u}(\mu)(Q_{u,u+\tau}(1)). \quad (17)$$

with

$$[s, t[_\tau := \{s + n\tau \in [s, t[: n \in \mathbb{N}\}.$$

To check this claim, observe that for any $t := s + n\tau$ we have

$$\Phi_{s,s+p\tau}(\mu)(Q_{s+p\tau,s+(p+1)\tau}(1)) = \mu Q_{s,s+(p+1)\tau}(1) / \mu Q_{s,s+p\tau}(1)$$

$$\text{and therefore } \prod_{0 \leq p < n} \Phi_{s,s+p\tau}(\mu)(Q_{s+p\tau,s+(p+1)\tau}(1)) = \mu Q_{s,s+n\tau}(1)$$

The above formula coincides with the product formula relating the unnormalised operators $Q_{s,t}$ with the normalised semigroup $\Phi_{s,t}$ discussed in [14, Section 1.3.2], see also [17, Proposition 2.3.1] and [19, Section 12.2.1].

We strengthen (14) and assume that for any $s \geq 0$ and $\tau > 0$, the integral operator $Q_{t,t+\tau}$ has a density $q_{s,s+\tau}$ with respect to some Radon positive measure ν_τ on E ; that is we have that

$$Q_{s,s+\tau}(x, dy) = q_{s,s+\tau}(x, y) \nu_\tau(dy). \quad (18)$$

We also assume there exists some $r_0 > 1$ such that for any $r \geq r_0$ we have

$$0 < \iota_r(\tau) := \inf_{s \in \mathcal{T}} \inf_{\mathcal{V}(r)^2} q_{s,s+\tau} \leq \sup_{s \in \mathcal{T}} \sup_{\mathcal{V}(r)^2} q_{s,s+\tau} < \infty \quad \text{and} \quad \nu_\tau(\mathcal{V}(r)) > 0. \quad (19)$$

In this situation, for any $r \geq r_0$ and $\bar{r} \geq r$ we have the uniform estimate

$$\inf_{\mathcal{V}(r)} Q_{s,s+\tau}(1) \geq J_{r,\bar{r}}(\tau) := \inf_{\mathcal{V}(r)} Q_{s,s+\tau}(1_{\mathcal{V}(\bar{r})}) \geq \iota_r(\tau) \nu_\tau(\mathcal{V}(r)) > 0.$$

We associate with a given $\mu \in \mathcal{P}_V(E)$ and some function $H \in \mathcal{B}_{0,V}(E)$ the finite rank (and hence compact) operator

$$f \in \mathcal{B}_V(E) \mapsto T_{s,t}^{\mu,H}(f) := \frac{Q_{s,t}(H)}{\mu_s(Q_{s,t}(1))} \mu_t(f) \in \mathcal{C}_V(E)$$

with the flow of measures $\mu_t = \Phi_{s,t}(\mu_s)$ starting at $\mu_0 = \mu$. With this notation at hand, one has the following theorem.

Theorem 2.3 ([21]). *Consider a V -positive semigroups $Q_{s,t}$ with a density (18) satisfying (19) for some parameter $\tau > 0$ and some $r_0 > 1$. In this situation, there exists a parameter $b > 0$ such that for any $\mu, \eta \in \mathcal{P}_V(E)$ and any $s \geq 0$ and $t \in [s, \infty[_\tau$ we have the local Lipschitz estimate*

$$\|\Phi_{s,t}(\mu) - \Phi_{s,t}(\eta)\|_V \leq c(\mu, \eta) e^{-b(t-s)} \|\mu - \eta\|_V. \quad (20)$$

For any $(\mu, H) \in (\mathcal{P}_V(E) \times \mathcal{B}_{0,V}(E))$ there exists some finite constant $c_H(\mu) < \infty$ such that for any $s \geq 0$ and $t \in [s, \infty[_\tau$ we have

$$\left\| \frac{Q_{s,t}}{\mu_s Q_{s,t}(1)} - T_{s,t}^{\mu,H} \right\|_V \leq c_H(\mu) e^{-b(t-s)}. \quad (21)$$

For continuous time semigroups, the above estimates also hold for any continuous time indices $s \leq t$ as soon as for any $r \geq r_0$ there exists some $\bar{r} \geq r$ such that $\inf_{\delta \in [0, \tau]} J_{r, \bar{r}}(\delta) > 0$.

The proof of Theorem 2.3 is based on discrete time type V -norm operator contraction techniques combining the geometric drift condition stated in the l.h.s. of (1) with the local minorization condition stated in (19). The condition $\inf_{\delta \in [0, \tau]} J_{r, \bar{r}}(\delta) > 0$ is a technical condition only made for continuous time semigroups to ensure that (20) and (21) also hold for continuous time indices.

Theses regularity conditions are rather flexible as we will now explain.

Absolutely continuous integral operators arise in a natural way in discrete time settings [17, 14, 27, 64] and in the analysis of continuous time elliptic diffusion absorption models [2, 29, 30, 61]. In connection to this, two-sided estimates for stable-like processes are provided in [6, 43, 59, 63]. Two sided Gaussian estimates can also be obtained for some classes of degenerate diffusion processes of rank 2, that is when the Poisson brackets of the first order span the whole space [44]. This class of diffusions includes frictionless Hamiltonian kinetic models.

Diffusion density estimates can be extended to sub-Markovian semigroups using the multiplicative functional methodology developed in [15]. Whenever the trajectories of these diffusion flows, say $t \mapsto X_t(x)$, where $x \in E$ is the initial position, are absorbed on the smooth boundary ∂E of a open connected domain E , for any $\tau > 0$ the densities $q_\tau(x, y)$ of the sub-Markovian semigroup Q_τ (with respect to the trace of the Lebesgue measure on E) associated with the non absorption event are null at the boundary. Nevertheless, whenever these densities are positive and continuous on the open set E^2 for some $\tau > 0$, they are uniformly positive and bounded on any compact subset of E ; thus condition (19) is satisfied.

In this context, whenever $T(x)$ stands for first exit time from E and $T_r(x)$ the first exit time from the compact level set $\mathcal{V}(r) \subset E$ starting from $x \in \mathcal{V}(r)$, for any $\delta \in [0, \tau]$ and $r_+ > r$ we have the estimate

$$Q_\delta(1_{\mathcal{V}(r_+)}) (x) := \mathbb{E} (1_{\mathcal{V}(r_+)}(X_\delta(x)) 1_{T(x) > \delta}) \geq \mathbb{P} (T_{r_+}(x) > \delta) \geq \mathbb{P} (T_{r_+}(x) > \tau).$$

In this context, we have

$$\inf_{x \in \mathcal{V}(r)} \mathbb{P} (T_{r_+}(x) > \tau) > 0 \implies \inf_{\delta \in [0, \tau]} \inf_{\mathcal{V}(r)} Q_\delta(1) \geq \inf_{\delta \in [0, \tau]} J_{r, r_+}(\delta) > 0. \quad (22)$$

Whenever the interior $E_+ := \mathcal{V}(r_+)^\circ$ is a connected domain, the l.h.s. estimate in (22) is met as soon as the sub-Markovian semigroup Q_τ^+ associated with the non absorption

event at the boundary ∂E_+ has a continuous density $(x, y) \in E_+^2 \mapsto q_\tau^+(x, y)$. To check this claim, observe that for any $x \in \mathcal{V}(r)$ we have

$$\begin{aligned} \mathbb{P}(T_{r_+}(x) > \tau) &= Q_\tau^+(1)(x) \geq Q_\tau^+(1_{\mathcal{V}(r)})(x) = \int q_\tau^+(x, y) 1_{\mathcal{V}(r)}(y) \nu_\tau(dy) \\ &\geq \nu_\tau(\mathcal{V}(r)) \inf_{\mathcal{V}(r)^2} q_\tau^+ > 0. \end{aligned}$$

It is out of the scope of this article to review the different classes of absolutely continuous operators and related two-sided Gaussian estimates arising in the analysis of continuous time elliptic diffusion and particle absorption models. For a more thorough discussion on this topic we refer to the series of reference pointers presented above.

Needless to say that the design of Lyapunov functions is a crucial and challenging problem in the stability analysis of positive semigroups. We have chosen to concentrate our review on presenting practical and general principles for designing Lyapunov functions.

2.3 Time homogenous models

For time homogeneous models we use the notation

$$(\Phi_t, Q_t, \bar{Q}_t) := (\Phi_{0,t}, Q_{0,t}, \bar{Q}_{0,t}).$$

As expected for time homogeneous semigroups a variety of results follow almost immediately from the estimates obtained in Theorem 2.3. Following [21], these results include the existence of an unique leading eigen-triple

$$(\rho, \eta_\infty, h) \in (\mathbb{R} \times \mathcal{P}_V(E) \times \mathcal{B}_{0,V}(E)) \quad \text{with} \quad \eta_\infty(h) = 1 \quad (23)$$

in the sense that for any $t \in \mathcal{T}$ we have

$$Q_t(h) = e^{\rho t} h \quad \text{and} \quad \eta_\infty Q_t = e^{\rho t} \eta_\infty \quad \text{or equivalently} \quad \Phi_t(\eta_\infty) = \eta_\infty. \quad (24)$$

The eigenfunction h is sometimes called the ground state and the fixed point measure η_∞ the quasi-invariant measure. For any $x \in E$ we also have the product series formulation

$$0 < h(x) := \prod_{n \geq 0} \{1 + e^{-\rho \tau} [\Phi_{n\tau}(\delta_x)(Q_\tau(1)) - \Phi_{n\tau}(\eta_\infty)(Q_\tau(1))]\}.$$

In this context, choosing $(\mu, H) = (\eta_\infty, h)$ in (21), we readily check that

$$T_{s,s+t}^{\eta_\infty, h}(f) = T(f) := \frac{h}{\eta_\infty(h)} \eta_\infty(f) \quad \text{and} \quad \|e^{-\rho t} Q_t - T\|_V \leq c_h(\eta_\infty) e^{-bt}.$$

For any $\eta \in \mathcal{P}_V(E)$ we have the conjugate formulae

$$\Psi_h(\Phi_t(\eta)) = \Psi_h(\eta) P_t^h \quad (25)$$

with the Doob h -transform of Q_t defined by the Markov semigroup

$$P_t^h : f \in \mathcal{B}_{V^h}(E) \mapsto P_t^h(f) := e^{-\rho t} \frac{1}{h} Q_t(hf) \in \mathcal{B}_{V^h}(E).$$

Observe that

$$\eta_\infty = \Phi_t(\eta_\infty) \iff \eta_\infty^h := \Psi_h(\eta_\infty) = \eta_\infty^h P_t^h.$$

The Markov semigroup P_t^h is sometimes called the transition semigroup of the h -process, a.k.a. the process evolving in the ground state.

We further assume that Q_t is a sub-Markov semigroup of *self-adjoint operators* on $\mathbb{L}_2(\nu)$ with respect to some locally finite measure ν on E . In addition, there exists an orthonormal basis $(\varphi_n)_{n \geq 1}$ associated with a decreasing sequence of eigenvalues $\rho_n \leq 0$ such that

$$Q_t(x, dy) = \sum_{n \geq 1} e^{\rho_n t} \varphi_n(x) \varphi_n(y) \nu(dy). \quad (26)$$

In this context, the formulae (24) are satisfied with the parameters

$$(\rho, h) = (\rho_1, \varphi_1) \quad \text{and} \quad \eta_\infty(dx) = \Psi_h(\nu)(dx) := \frac{1}{\nu(h)} h(x) \nu(dx).$$

Note that in this case h has unit norm $\nu(h^2) = 1$. The spectral resolution (26) yields for any $t \geq 0$ and $f \in \mathbb{L}_2(\nu)$ the following decomposition

$$e^{-\rho t} Q_t(f)(x) - \frac{h(x)}{\eta_\infty(h)} \eta_\infty(f) = \sum_{n \geq 2} e^{\rho_n t} \varphi_n(x) \nu(\varphi_n f) \quad \text{with} \quad \rho_n^h = \rho_n - \rho_1. \quad (27)$$

This yields the following result.

Proposition 2.4. *For any time horizon $t \geq 0$ and any $f \in \mathbb{L}_2(\nu)$ we have the exponential estimates*

$$\left\| e^{-\rho t} Q_t(f) - \frac{h}{\eta_\infty(h)} \eta_\infty(f) \right\|_{\mathbb{L}_2(\nu)} \leq e^{\rho_2^h t} (\nu(f^2) - \nu(hf)^2)^{1/2}. \quad (28)$$

Whenever Q_t is a positive semigroup of *self-adjoint operators* on $\mathbb{L}_2(\nu)$ the Doob h -transform P_t^h is a semigroup of self-adjoint operators on $\mathbb{L}_2(\eta_\infty^h)$ and we have the following spectral decomposition

Lemma 2.5. *For any $t \geq 0$ and $f \in \mathbb{L}_2(\eta_\infty^h)$ we have*

$$P_t^h(x, dy) = \eta_\infty^h(dy) + \sum_{n \geq 2} e^{\lambda_n t} h_n(x) h_n(y) \eta_\infty^h(dy) \quad (29)$$

with the $\mathbb{L}_2(\eta_\infty^h)$ orthonormal basis $(h_n)_{n \geq 2}$ defined for any $n \geq 2$ by

$$h_n := \varphi_n/h \quad \text{and} \quad \lambda_n = \rho_n - \rho_1 < 0 \quad \text{and} \quad \eta_\infty^h = \Psi_{h^2}(\nu).$$

Note that the density of the integral operator $P_t^h(x, dy)$ w.r.t. $\eta_\infty^h(dy)$ is given by

$$p_t^h(x, y) = e^{-\rho_1 t} \frac{q_t(x, y)}{h(x)h(y)} = 1 + \sum_{n \geq 2} e^{\lambda_n t} h_n(x) h_n(y). \quad (30)$$

We further assume that $h \in \mathcal{B}_0(E)$ and P_t^h is ultra contractive, in the sense that for any $t > 0$ we have

$$\|P_t^h\|_{\mathbb{L}_2(\eta_\infty^h) \rightarrow \mathbb{L}_\infty(\eta_\infty^h)} = e^{-\rho_1 t} \sup_{(x,y) \in E^2} \frac{q_t(x, y)}{h(x)h(y)} = \sup_{(x,y) \in E^2} p_t^h(x, y) < \infty. \quad (31)$$

Proposition 2.6. *Assume that $\nu(E) < \infty$ and $h \in \mathcal{B}_0(E)$. In addition, for any $t > 0$ (31) holds and the mapping $x \mapsto \int p_t^h(x, y) \nu(dy)$ is u.s.c. and locally lower bounded. In this situation, the function $V := 1/h \in \mathcal{B}_\infty(E)$ and for any $t > 0$ we have $Q_t(V)/V \in \mathcal{B}_0(E)$. In addition, for any $t > 0$ we have*

$$Q_t(V)/V \leq c_t/V^2 \in \mathcal{B}_0(E). \quad (32)$$

2.4 Markov diffusion semigroups

This section is mainly concerned with the design of Lyapunov functions for continuous time Markov semigroups. To simplify notation, we only consider time homogeneous models. All the semigroups discussed in this section satisfy condition (8). Thus, by (14) the contraction theorem, Theorem 2.1 applies to all the Markov semigroups discussed in this section as soon as the transition semigroups have a continuous density with respect to the Lebesgue measure.

Section 2.4.1 presents some elementary principles based on spectral conditions on the drift function and a simple way to design Lyapunov functions in terms of the generator of diffusion process. These generator-type techniques are illustrated in Section 2.4.2 in the context of overdamped Langevin diffusions. The design of Lyapunov functions for hypo-elliptic diffusions and Langevin diffusions are discussed respectively in Section 2.4.3 and Section 2.4.4.

2.4.1 Some general principles

Consider the Markov semigroup P_t of a diffusion flow $X_t(x)$ on $E = \mathbb{R}^n$ defined by

$$dX_t(x) = b(X_t(x)) dt + \sigma(X_t(x)) dB_t. \quad (33)$$

In the above display, B_t is a n_1 -dimensional Brownian motion starting at the origin for some $n \geq 1$, b is a differentiable drift function from \mathbb{R}^n into itself with gradient-matrix $\nabla b = (\partial_{x_i} b^j)_{1 \leq i, j \leq n}$, and σ stands for some diffusion function from \mathbb{R}^n into $\mathbb{R}^{n \times n_1}$. We set $\Sigma^2 := \sigma \sigma'$, where $\sigma'(x) := \sigma(x)'$ stands for the transposition of the matrix $\sigma(x)$, so that $\Sigma^2(x) := \sigma(x) \sigma'(x)$. The absolute continuity of the transition semigroup $P_t(x, dy) = \mathbb{P}(X_t(x) \in dy) = p_t(x, y) \nu(dy)$ for some continuous transition densities

$p_t(x, y)$ (w.r.t. the Lebesgue measure $\nu(dy)$) is ensured as soon as (b, σ) are globally Lipschitz continuous and the diffusion matrix is invertible or more generally satisfying a parabolic Hörmander condition (see for instance [53, 58, 56] and references therein). The generator L of the diffusion flow $X_t(x)$ and its carré du champ operator Γ_L are given respectively by the formula

$$L(f) := b' \nabla f + \frac{1}{2} \text{Tr} (\Sigma^2 \nabla^2 f) \quad \text{and} \quad \Gamma_L(f, g) := (\nabla f)' \Sigma^2 \nabla g. \quad (34)$$

The next proposition provides a rather elementary way to design a Lyapunov function.

Proposition 2.7. *Assume that $\sigma(x) = \sigma_0$ for some $\sigma_0 \in \mathbb{R}^{n \times n_1}$ and we have*

$$\nabla b + (\nabla b)' \leq -2\lambda I \quad \text{for some } \lambda > 0. \quad (35)$$

Then for any $v > 0$ and $t > 0$ there exists some $\delta_t > 0$ such that

$$V(x) := \exp(v\|x\|) \implies P_t(V)/V \leq c_t/V^{\delta_t}. \quad (36)$$

The proof of the above proposition is rather technical, thus it is provided in the appendix on page 63.

The next proposition is a slight extension of Theorem 2.6 [49] on reversible semi-groups to stochastic flows in Euclidean spaces. It provides a rather simple way to design Lyapunov functions in terms of generators.

Proposition 2.8. *Assume there exists some $\alpha > 0$, $\beta \in \mathbb{R}$ and $0 < \epsilon < 1$ such that*

$$\alpha W + \beta + L(W) \leq -\epsilon \Gamma_L(W, W). \quad (37)$$

In this situation, for any $t > 0$ we have

$$V := \exp(2\epsilon W) \implies P_t(V)/V \leq v_t/V^{\delta_t} \quad (38)$$

with the parameters

$$v_t = \exp(-2\beta\epsilon(1 - e^{-\alpha t})/\alpha) \quad \text{and} \quad \delta_t := (1 - e^{-\alpha t}).$$

The proof of the above proposition follows word-for-word the proof of Theorem 2.6 in [49], thus it is provided in the appendix on page 64.

We further assume that P_t satisfies for any $t > 0$ the sub-Gaussian estimate

$$P_t(x, dy) \leq c_t \exp\left(-\frac{1}{2\sigma_t^2} \|y - m_t(x)\|^2\right) dy \quad (39)$$

for some parameters $\sigma_t > 0$ and some function m_t on \mathbb{R}^n such that

$$\|m_t(x)\| \leq c_t (1 + \|x\|).$$

In this situation, for any $n \geq 1$ and $t \geq 0$ we have

$$V(x) := 1 + \|x\|^n \implies \|P_t(V)/V\| < \infty.$$

More refined estimates can be found when the function m_t is such that

$$|m_t(x)| \leq \epsilon_t |x| \quad \text{with} \quad \epsilon_t \in]0, 1[\quad (40)$$

for some norm $|\cdot|$ on \mathbb{R}^n . In this situation, observe that any $v \geq 0$ and any centered Gaussian random variable Y on \mathbb{R}^n with identity covariance matrix I_n we have

$$e^{-v|x|} \mathbb{E} \left(e^{v|m_t(x) + \sigma_t^2 Y|} \right) \leq c_t e^{-v(1-\epsilon_t)|x|}.$$

This yields the following lemma.

Lemma 2.9. *Consider a Markov semigroup P_t satisfying the sub-Gaussian estimate (39) as well as (40) for some norm $|\cdot|$ on \mathbb{R}^n . Then for any $v \geq 0$ and $t > 0$ there also exists some finite constant $\delta_t > 0$ such that*

$$V(x) := \exp(v|x|) \implies P_t(V)/V \leq c_t/V^{\delta_t}.$$

2.4.2 Overdamped Langevin diffusion

Let $W(x)$ be some twice differentiable potential function from \mathbb{R}^n into \mathbb{R} . The overdamped Langevin diffusion is defined by choosing in (33) the drift function

$$b(x) := -\gamma \nabla W(x) \quad \text{and} \quad (n_1, \sigma(x)) = (n, \rho I) \quad \text{for some} \quad \gamma, \rho > 0.$$

In this context, we have

$$(35) \iff \nabla^2 W \geq (\lambda/\gamma) I \quad \text{for some} \quad \lambda > 0.$$

Also observe that

$$(37) \iff \alpha W + \beta + \frac{\rho^2}{2} \text{Tr}(\nabla^2 W) \leq (\gamma - \epsilon \rho^2) \|\nabla W\|^2.$$

The above condition is clearly met when W behaves as $\|x\|^m$ with $m \geq 1$ at infinity; that is, there exists some sufficiently large radius r such that for any $\|x\| \geq r$ we have

$$|\text{Tr}(\nabla^2 W(x))| \leq c_1 \|x\|^{(m-2)+} \quad \text{and} \quad \|\nabla W(x)\|^2 \geq c_2 \|x\|^{2(m-1)}.$$

2.4.3 Hypo-elliptic diffusions

Consider the \mathbb{R}^n -valued diffusion (33) with $(b(x), \sigma(x)) = (Ax, \Sigma)$, for some matrices (A, Σ) with appropriate dimensions. We assume that A is stable (a.k.a. Hurwitz); that is its spectral abscissa $\zeta(A)$ defined below is negative

$$\zeta(A) := \sup \{ \operatorname{Re}(\lambda(A)) : \lambda(A) \in \operatorname{Spec}(A) \} < 0. \quad (41)$$

In the above display $\operatorname{Spec}(A)$ denotes the spectrum of the matrix A , and $\operatorname{Re}(\lambda(A))$ the real part of $\lambda(A)$. We also assume that $R := \Sigma\Sigma'$ is positive semi-definite and the pair of matrices $(A, R^{1/2})$ are controllable, in the sense that the $(n \times n^2)$ -matrices

$$[R^{1/2}, AR^{1/2}, \dots, A^{r-1}R^{1/2}] \quad \text{has rank } n. \quad (42)$$

Whenever $\zeta(A) < 0$ we have

$$P_t(x, dy) = \frac{1}{\sqrt{\det(2\pi C_t)}} \exp\left(-\frac{1}{2}(y - m_t(x))' C_t^{-1} (y - m_t(x))\right) dy \quad (43)$$

with the mean value function

$$x \mapsto m_t(x) := e^{tA}x \longrightarrow_{t \rightarrow \infty} 0$$

and the covariance matrices C_t defined for any $t > 0$ by

$$0 < C_t := \int_0^t e^{sA} R e^{sA'} ds \longrightarrow_{t \rightarrow \infty} C_\infty := \int_0^\infty e^{sA} R e^{sA'} ds.$$

Since A is stable, there exists some norm $|\cdot|$ on \mathbb{R}^n such that the corresponding operator norm satisfies $|e^{tA}| \leq e^{l(A)t}$ for some log-norm parameter $l(A) < 0$. This implies that

$$|m_t(x)| = |e^{tA}x| \leq e^{l(A)t} |x|. \quad (44)$$

This clearly shows that the semigroup P_t of the hypo-elliptic Ornstein-Uhlenbeck diffusion satisfies (39) and (40).

Let \mathcal{P}_t be the Markov semigroup of the \mathbb{R}^n -valued linear diffusion

$$d\mathcal{X}_t(x) = (A\mathcal{X}_t(x) + a(\mathcal{X}_t(x))) dt + \Sigma dB_t \quad (45)$$

with some bounded drift function a on \mathbb{R}^n , an $(n \times n)$ -matrix A satisfying (41), some n_1 -valued Brownian motion B_t starting at the origin and some $(n \times n_1)$ -matrix Σ satisfying the rank condition (42).

Using the stochastic interpolation formula (cf. Theorem 1.2 in [23]) given by

$$\mathcal{X}_t(x) - X_t(x) = \int_0^t e^{(t-s)A'} a(\mathcal{X}_s(x)) ds$$

we check the almost sure estimate

$$|X_t(x) - \mathcal{X}_t(x)| \leq c \quad \text{for some finite constant } c < \infty.$$

This yields the following proposition.

Proposition 2.10. *For any $v > 0$ and $t > 0$ there exists some $\delta_t > 0$ such that*

$$V(x) := \exp(v|x|) \implies \mathcal{P}_t(V)/V \leq c_t/V^{\delta_t}.$$

2.4.4 Langevin diffusion

Consider the Langevin diffusion flow $\mathcal{X}_t(z) = (X_t(z), Y_t(z)) \in (\mathbb{R}^r \times \mathbb{R}^r)$ starting at $z = (x, y) \in (\mathbb{R}^r \times \mathbb{R}^r)$ and given by

$$\begin{aligned} dX_t(z) &= Y_t(z)/m dt \\ dY_t(z) &= (b(X_t(z)) - \beta Y_t(z)/m) dt + \sigma dB_t. \end{aligned}$$

In the above display, B_t stands for an r -dimensional Brownian motion B_t starting at the origin, $\sigma, \beta, m > 0$ some parameters and b a function of the form

$$b(x) := -\gamma x + a(x) \quad \text{with} \quad \gamma > 0 \quad \text{and} \quad \|a\| < \infty.$$

In statistical physics, the above diffusion represents the evolution of N particles $X_t(z) = (X_t^i(z))_{1 \leq i \leq N} \in \mathbb{R}^{3N}$ with mass $m > 0$, position $X_t(z) \in \mathbb{R}^{3N}$ and momenta $Y_t(z)$. In this context, $\gamma > 0$ stands for some friction parameter, and the diffusion parameter $\sigma > 0$ is related to the Boltzmann constant and the temperature of the system. In this context, the function $b(x) = -\nabla W(x)$ is often described by the gradient of some potential function W . For instance, for a quadratic confinement we have

$$W(x) := \gamma \|x\|^2/2 + w(x) \quad \text{with} \quad \|\nabla w\| < \infty$$

$$\implies b(x) = -\nabla W(x) := -\gamma x + a(x) \quad \text{and} \quad a(x) = \nabla w(x).$$

Notice that $\mathcal{X}_t(z)$ can be rewritten in vector form as in (45) with $n = 2r$, $a(x, y) = \begin{pmatrix} 0 \\ a(x) \end{pmatrix}$ and the matrices

$$A = \begin{pmatrix} 0 & m^{-1} I_{n \times n} \\ -\gamma I_{n \times n} & -\beta m^{-1} I_{n \times n} \end{pmatrix} \quad \text{and} \quad \Sigma := \begin{pmatrix} 0 & 0 \\ 0 & \sigma I_{n \times n} \end{pmatrix}. \quad (46)$$

It is a simple exercise to check that A satisfies (41) and (42).

Consider the \mathbb{R}^2 -valued stochastic process $X_t = (q_t, p_t)$ defined by

$$\begin{cases} dq_t &= \beta \frac{p_t}{m} dt \\ dp_t &= -\beta \left(\frac{\partial W}{\partial q}(q_t) + \frac{\sigma^2}{2} \frac{p_t}{m} \right) dt + \sigma dB_t \end{cases} \quad (47)$$

with some positive constants β, m, σ , a Brownian motion B_t , and a smooth positive function W on \mathbb{R} such that for sufficiently large r we have

$$\forall |q| \geq r \quad q \frac{\partial W}{\partial q}(q) \geq \delta (W(q) + q^2)$$

for some positive constant δ . This condition is clearly met when W behaves as q^{2l} for certain $l \geq 1$ at infinity. We let $V(q, p)$ be the function on \mathbb{R}^2 defined by

$$V(q, p) = 1 + \frac{1}{2m} p^2 + W(q) + \frac{\epsilon}{2} \left(\frac{\sigma^2}{2} q^2 + 2pq \right) \quad \text{with} \quad \epsilon < \frac{\sigma^2}{2m}.$$

In this situation, there exists some $a > 0$ and $c < \infty$ such that

$$L(V) \leq -aV + c. \quad (48)$$

The proof of the above estimate is rather technical, thus it is provided in the appendix on page 67.

2.5 Sub-Markov semigroups

Sub-Markov semigroups are prototype-based models of positive integral operators. In time homogeneous settings, these stochastic models are defined in terms of a stochastic flow $X_t(x)$ evolving on some metric Polish space (\mathcal{E}, d) , some non negative absorption potential function U on some non necessarily bounded Borel subset $E \subset \mathcal{E}$. For a given $x \in E$ we denote by $T(x)$ the exit time of the flow $X_t(x)$ from E .

We associate with these objects, the sub-Markov semigroup $Q_t^{[U]}$ defined for any $f \in \mathcal{B}_b(E)$ and $x \in E$ by

$$Q_t^{[U]}(f)(x) = \mathbb{E} \left(f(X_t(x)) 1_{T(x) > t} \exp \left(- \int_0^t U(X_s(x)) ds \right) \right). \quad (49)$$

The above model can be interpreted as the distribution of a stochastic flow evolving in an absorbing medium with hard and soft obstacles. Before killing, the flow starts at $x \in E$ and evolves as $X_t(x)$. Then, it is killed at rate U or as soon as it exits the set E . In the case $E = \mathcal{E}$, the flow cannot exit the set E and it is only killed at rate U . This situation is sometimes referred a sub-Markov semigroup with soft obstacles represented by the absorbing potential function U on E . When the flow may exit the set $E \subset \mathcal{E}$, the complementary subset $C := \mathcal{E} - E$ is interpreted as an hard obstacle, a.k.a. an infinite energy barrier.

We illustrate the V -positive semigroup analysis developed in this article through three typical examples of solvable sub-Markov semigroups arising in physics and applied probability.

2.5.1 The harmonic oscillator

Consider the case $E = \mathcal{E} = \mathbb{R}$, and let $X_t(x) = B_t(x)$ be a Brownian motion starting at $x \in \mathbb{R}$ and let $U(x) = x^2/2$. In this situation, the semigroup $Q_t^{[U]} = Q_t$ defined in (49) coincides with the one dimensional harmonic oscillator. For any $t > 0$, the integral operator Q_t has a continuous density w.r.t. the uniform measure ν on E given by

$$q_t(x, y) = \sum_{n \geq 1} e^{\rho_n t} \varphi_n(x) \varphi_n(y) \quad (50)$$

with the $\mathbb{L}_2(\nu)$ orthonormal basis eigenstates

$$\varphi_n(x) = (2^{n-1}(n-1)!\sqrt{\pi})^{-1/2} e^{-x^2/2} \mathbb{H}_{n-1}(x)$$

associated with the eigenvalues

$$\rho_n = -(n - 1/2) \quad \text{and the Hermite polynomials} \quad \mathbb{H}_n(x) = (-1)^n e^{x^2} \partial^n e^{-x^2}.$$

In this context, the eigenstate associated with the top eigenvalue $\rho = \rho_1 = -1/2$ is given by the harmonic function

$$h(x) = \varphi_1(x) = \pi^{-1/4} e^{-x^2/2}. \quad (51)$$

The spectral resolution of integral operator $P_t^h(x, dy)$ and its density $p_t^h(x, y)$ with respect to the invariant measure

$$\eta_\infty^h(dy) = \frac{1}{\sqrt{\pi}} e^{-y^2} dy$$

are given as in (29) and (30) with $\mathbb{L}_2(\eta_\infty^h)$ orthonormal basis defined for any $n \geq 2$ by

$$h_n = (2^{n-1}(n-1)!)^{-1/2} \mathbb{H}_{n-1} \quad \text{and} \quad \rho_n^h = \rho_n - \rho_1 = -(n-1).$$

In this context, the h -process is given by the Ornstein-Uhlenbeck diffusion

$$dX_t^h(x) = \partial \log h(X_t^h(x)) dt + dB_t = -X_t^h(x) dt + dB_t. \quad (52)$$

In the above display, $B_t = B_t(0)$ stands for the one dimensional Brownian motion starting at the origin. The conjugate formula

$$Q_t(hf)/Q_t(h) = P_t^h(f) \iff Q_t(f) = e^{ot} h P_t^h(f/h) \quad (53)$$

yields the following proposition.

Proposition 2.11. *For any time horizon $t \geq 0$ we have*

$$Q_t(x, dy) = \frac{1}{\sqrt{\cosh(t)}} \exp\left(-\frac{x^2}{2} p_t\right) \frac{1}{\sqrt{2\pi p_t}} \exp\left(-\frac{(y - m_t(x))^2}{2p_t}\right) dy$$

with the mean and variance parameters $(m_t(x), p_t)$ defined by

$$m_t(x) = x/\cosh(t) \quad \text{and} \quad p_t = \tanh(t).$$

The proof of the above proposition is a direct consequence of the conjugate formula, thus it is provided in the appendix, on page 65.

Choosing $V(x) = 1 + |x|^n$, for some $n \geq 1$, we readily check that

$$V \in \mathcal{C}_\infty(E) \quad \text{and} \quad \mathcal{Q}_t(V)/V \leq v_t \mathcal{Q}_t(1) \in \mathcal{C}_0(E) \quad (54)$$

where v_t is a constant depending only on t .

2.5.2 The half-harmonic oscillator

Consider the case $E =]0, \infty[\subset \mathcal{E} = \mathbb{R}$, and let $X_t(x) = B_t(x)$ be a Brownian motion starting at $x \in \mathbb{R}$ and let $U(x) = x^2/2$. In this situation, the semigroup $Q_t^{[U]} = Q_t$ defined in (49) coincides with the harmonic oscillator with an infinite barrier at the origin $\partial E = \{0\}$ (a.k.a. the half-harmonic oscillator). Using the fact that

$$e^{x^2/2} \frac{1}{2} \partial^2 e^{-x^2/2} = U(x) - 1/2$$

we have the conjugate formula

$$Q_t(f)(x) = e^{-t/2} e^{-x^2/2} \mathbb{E} \left(f(Y_t(x)) e^{Y_t(x)^2/2} 1_{T^Y(x) > t} \right)$$

with the Ornstein-Uhlenbeck diffusion

$$dY_t(x) = -Y_t(x) dt + dB_t \quad \text{and} \quad T^Y(x) := \inf \{t \geq 0 : Y_t(x) \in \partial E\}. \quad (55)$$

Note that the stochastic flow $Y_t(x)$ coincides with the h -process of the harmonic oscillator discussed in (52). Thus, by reflection arguments we have

$$\begin{aligned} Q_t^Y(f)(x) &:= \mathbb{E} (f(Y_t(x)) 1_{T^Y(x) > t}) = \mathbb{E} (f(B_{\sigma_t}(\epsilon_t x)) 1_{T(\epsilon_t x) > t}) \\ &= \int_0^\infty f(y) q_t^Y(x, y) dy \quad \text{with} \quad q_t^Y(x, y) := (r_t(x, y) - r_t(x, -y)). \end{aligned}$$

In the above display, (ϵ_t, σ_t) stands for the parameters

$$(\epsilon_t, \sigma_t) := \left(e^{-t}, \sqrt{\frac{1 - \epsilon_t^2}{2}} \right) \quad \text{and} \quad r_t(x, y) = \frac{1}{\sqrt{2\pi\sigma_t^2}} \exp \left(-\frac{1}{2\sigma_t^2} (y - \epsilon_t x)^2 \right).$$

This yields the following proposition.

Proposition 2.12. *For any $t > 0$ and $x \in E =]0, \infty[$ we have*

$$\bar{Q}_t(x, dy) = \frac{\sinh(y m_t(x))}{\mathbb{P}(0 \leq Z \leq m_t(x)/\sqrt{p_t})} \times \frac{1}{\sqrt{2\pi p_t}} \exp \left(-\frac{y^2 + m_t(x)^2}{2p_t} \right) \nu(dy).$$

In the above display, $\nu(dy) := 1_{[0, \infty[}(y) dy$ stands for the trace of the Lebesgue measure on the half-line, Z is a centered Gaussian variable with unit variance and $(m_t(x), p_t)$ are the mean and variance parameters defined in Proposition 2.11. In addition, the total mass function $Q_t(1)(x)$ is given by the formula

$$Q_t(1)(x) = 2 \frac{e^{-\frac{x^2}{2} p_t}}{\sqrt{\cosh(t)}} \times \mathbb{P}(0 \leq Z \leq m_t(x)/\sqrt{p_t}) \in \mathcal{C}_0(E).$$

The proof of the above proposition follows the same lines of arguments as the proof of Proposition 2.11; it is provided in the appendix, on page 66.

Choosing $V(x) = x^n + 1/x$, for some $n \geq 1$, we readily check that

$$V \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t(V)/V \leq c_t/V \in \mathcal{C}_0(E). \quad (56)$$

The proof of the above estimate follows elementary but lengthy calculations, thus it is provided in the appendix on page 68.

For any $t > 0$, the integral operator Q_t has a continuous density w.r.t. the uniform measure ν on E given by

$$q_t(x, y) = \sum_{n \geq 1} e^{\rho_n t} \varphi_n(x) \varphi_n(y)$$

with the $\mathbb{L}_2(\nu)$ orthonormal basis eigenstates

$$\varphi_n(x) = \sqrt{2} (2^{2n-1} (2n-1)! \sqrt{\pi})^{-1/2} e^{-x^2/2} \mathbb{H}_{2n-1}(x)$$

associated with the eigenvalues

$$\rho_n = -((2n-1) + 1/2).$$

In this context, the eigenstate associated with the top eigenvalue $\rho = \rho_1 = -3/2$ is given for any $x \in]0, \infty[$ by the harmonic function

$$h(x) = \varphi_1(x) = 2\pi^{-1/4} x e^{-x^2/2} = h_0(x) \mathbb{H}_1(x)$$

with the ground state h_0 of the harmonic oscillator discussed in (51). Note that h coincides with the restriction on $]0, \infty[$ of the first excited state of the harmonic oscillator (negative on $] -\infty, 0]$ and crossing the origin at $x = 0$).

The spectral resolution of integral operator $P_t^h(x, dy)$ and its density $p_t^h(x, y)$ with respect to the invariant measure

$$\eta_\infty^h(dy) = \frac{4}{\sqrt{\pi}} y^2 e^{-y^2} 1_{]0, \infty[}(y) dy$$

are given for any $x, y \in]0, \infty[$ as in (29) and (30) with $\mathbb{L}_2(\eta_\infty^h)$ orthonormal basis defined for any $n \geq 2$ and $x \in]0, \infty[$ by the odd Hermite functions

$$h_n(x) = (2^{2n} (2n-1)!)^{-1/2} \mathbb{H}_{2n-1}(x)/x \quad \text{and} \quad \rho_n^h = -2(n-1).$$

In this context, the h -process is given by the diffusion

$$dX_t^h(x) = \partial \log h(X_t^h(x)) dt + dB_t = \left(\frac{1}{X_t^h(x)} - X_t^h(x) \right) dt + dB_t. \quad (57)$$

2.5.3 The Dirichlet heat kernel

Let $X_t(x) = B_t(x)$ be a Brownian motion starting at $x \in E :=]0, 1[\subset \mathcal{E} := \mathbb{R}$ and $T(x)$ be the first time $t \geq 0$ the process $B_t(x) \in \partial E := \{0, 1\}$. Choosing $U = 0$ in (49), the semigroup $Q_t^{[U]} = Q_t$ takes the following form

$$Q_t(f)(x) := \mathbb{E}(f(B_t(x)) \mathbf{1}_{T(x) > t}).$$

For any $t > 0$, the integral operator Q_t has a continuous density w.r.t. the uniform measure ν on E given by the Dirichlet heat kernel

$$q_t(x, y) = \sum_{n \geq 1} e^{\rho_n t} \varphi_n(x) \varphi_n(y) \quad (58)$$

with the $\mathbb{L}_2(\nu)$ orthonormal basis eigenstates

$$\varphi_n(x) = \sqrt{2} \sin(n\pi x) \quad \text{associated with the eigenvalues} \quad \rho_n = -(n\pi)^2/2.$$

In this context, the eigenstate $h(x) = \varphi_1(x) = \sqrt{2} \sin(\pi x)$ associated with the top eigenvalue $\rho = \rho_1 = -\pi^2/2$ is strictly positive except at the boundary $\{0, 1\}$. By removing the boundary, the semigroup P_t^h of the process evolving in the ground state $h(x)$ on the open interval $E :=]0, 1[$ is a self-adjoint operators on $\mathbb{L}_2(\eta_\infty^h)$ with

$$\eta_\infty^h(dx) = h^2(x) \nu(dx) = 2 \sin^2(\pi x) \mathbf{1}_E(x) dx.$$

In addition, we have the spectral decomposition (29) with the $\mathbb{L}_2(\eta_\infty^h)$ orthonormal basis eigenstates

$$h_n(x) := \sin(n\pi x) / \sin(\pi x)$$

associated with the eigenvalues

$$\lambda_n = -\pi^2(n^2 - 1)/2 < 0.$$

Our next objective is to estimate the density $p_t^h(x, y)$ of the integral operator $P_t^h(x, dy)$ w.r.t. η_∞^h defined in (30). Recalling that $|\sin(ny)| \leq n|\sin(y)|$, for any $n \geq 1$ and $y \in \mathbb{R}$, for any $x \in E$ we have the diagonal estimate

$$p_t^h(x, x) - 1 = \sum_{n \geq 2} e^{\rho_n t} h_n(x)^2$$

with

$$h_n(x)^2 = \left(\frac{\sin(n\pi x)}{\sin(\pi x)} \right)^2 \leq n^2 \quad \text{so that condition (31) is satisfied.}$$

Observe that the function

$$V : x \in E \mapsto V(x) := \sqrt{2}/h(x) \in [1, \infty[$$

is locally bounded with compact level sets given for any $0 < \epsilon \leq 1$ by the formulae

$$\mathcal{K}_\epsilon := \{x \in]0, 1[: V(x) \leq 1/\epsilon\} = \{x : \sin(\pi x) \geq \epsilon\} \subset E.$$

In any dimension we can use the intrinsic ultracontractivity to produce a Lyapunov function V . Let E be a bounded domain of \mathbb{R}^n for some $n \geq 1$ and assume that it is a $C^{1,\alpha}$ domain for some $\alpha > 0$. Denote by $q_t(x, y)$ the Dirichlet heat kernel on E . By [57] one has

$$q_t(x, y) \leq c_t d(x, \partial E) d(y, \partial E)$$

for some constant c_t independent of x and y . Here $d(x, \partial E)$ denotes the distance from x to the boundary of E . Set $V(x) = \frac{1}{d(x, \partial E)}$. The above intrinsic ultracontractivity implies

$$Q_t(V)(x) = \int_E q_t(x, y) V(y) dy \leq c_t |E| d(x, \partial E)$$

which in turn gives $Q_t(V)/V \leq c_t |E|/V^2 \in \mathcal{B}_0(E)$, where $|E|$ stands for the volume of the bounded set E .

3 Lyapunov design principles

The aim of this section is to present some general principles to construct Lyapunov functions for positive semigroups. Section 3.1 provides equivalent formulations of the Lyapunov condition in (1) encountered in the literature in terms of exhausting sequences of compact level sets. This section also presents simple ways to design Lyapunov functions for sub-Markov semigroups on normed spaces in terms of their generators. Section 3.2 presents some principles to construct Lyapunov functions for positive semigroups dominated by semigroups with known Lyapunov functions. Section 3.3 is dedicated to the design of Lyapunov functions for conjugate semigroups. All the principles discussed in this section are illustrated in Section 5 as well as in Section 6 in the context of conditional diffusions.

3.1 Foster-Lyapunov conditions

For time homogeneous models $Q_{s,s+t} := Q_t$, the l.h.s. condition in (1) takes the form $Q_\tau(V)/V \leq \Theta_\tau \in \mathcal{B}_0(E)$. In terms of the compact sets $K_\epsilon := \{\Theta_\tau \geq \epsilon\}$, the l.h.s Lyapunov condition in (1) yields for any $\tau > 0$ the estimate

$$Q_\tau(V)(x) \leq \epsilon V(x) + 1_{K_\epsilon}(x) c_\epsilon \tag{59}$$

for any $\epsilon > 0$ with the parameter $c_\epsilon := \sup_{K_\epsilon}(V\Theta_\tau) < \infty$. This implies that for any $n \geq 1$ we have

$$Q_\tau(V)(x) \leq \epsilon_n V(x) + 1_{K_{\epsilon_n}}(x) c_{\epsilon_n} \tag{60}$$

where $K_{\epsilon_n} \subset E$ stands for some increasing sequence of compact sets and c_{ϵ_n} some finite constants, indexed by a decreasing sequence of parameters $\epsilon_n \in [0, 1]$ such that $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$. In the reverse angle, assume that $Q_\tau(V)/V$ is locally lower bounded and lower semicontinuous. In this situation, condition (60) ensures that $Q_\tau(V)/V \in \mathcal{B}_0(E)$ for any $\tau > 0$. Indeed, for any $\delta > 0$, there exists some $n \geq 1$ such that $\epsilon_n < \delta$ and we have

$$\{Q_\tau(V)/V \geq \delta\} \subset \{Q_\tau(V)/V > \epsilon_n\} \subset K_{\epsilon_n}.$$

Since $\{Q_\tau(V)/V \geq \delta\}$ is a closed subset of a compact set it is also compact.

More generally, whenever (60) is met for some exhausting sequence of compact sets K_{ϵ_n} , in the sense that for any compact subset $K \subset E$ there exists some $n \geq 1$ such that $K \subset K_{\epsilon_n}$ we have

$$\inf_K Q_\tau(V)/V \geq \inf_{K_{\epsilon_n}} Q_\tau(V)/V \geq \epsilon_n.$$

This ensures that the function $Q_\tau(V)/V$ is necessarily locally lower bounded. In this situation, we have $Q_\tau(V)/V \in \mathcal{B}_0(E)$ as soon as $Q_\tau(V)/V$ is lower semicontinuous.

Notice that the sub-level set $\mathcal{V}(r) := \{V \leq r\}$ of the Lyapunov function $V \in \mathcal{B}_\infty(E)$ and the ϵ -super-level sets $K_\epsilon := \{\Theta_\tau \geq \epsilon\}$ of $\Theta_\tau \in \mathcal{B}_0(E)$ are equivalent compact exhausting sequences, in the sense that for any $r \geq 1$ we have

$$\mathcal{V}(r) \subset K_{\epsilon_r} \subset \mathcal{V}(r_\epsilon) \quad \text{with} \quad \epsilon_r := \inf_{\mathcal{V}(r)} \Theta_\tau \quad \text{and} \quad r_\epsilon := \sup_{K_{\epsilon_r}} V.$$

Whenever E is a locally compact Polish space, the abstract sequence $C_n := K_{\epsilon_n}$ in (60) is automatically exhausting; that is, we have that $E = \cup_{n \geq 0} C_n$ with C_n included in the interior C_{n+1}° of the compact set C_{n+1} . To check this claim, observe that for any $n \geq 1$ there exists some $m_n \geq n$ such that

$$C_n \subset \{\Theta_\tau \geq \inf_{C_n} \Theta\} \subset C_{m_n} \subset \{\Theta_\tau \geq \inf_{C_{m_n}} \Theta\}.$$

Thus, the exhausting sequence C_n is equivalent to the one defined by the super-level sets of Θ_τ .

The rather abstract condition (60) is often presented in the literature as an initial condition to check on a case-by-case basis to analyze the stability property of time homogenous sub-Markov semigroups (see for instance [31, 36], as well as Section 17.5 in [22] in the context of Markov semigroups and the references therein).

We end this section with a brief discussion on condition (60) in the context of the sub-Markov semigroup discussed in (49). Note that this semigroup can be turned into a Markov semigroup by sending the killed process into a cemetery state, say Δ , at the killing time. In this interpretation, functions on E are extended to $E_\Delta = E \cup \{\Delta\}$ by setting $f(\Delta) = 0$. More interestingly, whenever E is locally compact its topology coincides with the weak topology induced by $\mathcal{C}_0(E) := \mathcal{B}_0(E) \cap \mathcal{C}_b(E)$, and inversely (cf. Proposition 2.1 in [1]). In this context a continuous function f vanishes at infinity

if and only if its extension to the one point compactification (a.k.a. Alexandroff compactification) $E_\Delta := E \cup \{\Delta\}$ (obtained by setting $f(\Delta) = 0$) is continuous. For locally compact spaces, we also recall that the one point extension E_Δ is compact.

Whenever it exists, the generator L^U of these sub-Markov semigroups $Q_t^{[U]}$ are defined on domain of functions $D(L^U) \subset \mathcal{B}_0(E)$. As expected, the analysis of this class of models in terms of generators often requires to develop a sophisticated analysis taking into account the topological structure of the set E . To the best of our knowledge, there is no simple sufficient condition to check (60) in terms of these generators.

The situation is greatly simplified for sub-Markov semigroups with soft obstacles. When $E = \mathcal{E}$ is a locally compact normed space $(E, \|\cdot\|)$ we let L be the generator of the flow $X_t(x)$. In this situation, the generator of the sub-Markov semigroup $Q_t^{[U]}$ is given by $L^U = L - U$. We further assume that L and L^U are defined on some common domain $\mathcal{D}(L) \subset \mathcal{B}(E)$.

Lemma 3.1 ([31]). *Let $V, V_0 \in \mathcal{D}(L)$ be a couple of functions such that $V, V_0 \geq 1$ and*

$$V(x) \xrightarrow{\|x\| \rightarrow \infty} \infty \quad \text{and} \quad V(x)/V_0(x) \xrightarrow{\|x\| \rightarrow \infty} \infty. \quad (61)$$

In this situation, condition (60) is satisfied as soon as there exists some finite constant $c_0 < \infty$ such that

$$L^U(V_0)/V_0 \leq c_0 \quad \text{and} \quad L^U(V)(x)/V(x) \xrightarrow{\|x\| \rightarrow \infty} -\infty. \quad (62)$$

Note that in this context, the compact sets in (60) are given for some sufficiently large radii $r_\epsilon > 0$ by the closed balls:

$$\mathcal{K}_\epsilon = \overline{\mathbb{B}}(0, r_\epsilon) := \{x \in E : \|x\| \leq r_\epsilon\}. \quad (63)$$

3.2 Semigroup domination

For a given $p \geq 1$ we clearly have

$$V \in \mathcal{B}_\infty(E) \iff V^p \in \mathcal{B}_\infty(E) \quad \text{and} \quad \mathcal{B}_{V^{1/p}}(E) \subset \mathcal{B}_V(E) \subset \mathcal{B}_{V^p}(E).$$

We say that a V -positive semigroup $Q_{s,t}$ is p -dominated by a collection of integral operators $\mathcal{Q}_{s,t}$ on $\mathcal{B}_{V^p}(E)$ and we write $Q \ll_p \mathcal{Q}$ as soon as for any non negative function $f \in \mathcal{B}_V(E)$ and any $s \leq t$ we have

$$Q_{s,t}(f) \leq c_{t-s}(p) \mathcal{Q}_{s,t}(f^p)^{1/p}.$$

To simplify notation, when $p = 1$ we write $Q \ll \mathcal{Q}$ instead of $Q \ll_1 \mathcal{Q}$. Observe that

$$Q \ll_p \mathcal{Q} \implies \forall s \leq t \quad (Q_{s,t}(V)/V)^p \leq c_{t-s}(p)^p \mathcal{Q}_{s,t}(V^p)/V^p.$$

This yields for any $\tau > 0$ and $\theta_\tau \in \mathcal{B}_0(E)$ the Lyapunov estimate

$$Q_{s,s+\tau}(V^p)/V^p \leq \theta_\tau^p \implies Q_{s,s+\tau}(V)/V \leq c_\tau \theta_\tau. \quad (64)$$

We illustrate the above domination property with the Langevin diffusion flow $\mathcal{X}_t^{(a)}(z) = (X_t(z), Y_t(z)) \in (\mathbb{R}^n \times \mathbb{R}^n)$ starting at $z = (x, y) \in (\mathbb{R}^n \times \mathbb{R}^n)$ and defined by the hypo-elliptic diffusion

$$\begin{aligned} dX_t(z) &= Y_t(z)/m dt \\ dY_t(z) &= (a(X_t(z)) - \gamma X_t(z) - \beta Y_t(z)/m) dt + \sigma dB_t. \end{aligned} \quad (65)$$

In the above display, $\sigma, \gamma, \beta, m > 0$ stands for some parameters and a some Lipschitz function on \mathbb{R}^n , with $n \geq 1$. Notice that when $a = 0$, the flow $\mathcal{X}_t^{(0)}(z)$ resumes to an hypo-elliptic Ornstein-Uhlenbeck on \mathbb{R}^{2n} . Consider a bounded open connected domain $D \subset \mathbb{R}^n$ and set

$$\forall z \in E := D \times \mathbb{R}^n \quad T^{(a)}(z) := \inf \left\{ t \geq 0 : \mathcal{X}_t^{(a)}(z) \in \partial E \right\}.$$

We associate with these objects, the sub-Markov semigroup defined for any $f \in \mathcal{B}_b(E)$ and $z = (x, y) \in E$ by

$$\mathcal{Q}_t^{(a)}(f)(z) := \mathbb{E} \left(f(\mathcal{X}_t^{(a)}(z)) 1_{T^{(a)}(z) > t} \right).$$

In this situation, we have

$$\sup_D a < \infty \implies \forall p > 1 \quad \mathcal{Q}^{(a)} \ll_p \mathcal{Q}^{(0)}. \quad (66)$$

The proof of the above assertion is a direct consequence of Girsanov's theorem and Hölder's inequality. For the convenience of the reader, a detailed proof is provided in the appendix on page 70.

To emphasize the role of the absorption in sub-Markov semigroups we return to the class of models discussed in (49). We let P_t be the free evolution Markov semigroup associated with the stochastic flow $X_t(x)$. Assume that $Q_t^{[U]}(1) \in \mathcal{B}_0(E)$ and

$$\|Q_t^{[U]}(V)/V\| < \infty \quad \text{for some } t > 0 \text{ and } V \in \mathcal{B}_\infty(E). \quad (67)$$

Applying Hölder's inequality and choosing $V_p := V^{1/p} \in \mathcal{B}_\infty(E)$ with $p > 1$ we readily check the estimate

$$Q_t^{[U]}(V_p)/V_p \leq c_t(p) Q_t^{[U]}(1)^{1-1/p} \in \mathcal{B}_0(E). \quad (68)$$

The next lemma provides several practical conditions to check the uniform estimate (67) for sub-Markov semigroups associated with soft obstacles.

Lemma 3.2. *Consider the sub-Markov semigroup discussed in (49) when $E = \mathcal{E}$ is a locally compact normed space $(E, \|\cdot\|)$. Assume that the generators L and L^U of the flows P_t and $Q_t^{[U]}$ are defined on some common domain $\mathcal{D}(L) \subset \mathcal{B}(E)$. In this situation, for any $V \in \mathcal{B}_\infty(E) \cap \mathcal{D}(L)$ and parameter $a > 0$ we have*

$$L^U(V) \leq -aV + c \implies \forall t \geq 0 \quad \|Q_t^{[U]}(V)/V\| < \infty. \quad (69)$$

Whenever $U \in \mathcal{B}_\infty(E) \cap \mathcal{D}(L)$, for any $a_0 \geq 0$ and $a_1 \in \mathbb{R}$ we have

$$L(U) \leq a_0 + a_1 U \implies \forall t \geq 0 \quad \|Q_t^{[U]}(U)\| < \infty. \quad (70)$$

The proof of the above lemma follows essentially the same lines of arguments as the proof of Lemma 3.1; thus it is provided in the appendix, on page 69.

Whenever $E = \mathcal{E}$ and the absorption potential function U is bounded, we have $P \ll Q^{[U]} \ll P$. In this context, *there is no hope to have that $Q_t^{[U]}(1) \in \mathcal{B}_0(E)$ for some $t > 0$* . Nevertheless, for any $V \in \mathcal{B}_\infty(E)$ and any time horizon $t > 0$ we have

$$Q_t^{[U]}(V)/V \in \mathcal{B}_0(E) \iff P_t(V)/V \in \mathcal{B}_0(E).$$

In this situation, the design of Lyapunov functions V satisfying (1) or equivalently Foster-Lyapunov conditions of the form (60) is equivalent to the problem of finding a Lyapunov function for the Markov semigroup P_t .

Whenever P_t is stable, in the sense that it has a Lyapunov $V \in \mathcal{B}_\infty(E)$ such that $P_t(V)/V \in \mathcal{B}_0(E)$ for some $t > 0$, then the domination property $Q^{[U]} \ll P$ yields automatically a Lyapunov function for $Q_t^{[U]}$.

Whenever P_t is not necessarily stable but we have $\|P_t(V)/V\| < \infty$ for some $t > 0$ and $V \in \mathcal{B}_\infty(E)$, applying (68) the domination property $Q^{[U]} \ll P$ ensures that for any $p > 1$ we have $V_p := V^{1/p} \in \mathcal{B}_\infty(E)$ and

$$Q_t^{[U]}(1) \in \mathcal{B}_0(E) \implies Q_t^{[U]}(V_p)/V_p \in \mathcal{B}_0(E).$$

Last, but not least, note that the above discussion extends without difficulties to time varying models.

3.3 Some conjugacy principles

For any given $V \in \mathcal{B}_\infty(E)$, observe that for any positive function H ,

$$H \in \mathcal{B}_{0,V}(E) \iff V^H := V/H \in \mathcal{B}_\infty(E).$$

Thus, \mathcal{Q}_t is a V -positive semigroup on $\mathcal{B}_V(E)$ if and only if the H -conjugate semigroup $\mathcal{Q}_t^H(f) := \mathcal{Q}_t(fH)/H$ is a V^H -positive semigroup on $\mathcal{B}_{V^H}(E)$. In this situation, any semigroup $Q \ll \mathcal{Q}^H$ dominated by \mathcal{Q}^H yields for any $s \geq 0$ and $t > 0$ the Lyapunov estimate

$$Q_{s,s+t}(V^H)/V^H \leq c_t \mathcal{Q}_t(V)/V \in \mathcal{B}_0(E).$$

To get one step further, observe that

$$\mathcal{Q}_t(V)/V = \mathcal{Q}_t(1) \overline{\mathcal{Q}_t}(V)/V.$$

In this notation, for any $H \in \mathcal{B}_{0,V}(E)$ and any V -positive semigroup \mathcal{Q}_t on $\mathcal{B}_V(E)$ such $\mathcal{Q}_t(1) \in \mathcal{B}_0(E)$ and $\|\overline{\mathcal{Q}_t}\|_V < \infty$ we have

$$Q \ll \mathcal{Q}^H \implies Q_{s,s+t}(V^H)/V^H \leq c_t \mathcal{Q}_t(1) \in \mathcal{B}_0(E). \quad (71)$$

We illustrate the above comparison principles with an elementary example. Let $\mathcal{E} := \mathbb{R}$ and $W \in \mathcal{B}_\infty(\mathbb{R})$ be some non negative function. Consider the stochastic flow $X_t^W(x)$ of a one-dimensional Langevin diffusion on \mathcal{E} with generator

$$L(f) = \frac{1}{2} e^{2W} \partial (e^{-2W} \partial f). \quad (72)$$

We associate with a given open connected interval $E \subset \mathcal{E}$, the sub-Markov semigroup Q_t on $\mathcal{B}_b(E)$ defined by

$$Q_t(f)(x) := \mathbb{E}(f(X_t^W(x)) 1_{T_{\partial E}^W(x) > t}) \quad \text{with} \quad T_{\partial E}^W(x) := \inf \{t \geq 0 : X_t^W(x) \in \partial E\}. \quad (73)$$

Observe that

$$H := e^{-W} \implies U := H^{-1} \frac{1}{2} \partial^2 H = \frac{1}{2} ((\partial W)^2 - \partial^2 W). \quad (74)$$

When $W = 0$ the flow $X_t^0(x) = B_t(x)$ coincides with the Brownian flow $B_t(x)$ starting at x . Thus, by a change of probability we check that

$$Q_t = \mathcal{Q}_t^H \quad \text{with} \quad \mathcal{Q}_t(f)(x) := \mathbb{E} \left(f(B_t(x)) 1_{T_{\partial E}^0(x) > t} e^{-\int_0^t U(B_s(x)) ds} \right). \quad (75)$$

Whenever $E =]0, 1[$ the semigroup \mathcal{Q}_t is dominated by the Dirichlet heat kernel on $]0, 1[$. When $E = \mathbb{R}$, respectively $E =]0, \infty[$, and $U(x) \geq c + \varsigma x^2/2$, for some $c < \infty$ and $\varsigma > 0$, the semigroup \mathcal{Q}_t is dominated by the harmonic oscillator, respectively the half-harmonic oscillator. All of these dominating semigroups are completely solvable with $\mathcal{Q}_t(1) \in \mathcal{B}_0(E)$ and known Lyapunov functions.

4 Boundary problems

Let (\mathcal{E}, d) be a locally compact Polish space with a distinguish complete metric $d : (x, y) \in E^2 \mapsto d(x, y) \in \mathbb{R}_+$. We recall that these metric spaces are complete σ -compact and locally compact metric spaces, thus they have the Heine-Borel property, that is each closed and bounded subsets in \mathcal{E} are compact.

We also recall that a subspace $E \subset \mathcal{E}$ is Polish if and only if it is the intersection of a countable collection of open subsets. The distance from $x \in \mathcal{E}$ to a measurable subset $\mathcal{A} \subset \mathcal{E}$ is denoted by

$$d(x, \mathcal{A}) := \inf \{d(x, y) : y \in \mathcal{A}\}.$$

We also denote by $\partial E := \overline{E} - E^\circ$ the boundary of some domain (open and connected) $E \subset \mathcal{E}$, where \overline{E} and E° stand for the closure and the interior of a subset E .

In the further development of the article, χ stands for some decreasing positive function χ on $]0, \infty[$ such that for any $0 < \alpha < 1$ we have

$$\lim_{\alpha \rightarrow 0} \chi(\alpha) = +\infty \quad \chi(\alpha) < 1/\alpha \quad \text{and} \quad \bar{\chi}(\alpha) := \int_0^\alpha \chi(u) du < \infty.$$

Definition 4.1. We associate with χ the function $V_\partial \in \mathcal{C}(E)$ defined by

$$V_\partial : x \in E \mapsto V_\partial(x) := \chi(d(x, \partial E)) \in]0, \infty[. \quad (76)$$

For instance, we can choose $\chi(u) = 1/u^{1-\epsilon}$, for some $\epsilon \in]0, 1[$. For any $r > 0$ the r -sub-level sets of V_∂ are given by the closed subsets

$$\mathcal{V}_\partial(r) := \{x \in E : V_\partial(x) \leq r\} = \{x \in E : d(x, \partial E) \geq \chi^{-1}(r)\}.$$

Note that $V_\partial \in \mathcal{C}_\infty(E)$ as soon as \overline{E} is compact.

4.1 Bounded domains

Let $E \subset \mathcal{E} := \mathbb{R}^n$ be some bounded domain with locally Lipschitz boundary ∂E , for some $n \geq 1$. Consider a semigroup of integral operators

$$Q_t(x, dy) = q_t(x, y) dy \quad (77)$$

having for any $t > 0$ a bounded density $(x, y) \in E^2 \mapsto q_t(x, y) \in [0, \infty[$ w.r.t. the trace of the Lebesgue measure $\nu(dy) = dy$ on E . In this situation, we have the following lemma.

Lemma 4.2. For any $t > 0$ we have

$$V_\partial \in \mathcal{C}_\infty(E) \quad \text{and} \quad \|Q_t(V_\partial)\| \leq c_t \int_E \chi(d(x, \partial E)) dx < \infty. \quad (78)$$

The proof of the above lemma follows from an elementary change of variable formulae, thus it is provided in the appendix, on page 70.

The estimate (78) clearly applies to the class of sub-Markov semigroups $Q_t^{[U]}$ defined in (49) for any choice of the absorption potential function, as soon as the semigroup $Q^{[U]} \ll Q$ is dominated by a collection of integral operators $Q_t(x, dy)$ having a bounded density $q_t(x, y)$ on E^2 w.r.t. the Lebesgue measure on E . For instance, when the transition semigroup of the free evolution flow $X_t(x)$ in (49) has a density $p_t(x, y)$ for any non negative function f on E and any $x \in E$ we have

$$Q_t^{[U]}(f)(x) \leq \int q_t(x, y) f(y) dy \quad \text{with} \quad q_t(x, y) := p_t(x, y) 1_E(y).$$

We summarize the above discussion with the following proposition.

Proposition 4.3. Assume that $Q^{[U]} \ll Q$ is dominated by a collection of integral operators Q_t satisfying (77). Then,

$$Q_t^{[U]}(V_\partial)/V_\partial \leq c_t/V_\partial \in \mathcal{B}_0(E).$$

The choice of the Lyapunov function V is clearly not unique. For instance, when $E =]0, 1[$ instead of V_∂ we can choose $V(x) := 1/\sqrt{x} + 1/\sqrt{1-x}$. For the Dirichlet heat kernel discussed in Section 2.5.3 we can also choose $V(x) = 1/\sin(\pi x)$.

We emphasize that sub-Markov integral operators on the compact interval $E = [0, 1]$ with a *positive* continuous density w.r.t. the Lebesgue measure on E arise when the free evolution process is reflected at both sides of the interval. In this context the process is not conditioned by any type of non absorption at the boundaries. In this context, the unit function $V = 1$ belongs to $\mathcal{B}_\infty(E)$. In the same vein, sub-Markov integral operators with mixed boundary conditions on the left-closed interval $E = [0, 1[$, or respectively on the right-closed interval $E =]0, 1]$ arise when the free evolution process is reflected at the Neumann boundary $\partial_N E := \{0\}$ and non absorbed at the Dirichlet boundary $\partial_D E = \{1\}$, or respectively reflected at $\partial_N E := \{1\}$ and non absorbed at $\partial_D E = \{0\}$. More generally, consider a bounded domain $\Omega \subset \mathbb{R}^n$ with Lipschitz boundary $\partial\Omega = \partial_D\Omega \cup \partial_N\Omega$ consisting of two disjoint connected components $\partial_D\Omega$ and $\partial_N\Omega$ closed in \mathbb{R}^n , and set $E := \Omega \cup \partial_N\Omega$. In this notation, the function $V_\partial(x) := \chi(d(x, \partial_D E))$ belongs to $\mathcal{C}_\infty(E)$. In addition, for any bounded density $q_t(x, y)$ on E^2 we have the uniform estimate

$$\int_E q_t(x, y) V_\partial(y) dy \leq c_t \int_E V_\partial(y) dy < \infty.$$

The above estimate also holds for the function $V_\partial(x) = \chi(d(x, \partial E))$.

4.2 Unbounded domains

When the domain E is not bounded the function $V_\partial \notin \mathcal{B}_\infty(E)$. In this context, one natural way to design a Lyapunov function $V \in \mathcal{B}_\infty(E)$ is to consider an auxiliary function $V_\mathcal{E} \in \mathcal{C}_\infty(\mathcal{E})$ with $V_\mathcal{E}(x) \geq 1$ for any $x \in E$. In this situation, we have

$$V := V_\partial + V_\mathcal{E} \in \mathcal{C}_\infty(E).$$

To check this claim, observe that the sub-level sets of V_∂ are given by the closed subsets

$$\mathcal{V}_\partial(r) := \{V_\partial \leq r\} = \{x \in E : d(x, \partial E) \geq \chi^{-1}(r)\} \subset E$$

and we have the compact inclusion

$$\mathcal{V}(r) := \{V \leq r\} \subset \mathcal{V}_\mathcal{E}(r) \cap \mathcal{V}_\partial(r) \quad \text{with} \quad \mathcal{V}_\mathcal{E}(r) := \{x \in \mathcal{E} : V_\mathcal{E}(x) \leq r\}.$$

This yields the following easily checked proposition.

Proposition 4.4. *For any $t > 0$ we have*

$$\|Q_t(V_\partial)\| \vee \|Q_t(V_\mathcal{E})\| < \infty \implies Q_t(V)/V \leq c_t/V \in \mathcal{B}_0(E).$$

When $Q_t(1) \in \mathcal{B}_0(E)$ we also have

$$\|\overline{Q}_t(V_\partial)\| \vee \|\overline{Q}_t(V_\mathcal{E})/V_\mathcal{E}\| < \infty \implies Q_t(V)/V \leq c_t Q_t(1) \in \mathcal{B}_0(E).$$

The design of a function $V_{\mathcal{E}}$ is rather flexible. For instance, assume that $\overline{Q} \ll P$ is dominated by some Markov integral operators P_t on $\mathcal{B}_b(\mathcal{E})$ such that $\|P_t(V_{\mathcal{E}})/V_{\mathcal{E}}\| < \infty$ for some $V_{\mathcal{E}} \in \mathcal{B}_{\infty}(\mathcal{E})$. In this situation, we have $\|\overline{Q}_t(V_{\mathcal{E}})/V_{\mathcal{E}}\| < \infty$ as well as

$$\|V_{\mathcal{E}} Q_t(1)\| < \infty \implies \|Q_t(V_{\mathcal{E}})\| < \infty.$$

For instance, when P_t satisfies the sub-Gaussian estimates (39) on $\mathcal{E} = \mathbb{R}^n$ we can choose $V_{\mathcal{E}}(x) := 1 + \|x\|^k$, for some $k \geq 1$, as soon as the function $Q_t(1)(x) \xrightarrow{\|x\| \rightarrow \infty} 0$ faster than $\|x\|^{-k}$.

When the domain E and its boundary ∂E are both non necessarily bounded, it may happens that $Q_t(1) \in \mathcal{B}_0(E)$ but $\overline{Q}_t(V_{\partial}) \notin \mathcal{B}_b(E)$. In this situation, we can use the following proposition.

Proposition 4.5. *Assume there exists some $V_{\mathcal{E}} \in \mathcal{C}_{\infty}(\mathcal{E})$ with $V_{\mathcal{E}}(x) \geq 1$ for any $x \in E$ and such that*

$$\|\overline{Q}_t(V_{\partial})/V_{\mathcal{E}}\| \vee \|\overline{Q}_t(V_{\mathcal{E}})/V_{\mathcal{E}}\| \vee \|Q_t(1)V_{\mathcal{E}}\| < \infty.$$

Then we have

$$Q_t(V)/V \leq c_t/V \in \mathcal{B}_0(E).$$

Proof. Using the following decompositions

$$Q_t(V_{\partial}) = Q_t(1)V_{\mathcal{E}} \overline{Q}_t(V_{\partial})/V_{\mathcal{E}} \quad \text{and} \quad Q_t(V_{\mathcal{E}}) = Q_t(1)V_{\mathcal{E}} \overline{Q}_t(V_{\mathcal{E}})/V_{\mathcal{E}}$$

and applying Proposition 4.4 we have

$$\|Q_t(V_{\partial})\| \vee \|Q_t(V_{\mathcal{E}})\| < \infty \quad \text{and therefore} \quad Q_t(V)/V \leq c_t/V \in \mathcal{B}_0(E).$$

This ends the proof of the proposition. ■

The case $Q_t(1) \notin \mathcal{B}_0(E)$ can also be handle whenever the pair $(V_{\partial}, V_{\mathcal{E}})$ can be chosen so that

$$\forall \delta > 0 \quad V_{\partial} V_{\mathcal{E}}^{\delta} \in \mathcal{C}_{\infty}(E). \tag{79}$$

For instance we can choose for some $v > 0$ and $\epsilon \in]0, 1[$ the functions

$$V_{\mathcal{E}}(x) := \exp(v\|x\|) \quad \text{and} \quad \chi(u) := 1/u^{1-\epsilon}.$$

Observe that

$$d(x, \partial E) \leq \|x\| + d(0, \partial E) \quad \text{and} \quad V_{\partial}(x) \geq \chi(\|x\| + (1 \vee d(0, \partial E)))$$

and for any $m \geq 0$ and $\delta > 0$ we have

$$V_{\mathcal{E}}(x) \geq c_v(m, \delta) (1 + \|x\|)^{(m+1)/\delta}.$$

This implies that

$$V_{\mathcal{E}}(x)^{\delta} V_{\partial}(x) \geq c_2 \frac{(1 + \|x\|)^{m+1}}{(\|x\| + (1 \vee d(0, \partial E)))^{1-\epsilon}} \geq c (1 + \|x\|)^{m+\epsilon}.$$

Using the fact that $V_{\mathcal{E}}(x) \geq 1$ for any $x \in E$, this implies that

$$\{x \in E : V_{\mathcal{E}}(x)^{\delta} V_{\partial}(x) \leq r\} \subset \{x \in E : c (1 + \|x\|)^{m+\epsilon} \leq r\} \cap \{x \in E : V_{\partial}(x) \leq r\}.$$

We conclude that $V_{\mathcal{E}}^{\delta} V_{\partial}$ has compact level sets and (79) is satisfied.

In this context, we have the following proposition.

Proposition 4.6. *Consider a couple of functions $(V_{\partial}, V_{\mathcal{E}})$ satisfying (79). Assume there exists some parameters $t > 0$, $\delta_t > 0$ and $\epsilon \geq 0$ such that*

$$Q_t(V_{\mathcal{E}})/V_{\mathcal{E}} \leq c_t/V_{\mathcal{E}}^{\delta_t} \quad \text{and} \quad Q_t(V_{\partial}) \leq c_t V_{\mathcal{E}}^{\epsilon \delta_t}. \quad (80)$$

In this situation, for any $p > 1 + \epsilon$ we have

$$V := V_{\mathcal{E}}^{1-1/p} V_{\partial}^{1/p} \in \mathcal{C}_{\infty}(E)$$

as well as

$$Q_t(V)/V \leq c_t/(V_{\mathcal{E}}^{\delta_t \epsilon_p} V_{\partial}^{1/p}) \in \mathcal{C}_0(E) \quad \text{with} \quad \epsilon_p := (1 - (1 + \epsilon)/p) > 0.$$

Proof. Observe that for any $p > 1 + \epsilon$ we have

$$V_{\partial} V_{\mathcal{E}}^{p-1} \in \mathcal{C}_{\infty}(E) \quad \text{and therefore} \quad V := V_{\partial}^{1/p} V_{\mathcal{E}}^{1-1/p} \in \mathcal{C}_{\infty}(E).$$

In the same vein, for any $\epsilon \geq 0$ we have

$$(79) \implies V_{\partial} V_{\mathcal{E}}^{p \delta_t \epsilon_p} \in \mathcal{C}_{\infty}(E) \quad \text{and therefore} \quad V_{\partial}^{1/p} V_{\mathcal{E}}^{\delta_t \epsilon_p} \in \mathcal{C}_{\infty}(E).$$

On the other hand, using Hölder's inequality, we have

$$\begin{aligned} Q_t(V)/V &\leq (Q_t(V_{\mathcal{E}})/V_{\mathcal{E}})^{1-1/p} (Q_t(V_{\partial})/V_{\partial})^{1/p} \\ &\leq c_t(1)(Q_t(V_{\partial})/(V_{\mathcal{E}}^{\delta_t(p-1)} V_{\partial}))^{1/p} \leq c_t(2)(1/(V_{\mathcal{E}}^{p \delta_t(1-(1+\epsilon)/p)} V_{\partial}))^{1/p}. \end{aligned}$$

This ends the proof of the proposition. ■

The design of a function $V_{\mathcal{E}}$ satisfying (80) is rather flexible. For instance, (80) is automatically satisfied when $Q \ll P$ is dominated by some Markov integral operators P_t on $\mathcal{B}_b(\mathcal{E})$ such that

$$P_t(V_{\mathcal{E}})(x)/V_{\mathcal{E}}(x) \leq c_t(1)/V_{\mathcal{E}}(x)^{\delta_t}.$$

Section 2.4 discusses a variety of Lyapunov functions $V_{\mathcal{E}}$ satisfying the above condition for Markov diffusion semigroups. These Lyapunov functions can also be designed

using the domination principles presented in Section 3.2. For instance, consider the semigroup $Q_t := \mathcal{Q}_t^{(a)}$ associated with the Langevin diffusion flow on a cylinder discussed in (66). In this situation, combining (64) with Proposition 2.9, for any $v \geq 0$ and $t > 0$ there exists some finite constant $\delta_t > 0$ such that

$$V_{\mathcal{E}}(x) := \exp(v|x|) \implies Q_t(V_{\mathcal{E}})/V_{\mathcal{E}} \leq c_t/V_{\mathcal{E}}^{\delta_t}.$$

Next, we illustrate the r.h.s. condition in (80) when q_t are sub-Gaussian densities; in the sense that for any $x, y \in E$ we have

$$q_t(x, y) \leq c_t g_t(x, y) \quad \text{with} \quad g_t(x, y) := \frac{1}{(2\pi\sigma_t^2)^{n/2}} \exp\left(-\frac{1}{2\sigma_t^2} \|y - m_t(x)\|^2\right) \quad (81)$$

for some parameter $\sigma_t > 0$ and some non necessarily bounded function m_t on E .

Proposition 4.7. *Let φ be a Lipschitz function on \mathbb{R}^{n-1} with uniformly bounded gradient and set*

$$E := \{x = (x_i)_{1 \leq i \leq n} \in \mathbb{R}^n : x_n > \varphi(x_{-n})\} \quad \text{with} \quad x_{-n} := (x_i)_{1 \leq i < n} \in \mathbb{R}^{n-1}.$$

Then the r.h.s. condition in (80) is met with $\epsilon = 0$ for any positive semigroup satisfying (81). The same property holds when the boundary ∂E can be decomposed as a finite union of graphs of differentiable functions on \mathbb{R}^{n-1} with uniformly bounded gradients.

Proof. We choose $\alpha > 0$ sufficiently small so that for any

$$x \in D_{\alpha}(E) := \{x \in \bar{E} : d(x, \partial E) \leq \alpha\}$$

there exists a projection $\bar{x} \in \partial E$ with $d(x, \partial E) = \|x - \bar{x}\|$. Let $\mathcal{C}_{\varpi}(\bar{x})$ be an interior cone with a given base vertex $\bar{x} = (x_{-n}, \varphi(x_{-n})) \in \partial E$ and a given half-opening angle ϖ around the axis $\mathcal{A}(\bar{x}) := \{(x_{-n}, x_n) : x_n \geq \varphi(x_{-n})\}$. For any $x \in \mathcal{A}(\bar{x})$ there exists a projection $\hat{x} \in \partial \mathcal{C}_{\varpi}(\bar{x})$ on the boundary $\partial \mathcal{C}_{\varpi}(\bar{x})$ with

$$d(x, \partial \mathcal{C}_{\varpi}(\bar{x})) = d(x, \hat{x}) = \cos\left(\frac{\pi}{2} - \varpi\right) (x_n - \varphi(x_{-n})) \leq d(x, \bar{x})$$

On the other hand, for any $y \in \partial E$ we have

$$z := (y_{-n}, \varphi(x_{-n})) \implies 0 \leq \frac{\pi}{2} - \varpi \leq \widehat{y\bar{x}z} \quad \text{and} \quad \tan(\widehat{y\bar{x}z}) = \frac{|\varphi(x_{-n}) - \varphi(y_{-n})|}{\|x_{-n} - y_{-n}\|}.$$

This yields the estimate

$$\cos\left(\frac{\pi}{2} - \varpi\right) \geq \cos(\widehat{y\bar{x}z}) = \frac{1}{\sqrt{1 + \tan^2(\widehat{y\bar{x}z})}}$$

from which we conclude that

$$0 \leq x_n - \varphi(x_{-n}) \leq \kappa d(x, \bar{x})$$

with

$$\kappa := \sqrt{1 + \|\nabla\varphi\|^2} \quad \text{and} \quad \|\nabla\varphi\| := \sup_{y \in \mathbb{R}^{n-1}} \|\nabla\varphi(y)\| < \infty.$$

This implies that

$$\begin{aligned} & \int_{D_\alpha(E)} \chi(d(y, \partial E)) q_t(x, y) dy \\ & \leq c_t(1) \int_{D_\alpha(E)} \chi((y_n - \varphi(y_{-n}))/\kappa) \exp\left(-\frac{1}{2\sigma_t^2}((y_n - \varphi(y_{-n})) + (\varphi(y_{-n}) - (m_t(x))_n))^2\right) \\ & \quad \times \exp\left(-\frac{1}{2\sigma_t^2} \|y_{-n} - (m_t(x))_{-n}\|^2\right) dy_n dy_{-n}. \end{aligned}$$

Using the change of variables

$$z := (y_n - \varphi(y_{-n}))/\kappa \implies dy_n = \kappa dz$$

we find that

$$\begin{aligned} & \int_{D_\alpha(E)} \chi(d(y, \partial E)) q_t(x, y) dy \\ & \leq \kappa c_t(1) \bar{\chi}(\alpha) \int_{\mathbb{R}^{n-1}} \exp\left(-\frac{1}{2\sigma_t^2} \|y_{-n} - (m_t(x))_{-n}\|^2\right) dy_{-n} \leq c_t(2). \end{aligned}$$

On the other hand, for any $\alpha > 0$ we have

$$\int_{E - D_\alpha(E)} \chi(d(y, \partial E)) q_t(x, y) dy \leq \chi(\alpha) \|Q_t(1)\| \leq c_t(3).$$

This ends the proof of the proposition. ■

4.3 Smooth boundaries

Next, we illustrate the Lyapunov conditions on V_∂ in the context of absolutely continuous sub-Markov semigroup of the form (77) with a bounded density $q_t(x, y)$ on a non necessarily bounded domain $E \subset \mathbb{R}^n$ with smooth non necessarily bounded \mathcal{C}^2 -boundary with uniformly bounded interior curvature.

We assume that there exists $\alpha > 0$ sufficiently small so that every point of the α -offset of ∂E (a.k.a. α -tubular neighborhood) defined by

$$Tub_\alpha(\partial E) := \{x \in \mathbb{R}^n : d(x, \partial E) \leq \alpha\}$$

lies on some normal ray passing through a point on ∂E and no two normal rays passing through different points of ∂E intersect in $Tub_\alpha(\partial E)$. We let $N(z)$ be the unit normal vector at $z \in Tub_\alpha(\partial E)$ pointing inward E , and let $\mathcal{D}_r(E)$ the closed subset defined for any $r \leq \alpha$ by

$$\mathcal{D}_r(E) := \{x \in \bar{E} : d(x, \partial E) \leq r\} \quad \text{and} \quad \mathcal{D}_{-r}(E) := \{x \in \mathbb{R}^n - E : d(x, \partial E) \leq r\}.$$

In this notation, the inverse of the normal coordinate map

$$F : (z, r) \in \partial E \times [-\alpha, \alpha] \mapsto F(z, r) = z + r N(z) \in Tub_\alpha(\partial E) \quad (82)$$

is given for any $x \in Tub_\alpha(\partial E)$ by

$$F^{-1}(x) = (\text{proj}_{\partial E}(x), d_\alpha(x, \partial E))$$

where $\text{proj}_{\partial E}(x)$ stands for the projection of $x \in Tub_\alpha(\partial E)$ onto ∂E and $d_\alpha(x, \partial E)$ stands for the signed distance function

$$d_\alpha(x, \partial E) := d(x, \partial E) 1_{\mathcal{D}_\alpha(E)}(x) - d(x, \partial E) 1_{\mathcal{D}_{-\alpha}(E)}(x) \in [-\alpha, \alpha].$$

In addition, the inward normal $N(x)$ at any x on the \mathcal{C}^2 boundary ∂E is given by

$$\nabla d_\alpha(x, \partial E) = N(x).$$

The Hessian of the signed distance function on the boundary ∂E gives the Weingarten map $\mathcal{W}(x)$. With this notation at hand, we have

$$\int_{\mathcal{D}_\alpha(E)} f(d(y, \partial E)) q_t(x, y) dy = \int_0^\alpha f(r) q_t^\partial(x, r) dr$$

with the level-set density function

$$\begin{aligned} q_t^\partial(x, r) &:= \int_{\partial E_r} q_t(x, y) \sigma_{\partial, r}(dy) \\ &= \int_{\partial E} q_t(x, z + rN(z)) |\det(I - r\mathcal{W}(z))| \sigma_\partial(dz). \end{aligned} \quad (83)$$

In the above display, $\sigma_{\partial, r}(dz)$ stands for the Riemannian volume measure on the r -extended boundary

$$\partial E_r := \{x \in E : d(x, \partial E) = r\}.$$

Moreover, since E has uniformly bounded interior curvature, for any $r \leq \alpha$ we have

$$\kappa_\partial(\alpha) := \sup |\det(I - r \mathcal{W}(z))| < \infty \quad \text{and} \quad \kappa_\partial^-(\alpha) := \sup |\det(I + r \mathcal{W}(y))| < \infty.$$

In the above display, the supremum is taken over all $z \in \partial E$, $y \in \partial E_r$, and $r \leq \alpha$. Several examples of hypersurface boundaries satisfying the above conditions are discussed in Section 7 (cf. for instance Proposition 7.4).

We denote by $\bar{q}_t^\partial \geq q_t^\partial$ the function defined as q_t^∂ by replacing q_t by \bar{q}_t . Using the fact that

$$\bar{Q}_t(V_\partial)(x) \leq \chi(\alpha) + \int_0^\alpha \chi(r) \bar{q}_t^\partial(x, r) dr$$

we readily check the following proposition.

Proposition 4.8. *For any $t > 0$ we have*

$$\begin{aligned} \sup_{0 \leq r \leq \alpha} \sup_{x \in E} q_t^\partial(x, r) < \infty &\implies Q_t(V_\partial) \leq \chi(\alpha) Q_t(1) + c_t(\alpha) \bar{\chi}(\alpha) \\ \sup_{0 \leq r \leq \alpha} \sup_{x \in E} \bar{q}_t^\partial(x, r) < \infty &\implies \bar{Q}_t(V_\partial) \leq \chi(\alpha) + c_t(\alpha) \bar{\chi}(\alpha). \end{aligned} \quad (84)$$

When the boundary ∂E is bounded, for any $t > 0$ we have the estimate

$$\|\bar{Q}_t(V_\partial)\| \leq c_t(\alpha) \left(\chi(\alpha) + \bar{\chi}(\alpha) \sup_{0 \leq r \leq \alpha} \sigma_{\partial, r}(\partial E_r) \right). \quad (85)$$

We end this section with some practical tools to estimate the level-set density functions discussed in Section 4.3. Most of our estimates are based on the following technical lemma.

Lemma 4.9. *Consider a couple of non negative functions f, g on \mathbb{R}^n and some parameter $\alpha > 0$ such that*

$$\sup_{\|u\| \leq \alpha} f(z + u) \leq \iota(\alpha) g(z) \quad \text{for some } \iota(\alpha) < \infty.$$

In this situation, we have the uniform estimate

$$\sup_{0 \leq r \leq \alpha} \int_{\partial E_r} f(z) \sigma_{\partial, r}(dz) \leq \iota(\alpha) \kappa_\partial(\alpha) \int_{\partial E} g(z) \sigma_\partial(dz)$$

as well as the co-area estimate

$$\int_{\partial E} f(z) \sigma_\partial(dz) \leq \frac{1}{\alpha} \iota(\alpha) \kappa_\partial^-(\alpha) \int_{\mathcal{D}_\alpha(E)} g(z) dz.$$

The proof of the above lemma is provided in the appendix, on page 72.

Note that the level-set density function defined in (83) can be estimated for any $0 \leq r \leq \alpha$ by the formula

$$q_t^\partial(x, r) \leq \kappa_\partial(\alpha) \int_{\partial E} q_t(x, z + rN(z)) \sigma_\partial(dz).$$

Proposition 4.10. *Assume that $q_t(x, y) \leq \varpi_t g_t(x, y)$ is dominated by some probability density $y \mapsto g_t(x, y)$ on \mathbb{R}^n for some $t > 0$ and some parameter $\varpi_t < \infty$. In addition, we have*

$$\sup_{\|u\| \leq \alpha} g_t(x, y + u) \leq \iota_t(\alpha) g_{\alpha, t}(x, y) \quad (86)$$

for some probability density $y \mapsto g_{\alpha, t}(x, y)$ and some $\iota_t(\alpha) < \infty$. In this situation, we have the uniform density estimates

$$\sup_{0 \leq r \leq \alpha} \sup_{x \in E} q_t^\partial(x, r) \leq \varpi_t \iota_t(\alpha) \kappa_\partial^-(\alpha) \kappa_\partial(\alpha) / \alpha. \quad (87)$$

Proof. By (86) for any $0 \leq r \leq \alpha$ we have

$$q_t^\partial(x, r) \leq \varpi_t \kappa_\partial(\alpha) \int_{\partial E} g_t(x, z + rN(z)) \sigma_\partial(dz).$$

On the other hand, we have

$$\int_{\mathcal{D}_\alpha(E)} g_{t, \alpha}(x, y) dy \leq 1.$$

The estimate (87) is now a direct consequence of the co-area estimate stated in Lemma 4.9. This ends the proof of the proposition. \blacksquare

We illustrate the above condition when q_t are the sub-Gaussian densities discussed in (81). In this situation, using the fact that $2a'b \leq \frac{1}{\epsilon} \|a\|^2 + \epsilon \|b\|^2$ for any $0 < \epsilon < 1$ and $\|u\| \leq \alpha$ we check that

$$-\frac{1}{2\sigma_t^2} \|(y + u) - m_t(x)\|^2 \leq -\frac{(1 - \epsilon)}{2\sigma_t^2} \|y - m_t(x)\|^2 + \frac{1}{2\sigma_t^2} \left(\frac{1}{\epsilon} - 1\right) \alpha^2.$$

In this context, condition (86) is met with the gaussian density

$$g_{\alpha, t}(x, y) := \frac{e^{-\frac{1}{2\sigma_t(\epsilon)^2} \|y - m_t(x)\|^2}}{(2\pi\sigma_t(\epsilon)^2)^{n/2}}$$

with

$$\iota_t(\alpha) := c_t e^{\frac{\alpha^2/\epsilon}{2\sigma_t(\epsilon)^2}} \quad \text{and} \quad \sigma_t(\epsilon)^2 := \sigma_t^2/(1 - \epsilon).$$

5 Riccati type processes

5.1 Positive diffusions

Consider the Riccati type diffusion on $E =]0, +\infty[$ defined for any $x \in E$ by

$$dX_t(x) = (a_0 + a_1 X_t(x) - b X_t(x)^2) dt + \sigma_1(X_t(x)) dB_t^1 + \sigma_2(X_t(x)) dB_t^2, \quad X_0(x) = x$$

for some Brownian motion (B_t^1, B_t^2) on \mathbb{R}^2 , the diffusion functions

$$\sigma_1(x) := \varsigma_1 \sqrt{x} \quad \sigma_2(x) := \varsigma_2 x$$

and the parameters

$$a_1 \in \mathbb{R} \quad a_0 > \varsigma_1^2 \quad b > 0 \quad \text{and} \quad \varsigma_1, \varsigma_2 \geq 0.$$

Applying Itô's formula, we readily check that

$$\partial_t \mathbb{E}(X_t(x)) \leq \text{Ricc}(\mathbb{E}(X_t(x))) \quad \text{and} \quad \partial_t \mathbb{E}(1/X_t(x)) \leq \text{Ricc}^-(\mathbb{E}(1/X_t(x)))$$

with the Riccati drift functions defined by

$$\text{Ricc}(z) := a_0 + a_1 z - b z^2 \quad \text{and} \quad \text{Ricc}^-(z) := a_0^- + a_1^- z - b^- z^2 \quad (88)$$

with the parameters

$$a_0^- := b \quad a_1^- := (\varsigma_2^2 - a_1) \quad \text{and} \quad b^- := a_0 - \varsigma_1^2.$$

Consider the Lyapunov function $V \in \mathcal{B}_\infty(E)$ defined by $V(x) := x + 1/x$. By well known properties of Riccati flows, for any $t > 0$ we have $\|P_t(V)\| < \infty$. For a more thorough discussion on this class of one-dimensional Riccati diffusions, we refer to the article [4].

5.2 Matrix valued diffusions

Let \overline{E} and E be the space of $(n \times n)$ -positive semi-definite and definite matrices respectively. Also let $\lambda_1(x) \geq \dots \geq \lambda_n(x)$ denote the ordered eigenvalues of $x \in \overline{E}$. Let \mathcal{W}_t denotes an $(n \times n)$ -matrix with independent Brownian entries. Also let A be an $(n \times n)$ -matrix with real entries and let $R, S \in E$. We associate with these objects the E -valued diffusion

$$dX_t = (AX_t + X_t A' + R - X_t S X_t) dt + \frac{\epsilon}{2} \left[X_t^{1/2} d\mathcal{W}_t R^{1/2} + R^{1/2} d\mathcal{W}_t' X_t^{1/2} \right].$$

Whenever $\epsilon \leq 2/\sqrt{n+1}$, the diffusion X_t has a unique strong solution that never hits the boundary $\partial E = \overline{E} - E$. In addition, the transition semigroup P_t of X_t is strongly Feller and admits a smooth density w.r.t. the Lebesgue measure on E , thus it is irreducible. Furthermore, when $\epsilon^2(1+n)/2 \leq \lambda_n(R)/\lambda_1(R)$ then the function $V(x) = \text{Tr}(x) + \text{Tr}(x^{-1})$ is a Lyapunov function with compact level subsets. For a detailed proof of the above assertion for more general classes of Riccati matrix valued diffusions we refer to [5] (see for instance the stability Theorem 2.4 and Section 5.4 in [5]).

5.3 Logistic birth and death process

Let $X_t(x)$ be the stochastic flow on $E := \mathbb{N} - \{0\}$ with generator L defined for any $f \in \mathcal{B}_b(E)$ and $x \geq 2$ by

$$L(f)(x) = J(x, x-1) (f(x-1) - f(x)) + J(x, x+1) (f(x+1) - f(x))$$

and for $x = 1$ by

$$L(f)(1) = J(1, 2)(f(2) - f(1)).$$

In the above display, the birth and death rates in the above display are given by

$$J(x, x+1) := \lambda_b x + v_b \quad \text{and} \quad J(x, x-1) := \lambda_d x + \lambda_l x(x-1) + v_d \quad (89)$$

for some non negative parameters $\lambda_d, \lambda_b, v_b, v_d \geq 0$ and $\lambda_l > 0$. Consider the identity function $V : x \in E \mapsto V(x) = x$. For any $x \geq 2$ we have

$$L(V)(x) = J(x, x+1) - J(x, x-1) = (F \circ V)(x)$$

with the concave function

$$z \in \mathbb{R}_+ \mapsto F(z) := (v_b - v_d) + (\lambda_b + \lambda_l - \lambda_d) z - \lambda_l z^2 \in \mathbb{R}. \quad (90)$$

Observe that

$$L(V)(1) - F(V(1)) = J(1, 2) - F(1) = J(1, 0) = v_d + \lambda_d.$$

This yields the estimate

$$\begin{aligned} P_t(L(V))(x) &= P_t(1_{[2, \infty[} L(V))(x) + P_t(1_{\{1\}} L(V))(x) \\ &= P_t((F \circ V))(x) + P_t(1_{\{1\}})(x) (L(V)(1) - F(V(1))) \\ &\leq F(P_t(V))(x) + J(1, 0) \end{aligned}$$

from which we check that

$$\partial_t P_t(V)(x) \leq \text{Ricc}(P_t(V)(x))$$

with the Riccati drift function defined in (88) with the parameters

$$a_0 := v_b + \lambda_d \quad a_1 := \lambda_b + \lambda_l - \lambda_d \quad \text{and} \quad b := \lambda_l > 0.$$

By well known properties of Riccati flows, for any $t > 0$ we conclude that $\|P_t(V)\| < \infty$.

5.4 Multivariate birth and death processes

We denote by $e := \{e_i, 1 \leq i \leq n\}$ the collection of column vector e_i on $\{0, 1\}^n$ with entries $e_i(j) = 1_{i=j}$ and with a slight abuse of notation we denote by 0 the null state in \mathbb{N}^n . Let $X_t(x)$ be a stochastic flow on $E = \mathbb{N}^n - \{0\}$ with generator L defined by

$$L(f)(x) := \sum_{y \in E} J(x, y) (f(y) - f(x)). \quad (91)$$

Let $\lambda, \mu, v, \varsigma$ be some column vectors and let C, D some $(d \times d)$ -matrices with real entries such that for any $1 \leq i \leq d$ and any $x \in E$ we have

$$J(x, x + e_i) := v_i + x_i (\lambda_i + (Cx)_i) \geq 0 \quad \text{and} \quad J(x, x - e_i) := \varsigma_i + x_i (\mu_i + (Dx)_i) \geq 0.$$

We also set

$$J(x, y) = 0 \quad \text{as soon as} \quad |x - y| \geq 2.$$

We further assume that

$$|v| \geq |\varsigma| \quad B := (D - C) \geq b I > 0 \quad \text{for some } b > 0.$$

and we set

$$a_0 := |v| - |\varsigma| \geq 0 \quad a_1 := \vee_{1 \leq i \leq n} (\lambda_i - \mu_i)$$

and for any $x \in \mathbb{N}^n$

$$\|x\| := \left(\sum_{1 \leq i \leq n} x_i^2 \right)^{1/2} \geq |x| := \sum_{1 \leq i \leq n} x_i.$$

Consider the Lyapunov function

$$x \in E \mapsto V(x) = |x| \in \mathbb{N}_+.$$

Note that V is locally bounded with finite level sets and for any $x \in E - e$ we have

$$L(V)(x) = \sum_{1 \leq i \leq n} ((v_i + x_i (\lambda_i + (Cx)_i)) - (\varsigma_i + x_i (\mu_i + (Dx)_i))).$$

In this situation, we have the formula

$$L(V)(x) = a_0 + (\lambda - \mu)'x - x' B x \leq a_0 + a_1 |x| - b \|x\|^2. \quad (92)$$

On the other hand, for any $y = e_j$ we have

$$L(V)(y) = \sum_{1 \leq i \leq n} J(y, y + e_i).$$

This implies that

$$\begin{aligned}
P_t(L(V)) &= P_t(1_{E-e} L(V)) + P_t(1_e L(V)) \\
&= a_0 + a_1 P_t(V) - b P_t(V^2) \\
&\quad + \sum_{1 \leq j \leq n} P_t(1_{e_j}) (L(V)(e_j) - (a_0 + (\lambda - \mu)'e_j - e_j' B e_j))
\end{aligned}$$

from which we readily check that

$$\partial_t \mathbb{E}(V(X_t(x))) \leq a_0^+ + a_1 \mathbb{E}(V(X_t(x))) - b (\mathbb{E}(V(X_t(x))))^2$$

with

$$\begin{aligned}
a_0^+ &:= a_0 + \sum_{1 \leq j \leq d} \left(\sum_{1 \leq i \leq d} J(e_j, e_j + e_i) - (|v| - |\varsigma| + (\lambda - \mu)'e_j - e_j'(D - C)e_j) \right) \\
&= a_0 + \sum_{1 \leq j \leq d} (|\varsigma| + \mu' e_j + e_j' D e_j).
\end{aligned}$$

We conclude that $\|P_t(V)\| < \infty$. The semigroup analysis discussed above can be extended without difficulties to more general process on countable spaces models satisfying condition (92). The extension to time varying models can also be handle using a more refined analysis on time varying Riccati equations.

We also mention, that the case $|v| = 0 = |\varsigma|$ coincides with the competitive and multivariate Lotka-Volterra birth and death process discussed in Theorem 1.1 in [10].

6 Some conditional diffusions

6.1 Coupled harmonic oscillators

Consider the \mathbb{R}^n -valued diffusion (33) with $(b(x), \sigma(x)) = (Ax, \Sigma)$, for some *non necessarily stable* drift matrix A and some diffusion matrix Σ with appropriate dimensions. We associate with a given semi-definite positive $(n \times n)$ matrix $S \geq 0$ the potential function

$$U(x) := \frac{1}{2} x' S x \quad \text{and we set} \quad R = \Sigma \Sigma'. \quad (93)$$

We assume that the pairs $(A, R^{1/2})$ and $(A', S^{1/2})$ are both controllable. Let $Q_t = Q_t^{[U]}$ be the sub-Markov semigroup defined in (49) on the Euclidean space $\mathcal{E} = E = \mathbb{R}^n$. As shown in [13], the leading-triple (ρ, h, η_∞) discussed in (24) is given by

$$\begin{aligned}
\rho &= -\text{Tr}(R q_\infty)/2 = -\text{Tr}(p_\infty S)/2 \\
h(x) &= \exp(-x' q_\infty x/2) \quad \text{and} \quad \eta_\infty(dx) = \frac{\exp(-x' p_\infty^{-1} x/2)}{\sqrt{\det(2\pi p_\infty)}} dx, \quad (94)
\end{aligned}$$

with the positive fixed points p_∞ and q_∞ of the dual algebraic Riccati matrix equation

$$Ap_\infty + p_\infty A' + R - p_\infty S p_\infty = 0 \quad \text{and} \quad A' q_\infty + q_\infty A + S - q_\infty R q_\infty = 0.$$

In this context, the h -process, denoted $(X_t^h(x))_{t \geq 0}$ and defined by the stochastic differential equation

$$dX_t^h(x) = A^h X_t^h(x) dt + \Sigma dB_t \quad \text{with} \quad A^h := A - R q_\infty. \quad (95)$$

Our controllability conditions ensures that A^h is a stable matrix. Note that $X_t^h(x)$ is an \mathbb{R}^n -valued Gaussian random variable with mean $m_t^h(x)$ and covariance matrix $p_t^h \in \mathbb{R}^{n \times n}$ given for any $t > 0$ by

$$m_t^h(x) = \exp(A^h t) x \quad \text{and} \quad p_t^h = \int_0^t \exp(A^h s) R \exp((A^h)' s) ds > 0.$$

This yields the explicit formula

$$P_t^h(x, dy) = \frac{1}{\sqrt{\det(2\pi p_t^h)}} \exp\left(-\frac{1}{2}(y - m_t^h(x))'(p_t^h)^{-1}(y - m_t^h(x))\right) dy.$$

Moreover the invariant measure $\eta_\infty^h = \eta_\infty^h \mathcal{P}_t^h$ is unique and given by

$$\eta_\infty^h(dx) = \frac{1}{\sqrt{\det(2\pi p_\infty^h)}} \exp\left(-\frac{1}{2}y'(p_\infty^h)^{-1}y\right) dy$$

with the limiting covariance matrix

$$p_\infty^h := \int_0^\infty \exp(A^h s) \Sigma^2 \exp((A^h)' s) ds = (p_\infty^{-1} + q_\infty)^{-1} > 0.$$

For any time horizon $t \geq 0$ and any measurable function F on the set $\mathcal{C}([0, t], \mathbb{R}^n)$ of continuous paths from $[0, t]$ into \mathbb{R}^n we have the path space exponential change of measure Feynman-Kac formula

$$\mathbb{E}\left(F(\mathbb{X}_t(x)) \exp\left(\int_0^t \mathbb{U}_s(\mathbb{X}_s(x)) ds\right)\right) = e^{\rho t} h(x) \mathbb{E}\left(F(\mathbb{X}_t^h(x))/h(X_t^h(x))\right)$$

with the historical processes

$$\mathbb{X}_t(x) := (X_s(x))_{0 \leq s \leq t}, \quad \mathbb{X}_t^h(x) := (X_s^h(x))_{0 \leq s \leq t} \quad \text{and} \quad \mathbb{U}_s(\mathbb{X}_s(x)) := U_s(X_s(x)).$$

This yields the conjugate formulae

$$Q_t(f) = e^{\rho t} h P_t^h(f/h).$$

We denote by $(m_t(x), p_t) \in (\mathbb{R}^n \times \mathbb{R}^{n \times n})$ the mean and covariance parameters satisfying the linear evolution and the Riccati matrix differential equations

$$\begin{cases} \partial_t m_t(x) &= (A - p_t S) m_t(x) \\ \partial_t p_t &= A p_t + p_t A' + \Sigma^2 - p_t S p_t \quad \text{with} \quad (m_0(x), p_0) = (x, 0). \end{cases} \quad (96)$$

The next proposition provides an explicit description of these semigroups.

Proposition 6.1 ([13]). *For any time horizon $t > 0$ we have $p_t > 0$ and*

$$\overline{\mathcal{Q}}_t(x, dy) = \frac{1}{\sqrt{\det(2\pi p_t)}} \exp\left(-\frac{1}{2}(y - m_t(x))' p_t^{-1} (y - m_t(x))\right) dy \quad (97)$$

as well as

$$-2 \log \mathcal{Q}_t(1)(x) = x' \left(\int_0^t F'_s S F_s ds \right) x + \int_0^t \text{Tr}(S p_s) ds$$

with the fundamental matrix semigroup F_t starting at $F_0 = I$ given by

$$\partial_t F_t = (A - p_t S) F_t.$$

Observe that the normalized Markov operator $\overline{\mathcal{Q}}_t$ satisfies (39) and (40) with the parameters

$$c_t = \frac{1}{\sqrt{\det(2\pi p_t)}}, \sigma_t^2 = \lambda_{\max}(p_t) \quad \text{and} \quad \epsilon_\tau = |e^{\tau(A - p_\infty S)}| \longrightarrow 0 \text{ as } \tau \rightarrow \infty \quad (98)$$

for some matrix norm $|\cdot|$. The r.h.s. assertion is a direct consequence of the Floquet representation theorem presented in [3] (cf. (1.3) and Theorem 1.1) and the fact that $(A - p_\infty S)$ is a stable matrix. Applying Lemma 2.9 for any $v \geq 0$ and $t > 0$ there also exists some finite constant $\delta_t > 0$ such that

$$V(x) := \exp(v|x|) \implies \overline{\mathcal{Q}}_t(V)/V \leq c_t/V^{\delta_t}.$$

Using Proposition 6.1, for any $k \geq 0$ and $t \geq 0$ it is also readily checked that

$$V(x) := (1 + \|x\|)^k \implies \|\overline{\mathcal{Q}}_t(V)/V\| < \infty \quad \text{and} \quad \|\mathcal{Q}_t(V)\| < \infty.$$

6.2 Half-harmonic linear diffusions

For one dimensional models, the coupled harmonic oscillator discussed in Section 6.1 resumes to one dimensional linear diffusion

$$dX_t(x) = a X_t(x) dt + dB_t \quad \text{and the potential} \quad U(x) = \varsigma x^2/2 \quad (99)$$

for some parameters $\varsigma > 0$ and $a \in \mathbb{R}$. We set $\beta := a + \sqrt{a^2 + \varsigma}$. In this notation, the leading pair $(\rho, h) = (\rho_1, \varphi_1)$ is given by

$$\rho = -\beta/2 \quad \text{and} \quad h(x) = ((\beta - a)/\pi)^{1/4} \exp(-\beta x^2/2). \quad (100)$$

The quasi-invariant measure is therefore given by

$$\eta_\infty(dx) = \sqrt{\frac{\varsigma}{2\pi\beta}} \exp(-\varsigma x^2/(2\beta)) dx.$$

Therefore, the h -process resumes to the Ornstein-Uhlenbeck diffusion

$$dX_t^h(x) = -b X_t^h(x) dt + dB_t \quad (101)$$

with the invariant measure

$$\eta_\infty^h(dx) := \sqrt{\frac{b}{\pi}} \exp(-b x^2) dx \quad \text{with} \quad b := (\beta - a) = \sqrt{a^2 + \varsigma} > 0.$$

Note that any Ornstein-Uhlenbeck process can be seen as the h -process associated with a non absorbed (possibly transient) linear diffusion evolving in some quadratic potential well.

In this context, Proposition 6.1 is also satisfied with the mean and variance parameters

$$\begin{cases} \partial_t m_t(x) &= (a - p_t \varsigma) m_t(x) \\ \partial_t p_t &= 2ap_t + 1 - \varsigma p_t^2 \quad \text{with} \quad (m_0(x), p_0) = (x, 0). \end{cases} \quad (102)$$

We also have

$$-\frac{2}{\varsigma} \log \mathcal{Q}_t(1)(x) = \bar{p}_t + \chi_t x^2 \quad (103)$$

with

$$\chi_t := \int_0^t \exp\left(-2 \int_0^s (a - p_u \varsigma) du\right) ds \quad \text{and} \quad \bar{p}_t := \int_0^t p_s ds.$$

The half-harmonic semigroup associated with the flow $X_t(x)$ is defined for any $x \in E :=]0, \infty[$ and $f \in \mathcal{B}_b(E)$ by the formulae

$$\mathcal{Q}_t(f)(x) := \mathbb{E} \left(f(X_t(x)) 1_{T(x) > t} \exp \left\{ - \int_0^t U(X_s(x)) ds \right\} \right). \quad (104)$$

In the above display, $T(x)$ stands for the hitting time of the origin. In terms of the h -process of the flow in the harmonic potential (101) we also have the conjugate formula

$$\mathcal{Q}_t(f)(x) = e^{t\rho} e^{-\beta x^2/2} \mathbb{E} \left(f(Y_t(x)) e^{\beta Y_t(x)^2/2} 1_{T^Y(x) > t} \right) \quad (105)$$

with the parameters (ρ, β) defined in (100) and the Ornstein-Uhlenbeck diffusion flow defined by

$$dY_t(x) = -b Y_t(x) dt + dB_t \quad \text{with} \quad b := (\beta - a) > 0.$$

In the above display, $T^Y(x)$ stands for the hitting time of the origin by the flow $Y_t(x)$ starting at $x > 0$. Arguing as in Section 2.5.2 we check that

$$\begin{aligned} & \mathcal{Q}_t(x, dy) \\ &= \sinh(y m_t(x)) \exp\left(-\frac{\varsigma}{2} (\chi_t x^2 + \bar{p}_t)\right) \times \sqrt{\frac{2}{\pi p_t}} \exp\left(-\frac{y^2 + m_t(x)^2}{2p_t}\right) 1_{]0, \infty[}(y) dy \end{aligned}$$

with the parameters $(m_t(x), p_t)$ and (χ_t, \bar{p}_t) defined in (102) and (103).

Arguing as in (56), choosing the Lyapunov function $V(x) = x^n + 1/x$, for some $n \geq 1$, we readily check that

$$V \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t(V)/V \leq c_t/V \in \mathcal{C}_0(E). \quad (106)$$

6.3 Linear diffusions in some domains

Consider the one-dimensional stochastic flow $Y_t(x)$ of an Ornstein-Uhlenbeck

$$dY_t(x) = -b Y_t(x) dt + dB_t \quad \text{for some } b > 0.$$

In the above display, B_t is a one-dimensional Brownian motion starting at the origin. For a given $x \in E :=]0, \infty[$, we let $T^Y(x)$ be the hitting time of the origin by the flow $Y_t(x)$ starting at $x > 0$. Consider the semigroup

$$Q_t^Y(f)(x) := \mathbb{E}(f(Y_t(x)) 1_{T^Y(x) > t}).$$

Choosing $(a, \varsigma, \beta, \rho) = (0, b^2, b, -b/2)$ in (100), formula (105) takes the form

$$Q_t^Y(f)(x) = e^{-\rho t} H(x)^{-1} Q_t(fH)(x) \quad \text{with } H(x) = \exp(-bx^2/2)$$

with the semigroup Q_t defined in (104) with $U(x) = b^2x/2$.

For any given $n \geq 1$ we have

$$V(x) := x^n + 1/x \implies V \in \mathcal{C}_\infty(E) \quad \text{and} \quad V^H := V/H \in \mathcal{C}_\infty(E).$$

Using (106) we conclude that

$$V^H \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t^Y(V^H)/V^H = e^{-\rho t} Q_t(V)/V \leq c_t/V \in \mathcal{C}_0(E).$$

The long time behavior of the positive semigroup Q_t^Y is also studied in [46], and more recently in [55] in terms of the tangent of the h -process.

More generally, consider the \mathbb{R}^n -valued diffusion flow $X_t(x)$ defined in (33) with $(b(x), \sigma(x) = (Ax, \Sigma))$, for some matrices (A, Σ) with appropriate dimensions. Assume that $R := \Sigma\Sigma'$ is positive semi-definite and the pair of matrices $(A, R^{1/2})$ are controllable. In this situation, the Markov semigroup P_t of the stochastic flow $X_t(x)$ satisfies the sub-Gaussian estimate (81) for some parameters $(\sigma_t, m_t(x))$.

Consider a domain $E \subset \mathbb{R}^n$ with \mathcal{C}^2 -boundary with uniformly bounded interior curvature. For any given $x \in E$, let Q_t be the sub-Markov semigroup

$$Q_t(f)(x) := \mathbb{E}(f(X_t(x)) 1_{T(x) > t}) \quad \text{with } T(x) := \inf \{t \geq 0 : X_t(x) \in \partial E\}. \quad (107)$$

We clearly have $Q_t(V_\partial) \leq P_t(V_\partial)$, with the function V_∂ defined in (76). When E is non necessarily bounded but its boundary ∂E is bounded we known from (85) that $\|Q_t(V_\partial)\| < \infty$. For non necessarily bounded boundaries the sub-gaussian property (81) ensures that $\|Q_t(V_\partial)\| < \infty$.

When E is bounded, applying Lemma 4.2 (see also Proposition 4.3) we have

$$V_\partial \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t(V_\partial)/V_\partial \leq c_t/V_\partial \in \mathcal{C}_0(E).$$

For unbounded domains we need to ensure that A is stable so that (44) is satisfied for some norm $|\cdot|$ on \mathbb{R}^n . In this situation, applying Proposition 2.10 for any $t > 0$ there exists some $\delta_t > 0$ such that

$$V_\mathcal{E}(x) := \exp(v|x|) \implies Q_t(V_\mathcal{E})/V_\mathcal{E} \leq c_t/V_\mathcal{E}^{\delta_t}.$$

Applying Proposition 4.6 with $\epsilon = 0$, for any $p > 1$ we conclude that

$$V_p := V_\mathcal{E}^{1-1/p} V_\partial^{1/p} \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t(V_p)/V_p \leq c_t \Theta_{p,t} \quad (108)$$

with the function

$$\Theta_{p,t} := 1/(V_\mathcal{E}^{\delta_t(1-1/p)} V_\partial^{1/p}) \in \mathcal{C}_0(E).$$

6.4 Langevin diffusions in some domains

Consider the semigroup Q_t of the one-dimensional Langevin diffusion defined in (73) with $E =]0, \infty[$ and a quadratic confinement potential

$$W(x) = x^2/2 \implies H(x) := e^{-W(x)} = e^{-x^2/2} \quad \text{and} \quad U := \frac{1}{2} (x^2 + 1).$$

In this case, the semigroup \mathcal{Q}_t defined in (75) coincides with the semigroup of the half-harmonic oscillator discussed in Section 2.5.2. By (56) for any $n \geq 1$ we have

$$V(x) := x^n + 1/x \implies \mathcal{Q}_t(V)/V \leq c_t/V \in \mathcal{C}_0(E).$$

Notice that

$$V^H(x) := V(x)/H(x) = x^n e^{x^2/2} + \frac{e^{x^2/2}}{x}. \quad (109)$$

Using (75) we conclude that

$$V^H \in \mathcal{C}_\infty(E) \quad \text{and} \quad \mathcal{Q}_t(V^H)/V^H = \mathcal{Q}_t(V)/V \leq c_t/V \in \mathcal{C}_0(E).$$

More generally, consider the case $E =]0, \infty[$ with at least a quadratic confinement potential U , in the sense that

$$U(x) = \frac{1}{2} ((\partial W)^2 - \partial^2 W)(x) \geq U_2(x) := c + \varsigma x^2/2 \quad \text{for some } \varsigma > 0.$$

In this situation, $\mathcal{Q}_t \ll \mathcal{Q}^{[U_2]}$ is dominated by the semigroup $\mathcal{Q}^{[U_2]}$ of the half-harmonic oscillator discussed in Section 2.5.2. Arguing as in (109) we have

$$H := e^{-W} \quad V^H := V/H \in \mathcal{C}_\infty(E) \quad \text{and} \quad \mathcal{Q}_t(V^H)/V^H \leq c_t/V \in \mathcal{C}_0(E).$$

For instance, whenever the confinement potential W is chosen so that

$$W(x) \geq \epsilon_0 \log x + W_1(x) \quad \text{for some } 0 \leq \epsilon_0 < 1$$

and some function $W_1 \geq 1$ such that $W_1(x) \xrightarrow{x \rightarrow \infty} \infty$ we have

$$H = e^{-W} \implies V^H(x) := V(x)/H(x) = x^n e^{W(x)} + \frac{e^{W(x)}}{x} \geq x^n e^{W(x)} + \frac{e^{W_1(x)}}{x^{1-\epsilon_0}}.$$

Using (75) we conclude that

$$V^H \in \mathcal{C}_\infty(E) \quad \text{and} \quad \mathcal{Q}_t(V^H)/V^H = \mathcal{Q}_t(V)/V \leq c_t/V \in \mathcal{C}_0(E).$$

We illustrate the above result, with the logistic diffusion discussed in [9]. Consider the generalized Feller diffusion

$$dY_t(x) := (2a Y_t(x) - (8b/\sigma^2) Y_t(x)^2) dt + \sigma \sqrt{Y_t(x)} dB_t$$

starting at $x \in E :=]0, \infty[$. In the above display, B_t is a one dimensional Brownian motion starting at the origin and $a, b, \sigma > 0$ some parameters. Observe that

$$X_t(x) := (2/\sigma)\sqrt{Y_t(x)} \implies dX_t(x) = -\partial W(X_t(x)) dt + dB_t$$

with the potential function

$$\partial W(x) = \frac{1}{2x} - a x + b x^3 \quad \text{with } a, b > 0.$$

Thus, choosing

$$W(x) = \frac{1}{2} \log x + b \frac{x^4}{4} - a \frac{x^2}{2}$$

we readily check that

$$V^H(x) := e^{W(x)}(x^n + 1/x) = (x^{n-1/2} + 1/\sqrt{x}) e^{b\frac{x^4}{4} - a\frac{x^2}{2}} \implies V^H \in \mathcal{C}_\infty(E).$$

More generally, consider the Langevin diffusion flow

$$\mathcal{X}_t(z) = (X_t(z), Y_t(z)) \in (\mathbb{R}^n \times \mathbb{R}^n)$$

starting at $z = (x, y) \in (\mathbb{R}^n \times \mathbb{R}^n)$ and defined by the hypo-elliptic diffusion (65). We further assume that $\sup_D a < \infty$ for some bounded open connected domain $D \subset \mathbb{R}^n$ with \mathcal{C}^2 -boundary, and for any $z \in E := D \times \mathbb{R}^n$ and $f \in \mathcal{B}_b(E)$ we set

$$\mathcal{Q}_t(f)(z) := \mathbb{E}(f(\mathcal{X}_t(z)) 1_{T^{(a)}(z) > t}) \quad \text{with } T(z) := \inf\{t \geq 0 : X_t(z) \in \partial D\}.$$

We know from (66) that for any $q > 1$ we have $\mathcal{Q} \ll_q Q$ is q -dominated by the sub-Markov semigroup Q_t associated with the Ornstein-Uhlenbeck diffusion on E defined in (107), with the matrices (A, Σ) defined in (46). In terms of the functions $(V_p, \Theta_{p,t})$ defined in (108), combining (64) with (108) for any $p, q > 1$ we conclude that

$$\mathcal{Q}_t(V_{p,q})/V_{p,q} \leq c_t(p, q) \Theta_{p,q,t}$$

with the collection of Lyapunov functions

$$V_{p,q} := V_p^{1/q} \in \mathcal{C}_\infty(E) \quad \text{and the function } \Theta_{p,q,t} := \Theta_{p,t}^{1/q} \in \mathcal{C}_0(E).$$

6.5 Coupled oscillators in some domains

Consider the \mathbb{R}^n -valued diffusion $X_t(x)$ and the quadratic potential function U discussed in Section 6.1, for some $n \geq 2$ and set $E :=]0, \infty[\times \mathbb{R}^{n-1}$. Let Q_t be the semigroup defined for any $f \in \mathcal{B}_b(E)$ and $x \in E$ by the formulae

$$Q_t(f)(x) := \mathbb{E} \left(f(X_t(x)) 1_{T(x) > t} \exp \left(- \int_0^t U(X_s(x)) ds \right) \right) \quad (110)$$

with the quadratic function U in (93) and the exit time $T(x)$ given by

$$T(x) := \inf \{ t \geq 0 : X_t(x) \in \partial E \} \quad \text{with} \quad \partial E = \{0\} \times \mathbb{R}^{n-1}.$$

In terms of the h -process $Y_t(x) := X_t^h(x)$ associated with the leading pair (ρ, h) defined in (95) we also have the conjugate formula

$$Q_t(f) = e^{\rho t} h Q_t^Y(f/h) \quad \text{with} \quad Q_t^Y(f)(x) := \mathbb{E} (f(Y_t(x)) 1_{T^Y(x) > t}).$$

In the above display, $T^Y(x)$ stands for the boundary hitting time

$$T^Y(x) := \inf \{ t \geq 0 : Y_t(x) \in \partial E \}.$$

When $n = 2$, the linear diffusion $X_t(x)$ associated to the matrices $A_{1,2} = A_{2,1} = A_{2,2} = 0$ and $A_{1,2} = 1$ and $\Sigma_{1,1} = \Sigma_{1,2} = \Sigma_{2,1} = 0$ and $\Sigma_{2,2} = 1$ coincides with the integrated Wiener process model discussed in [45, 35, 48]. In a seminal article [48], McKean obtained the joint distribution of the pair $(T(x), X_{T(x)}^2)$ in the absence of soft absorption, that is when $U = 0$. To the best of our knowledge, an explicit description of the distribution of this pair and the corresponding sub-Markov semigroup is unknown in more general situations.

Observe that for any $x \in E$ and any non negative function $f \in \mathcal{B}_b(\mathbb{R}^n)$ we have

$$Q_t(f)(x) \leq \mathcal{Q}_t(f)(x) := e^{\rho t} h(x) \mathbb{E} (f(Y_t(x))/h(Y_t(x))).$$

The semigroup \mathcal{Q}_t defined above coincides with the semigroup of the coupled harmonic oscillator discussed in Section 6.1. We know from (98) that $\overline{\mathcal{Q}}_t$ satisfies the sub-Gaussian estimates (39) with

$$c_t = \frac{1}{\sqrt{\det(2\pi p_t)}} \quad \text{and} \quad \sigma_t^2 = \lambda_{\max}(p_t)$$

with the solution p_t of the Riccati-matrix equation (96). Using Proposition 6.1 for any $k \geq 1$ we have

$$V_{\mathcal{E}}(x) := 1 + \|x\|^k \implies \|\overline{\mathcal{Q}}_t(V_{\mathcal{E}})/V_{\mathcal{E}}\| < \infty \implies Q_t(V_{\mathcal{E}}) \leq c_t \mathcal{Q}_t(1)V_{\mathcal{E}}.$$

Recalling that $\mathcal{Q}_t(1)(x)$ tends to 0 exponentially fast as $\|x\| \rightarrow \infty$, this implies that

$$\forall t > 0 \quad \|Q_t(V_{\mathcal{E}})\| < \infty.$$

On the other hand for any $y = (y_1, y_{-1}) \in E =]0, \infty[\times \mathbb{R}^{n-1}$, the distance to the boundary is given by $d(y, \partial E) = y_1$. In terms of the function V_∂ defined in (76) this implies that

$$Q_t(V_\partial)(x) \leq \int Q_t(x, dy) \mathbb{1}_{]0,1[}(y_1) \chi(y_1) + \chi(1) Q_t(1)(x)$$

from which we check that $\|Q_t(V_\partial)\| < \infty$. Applying Proposition 4.4 we conclude that

$$V := V_\partial + V_\mathcal{E} \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t(V)/V \leq c_t/V \in \mathcal{C}_0(E).$$

The same analysis applies by replacing the half line E_1 by the unit interval $E_1 :=]0, 1[$. In this context, the boundary is given by the two infinite potential walls

$$\partial E = (\{0\} \times \mathbb{R}^{n-1}) \cup (\{1\} \times \mathbb{R}^{n-1}) \quad \text{and} \quad d(x, \partial E) = x_1 \wedge (1 - x_1).$$

More generally, consider a domain $E \subset \mathbb{R}^n$ with \mathcal{C}^2 -boundary with uniformly bounded interior curvature. In this situation, the sub-Gaussian property (81) ensures that $\|\overline{Q}_t(V_\partial)\| < \infty$ and therefore

$$\|Q_t(V_\partial)\| \leq \|Q_t(V_\partial)\| = \|Q_t(1)\overline{Q}_t(V_\partial)\| < \infty.$$

Applying Proposition 4.4, we conclude that

$$V := V_\partial + V_\mathcal{E} \in \mathcal{C}_\infty(E) \quad \text{and} \quad Q_t(V)/V \leq c_t/V \in \mathcal{C}_0(E).$$

7 Some hypersurface boundaries

7.1 Defining functions and charts

Consider a smooth function $y \in \mathbb{R}^{n-1} \mapsto \varphi(y) \in \mathbb{R}$ with non empty and connected level set, for some $n \geq 2$. Consider a domain E in \mathbb{R}^n with a smooth boundary $\partial E = \overline{\varphi}^{-1}(\{0\})$ defined as the null level set of the function

$$x = (x_i)_{1 \leq i \leq n} \in \mathbb{R}^n \mapsto \overline{\varphi}(x) := \varphi(x_{-n}) - x_n \quad \text{with} \quad x_{-n} := (x_i)_{1 \leq i < n} \in \mathbb{R}^{n-1}.$$

Consider the column vectors $\nabla \varphi(x_{-n}) := (\partial_{x_i} \varphi(x_{-n}))_{1 \leq i < n}$. In this notation, the unit normal vector $N(x)$ at $x \in \partial E$ is given by the column vectors

$$N(x) = \frac{\nabla \overline{\varphi}(x)}{\|\nabla \overline{\varphi}(x)\|} = \frac{1}{\sqrt{1 + \|\nabla \varphi(x_{-n})\|^2}} \begin{pmatrix} \nabla \varphi(x_{-n}) \\ -1 \end{pmatrix}.$$

Observe that the vector $N(x)$ is the outward-pointing normal direction to E as soon as $E = \overline{\varphi}^{-1}(]-\infty, 0[)$ and the inward-pointing normal direction to E when $E = \overline{\varphi}^{-1}(]0, +\infty[)$.

Consider the column vectors $e_i := (1_i(j))_{1 \leq i < n}$, with $1 \leq i < n$. In this notation, the $(n-1)$ tangential column vectors $T_i(x)$ at $x \in \partial E$ are given for any $1 \leq i < n$ by the column vectors

$$T_i(x) := \begin{pmatrix} e_i \\ \partial_{x_i} \varphi(x_{-n}) \end{pmatrix}.$$

The inner product $g(x)$ on the tangent space $\mathbb{T}_x(\partial E)$ (a.k.a. the first fundamental form on ∂E) is given by the Gramian matrix

$$g(x) = (T_i(x)' T_j(x))_{1 \leq i, j < n} = T(x) T(x)' \quad \text{with} \quad T(x)' := (T_1(x)', \dots, T_{n-1}(x)').$$

This yields the matrix formula

$$g(x) = \begin{pmatrix} I, \nabla \varphi(x_{-n}) \end{pmatrix} \begin{pmatrix} I \\ \nabla \varphi(x_{-n})' \end{pmatrix} = I + \nabla \varphi(x_{-n}) \nabla \varphi(x_{-n})'.$$

In this notation, the projection $\text{proj}_{\mathbb{T}_x(\partial E)}$ on the tangent space $\mathbb{T}_x(\partial E)$ is defined for any column vector $V = (V^i)_{1 \leq i \leq n} \in \mathbb{R}^n$ by

$$\text{proj}_{\mathbb{T}_x(\partial E)}(V) := (T_1(x)', \dots, T_{n-1}(x)') g(x)^{-1} \begin{pmatrix} T_1(x)' \\ \vdots \\ T_{n-1}(x)' \end{pmatrix} \begin{pmatrix} V^1 \\ \vdots \\ V^n \end{pmatrix}.$$

In matrix notation, the projection of m column vectors $V_i \in \mathbb{R}^n$, with $i \in \{1, \dots, m\}$ and any $m \geq 1$ takes the synthetic form

$$\begin{aligned} \text{proj}_{\mathbb{T}_x(\partial E)}(V_1, \dots, V_m) &= (T(x)' g(x)^{-1} T(x)) (V_1, \dots, V_m) \\ &= (\text{proj}_{\mathbb{T}_x(\partial E)}(V_1), \dots, \text{proj}_{\mathbb{T}_x(\partial E)}(V_m)). \end{aligned}$$

Equivalently, if $g(x)^{i,j}$ denotes the (i, j) -entry of the inverse matrix $g(x)^{-1}$, the projection of a column vector $V \in \mathbb{R}^n$ onto $\mathbb{T}_x(\partial E)$ is defined by

$$\text{proj}_{\mathbb{T}_x(\partial E)}(V) = \sum_{1 \leq i, j < n} g(x)^{i,j} (T_j(x)' V) T_i(x).$$

7.2 The shape matrix

Consider the Monge parametrization

$$\psi : \theta = (\theta_i)_{1 \leq i < n} \in \mathcal{S} := \mathbb{R}^{n-1} \mapsto \psi(\theta) = \begin{pmatrix} \theta \\ \varphi(\theta) \end{pmatrix} \in \partial E \subset \mathbb{R}^n. \quad (111)$$

In this chart, the tangent vectors and the normal unit vector at $x = \psi(\theta)$ are given for any $1 \leq i < n$ by

$$T_i^\psi(\theta) := \partial_{\theta_i} \psi(\theta) = T_i(\psi(\theta)) \in \mathbb{T}_x(\partial E) \quad \text{and} \quad N^\psi(\theta) := N(\psi(\theta)) \in \mathbb{T}_x^\perp(\partial E).$$

For any $1 \leq i, j < n$ we have

$$\begin{aligned} (\partial_{\theta_i} \psi(\theta))' N^\psi(\theta) &= 0 \\ \implies \Omega(\psi(\theta))_{i,j} &:= (\partial_{\theta_i, \theta_j} \psi(\theta))' N^\psi(\theta) = -(\partial_{\theta_i} \psi(\theta))' \partial_{\theta_j} N^\psi(\theta). \end{aligned}$$

Observe that for $x = \psi(\theta)$,

$$\partial_{\theta_i} N^\psi(\theta) = \sum_{1 \leq k \leq n} (\partial_{x_k} N)(x) \partial_{\theta_i} \psi^k(\theta) = (\nabla N(x))' \partial_{\theta_i} \psi(\theta)$$

from which we check that for any $1 \leq i, j < n$ the coefficients of the second fundamental form can be computed as follows:

$$\Omega(x)_{i,j} = -(\partial_{\theta_i} \psi(\theta))' (\nabla N(x))' \partial_{\theta_j} \psi(\theta).$$

We set

$$(\partial N^\psi(\theta))' := (\partial_{\theta_1} N^\psi(\theta), \dots, \partial_{\theta_{n-1}} N^\psi(\theta)) \in (\mathbb{T}_{\psi(\theta)}(\partial E))^{n-1}.$$

In this notation, for any $x = \psi(\theta)$ we have the matrix formulation

$$\begin{aligned} \Omega(x) &:= -\partial\psi(\theta) (\partial N^\psi(\theta))' = \left((\partial_{\theta_i, \theta_j} \psi(\theta))' N(x) \right)_{1 \leq i, j < n} \\ &= -\frac{\nabla^2 \varphi(\theta)}{\sqrt{1 + \|\nabla \varphi(\theta)\|^2}} \quad \text{with} \quad \nabla^2 \varphi(\theta) := (\partial_{\theta_i, \theta_j} \varphi(\theta))_{1 \leq i, j < n}. \end{aligned}$$

We also readily check the matrix formulation of the Weingarten's equations

$$(\partial N^\psi(\theta))' = ((\partial\psi(\theta))' g(\psi(\theta))^{-1} (\partial\psi(\theta)) (\partial N^\psi(\theta))' = -(\partial\psi(\theta))' \mathcal{W}(x).$$

In the above display, $\mathcal{W}(x)$ stands for the shape matrix (a.k.a. the Weingarten map or the mixed second fundamental form) defined by

$$\begin{aligned} \mathcal{W}(x) &:= g(x)^{-1} \Omega(x) \\ &= -\frac{1}{\sqrt{1 + \|\nabla \varphi(x_{-n})\|^2}} (I + \nabla \varphi(x_{-n}) \nabla \varphi(x_{-n})')^{-1} \nabla^2 \varphi(x_{-n}). \end{aligned}$$

We summarize the above discussion in the following proposition.

Proposition 7.1. *For any $1 \leq i < n$ we have the Weingarten's equations*

$$\partial_{\theta_i} N^\psi(\theta) = - \sum_{1 \leq k < n} \mathcal{W}(\psi(\theta))_{k,i} \partial_{\theta_k} \psi(\theta) \in \mathbb{T}_{\psi(\theta)}(\partial E).$$

Example 7.2. *For $n = 2$ we have $x \in \mathbb{R} \mapsto \bar{\varphi}(x) = \varphi(x) - x$, so that the boundary $\partial E = \bar{\varphi}^{-1}(\{0\})$ coincides with the graph of the function φ . In this context, the metric and Weingarten map at $x \in \partial E = \{x = (x_1, x_2) \in \mathbb{R}^2 : x_2 = \varphi(x_1)\}$ take the form*

$$g(x) = 1 + \|\partial\varphi(x_1)\|^2 \quad \text{and} \quad \mathcal{W}(x) = -\frac{1}{(1 + \|\partial\varphi(x_1)\|^2)^{3/2}} \partial^2 \varphi(x_1).$$

Example 7.3. For $n = 3$, the boundary ∂E is given by the surface in \mathbb{R}^3 defined

$$\partial E := \{x = (x_i)_{1 \leq i \leq 3} \in \mathbb{R}^3 : x_3 = \varphi(x_1, x_2)\}.$$

The Monge parametrization is given by

$$\psi : \theta = (\theta_1, \theta_2) \in \mathbb{R}^2 \mapsto \psi(\theta) = \begin{pmatrix} \theta_1 \\ \theta_2 \\ \varphi(\theta_1, \theta_2) \end{pmatrix} \in \partial E \subset \mathbb{R}^3.$$

In this situation, the tangent vectors at $x \in \partial E$ are given by

$$T_1(x) = \begin{pmatrix} 1 \\ 0 \\ \partial_{x_1}\varphi(x) \end{pmatrix} \quad \text{and} \quad T_2(x) = \begin{pmatrix} 0 \\ 1 \\ \partial_{x_2}\varphi(x) \end{pmatrix}.$$

In the same vein, whenever $E = \{x \in \mathbb{R}^3 : \varphi(x_1, x_2) \leq x_3\}$ the outward pointing unit normal at $x \in \partial E$ is given by

$$N(x) = \frac{1}{\sqrt{1 + (\partial_{x_1}\varphi(x))^2 + (\partial_{x_2}\varphi(x))^2}} \begin{pmatrix} \partial_{x_1}\varphi(x) \\ \partial_{x_2}\varphi(x) \\ -1 \end{pmatrix}.$$

The inner product $g(x)$ is easily computed and given by

$$g(x) = \begin{pmatrix} 1 + (\partial_{x_1}\varphi(x))^2 & (\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x)) \\ (\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x)) & 1 + (\partial_{x_2}\varphi(x))^2 \end{pmatrix}.$$

The inverse metric is given by

$$g(x)^{-1} = \frac{1}{\det(g(x))} \begin{pmatrix} 1 + (\partial_{x_2}\varphi(x))^2 & -(\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x)) \\ -(\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x)) & 1 + (\partial_{x_1}\varphi(x))^2 \end{pmatrix}$$

with

$$\det(g(x)) = 1 + (\partial_{x_1}\varphi(x))^2 + (\partial_{x_2}\varphi(x))^2 = 1 + \|\nabla\varphi(x)\|^2.$$

The second fundamental form is also given by

$$\Omega(x) = -\frac{1}{\sqrt{1 + \|\nabla\varphi(x)\|^2}} \begin{pmatrix} \partial_{x_1}^2\varphi(x) & \partial_{x_1, x_2}\varphi(x) \\ \partial_{x_1, x_2}\varphi(x) & \partial_{x_2}^2\varphi(x) \end{pmatrix}$$

and the Weingarten map is defined by

$$\mathcal{W}(x) = -\frac{1}{(1 + \|\nabla\varphi(x)\|^2)^{3/2}} \times \begin{pmatrix} (1 + (\partial_{x_2}\varphi(x))^2)\partial_{x_1}^2\varphi(x) - (\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x))\partial_{x_1, x_2}\varphi(x) & (1 + (\partial_{x_2}\varphi(x))^2)\partial_{x_1, x_2}\varphi(x) - (\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x))\partial_{x_2}^2\varphi(x) \\ -(\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x))\partial_{x_1}^2\varphi(x) + (1 + (\partial_{x_1}\varphi(x))^2)\partial_{x_1, x_2}\varphi(x) & -(\partial_{x_1}\varphi(x))(\partial_{x_2}\varphi(x))\partial_{x_1, x_2}\varphi(x) + (1 + (\partial_{x_1}\varphi(x))^2)\partial_{x_2}^2\varphi(x) \end{pmatrix}.$$

7.3 Surface and volume forms

The surface form σ_∂ on the boundary ∂E expressed in the chart ψ introduced in (111) is given by

$$(\sigma_\partial \circ \psi^{-1})(d\theta) = \sqrt{\det(g(\psi(\theta)))} d\theta$$

with the Gramian of the coordinate chart

$$\begin{aligned} g(\psi(\theta)) &:= \text{Gram}(\partial_{\theta_1}\psi(\theta), \dots, \partial_{\theta_{n-1}}\psi(\theta)) \\ &:= (\partial\psi(\theta))(\partial\psi(\theta))' = I + \nabla\varphi(\theta)(\nabla\varphi(\theta))' \end{aligned}$$

with the coordinates tangent vectors $\partial\psi(\theta) = T^\psi(\theta) := T(\psi(\theta))$. To check this claim recall that the surface area spanned by the column vectors

$$\partial\psi(\theta)' := (\partial_{\theta_1}\psi(\theta), \dots, \partial_{\theta_{n-1}}\psi(\theta))$$

is equal to the volume of the parallelepiped generated by the column vectors

$$(\partial\psi(\theta)', N(\psi(\theta))) := (\partial_{\theta_1}\psi(\theta), \dots, \partial_{\theta_{n-1}}\psi(\theta), N(\psi(\theta)))$$

which is given by the determinant of the column vectors, so that

$$(\sigma_\partial \circ \psi^{-1})(d\theta) = |\det(\partial\psi(\theta)', N(\psi(\theta)))| d\theta.$$

On the other hand, we have

$$\begin{aligned} &\begin{pmatrix} \partial\psi(\theta) \\ N(\psi(\theta))' \end{pmatrix} (\partial\psi(\theta)', N(\psi(\theta))) \\ &= \begin{pmatrix} (\partial_{\theta_1}\psi(\theta))' \\ \vdots \\ (\partial_{\theta_{n-1}}\psi(\theta))' \\ N(\psi(\theta))' \end{pmatrix} (\partial_{\theta_1}\psi(\theta), \dots, \partial_{\theta_{n-1}}\psi(\theta), N(\psi(\theta))) = \begin{pmatrix} (\partial\psi(\theta))(\partial\psi(\theta))' & 0_{n-1,1} \\ 0_{1,n-1} & 1 \end{pmatrix}. \end{aligned}$$

This implies that

$$\begin{aligned} |\det(\partial\psi(\theta)', N(\psi(\theta)))| &= \sqrt{|\det((\partial\psi(\theta)', N(\psi(\theta)))'(\partial\psi(\theta)', N(\psi(\theta))))|} \\ &= \sqrt{\det((\partial\psi(\theta))(\partial\psi(\theta))')}. \end{aligned}$$

Using the determinant perturbation formula w.r.t. rank-one matrices $\det(I + uv') = 1 + v'u$ which is valid for any column vectors $u, v \in \mathbb{R}^n$ we check that

$$\det(I + \nabla\varphi(\theta)\nabla\varphi(\theta)') = 1 + \|\nabla\varphi(\theta)\|^2.$$

This yields the formula

$$(\sigma_\partial \circ \psi^{-1})(d\theta) = \sqrt{1 + \|\nabla\varphi(\theta)\|^2} d\theta.$$

The mapping F defined in (82) can also be rewritten as a chart $\bar{\psi}$ on $\mathcal{D}_r(E)$ defined for any $(\theta, u) \in (\mathcal{S} \times [0, r])$ defined by

$$\bar{\psi}(\theta, u) := F(\psi(\theta), u) = \psi(\theta) + u N(\psi(\theta)) \in \mathcal{D}_r(E).$$

The Jacobian matrix of $\bar{\psi}$ is given by

$$\text{Jac}(\bar{\psi})(\theta, u) = (\partial_{\theta_1} \bar{\psi}(\theta, u), \dots, \partial_{\theta_{n-1}} \bar{\psi}(\theta, u), N(\psi(\theta))).$$

By Proposition 7.1 we have

$$\begin{aligned} \partial_{\theta_i} \bar{\psi}(\theta, u) &= \partial_{\theta_i} \psi(\theta) + u \partial_{\theta_i} N(\psi(\theta)) \\ &= \partial_{\theta_i} \psi(\theta) - u \sum_{1 \leq k < n} \partial_{\theta_k} \psi(\theta) \mathcal{W}(\psi(\theta))_{k,i}. \end{aligned}$$

This yields the formula

$$(\partial_{\theta_1} \bar{\psi}(\theta, u), \dots, \partial_{\theta_{n-1}} \bar{\psi}(\theta, u)) = (\partial_{\theta_1} \psi(\theta), \dots, \partial_{\theta_{n-1}} \psi(\theta)) (I - u \mathcal{W}(\psi(\theta)))$$

from which we check that

$$|\det(\text{Jac}(\bar{\psi})(\theta, u))| = \sqrt{\det(g(\psi(\theta)))} |\det(I - u \mathcal{W}(\psi(\theta)))|.$$

Note that $\bar{\psi}(\theta, 0) = \psi(\theta)$, and for any given $u < r$, the mapping $\theta \mapsto \bar{\psi}(\theta, u)$ is a chart on ∂E_u . This yields the following proposition. For the convenience of the reader, a more detailed proof of the next proposition is provided in the appendix on page 72.

Proposition 7.4. *For any $u \leq r$, the surface form $\sigma_{\partial, u}$ on the boundary ∂E_u expressed in the chart $\theta \in \mathcal{S} \mapsto \bar{\psi}(\theta, u) := F(\psi(\theta), u)$ is given by the formula*

$$(\sigma_{\partial, u} \circ \bar{\psi}(\cdot, u)^{-1})(d\theta) = |\det(I - u \mathcal{W}(\psi(\theta)))| (\sigma_{\partial} \circ \psi^{-1})(d\theta)$$

with

$$\begin{aligned} &|\det(I - u \mathcal{W}(\psi(\theta)))| \\ &= \left| \det \left(I + \frac{u}{\sqrt{1 + \|\nabla \varphi(\theta)\|^2}} (I + \nabla \varphi(\theta) \nabla \varphi(\theta)')^{-1} \nabla^2 \varphi(\theta) \right) \right|. \end{aligned}$$

In addition, the volume form $\sigma_{\mathcal{D}_r(E)}$ on $\mathcal{D}_r(E)$ expressed in the chart $\bar{\psi}$ is given by

$$\left(\sigma_{\mathcal{D}_r(E)} \circ \bar{\psi}^{-1} \right) (d(\theta, u)) = |\det(I - u \mathcal{W}(\psi(\theta)))| (\sigma_{\partial} \circ \psi^{-1})(d\theta) du.$$

Using Jacobi's formula for the derivative of determinants, we also have

$$\partial_u \log \det(I - u \mathcal{W}(x)) = -\text{Tr}((I - u \mathcal{W}(x))^{-1} \mathcal{W}(x)).$$

The level-set density function defined in (83) expressed in the chart ψ is given by the formula

$$\begin{aligned} &q_t^\partial(x, r) \\ &= \int_{\mathcal{S}} q_t(x, \psi(\theta) + r N(\psi(\theta))) |\det(I - r \mathcal{W}(\psi(\theta)))| \sqrt{\det(g(\psi(\theta)))} d\theta. \end{aligned}$$

7.4 Boundary decompositions

For some given coordinate index $k \in \{1, \dots, n\}$ and $x = (x_i)_{1 \leq i \leq n} \in \mathbb{R}^n$ we set

$$x_{-k} := (x_i)_{i \in \mathcal{I}} \quad \text{with} \quad \mathcal{I} := \{1, \dots, n\} - \{k\}$$

We further assume that

$$\partial E = \{x \in \mathbb{R}^n : x_{-k} \in \mathcal{S} \quad \text{and} \quad \varphi(x_{-k}) = x_k\} = \overline{\varphi}^{-1}(\{0\})$$

is defined as the null level set of some global defining function of the form

$$\overline{\varphi} : x \in \{(x_i)_{1 \leq i \leq n} \in \mathbb{R}^n : x_{-k} \in \mathcal{S}\} \mapsto \overline{\varphi}(x) := \varphi(x_{-k}) - x_k \in \mathbb{R}$$

for some open domain $\mathcal{S} \subset \mathbb{R}^{n-1}$.

Example 7.5 (Cylindrical boundaries). *Let $1 \leq k \leq n_1$ and $n = n_1 + n_2$ for some $n_1 > 1$ and $n_2 \geq 1$. Consider a domain \mathcal{S} of the form $\mathcal{S} = (\widehat{\mathcal{S}} \times \mathbb{R}^{n_2})$ with $\widehat{\mathcal{S}} \subset \mathbb{R}^{n_1-1}$ and assume that*

$$\forall y \in \mathbb{R}^{n_1} \text{ s.t. } y_{-k} \in \widehat{\mathcal{S}} \quad \text{and} \quad \forall z \in \mathbb{R}^{n_2} \quad \text{we have} \quad \varphi(y_{-k}, z) := \widehat{\varphi}(y_{-k}).$$

In this situation, the set ∂E is a cylindrical boundary given by the formula

$$\partial E = \partial \widehat{E} \times \mathbb{R}^{n_2} \quad \text{with} \quad \partial \widehat{E} := \left\{ y \in \mathbb{R}^{n_1} : y_{-k} \in \widehat{\mathcal{S}} \quad \text{and} \quad \widehat{\varphi}(y_{-k}) = y_k \right\}.$$

In this context, the coordinates of the outward normal by

$$N^j(x) = \frac{\epsilon}{\sqrt{1 + \|\nabla \varphi(x_{-k})\|^2}} \left(1_{\mathcal{I}}(j) \partial_{x_j} \varphi(x_{-k}) + 1_k(j) (-1) \right)$$

with the orientation parameter $\epsilon = 1$ when $E = \overline{\varphi}^{-1}(] - \infty, 0[)$; and $\epsilon = -1$ when $E = \overline{\varphi}^{-1}(]0, +\infty[)$. In the same vein, the entries $T_i^j(x)$ of the tangent vectors $T_i(x)$ indexed by $i \in \mathcal{I}$ are given for any $1 \leq j \leq n$ by

$$T_i^j(x) = 1_i(j) + 1_k(j) \partial_{x_i} \varphi(x_{-k}).$$

Consider the $(n \times (n - 1))$ -matrix

$$T(x)' := (T_1(x), \dots, T_{k-1}(x), T_{k+1}(x), \dots, T_n(x)).$$

In this notation, the inner product $g(x)$ on the tangent space $\mathbb{T}_x(\partial E)$ is given by the $((n - 1) \times (n - 1))$ -square Gramian matrix

$$g(x) = T(x)T(x)' = I + \nabla \varphi(x_{-k}) \nabla \varphi(x_{-k})'$$

with the gradient column vector

$$\nabla\varphi(x_{-k}) := (\partial_{x_i}\varphi(x_{-k}))_{i\in\mathcal{I}} = \begin{pmatrix} \partial_{x_1}\varphi(x_{-k}) \\ \vdots \\ \partial_{x_{k-1}}\varphi(x_{-k}) \\ \partial_{x_{k+1}}\varphi(x_{-k}) \\ \vdots \\ \partial_{x_n}\varphi(x_{-k}) \end{pmatrix} \in \mathbb{R}^{n-1}.$$

We check this claim using the fact that for any $i_1, i_2 \in \mathcal{I}$ we have

$$\begin{aligned} T_{i_1}(x)'T_{i_2}(x) &= \sum_{1\leq j\leq n} (1_{i_1}(j) + 1_k(j) \partial_{x_{i_1}}\varphi(x_{-k})) (1_{i_2}(j) + 1_k(j) \partial_{x_{i_2}}\varphi(x_{-k})) \\ &= 1_{i_1=i_2} + \partial_{x_{i_1}}\varphi(x_{-k}) \partial_{x_{i_2}}\varphi(x_{-k}). \end{aligned}$$

The parametrization of the hyper surface ∂E is now given by the chart function

$$\psi : \theta = (\theta_i)_{i\in\mathcal{I}} \in \mathcal{S} \mapsto \psi(\theta) \in \partial E$$

with

$$\forall 1 \leq j \leq n \quad \psi^j(\theta) := 1_{\mathcal{I}}(j) \theta_j + 1_k(j) \varphi(\theta).$$

For any $1 \leq j \leq n$ and $i_1, i_2 \in \mathcal{I}$ observe that

$$\partial_{\theta_{i_1}}\psi(\theta) = T_{i_1}^\psi(\theta) := T_{i_1}(\psi(\theta)) \quad \text{and} \quad \partial_{\theta_{i_1}, \theta_{i_2}}\psi^j(\theta) = 1_{k_i}(j) \partial_{\theta_{i_1}, \theta_{i_2}}\varphi(\theta).$$

This implies that

$$\left(\partial_{\theta_{i_1}, \theta_{i_2}}\psi(\theta) \right)' N(\psi(\theta)) = -\epsilon \frac{(\nabla^2\varphi(\theta))_{i_1, i_2}}{\sqrt{1 + \|\nabla\varphi(x_{-k})\|^2}} \quad \text{with} \quad \nabla^2\varphi(\theta) := (\partial_{\theta_{i_1}, \theta_{i_2}}\varphi(\theta))_{(i_1, i_2) \in \mathcal{I}^2}.$$

We set $(\partial\psi(\theta))' := T(\psi(\theta))'$ and $N^\psi(\theta) := N(\psi(\theta))$. In this notation, we also have

$$\begin{aligned} & (\partial N^\psi(\theta))' \\ & := (\partial_{\theta_1}N^\psi(\theta), \dots, \partial_{\theta_{k_\ell-1}}N^\psi(\theta), \partial_{\theta_{k_\ell+1}}N^\psi(\theta), \dots, \partial_{\theta_n}N^\psi(\theta)) \\ & = -\mathcal{W}(\psi(\theta)) := -g(\psi(\theta))^{-1}\Omega(\psi(\theta)) \quad \text{with} \quad \Omega(\psi(\theta)) = -\epsilon \frac{\nabla^2\varphi(\theta)}{\sqrt{1 + \|\nabla\varphi(\theta)\|^2}}. \end{aligned}$$

Example 7.6. For the cylindrical boundary discussed in Example 7.5, the inner product and the Weingarten map on the boundary $\partial\hat{E}$ are given for any $y \in \partial\hat{E}$ by the matrices

$$\hat{g}(y) = I_{(n_1-1, n_1-1)} + \nabla\hat{\varphi}(y_{-k})\nabla\hat{\varphi}(y_{-k})' \quad \text{and} \quad \hat{\mathcal{W}}(y) := \epsilon \hat{g}(y)^{-1} \frac{\nabla^2\hat{\varphi}(y_{-k})}{\sqrt{1 + \|\nabla\hat{\varphi}(y_{-k})\|^2}}$$

with the gradient column vector and the Hessian matrix given by

$$\begin{aligned}\nabla \widehat{\varphi}(y_{-k}) &:= (\partial_{y_i} \widehat{\varphi}(y_{-k}))_{i \in \widehat{\mathcal{I}}} \\ \nabla^2 \widehat{\varphi}(y_{-k}) &:= (\partial_{y_{i_1}, y_{i_2}} \widehat{\varphi}(y_{-k}))_{i_1, i_2 \in \widehat{\mathcal{I}}} \quad \text{with } \widehat{\mathcal{I}} := \{1, \dots, n_1\} - \{k\}.\end{aligned}$$

Observe that

$$\det(\widehat{g}(y)) = 1 + \|\nabla \widehat{\varphi}(y_{-k})\|^2 \quad \text{and} \quad \nabla^2 \varphi(y_{-k}, z) = \begin{pmatrix} \nabla^2 \widehat{\varphi}(y_{-k}) & 0_{(n_1-1, n_2)} \\ 0_{(n_2, n_1-1)} & 0_{(n_2, n_2)} \end{pmatrix}.$$

In this case, the inner product and the Weingarten map on the boundary ∂E are given for any point $x = (y, z) \in (\partial \widehat{E} \times \mathbb{R}^{n_2})$ by the matrices

$$g(x) = \begin{pmatrix} \widehat{g}(y) & 0_{(n_1-1, n_2)} \\ 0_{(n_2, n_1-1)} & I_{(n_2, n_2)} \end{pmatrix} \quad \text{and} \quad \mathcal{W}(x) = \begin{pmatrix} \widehat{\mathcal{W}}(y) & 0_{(n_1-1, n_2)} \\ 0_{(n_2, n_1-1)} & 0_{(n_2, n_2)} \end{pmatrix}.$$

Observe that the above matrices are bounded (w.r.t. any matrix norm) as soon as $\partial \widehat{E}$ is bounded.

More generally, assume that the boundary $\partial E \subset \cup_{\iota \in \mathcal{J}} \mathcal{O}(\iota) \subset \mathbb{R}^n$ admits a finite covering by open connected subsets $\mathcal{O}(\iota) \subset \mathbb{R}^n$ indexed by some finite set \mathcal{J} . In addition, there exists some local defining smooth functions $\overline{\varphi}_\iota$ with non vanishing gradients on $\mathcal{O}(\iota)$ such that

$$\partial E(\iota) := \partial E \cap \mathcal{O}(\iota) = \overline{\varphi}_\iota^{-1}(\{0\}) \quad \text{and} \quad E(\iota) := E \cap \mathcal{O}(\iota) = \overline{\varphi}_\iota^{-1}(]0, \infty[).$$

Up to shrinking the set $\mathcal{O}(\iota)$, by the implicit function theorem there is no loss of generality to assume that the defining functions are given by

$$\overline{\varphi}_\iota : x = (x_i)_{1 \leq i \leq n} \in \mathcal{O}(\iota) \mapsto \overline{\varphi}_\iota(x) = \varphi_\iota(x_{-k_\iota}) - x_{k_\iota}$$

for some parameter $1 \leq k_\iota \leq n$ and some smooth function φ_ι on some ball $\mathcal{S}(\iota) \subset \mathbb{R}^{n-1}$. We set $\mathcal{I}_\iota := \{1, \dots, n\} - \{k_\iota\}$. In this notation, the parametrization of the hyper surface $\partial E(\iota)$ is now given by the smooth homeomorphism

$$\psi_\iota : \theta = (\theta_i)_{i \in \mathcal{I}_\iota} \in \mathcal{S}(\iota) \mapsto \psi_\iota(\theta) \in \partial E(\iota) \quad \text{with} \quad \psi_\iota^j(\theta) := 1_{\mathcal{I}_\iota}(j) \theta_j + 1_{k_\iota}(j) \varphi_\iota(\theta). \quad (112)$$

The first and second fundamental forms on $\mathbb{T}_x(\partial E(\iota))$ as well as the Weingarten map at $x \in \partial E(\iota)$ are given by

$$\begin{aligned}g_\iota(x) &= I + \nabla \varphi_\iota(x_{-k_\iota}) \nabla \varphi_\iota(x_{-k_\iota})' \\ \Omega_\iota(x) &= -\frac{\nabla^2 \varphi_\iota(x_{-k_\iota})}{\sqrt{1 + \|\nabla \varphi_\iota(x_{-k_\iota})\|^2}} \quad \text{and} \quad \mathcal{W}_\iota(x) := g_\iota(x)^{-1} \Omega_\iota(x).\end{aligned}$$

The atlas $\mathcal{A} = (\psi_\iota, \mathcal{S}_\iota)_{\iota \in \mathcal{J}}$ represents a collection of local coordinate systems of the boundary $\partial E = \cup_{\iota \in \mathcal{J}} \partial E(\iota)$. In this situation, the surface form on ∂E and the volume form $\sigma_{\mathcal{D}_r(E)}$ on $\mathcal{D}_r(E)$ expressed in the atlas \mathcal{A} are defined by the formulae

$$\begin{aligned}\sigma_\partial^{\mathcal{A}}(d\theta) &:= \sum_{\iota \in \mathcal{J}} \pi_\iota(\psi_\iota(\theta)) \, 1_{\mathcal{S}(\iota)}(\theta) \, \sqrt{1 + \|\nabla \varphi_\iota(\theta)\|^2} \, d\theta \\ \sigma_{\mathcal{D}_r(E)}^{\mathcal{A}}(d(\theta, u)) &:= \sum_{\iota \in \mathcal{J}} \pi_\iota(\psi_\iota(\theta)) \, 1_{\mathcal{S}(\iota)}(\theta) \, |\det(I - u \mathcal{W}_\iota(\psi_\iota(\theta)))| \, \sqrt{1 + \|\nabla \varphi_\iota(\theta)\|^2} \, du.\end{aligned}$$

In the above display, $\pi_\iota : \partial E \mapsto [0, 1]$ stands for some partition of unity subordinate to the open cover of the boundary induced by the atlas.

Example 7.7. *Observe that the metric in the graph model discussed in Example 7.2 is not necessarily bounded. In this context, we can also use for any $a < a_+ < b_- < b$ a covering of the form*

$$\mathcal{O}(0) =]a, b[\times \mathbb{R} \quad \mathcal{O}(-1) =]b_-, +\infty[\times \mathbb{R} \quad \text{and} \quad \mathcal{O}(1) =]-\infty, a_+[\times \mathbb{R}.$$

For instance when $\varphi(z) = z^2$ and $(a, a_+, b_-, b) = (-2, -1, 1, 2)$ we have

$$\begin{aligned}\partial E(0) &= \{(x_1, x_2) \in]-2, 2[\times]4, \infty[: x_2 = \varphi_0(x_1)\} \\ \partial E(1) &= \{(x_1, x_2) \in]-\infty, -1[\times]1, +\infty[: x_1 = \varphi_1(x_2)\} \\ \partial E(-1) &= \{(x_1, x_2) \in]1, \infty[\times]1, +\infty[: x_1 = \varphi_{-1}(x_2)\}\end{aligned}$$

with the functions

$$\varphi_0(z) = z^2 \quad \text{and} \quad \forall \epsilon \in \{-1, 1\} \quad \varphi_\epsilon(z) = -\epsilon \sqrt{z}.$$

Whenever E is the sub-graph of φ , the parameter $\epsilon \in \{-1, 1\}$ plays the role of the orientation and the outward pointing unit normal vector at $x \in \partial E(0)$ and $y \in \partial E(\epsilon)$ are given by

$$N_0(x) = \frac{1}{\sqrt{1 + 4x_1^2}} \begin{pmatrix} 2x_1 \\ -1 \end{pmatrix} \quad \text{and} \quad N_\epsilon(y) = \frac{\epsilon}{\sqrt{1 + 1/(4y_2)}} \begin{pmatrix} -1 \\ -\epsilon/\sqrt{4y_2} \end{pmatrix}.$$

The tangent vectors at $x \in \partial E(0)$ and at $y \in \partial E(\epsilon)$ are defined by

$$T_0(x) = \begin{pmatrix} 1 \\ 2x_1 \end{pmatrix} \quad \text{and} \quad T_\epsilon(y) = \begin{pmatrix} -\epsilon/(\sqrt{4y_2}) \\ 1 \end{pmatrix}.$$

The above sub-graphs can be described with 3 charts $\{\psi_0, \psi_{+1}, \psi_{-1}\}$ defined for any $\epsilon \in \{-1, 1\}$ by

$$\psi_0 : \theta \in]-2, 2[\mapsto \psi_0(\theta) = \begin{pmatrix} \theta \\ \theta^2 \end{pmatrix} \quad \text{and} \quad \psi_\epsilon : \theta \in]1, \infty[\mapsto \psi_\epsilon(\theta) := \begin{pmatrix} -\epsilon\sqrt{\theta} \\ \theta \end{pmatrix}.$$

In this situation, the tangent vectors are given by

$$\partial_\theta \psi_0(\theta) = T_0(\psi_0(\theta)) = \begin{pmatrix} 1 \\ 2\theta \end{pmatrix} \quad \text{and} \quad \partial_\theta \psi_\epsilon(\theta) = T_\epsilon(\psi_\epsilon(\theta)) = \begin{pmatrix} -\epsilon/(\sqrt{4\theta}) \\ 1 \end{pmatrix}.$$

In this context, for any $\theta \in]-2, 2[$ we have

$$g(\psi_0(\theta)) = 1 + 4\theta^2 \quad \text{and} \quad \mathcal{W}(\psi_0(\theta)) = -2 (1 + 4\theta^2)^{-3/2}.$$

In addition, for any $\theta \in]1, \infty[$ we have

$$g(\psi_\epsilon(\theta)) = 1 + 1/(4\theta) \quad \text{and} \quad \mathcal{W}(\psi_\epsilon(\theta)) = -2\epsilon (1 + 4\theta)^{-3/2}.$$

Observe that the metric expressed in the chart $\{\psi_0, \psi_{+1}, \psi_{-1}\}$ is defined in terms of bounded functions.

Example 7.8. Consider the hyperbolic paraboloid boundary

$$\begin{aligned} \partial E &= \{(y_1, y_2, y_3) \in \mathbb{R}^3 : y_3 = y_1^2 + y_2^2\} \\ &= \partial E(0) \cup \partial E(1, 1) \cup \partial E(1, -1) \cup \partial E(2, 1) \cup \partial E(2, -1). \end{aligned}$$

In the above display, $\partial E(0)$ and $\partial E(i, \epsilon)$ with $i \in \{1, 2\}$ and $\epsilon \in \{-1, 1\}$ stands for the partition defined for any $\epsilon \in \{-1, 1\}$ by

$$\begin{aligned} \partial E(0) &:= \{y \in \mathbb{R}^3 : (y_1, y_2) \in \mathcal{S}_0 \quad y_3 = \varphi_0(y_1, y_2) := y_1^2 + y_2^2\} \\ \partial E(1, \epsilon) &:= \{y \in \mathbb{R}^3 : (y_1, y_3) \in \mathcal{S} \quad y_2 = \varphi_{1, \epsilon}(y_1, y_3) := \epsilon \sqrt{y_3 - y_1^2}\} \\ \partial E(2, \epsilon) &:= \{y \in \mathbb{R}^3 : (y_2, y_3) \in \mathcal{S} \quad y_1 = \varphi_{2, \epsilon}(y_2, y_3) := \epsilon \sqrt{y_3 - y_2^2}\} \end{aligned}$$

with the sets

$$\begin{aligned} \mathcal{S}_0 &:= \{(y_1, y_2) \in \mathbb{R}^2 : y_1^2 + y_2^2 < 2\} \\ \mathcal{S} &:= \{(y_2, y_3) \in \mathbb{R}^2 : y_3 > 1 \quad \& \quad |y_2| < \sqrt{3y_3/4}\}. \end{aligned}$$

On the truncated boundary $\partial E(0)$ we use a single chart defined by

$$\psi_0 : \theta = (\theta_1, \theta_2) \in \mathcal{S}_0 \mapsto \psi_0(\theta) = \begin{pmatrix} \theta_1 \\ \theta_2 \\ \theta_1^2 + \theta_2^2 \end{pmatrix} \in \partial E(0).$$

On $\partial E(1, \epsilon)$ we use the chart defined by

$$\psi_{1, \epsilon} : \theta = (\theta_1, \theta_3) \in \mathcal{S} \mapsto \psi_{1, \epsilon}(\theta) = \begin{pmatrix} \theta_1 \\ \epsilon \sqrt{\theta_3 - \theta_1^2} \\ \theta_3 \end{pmatrix} \in \partial E(1, \epsilon).$$

Finally, on $\partial E(2)$ we use the chart defined by

$$\psi_{2,\epsilon} : \theta = (\theta_2, \theta_3) \in \mathcal{S} \mapsto \psi_{2,\epsilon}(\theta) = \begin{pmatrix} \epsilon\sqrt{\theta_3 - \theta_2^2} \\ \theta_2 \\ \theta_3 \end{pmatrix} \in \partial E(2, \epsilon).$$

For any $\theta = (\theta_1, \theta_2) \in \mathcal{S}_0$ we have

$$\partial_{\theta_1}\psi_0(\theta) = \begin{pmatrix} 1 \\ 0 \\ 2\theta_1 \end{pmatrix} \quad \text{and} \quad \partial_{\theta_2}\psi_0(\theta) = \begin{pmatrix} 0 \\ 1 \\ 2\theta_2 \end{pmatrix}.$$

In this chart, the metric is given by

$$g(\psi_0(\theta)) = \begin{pmatrix} 1 + 4\theta_1^2 & 4\theta_1\theta_2 \\ 4\theta_1\theta_2 & 1 + 4\theta_2^2 \end{pmatrix} \quad \text{and} \quad g(\psi_0(\theta))^{-1} = \frac{1}{1 + 4(\theta_1^2 + \theta_2^2)} \begin{pmatrix} 1 + 4\theta_2^2 & -4\theta_1\theta_2 \\ -4\theta_1\theta_2 & 1 + 4\theta_1^2 \end{pmatrix}.$$

In addition, the outward pointing unit normal at $\psi_0(\theta) \in \partial E(0)$ is given by

$$N_0(\psi_0(\theta)) = \frac{1}{\sqrt{1 + 4(\theta_1^2 + \theta_2^2)}} \begin{pmatrix} 2\theta_1 \\ 2\theta_2 \\ -1 \end{pmatrix} \quad \text{and} \quad \Omega_0(\psi_0(\theta)) = \frac{1}{\sqrt{1 + 4(\theta_1^2 + \theta_2^2)}} \begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix}.$$

For any $\theta = (\theta_1, \theta_3) \in \mathcal{S}$ we have

$$\partial_{\theta_1}\psi_{1,\epsilon}(\theta) = \begin{pmatrix} 1 \\ \frac{-\epsilon\theta_1}{\sqrt{\theta_3 - \theta_1^2}} \\ 0 \end{pmatrix} \quad \text{and} \quad \partial_{\theta_3}\psi_{1,\epsilon}(\theta) = \begin{pmatrix} 0 \\ \frac{\epsilon}{2\sqrt{\theta_3 - \theta_1^2}} \\ 1 \end{pmatrix}.$$

In this chart, the metric is given by

$$g(\psi_{1,\epsilon}(\theta)) = \begin{pmatrix} 1 + \frac{\theta_1^2}{\theta_3 - \theta_1^2} & -\frac{\theta_1}{2(\theta_3 - \theta_1^2)} \\ -\frac{\theta_1}{2(\theta_3 - \theta_1^2)} & 1 + \frac{1}{4(\theta_3 - \theta_1^2)} \end{pmatrix} \quad \text{and} \quad g(\psi_{1,\epsilon}(\theta))^{-1} = \frac{1}{1 + \frac{\theta_1^2}{\theta_3 - \theta_1^2} + \frac{1}{4(\theta_3 - \theta_1^2)}} \begin{pmatrix} 1 + \frac{1}{4(\theta_3 - \theta_1^2)} & \frac{\theta_1}{2(\theta_3 - \theta_1^2)} \\ \frac{\theta_1}{2(\theta_3 - \theta_1^2)} & 1 + \frac{\theta_1^2}{\theta_3 - \theta_1^2} \end{pmatrix}.$$

In addition, the outward pointing unit normal at $\psi_{1,\epsilon}(\theta) \in \partial E(1, \epsilon)$ is given by

$$N_{1,\epsilon}(\psi_{1,\epsilon}(\theta)) = \frac{-\epsilon}{\sqrt{1 + \frac{\theta_1^2}{\theta_3 - \theta_1^2} + \frac{1}{4(\theta_3 - \theta_1^2)}}} \begin{pmatrix} \frac{-\epsilon\theta_1}{\sqrt{\theta_3 - \theta_1^2}} \\ -1 \\ \frac{\epsilon}{2\sqrt{\theta_3 - \theta_1^2}} \end{pmatrix} \quad \text{and} \quad \Omega_{1,\epsilon}(\psi_{1,\epsilon}(\theta)) = \frac{-\epsilon}{\sqrt{1 + \frac{\theta_1^2}{\theta_3 - \theta_1^2} + \frac{1}{4(\theta_3 - \theta_1^2)}}} \begin{pmatrix} \frac{-\epsilon\theta_3}{(\theta_3 - \theta_1^2)^{3/2}} & \frac{-\epsilon\theta_1}{(\theta_3 - \theta_1^2)^{3/2}} \\ \frac{-\epsilon\theta_1}{(\theta_3 - \theta_1^2)^{3/2}} & -\frac{\epsilon}{4(\theta_3 - \theta_1^2)^{3/2}} \end{pmatrix}.$$

Finally, for any $\theta = (\theta_2, \theta_3) \in \mathcal{S}$ we have

$$\partial_{\theta_2}\psi_{2,\epsilon}(\theta) = \begin{pmatrix} \frac{-\epsilon\theta_2}{\sqrt{\theta_3 - \theta_2^2}} \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad \partial_{\theta_3}\psi_{2,\epsilon}(\theta) = \begin{pmatrix} \frac{\epsilon}{2\sqrt{\theta_3 - \theta_2^2}} \\ 0 \\ 1 \end{pmatrix}.$$

In this chart, the metric and the outward pointing unit normal at $\psi_{2,\epsilon}(\theta) \in \partial E(2, \epsilon)$ are given by

$$g(\psi_{2,\epsilon}(\theta)) = \begin{pmatrix} 1 + \frac{\theta_2^2}{\theta_3 - \theta_2^2} & -\frac{\theta_2}{2(\theta_3 - \theta_2^2)} \\ -\frac{\theta_2}{2(\theta_3 - \theta_2^2)} & 1 + \frac{1}{4(\theta_3 - \theta_2^2)} \end{pmatrix} \quad \text{and} \quad N_{2,\epsilon}(\psi_{2,\epsilon}(\theta)) = -\frac{\epsilon}{\sqrt{1 + \frac{\theta_2^2}{\theta_3 - \theta_2^2} + \frac{1}{4(\theta_3 - \theta_2^2)}}} \begin{pmatrix} -1 \\ \frac{-\epsilon\theta_2}{\sqrt{\theta_3 - \theta_2^2}} \\ \frac{\epsilon}{2\sqrt{\theta_3 - \theta_2^2}} \end{pmatrix}.$$

Appendix

Proof of (9)

We have

$$P_{s,s+t}(V) \leq V + \int_s^{s+t} (-aP_{s,u}(V) + c) du = V + ct - a \int_0^t P_{s,s+u}(V) du$$

and

$$\int_0^t P_{s,s+u}(V) du = tP_{s,s+t}(V) - \int_0^t uP_{s,s+u}(L_{s+u}(V)) du \geq tP_{s,s+t}(V) - ct^2/2.$$

Combining the above estimates, we readily check that

$$P_{s,s+t}(V) \leq (1 + at)^{-1}V + ct \frac{1 + at/2}{1 + at} \leq (1 + at)^{-1}V + ct.$$

This ends the proof of (9). ■

Proof of Proposition 2.7

We have the following almost sure estimate $\|\nabla X_\tau(x)\|_2 \leq e^{-\lambda\tau}$, where $\|A\|_2$ stands for the spectral norm of a matrix A . This yields for any $x, y \in \mathbb{R}^n$ the almost sure estimate

$$\|X_\tau(x) - X_\tau(y)\| \leq e^{-\lambda\tau} \|x - y\|. \quad (113)$$

Applying the above to $y = 0$ we find that

$$P_\tau(V)(x) \leq P_\tau(V)(0) V(x)^{1-\delta} \quad \text{with} \quad \delta = 1 - e^{-\lambda\tau}.$$

Next, we check that $P_\tau^X(V)(0) < \infty$. We have

$$X_u(0) = \int_0^u (b(0) ds + \sigma dB_s) + \int_0^u \int_0^1 \nabla b(\epsilon X_s(0))' X_s(0) d\epsilon ds.$$

This implies that

$$\|X_u(0)\| \leq \beta (u + \|B_u\|) + \beta \int_0^u \|X_s(0)\| ds$$

with $\beta := \sigma \vee \|b(0)\| \vee \|\nabla b\|$. Applying Grönwall lemma we check that

$$\|X_u(0)\| \stackrel{\text{law}}{\leq} \beta (u + \|B_u\|) + \beta^2 \int_0^u (s + \|B_s\|) e^{\beta(u-s)} ds.$$

On the other hand, we have

$$\int_0^u \|B_s\| ds = u \int_0^1 \|B_{us}\| ds \stackrel{\text{law}}{=} u^{3/2} \int_0^1 \|B_s\| ds.$$

This yields the rather crude estimate

$$\|X_u(0)\| \stackrel{\text{law}}{\leq} \beta (u + u^{1/2} \|B_1\|) + \beta^2 u^2/2 + \beta^2 e^{\beta u} u^{3/2} \int_0^1 \|B_s\| ds.$$

For any $a \geq 0$ by Jensen's inequality

$$\mathbb{E} \left(e^{a \int_0^1 \|B_s\| ds} \right) \leq \int_0^1 \mathbb{E} \left(e^{a \|B_s\|} \right) ds \leq e^{a^2 r/2}.$$

It is now an elementary exercise to check that $\mathbb{E}(e^{v\|X_\tau(0)\|}) < \infty$. This ends the proof of the proposition. \blacksquare

Proof of Proposition 2.8

Consider the function

$$\begin{aligned} f_t(x) &:= \exp \left(2\epsilon \left(e^{-\alpha t} W(x) - \beta \frac{1 - e^{-\alpha t}}{\alpha} \right) \right) \\ &\implies -\partial_t \log f_t(x) = 2\epsilon e^{-\alpha t} (\alpha W + \beta). \end{aligned}$$

In the same vein, we check that

$$\begin{aligned} \partial_{x_i} f_t(x) / f_t(x) &= 2\epsilon e^{-\alpha t} \partial_{x_i} W \\ \partial_{x_i x_j} f_t(x) / f_t(x) &= 2\epsilon e^{-\alpha t} (2\epsilon e^{-\alpha t} \partial_{x_i} W \partial_{x_j} W + \partial_{x_i x_j} W). \end{aligned}$$

This implies that

$$\begin{aligned} (L(f_t) - \partial_t f_t) / f_t \\ = 2\epsilon e^{-\alpha t} ((\alpha W + \beta) + L(W) + \epsilon e^{-\alpha t} \Gamma_L(W, W)). \end{aligned}$$

Combining the above with (37) we find that

$$(L(f_t)(x) - \partial_t f_t(x)) \leq -2 \epsilon^2 e^{-\alpha t} (1 - e^{-\alpha t}) \Gamma_L(W, W) f_t(x) \leq 0.$$

This yields the interpolation formula

$$\mathbb{E}(f_0(X_t(x))) - f_t(x) = \int_0^t \mathbb{E}(\partial_s f_{t-s}(X_s(x))) ds \leq 0.$$

We check (38) after some elementary manipulations, thus there are skipped. This ends the proof of the proposition. \blacksquare

Proof of Proposition 2.11

Notice that

$$X_t^h(x) \stackrel{\text{law}}{=} \epsilon_t x + \sigma_t Z = B_{\sigma_t}(\epsilon_t x) \quad \text{with} \quad \epsilon_t := e^{-t} \quad \text{and} \quad \sigma_t := \sqrt{\frac{1 - \epsilon_t^2}{2}}$$

and some centered Gaussian random variable Z with unit variance. The conjugate formula (53) yields the integral operator equation

$$Q_t(x, dy) = e^{-t/2} e^{-x^2/2} \frac{1}{\sqrt{2\pi\sigma_t^2}} \exp\left(-\frac{(y - \epsilon_t x)^2}{2\sigma_t^2} + \frac{y^2}{2}\right) dy.$$

Observe that

$$-\frac{(y - \epsilon_t x)^2}{\sigma_t^2} + y^2 = -\frac{1}{p_t} \left(y - \frac{\epsilon_t}{1 - \sigma_t^2} x\right)^2 + x^2 \frac{\epsilon_t^2}{1 - \sigma_t^2}$$

with

$$p_t := \frac{1 - \epsilon_t^2}{1 + \epsilon_t^2} = \tanh(t) \quad \iff \quad \partial_t p_t = 1 - p_t^2 \quad \text{with} \quad p_0 = 0. \quad (114)$$

We check this claim using the fact that

$$\frac{1}{\sigma_t^2} = \frac{2}{1 - \epsilon_t^2} = 1 + \frac{1 + \epsilon_t^2}{1 - \epsilon_t^2} = 1 + \frac{1}{p_t}.$$

On the other hand, we have

$$\frac{1 - \sigma_t^2}{\epsilon_t} = \cosh(t) \quad \text{and} \quad \partial_t \log \cosh(t) = p_t = \tanh(t).$$

This implies that

$$\int_0^t p_s ds = \log \cosh(t) \quad \text{and} \quad \frac{\epsilon_t}{1 - \sigma_t^2} = \frac{1}{\cosh(t)} = \exp\left(-\int_0^t p_s ds\right).$$

We also have

$$1 - \sigma_t^2 = 1 - \frac{1 - \epsilon_t^2}{2} = \frac{1 + \epsilon_t^2}{2} \implies 1 - \frac{\epsilon_t^2}{1 - \sigma_t^2} = \frac{1 - \epsilon_t^2}{1 + \epsilon_t^2} = p_t.$$

This implies that

$$Q_t(1)(x) = e^{-t/2} \frac{\sqrt{p_t}}{\sigma_t} \exp\left(-\frac{x^2}{2} p_t\right) = e^{-t/2} h(x) P_t^h(1/h)(x).$$

Notice that

$$e^{-t/2} \frac{\sqrt{p_t}}{\sigma_t} = \sqrt{\epsilon_t \frac{1 - \epsilon_t^2}{1 + \epsilon_t^2} \frac{2}{1 - \epsilon_t^2}} = \sqrt{\frac{2}{1/\epsilon_t + \epsilon_t}} = \frac{1}{\sqrt{\cosh(t)}}$$

and

$$\partial_t m_t(x) = -p_t m_t(x) \quad \text{and} \quad \partial_t p_t = 1 - p_t^2 \quad \text{with} \quad (m_0(x), p_0) = (x, 0).$$

This ends the proof of the proposition. ■

Proof of Proposition 2.12

Notice that

$$\begin{aligned} & e^{t/2} e^{x^2/2} Q_t(x, dy) \\ &= \frac{1}{\sqrt{2\pi\sigma_t^2}} \left(\exp\left(-\frac{(y - \epsilon_t x)^2}{2\sigma_t^2} + \frac{y^2}{2}\right) - \exp\left(-\frac{(y + \epsilon_t x)^2}{2\sigma_t^2} + \frac{y^2}{2}\right) \right) 1_{[0, \infty[}(y) dy. \end{aligned}$$

This implies that

$$\begin{aligned} Q_t(1)(x) &= \frac{e^{-\frac{x^2}{2} \tanh(t)}}{\sqrt{\cosh(t)}} \\ &\quad \times \int_0^\infty \frac{1}{\sqrt{2\pi p_t}} \left(\exp\left(-\frac{(y - m_t(x))^2}{2p_t}\right) - \exp\left(-\frac{(y + m_t(x))^2}{2p_t}\right) \right) dy. \end{aligned}$$

We conclude that

$$\begin{aligned} Q_t(1)(x) &= \frac{e^{-\frac{x^2}{2} \tanh(t)}}{\sqrt{\cosh(t)}} \times \mathbb{P}(-m_t(x)/\sqrt{p_t} \leq Z \leq m_t(x)/\sqrt{p_t}) \\ &= 2 \frac{e^{-\frac{x^2}{2} \tanh(t)}}{\sqrt{\cosh(t)}} \times \mathbb{P}\left(0 \leq Z \leq \frac{x}{\sqrt{\sinh(t) \cosh(t)}}\right) \\ &\longrightarrow 0 \quad \text{as } x \rightarrow \infty \text{ or } x \rightarrow 0 \text{ or as } t \rightarrow \infty. \end{aligned}$$

In the above display, Z stands for some centered Gaussian random variable with unit variance. Note that we have used the fact that

$$m_t(x)/\sqrt{p_t} = \frac{x}{\cosh(t) \sqrt{\tanh(t)}} = \frac{x}{\sqrt{\sinh(t) \cosh(t)}}.$$

In addition, we have

$$\begin{aligned} \bar{Q}_t(x, dy) &= \frac{1}{\mathbb{P}\left(0 \leq Z \leq x/\sqrt{\sinh(t) \cosh(t)}\right)} \\ &\quad \times \frac{1}{2\sqrt{2\pi p_t}} \left(\exp\left(-\frac{(y - m_t(x))^2}{2p_t}\right) - \exp\left(-\frac{(y + m_t(x))^2}{2p_t}\right) \right) 1_{[0, \infty[}(y) dy. \end{aligned}$$

This ends the proof of the proposition. ■

Proof of (48)

The generator of the process (47) is defined by

$$L(f)(q, p) = \beta \frac{p}{m} \frac{\partial f}{\partial q} - \beta \left(\frac{\partial W}{\partial q} + \frac{\sigma^2}{2} \frac{p}{m} \right) \frac{\partial f}{\partial p} + \frac{\sigma^2}{2} \frac{\partial^2 f}{\partial p^2}.$$

Recalling that $2pq \leq p^2 + q^2$, we prove that

$$\begin{aligned} V(q, p) &\leq \frac{1}{2} \left(\frac{1}{m} + \epsilon \right) p^2 + \frac{\epsilon}{2} \left(\frac{\sigma^2}{2} + 1 \right) q^2 + W(q) \\ &\leq C^\star(\epsilon) (1 + p^2 + q^2 + W(q)) \end{aligned}$$

with

$$C^\star(\epsilon) := \max \left\{ \frac{1}{2} \left(\frac{1}{m} + \epsilon \right), \frac{\epsilon}{2} \left(\frac{\sigma^2}{2} + 1 \right), 1 \right\}.$$

On the other hand, we have

$$\begin{aligned} L(V) &= \beta \frac{p}{m} \left(\frac{\partial W}{\partial q} + \epsilon \frac{\sigma^2}{2} q + \epsilon p \right) \\ &\quad - \beta \left(\frac{\partial W}{\partial q} + \frac{\sigma^2}{2} \frac{p}{m} \right) \left(\frac{p}{m} + \epsilon q \right) + \frac{\sigma^2}{2m} \\ &= -\beta \left[\frac{1}{m} \left(\frac{\sigma^2}{2m} - \epsilon \right) p^2 + \epsilon q \frac{\partial W}{\partial q} \right] + \frac{\sigma^2}{2m}. \end{aligned}$$

Under our assumptions, this implies that for any $|q| \geq r$ we have

$$\begin{aligned} L(V) &\leq -\beta \left[\frac{1}{m} \left(\frac{\sigma^2}{2m} - \epsilon \right) p^2 + \epsilon \delta (W(q) + q^2) \right] + \frac{\sigma^2}{2m} \\ &\leq -C_\star(\epsilon, \delta) (1 + p^2 + q^2 + W(q)) + c_m(\epsilon, \delta) \end{aligned}$$

with

$$C_\star(\epsilon, \delta) := \beta \min \left\{ \left(\frac{1}{m} \left(\frac{\sigma^2}{2m} - \epsilon \right), \epsilon \delta \right) \right\} \quad \text{and} \quad c_m(\epsilon, \delta) := C_\star(\epsilon, \delta) + \frac{\sigma^2}{2m}.$$

We conclude that for any $|q| > r$,

$$\begin{aligned} (V^{-1}L(V))(q, p) &\leq -\frac{C_\star(\epsilon, \delta) (1 + p^2 + q^2 + W(q)) - c_m(\epsilon, \delta)}{V(q, p)} \\ &\leq -\frac{C_\star(\epsilon, \delta) (1 + p^2 + q^2 + W(q)) - c_m(\epsilon, \delta)}{C^\star(\epsilon) (1 + p^2 + q^2 + W(q))} \\ &= -\frac{C_\star(\epsilon, \delta)}{C^\star(\epsilon)} + \frac{c_m(\epsilon, \delta)}{C^\star(\epsilon)} \frac{1}{1 + p^2 + q^2 + W(q)} \\ &\leq -\left[\frac{C_\star(\epsilon, \delta)}{C^\star(\epsilon)} - \frac{c_m(\epsilon, \delta)}{C^\star(\epsilon)} \frac{1}{1 + p^2 + q^2} \right]. \end{aligned}$$

We choose r sufficiently large to satisfy

$$|p| > r \quad \text{or} \quad |q| > r$$

$$\Rightarrow \frac{C_\star(\epsilon, \delta)}{C^\star(\epsilon)} - \frac{c_m(\epsilon, \delta)}{C^\star(\epsilon)} \frac{1}{p^2 + q^2} \geq \frac{C_\star(\epsilon, \delta)}{C^\star(\epsilon)} - \frac{c_m(\epsilon, \delta)}{C^\star(\epsilon)} \frac{1}{r^2} \geq a := \frac{C_\star(\epsilon, \delta)}{2C^\star(\epsilon)} > 0,$$

and we set

$$K_r := \{(q, p) \in \mathbb{R}^2 : |p| \vee |q| \leq r\}.$$

In this notation, we have

$$L(V) \leq -aV \mathbf{1}_{E-K_r} + \sup_{K_r} L(V) \leq -aV + c \quad \text{with} \quad c := \sup_{K_r} L(V) + a \sup_{K_r} V.$$

Proof of (56)

Observe that for any $0 < y \leq 1$ and $z \in E =]0, \infty[$ we have

$$\sinh(yz) \leq y \sinh(z) \quad \text{and} \quad \sinh(z) \leq \frac{1}{2} e^z.$$

This implies that

$$\int_0^\infty \mathcal{Q}_t(x, dy) \frac{1}{y} \leq \sinh(m_t(x)) \frac{e^{-\frac{x^2}{2}(p_t + e^{-2t}/p_t)}}{\sqrt{\cosh(t)}} \times \int_0^\infty \sqrt{\frac{2}{\pi p_t}} \exp\left(-\frac{y^2}{2p_t}\right) dy.$$

from which we check that

$$\int_0^\infty \mathcal{Q}_t(x, dy) \frac{1}{y} \mathbf{1}_{]0,1]}(y) \leq \frac{\exp\left(-\left(\frac{x^2}{2}(p_t + \frac{e^{-2t}}{p_t}) - e^{-t}x\right)\right)}{2\sqrt{\cosh(t)}}.$$

On the other hand, for any $n \geq 1$ we have

$$\begin{aligned} & \int_0^\infty \mathcal{Q}_t(x, dy) y^n \\ & \leq \frac{1}{2} \frac{1}{\sqrt{\cosh(t)}} \sqrt{\frac{2}{\pi p_t}} \exp\left(-\frac{\epsilon_t^2 x^2}{2p_t} - \frac{x^2}{2} p_t\right) \int_0^\infty y^n \exp\left(y\epsilon_t x - \frac{y^2}{2p_t}\right) dy. \end{aligned}$$

Notice that

$$y\epsilon_t x - \frac{y^2}{2p_t} = -\frac{1}{2p_t} (y - \epsilon_t x p_t)^2 + \frac{x^2}{2} \epsilon_t^2 p_t$$

so that

$$\begin{aligned} & \int_0^\infty \mathcal{Q}_t(x, dy) y^n \leq \frac{1}{2} \frac{1}{\sqrt{\cosh(t)}} \sqrt{\frac{2}{\pi p_t}} \\ & \times \exp\left(-\frac{x^2}{2} \left((1 - \epsilon_t^2) p_t + \frac{\epsilon_t^2}{p_t}\right)\right) \int_0^\infty y^n \exp\left(-\frac{1}{2p_t} (y - \epsilon_t x p_t)^2\right) dy. \end{aligned}$$

For any $n \geq 1$, we conclude that

$$V(x) := x^n + 1/x \implies V \in \mathcal{C}_\infty(E) \quad \text{and} \quad \|Q_t(V)\| < \infty.$$

This ends the proof of (56). ■

Proof of Lemma 3.2

To simplify notation, we write Q_t instead of $Q_t^{[U]}$. For any $V \in \mathcal{B}_\infty(E) \cap \mathcal{D}(L)$ we have

$$\begin{aligned} Q_t(V) &= V + \int_0^t Q_s(L(V) - UV) ds \\ &\leq V + \int_0^t [-a Q_s(V) + c Q_s(1)] ds = V + c \int_0^t Q_s(1) ds - a \int_0^t Q_s(V) ds. \end{aligned}$$

On the other hand, through integration by parts we have

$$\begin{aligned} \int_0^t Q_s(V) ds &= [s Q_s(V)]_0^t - \int_0^t s \frac{d}{ds} Q_s(V) ds \\ &= t Q_t(V) - \int_0^t s \underbrace{Q_s(L(V) - UV)}_{\leq c} ds \geq t Q_t(V) - c \int_0^t s Q_s(1) ds. \end{aligned}$$

This implies that

$$Q_t(V) \leq V + c \int_0^t Q_s(1) ds - a \left(t Q_t(V) - c \int_0^t s Q_s(1) ds \right)$$

from which we conclude that

$$Q_t(V) \leq \frac{V}{1+at} + c \int_0^t Q_s(1) ds \implies Q_t(V) \leq \frac{V}{1+at} + ct.$$

This ends the proof of (69). Now, we come to the proof of (70). We have the forward evolution equation given for any $f \in \mathcal{D}(L)$ by

$$\partial_t Q_t(f) = Q_t(L^U(f)).$$

Applying the above to $f = U$ we readily check that

$$\partial_t Q_t(U) \leq a_0 + a_1 Q_t(U) - Q_t(U)^2 \leq a_0 + a_1 Q_t(U) - (Q_t(U))^2 / Q_t(1)$$

from which we find the Riccati estimates

$$\partial_t Q_t(U) \leq a_0 + a_1 Q_t(U) - (Q_t(U))^2 \implies \forall t > 0 \quad \|Q_t(U)\| < \infty.$$

This ends the proof of the lemma. ■

Proof of (66)

By Girsanov theorem we have

$$\mathcal{Q}_t^{(a)}(f)(z) = \mathbb{E} \left(f(\mathcal{X}_t^0(z)) Z_t(z) 1_{T^0(z) > t} \right)$$

with the exponential martingale

$$Z_t(z) = \exp \left(\frac{1}{\sigma} \int_0^t a(X_s(z))' dB_u - \frac{1}{2\sigma^2} \int_0^t \|a(X_s(z))\|^2 du \right).$$

By Hölder's inequality, for any non negative function f on E , any $z \in E$ and any conjugate parameters $p, q > 1$ with $1/p + 1/q = 1$ we have

$$\mathcal{Q}_t^{(a)}(f)(z) \leq \mathbb{E} \left(Z_t(z)^q 1_{T^0(z) > t} \right)^{1/q} \mathcal{Q}_t^{(a)}(f^p)(z)^{1/p}.$$

On the other hand, we have

$$\begin{aligned} \mathbb{E} \left(Z_t(z)^q 1_{T^0(z) > t} \right) &= \mathbb{E} \left(\bar{Z}_t(z) \exp \left(\frac{q(q-1)}{2\sigma^2} \int_0^t \|a(X_s(z))\|^2 du \right) 1_{T^0(z) > t} \right) \\ &\leq c_t(p) := \exp \left(\frac{pt}{2((p-1)\sigma)^2} \sup_D a \right) \end{aligned}$$

with the exponential martingale

$$\bar{Z}_t(z) = \exp \left(\frac{q}{\sigma} \int_0^t a(X_s(z))' dB_u - \frac{q^2}{2\sigma^2} \int_0^t \|a(X_s(z))\|^2 du \right).$$

This ends the proof of (66). ■

Proof of Lemma 4.2

For any $z \in \partial E$ there exists some open ball $\mathbb{B}(z, r) \subset \mathbb{R}^n$ with $r > 0$ and some \mathcal{C}^1 -mapping g from \mathbb{R}^{n-1} into \mathbb{R} such that

$$\begin{aligned} E \cap \mathbb{B}(z, r) &= \{x \in \mathbb{B}(z, r) : x_n < g(x_{-n})\} \\ \partial E \cap \mathbb{B}(z, r) &= \{x \in \mathbb{B}(z, r) : x_n = g(x_{-n})\} \quad \text{with } x_{-n} := (x_1, \dots, x_{n-1}). \end{aligned}$$

We make the change of variables

$$\mathcal{E}(z, r) := E \cap \mathbb{B}(z, r)$$

$$\mapsto \varsigma(x) := (x_{-n}, x_n - g(x_{-n})) \in \mathcal{O}(z, r) := \varsigma(\mathcal{E}(z, r)) \subset (\mathbb{R}^{n-1} \times \mathbb{R}_+)$$

with Jacobian

$$\nabla \varsigma(x) = \begin{pmatrix} I_{(n-1) \times (n-1)} & -\nabla g(x_{-n}) \\ 0 & 1 \end{pmatrix}.$$

Observe that

$$\begin{aligned} \varsigma &: x \in \mathcal{E}_0(z, r) := (\partial E \cap \mathbb{B}(z, r)) \\ \implies \varsigma(x) &= (x_{-n}, 0) \in \mathcal{O}_0(z, r) := \varsigma(\mathcal{E}_0(z, r)) \subset (\mathbb{R}^{n-1} \times \{0\}). \end{aligned}$$

The inverse is given by

$$\begin{aligned} y \in \mathcal{O}(z, r) &\mapsto \varsigma^{-1}(y) = (y_{-n}, y_n + g(y_{-n})) \in \mathcal{E}(z, r) \\ \implies \nabla \varsigma^{-1}(y) &= \begin{pmatrix} I_{(n-1) \times (n-1)} & \nabla g(y_{-n}) \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

On the other hand we have

$$\begin{aligned} \|\varsigma(x) - \varsigma(\bar{x})\| &= (\|x_{-n} - \bar{x}_{-n}\|^2 + (|x_n - \bar{x}_n| + |g(x_{-n}) - g(\bar{x}_{-n})|)^2)^{1/2} \\ &\leq (\|x_{-n} - \bar{x}_{-n}\|^2 + 2|x_n - \bar{x}_n|^2 + 2\|\nabla g\|^2 \|x_{-n} - \bar{x}_{-n}\|^2)^{1/2} \\ &\leq c(g) \|x - \bar{x}\| \quad \text{with } c(g) := \sqrt{2 \vee (1 + 2\|\nabla g\|^2)} \geq 1. \end{aligned}$$

In the same vein, we have

$$\|\varsigma^{-1}(y) - \varsigma^{-1}(\bar{y})\| \leq c(g) \|y - \bar{y}\| \quad \text{so that} \quad \frac{1}{c(g)} \|y - \bar{y}\| \leq \|\varsigma^{-1}(y) - \varsigma^{-1}(\bar{y})\|.$$

For any $x \in \mathcal{E}(z, r)$ and $\bar{x} \in \mathcal{E}_0(z, r)$ we have $\varsigma(\bar{x}) \in \mathcal{O}_0(z, r)$ and

$$\|x - \bar{x}\| = \|\varsigma^{-1}(\varsigma(x)) - \varsigma^{-1}(\varsigma(\bar{x}))\| \geq \frac{1}{c(g)} \|\varsigma(x) - \varsigma(\bar{x})\| \geq \frac{1}{c(g)} |\varsigma(x)_n|.$$

Taking the infimum of all $\bar{x} \in \mathcal{E}_0(z, r)$ this implies that

$$d(x, \mathcal{E}_0(z, r)) \geq \frac{1}{c(g)} |\varsigma(x)_n| \quad \text{and} \quad d(\varsigma^{-1}(y), \mathcal{E}_0(z, r)) \geq \frac{1}{c(g)} |y_n|$$

for any $x \in \mathcal{E}(z, r)$ and $y \in \mathcal{O}(z, r)$. We conclude that

$$\begin{aligned} &\int_{\mathcal{E}(z, r)} \chi(d(x, \mathcal{E}_0(z, r))) \, dx \\ &= \int_{\mathcal{O}(z, r)} \chi(d(\varsigma^{-1}(y), \mathcal{E}_0(z, r))) \, |\det(\varsigma^{-1}(y))| \, dy \\ &\leq \frac{1}{c(g)} \sup_{y \in \mathcal{O}(z, r)} |\det(\varsigma^{-1}(y))| \int_{\mathcal{O}(z, r)} \chi(y_n) \, dy < \infty. \end{aligned}$$

We end the proof of the lemma by covering ∂E by finitely many boundary coordinates patches $(\mathcal{E}(z_i, r_i), g_i)_{1 \leq i \leq n}$, for some $z_i \in \partial E$, $r_i > 0$ and some local defining functions g_i . ■

Proof of Lemma 4.9

Using the change of variable formulae

$$\int_{\partial E_r} f(z) \sigma_{\partial,r}(dz) = \int_{\partial E} f(z + rN(z)) |\det(I - r\mathcal{W}(z))| \sigma_{\partial}(dz)$$

and

$$\int_{\partial E} f(z) \sigma_{\partial}(dz) = \int_{\partial E_r} f(z - rN(z)) |\det(I + r\mathcal{W}(z))| \sigma_{\partial,r}(dz)$$

we check that

$$\int_{\partial E_r} f(z) \sigma_{\partial,r}(dz) \leq \kappa_{\partial}(\alpha) \int_{\partial E} f(z + rN(z)) \sigma_{\partial}(dz)$$

and

$$\int_{\partial E} f(z) \sigma_{\partial}(dz) \leq \kappa_{\partial}^{-}(\alpha) \int_{\partial E_r} f(z - rN(z)) \sigma_{\partial,r}(dz).$$

This yields the estimate

$$\int_{\partial E_r} f(z) \sigma_{\partial,r}(dz) \leq \iota(\alpha) \kappa_{\partial}(\alpha) \int_{\partial E} g(z) \sigma_{\partial}(dz).$$

In the same vein, we have

$$\int_{\partial E} f(z) \sigma_{\partial}(dz) \leq \iota(\alpha) \kappa_{\partial}^{-}(\alpha) \int_{\partial E_r} g(z) \sigma_{\partial,r}(dz).$$

Integrating w.r.t. the parameter $r \in [0, \alpha]$ we check the co-area estimate

$$\begin{aligned} \alpha \int_{\partial E} f(z) \sigma_{\partial}(dz) &\leq \iota(\alpha) \kappa_{\partial}^{-}(\alpha) \int_0^{\alpha} dr \int_{\partial E_r} g(z) \sigma_{\partial,r}(dz) \\ &= \iota(\alpha) \kappa_{\partial}^{-}(\alpha) \int_{\mathcal{D}_{\alpha}(E)} g(z) dz. \end{aligned}$$

This ends the proof of the lemma. ■

Proof of Proposition 7.4

For any given $\bar{\theta} := (\bar{\theta}_1, \dots, \bar{\theta}_n) \in (\mathbb{R}^{n-1} \times [0, r])$ we set $\bar{\theta}_{-n} := (\theta_1, \dots, \theta_{n-1})$. In this notation, we have

$$\begin{aligned} \bar{\psi} : \bar{\theta} \in (\mathbb{R}^{n-1} \times [0, r]) &\mapsto \bar{\psi}(\bar{\theta}) := F(\psi(\bar{\theta}_{-n}), \bar{\theta}_n) \\ &= \psi(\bar{\theta}_{-n}) + \bar{\theta}_n N(\psi(\bar{\theta}_{-n})) \in \mathcal{D}_r(E). \end{aligned}$$

The volume form $\sigma_{\mathcal{D}_r(E)}$ on $\mathcal{D}_r(E)$ expressed in the chart $\bar{\psi}$ is given by

$$\left(\sigma_{\mathcal{D}_r(E)} \circ \bar{\psi}^{-1}\right)(d\bar{\theta}) = |\det(\text{Jac}(\bar{\psi})(\theta))| = \sqrt{\det\left((\partial\bar{\psi}(\bar{\theta}))(\partial\bar{\psi}(\bar{\theta}))'\right)} d\bar{\theta}.$$

Arguing as above, we have

$$(\partial\bar{\psi}(\bar{\theta}))' = \left(\left(\partial_{\bar{\theta}_{-n}}\bar{\psi}(\bar{\theta})\right)', \partial_{\bar{\theta}_n}\bar{\psi}(\bar{\theta})\right) \in \left(\mathbb{T}_{\bar{\psi}(\bar{\theta})}(\mathcal{D}_r(E))\right)^n$$

with the tangent vectors

$$\left(\partial_{\bar{\theta}_{-n}}\bar{\psi}(\bar{\theta})\right)' := \left(\partial_{\bar{\theta}_1}\bar{\psi}(\bar{\theta}), \dots, \partial_{\bar{\theta}_{n-1}}\bar{\psi}(\bar{\theta})\right) \quad \text{and} \quad \partial_{\bar{\theta}_n}\bar{\psi}(\bar{\theta}) = N(\psi(\bar{\theta}_{-n})).$$

In addition, we have

$$\left(\partial_{\bar{\theta}_{-n}}\bar{\psi}(\bar{\theta})\right)' = \left(\partial\psi(\bar{\theta}_{-n})\right)' + \bar{\theta}_n \left(\partial(N(\psi(\bar{\theta}_{-n})))\right)' = \left(\partial\psi(\bar{\theta}_{-n})\right)' (I - \bar{\theta}_n \mathcal{W}(\psi(\bar{\theta}_{-n}))).$$

This yields the formula

$$\left(\partial_{\bar{\theta}_{-n}}\bar{\psi}(\bar{\theta})\right) \left(\partial_{\bar{\theta}_{-n}}\bar{\psi}(\bar{\theta})\right)' = g(\psi(\bar{\theta}_{-n})) (I - \bar{\theta}_n \mathcal{W}(\psi(\bar{\theta}_{-n})))^2$$

from which we check that

$$\left(\partial\bar{\psi}(\bar{\theta})\right) \left(\partial\bar{\psi}(\bar{\theta})\right)' = \begin{pmatrix} g(\psi(\bar{\theta}_{-n})) (I - \bar{\theta}_n \mathcal{W}(\psi(\bar{\theta}_{-n})))^2 & 0_{n-1} \\ 0 & 1 \end{pmatrix}.$$

We conclude that

$$\sqrt{\det\left((\partial\bar{\psi}(\bar{\theta}))(\partial\bar{\psi}(\bar{\theta}))'\right)} = \sqrt{\det(g(\psi(\bar{\theta}_{-n})) |\det(I - \bar{\theta}_n \mathcal{W}(\psi(\bar{\theta}_{-n})))|)}$$

and therefore

$$\left(\sigma_{\mathcal{D}_r(E)} \circ \bar{\psi}^{-1}\right)(d\bar{\theta}) = |\det(I - \bar{\theta}_n \mathcal{W}(\psi(\bar{\theta}_{-n})))| d\bar{\theta}_n (\sigma_{\partial,0} \circ \psi^{-1})(d\bar{\theta}_{-n}).$$

For any given $\bar{\theta}_n = u \in [0, r]$, the volume form $\sigma_{\partial,u}$ on the boundary ∂E_u expressed in the boundary chart

$$\bar{\psi}(\cdot, u) : \theta \in \mathbb{R}^{n-1} \mapsto \bar{\psi}(\theta, u) = F(\psi(\theta), u) \in \partial E_u$$

is given by

$$\left(\sigma_{\partial,u} \circ \bar{\psi}(\cdot, u)^{-1}\right)(d\theta) = |\det(I - u \mathcal{W}(\psi(\theta)))| (\sigma_{\partial,0} \circ \psi^{-1})(d\theta).$$

This ends the proof of the proposition. ■

References

- [1] A.R. Aliabad, F. Azarpanah and M. Namdari. Rings of continuous functions vanishing at infinity. *Commentationes Mathematicae Universitatis Carolinae*, vol. 45, no. 3, pp. 519-533 (2004).
- [2] D. G. Aronson. Bounds for the fundamental solution of a parabolic equation. *Bulletin of American Math. Soc.*, vol. 73, pp. 890–896 (1967).
- [3] A.N. Bishop and P. Del Moral. An explicit Floquet-type representation of Riccati aperiodic exponential semigroups. *International Journal of Control*, vol. 94, no. 1, pp. 258–266 (2021).
- [4] A.N. Bishop, P. Del Moral, K. Kamatani and B. Remillard. On one-dimensional Riccati diffusions. *Annals of Applied Probability*. volume 29, no. 2, pp. 1127–1187 (2019).
- [5] A.N. Bishop and P. Del Moral. On the stability of matrix-valued Riccati diffusions. *Electronic Journal of Probability*, vol. 24, pp. 1–40 (2019).
- [6] K. Bogdan, K. Szczypkowski. Gaussian estimates for Schrödinger perturbations. arXiv preprint arXiv:1301.4627 (2013).
- [7] P. Brémaud. Lyapunov Functions and Martingales. *Markov Chains. Texts in Applied Mathematics*, book series vol.31, pp. 167–193 (1999).
- [8] P. Cattiaux. Stochastic calculus and degenerate boundary value problems. *Ann. Institut Fourier*, vol. 42, no. 3, pp. 541–624 (1992).
- [9] P. Cattiaux, P. Collet, A. Lambert, S. Martínez, S. Méléard and J. San Martín. Quasi-stationary distributions and diffusion models in population dynamics. *The Annals of Probability*, vol. 37, no. 5, pp. 1926–1969 (2009).
- [10] N. Champagnat and D. Villemonais. Lyapunov criteria for uniform convergence of conditional distributions of absorbed Markov processes. *Stochastic Processes and their Applications*, (Arxiv) vol. 135, pp 51-74 (2021).
- [11] P. Del Moral and A. Guionnet. On the stability of interacting processes with applications to filtering and genetic algorithms. *Publication du Laboratoire de Statistique et Probabilités 3-98*, Université Paul Sabatier, Toulouse (1998). *Annales de l’Institut Henri Poincaré Probab. & Statist.* vol. 37, no. 2, pp. 155–194 (2001).
- [12] P. Del Moral and A. Guionnet. On the stability of measure valued processes with applications to filtering. *Comptes Rendus de l’Académie des Sciences-Series I-Mathematics*, vol. 329, no. 5, pp. 429–434 (1999).

- [13] P. Del Moral and E. Horton. Quantum harmonic oscillators and Feynman-Kac path integrals for linear diffusive particles. arXiv preprint arXiv:2106.14592 (2021).
- [14] P. Del Moral and L. Miclo. Branching and interacting particle systems approximations of Feynman-Kac formulae with applications to non-linear filtering. In *Séminaire de Probabilités, XXXIV*, volume 1729, pp. 1–145 (2000).
- [15] P. Del Moral and L. Miclo. On the stability of nonlinear Feynman-Kac semigroups. *Annales de la Faculté des Sciences de Toulouse: Mathématiques, serie 6*, t. 11, no.2, pp. 135–175 (2002).
- [16] P. Del Moral, M. Ledoux and L. Miclo. On contraction properties of Markov kernels. *Probability Theory and Related Fields*, vol. 126, no. 3, pp. 395–420 (2003).
- [17] P. Del Moral. *Feynman-Kac Formulae*. Springer, New York (2004).
- [18] P. Del Moral and A. Doucet. Particle motions in absorbing medium with hard and soft obstacles. *Stochastic Analysis and Applications*. vol. 22, no. 5, pp. 1175–1207 (2004).
- [19] P. Del Moral. *Mean Field Simulation for Monte Carlo Integration*. CRC press (2013).
- [20] P. Del Moral and A. Jasra. A Note on Random Walks with Absorbing barriers and Sequential Monte Carlo Methods. *Stochastic Analysis and Applications*, vol. 36, no. 3, pp. 413–442 (2018).
- [21] P. Del Moral, E. Horton and A. Jasra. On the Stability of Positive Semigroups. arXiv preprint arXiv:2112.03751 (2021), to appear in the *Annals of Applied Probability* (2023).
- [22] P. Del Moral and S. Penev. *Stochastic Processes: From Applications to Theory*. CRC-Press (2017).
- [23] P. Del Moral and S.S. Singh. Backward Itô-Ventzell and stochastic interpolation formulae. arXiv preprint arXiv:1906.09145 (2019).
- [24] P. Diaconis and D. Freedman. On the consistency of Bayes estimates (with discussion). *The Annals of Statistics*, vol. 14, no. 1, pp. 1–67, (1986).
- [25] P. Diaconis and L. Miclo. On quantitative convergence to quasi-stationarity. *Annales de la Faculté des Sciences de Toulouse: Mathématiques*, vol. 24, no. 4, pp. 973–1016 (2015).

- [26] R. Douc, E. Moulines and J.S. Rosenthal. Quantitative bounds on convergence of time-inhomogeneous Markov chains. *Ann. Appl. Probab.* 14, pp. 1643–1665 (2004).
- [27] R. Douc, E. Moulines and Y. Ritov. Forgetting of the initial condition for the filter in general state-space hidden Markov chain: a coupling approach. *Electronic Journal of Probability*, vol. 14, pp. 27–49 (2009).
- [28] Y. Du. Order structure and topological methods in nonlinear partial differential equations: Vol. 1: Maximum principles and applications (Vol. 2). World Scientific (2006).
- [29] G. Ferré. Large Deviations Theory in Statistical Physics: Some Theoretical and Numerical Aspects. Diss. Université Marne La Vallée (2019).
- [30] G. Ferré, M. Rousset, and G. Stoltz. Large deviations of empirical measures of diffusions in weighted topologies. *Electronic Journal of Probability*, vol. 25, pp. 1–52 (2020).
- [31] G. Ferré, M. Rousset and G. Stoltz. More on the long time stability of Feynman-Kac semigroups. *Stochastics and Partial Differential Equations: Analysis and Computations*, vol. 9, no. 3, pp. 630–673 (2021).
- [32] F.G. Foster. On the Stochastic Matrices Associated with Certain Queuing Processes. *The Annals of Mathematical Statistics*, vol. 24, no. 3; pp. 355–360 (1953).
- [33] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Springer reprint of the 1998 edition, vol 224. Springer-Verlag Berlin Heidelberg New York (2001).
- [34] F. Le Gland and L. Mevel. Stability and Approximation of nonlinear filters: an information theoretic approach. *Proceedings of the 38th Conference on Decision and Control*, Phoenix Arizona (1999).
- [35] M. Goldman. On the first passage of the integrated Wiener process. *Annals of Mathematical Statistics*, vol.42, no. 6, pp 2150–2155 (1971).
- [36] A. Guillin, B. Nectoux and L. Wu. Quasi-stationary distribution for strongly Feller Markov processes by Lyapunov functions and applications to hypoelliptic Hamiltonian. 2020 (<https://hal.archives-ouvertes.fr/hal-03068461>).
- [37] M. Hairer and J. C. Mattingly. Yet another look at Harris’ ergodic theorem for Markov chains. *Seminar on Stochastic Analysis, Random Fields and Applications VI*, 109–117, *Progr. Probab.*, 63, Birkhäuser/Springer Basel AG, Basel, 2011.

- [38] T. E. Harris. The existence of stationary measures for certain Markov processes. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954-1955*, vol. II, pp. 113–124. University of California Press, Berkeley and Los Angeles (1956).
- [39] R. Z. Hasminskii. *Stochastic Stability of Differential Equations*, vol. 7 of *Monographs and Textbooks on Mechanics of Solids and Fluids: Mechanics and Analysis*. Sijthoff & Noordhoff, Alphen aan den Rijn. Translated from the Russian by D. Louvish (1980).
- [40] M. Hürzeler. *Statistical Methods for General State-Space Models*. Ph.D. Thesis, Department of Mathematics, ETH Zurich, Zürich (1998).
- [41] M.L. Kleptsyna and A. Yu Veretennikov. On continuous time ergodic filters with wrong initial data. *Theory of Probability & Its Applications* vol. 53, no. 2, pp. 269–300 (2009).
- [42] M.L. Kleptsyna and A. Yu Veretennikov. On discrete time ergodic filters with wrong initial data. *Probab. Theory Related Fields*, vol. 141, pp. 411–444 (2008).
- [43] V. Knopova. On the Feynman-Kac semigroup for some Markov processes. *Modern Stochastics: Theory and Applications*, vol. 2, n.2 pp. 107–129 (2015).
- [44] V. Konakov, S. Menozzi, S. Molchanov. Explicit parametrix and local limit theorems for some degenerate diffusion processes. *Annales de l’IHP Probabilités et statistiques*. Vol. 46. No. 4., pp. 908–923 (2010).
- [45] A. Lachal. Sur le premier instant de passage de l’intégrale du mouvement brownien *Annales de l’I.H.P., Section B*, vol. 27, no. 3, pp. 385–405 (1991).
- [46] M. Lladser and J. San Martìn. Domain of attraction of the quasi-stationary distributions for the Ornstein-Uhlenbeck process. *Journal of Applied Probability*, vol. 37, no. 2, pp. 511–520 (2000).
- [47] T. Lelièvre, M. Ramil and J. Reygner. A probabilistic study of the kinetic Fokker-Planck equation in cylindrical domains. *arXiv preprint arXiv:2010.10157* (2020).
- [48] H. P. Jr. McKean. A winding problem for a resonator driven by a white noise. *J. Math. Kyoto Univ.* vol. 2, pp. 227–235 (1963).
- [49] S. Kusuoka and D. Stroock. Some boundedness properties of certain stationary diffusion semigroups. *Journal of Functional Analysis*, vol. 60, no. 2, pp. 243–264 (1985).
- [50] S.P. Meyn and R. L. Tweedie. A survey of Foster-Lyapunov techniques for general state space Markov processes. In *Proceedings of the Workshop on Stochastic Stability and Stochastic Stabilization*, Metz, France (1993).

- [51] S.P. Meyn and R. L. Tweedie. Markov Chains and Stochastic Stability. Springer Science & Business Media (2012).
- [52] J. Nash. Continuity of solutions of parabolic and elliptic equations. American J. Math. vol. 80, pp 931–954 (1958).
- [53] D. Nualart. Malliavin Calculus and Related Topics. Second Edition. Springer New York (2006).
- [54] R. D. Nussbaum. Eigenvectors of order-preserving linear operators. J. London Math. Soc., vol. 58, no. 2, pp. 480–496 (1998).
- [55] W. Oçafrain. Convergence to quasi-stationarity through Poincaré inequalities and Bakry-Emery criteria. Electronic Journal of Probability, vol. 26, pp. 1–30 (2021).
- [56] C. Olivera, C. Tudor. Density for solutions to stochastic differential equations with unbounded drift. Brazilian Journal of Probability and Statistics, vol. 33, no. 3, pp. 520–531 (2019).
- [57] E.M. Ouhabaz and F.Y. Wang. Sharp estimates for intrinsic ultracontractivity on $C^{1,\alpha}$ -domains. Manuscripta Math. 122 no. 2, pp. 229–244 (2007).
- [58] M. Sanz-Solé. Malliavin Calculus. With applications to stochastic partial differential equations. Fundamental Sciences, EPFL Press, Lausanne (1995).
- [59] R. Song. Two-sided estimates on the density of the Feynman-Kac semigroups of stable-like processes. Electronic Journal of Probability, vol. 11, pp. 146–161 (2006).
- [60] F.M. Spieksma. Countable state space Markov processes: Non explosiveness and moment function. Probability in the Engineering and Informational Sciences, vol. 29, pp. 623–637 (2015).
- [61] D. W. Stroock. Diffusion semigroups corresponding to uniformly elliptic divergence form operators. Séminaire de Probabilités, XXII, pp. 316– 347. Springer, Berlin (1988).
- [62] N.T. Varopoulos. Small time gaussian estimates of heat diffusion kernels. II. The theory of large deviations. Journal of Functional Analysis, vol. 93, pp. 1-33 (1990).
- [63] C. Wang. On estimates of the density of Feynman-Kac semigroups of α stable-like processes. Journal of mathematical analysis and applications, vol. 348, no.2, pp. 938–970 (2008).
- [64] N. Whiteley. Stability properties of some particle filters. The Annals of Applied Probability, vol. 23, no. 6, pp. 2500–2537 (2013).

- [65] Y. Zhang. On the non explosion and explosion for non homogeneous Markov pure jump processes. *Journal of Theoretical Probability*, vol. 31, no. 3, 1322–1355 (2018).