



HAL
open science

The Potential Implication of the Autonomic Nervous System in Hepatocellular Carcinoma

Romain Parent, Yori Gidron, Fanny Lebossé, Thomas Decaens, Fabien Zoulim

► **To cite this version:**

Romain Parent, Yori Gidron, Fanny Lebossé, Thomas Decaens, Fabien Zoulim. The Potential Implication of the Autonomic Nervous System in Hepatocellular Carcinoma. Cellular and Molecular Gastroenterology and Hepatology, 2019, 8 (1), pp.145-148. 10.1016/j.jcmgh.2019.03.002 . hal-03932347

HAL Id: hal-03932347

<https://hal.science/hal-03932347>

Submitted on 19 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMMENTARY

The Potential Implication of the Autonomic Nervous System in Hepatocellular Carcinoma



Liver cancer casualties are increasing worldwide despite the recent advent of adequate hepatitis B virus and hepatitis C virus therapies. Though extensive effort has been made in the past 20 years to study the effects of nonviral comorbidities on hepatocellular carcinoma (HCC), the unexplained heterogeneity of clinical evolution among patients in terms of traits and speed remains problematic for clinicians. After almost two decades of intense molecular profiling, the overall understanding of HCC development at the level of cells and tissues remains incomplete. Despite many potential therapeutic options, several drugs have failed to exceed the efficacy of the currently available compounds in phase III trials, indicating a need for novel targets¹ and renewed approaches for studying the disease.²

Indeed, although the major contribution of hepatocytes, and the implication of activated hepatic stellate cells and immune cells, is well documented in HCC, other cell types including intrahepatic neurons have seldom been studied. These latter lie at the interface between neurobehavioral traits of the patient and the pathology of the organ, and may thus improve prediction of disease prognosis and response to personalized (ie, tailor-made) therapy. Furthermore, keeping in mind that, in general oncology, 60% of patients develop cancers for reasons as yet unknown, once likely risk factors (eg, predisposing congenital mutations, smoking, alcohol consumption) have been considered,³ this novel research approach may identify as yet unsuspected HCC risk factors.

Brief Neurobiological Considerations in the Liver: Potential Influence on HCC-Predisposing Chronic Liver Disease

Succinct Anatomical Notions Pertaining to the Hepatic Autonomic Nervous System

The autonomic nervous system (ANS) (ie, visceral and involuntary sympathetic and parasympathetic nervous systems, or sympathetic nervous system and parasympathetic nervous system, respectively) is part of the peripheral nervous system, which plays a key role in regulating numerous physiological events, in synergy with the central nervous system (CNS). The sympathetic and parasympathetic nervous systems relay signals along the brain-liver neural axis. In 1924, it was shown that branches of the vagal nerve innervate the liver via the portal area, and are closely associated with the portal vein and bile ducts.⁴ In most species, a number of markers of sympathetic and parasympathetic nerves have been

observed surrounding the hepatic artery, portal vein and bile ducts, indicating general innervation of these structures by a variety of neuron subtypes. Intriguingly, the expression of sympathetic nerve fibers in species such as the rat is restricted to these areas, whereas in humans the expression of markers of ANS fibers extends deep into hepatic lobules and reaches liver parenchymal cells.⁵⁻⁷ This suggests a bidirectional and permanent connection between liver pathophysiology and the CNS, through afferent and efferent ANS fibers that sense and respond to relevant signals.

Pathophysiological Implication of the ANS in Chronic Liver Disease

CNS- or neural-derived events have recently been associated with chronic disease⁸ and cancer survival rates.⁹ Psychosocial factors independently and significantly predicted cancer prognosis in a meta-analysis of over 150 studies.¹⁰ Systematic hepatic neural ablation highlighted the importance of autonomic innervation in both physiological and pathological contexts, including metabolism, liver regeneration, and liver fibrosis. Indeed, afferent hepatic nerves of the ANS ensure liver sensing of glucose (via the GLUT2 receptor), lipids (through several protein kinase C isoforms), hormones (eg, GLP-1 and CCK), and owing to its well-known implication in liver pathology, interleukin (IL)-1 β , which in turn increases hepatic vagal activity.¹¹ Efferent hepatic nerves, linking the CNS to hepatic physiology, regulate the hepatic vasculature through its adrenergic and cholinergic arms.¹² The impact of the parasympathetic system on sinusoidal contractility, through its neurotransmitter acetylcholine, has also been reported¹² with likely consequences on blood supply regulation.

The influence of such processes on tumor metabolism is very likely, as nutrients and metabolites are delivered and excreted via the bloodstream, and their implication on the evolution of cirrhosis, the gateway to HCC, has been experimentally established.¹³ Such efferent nerves also alter hepatic metabolism in real time, after CNS sensing of circulating amino acids. This occurs through the implication of hypothalamic nuclei and the brainstem, themselves influencing the local secretion of different neurohormones, such as neuropeptide Y, pro-opiomelanocortin, pituitary adenylate cyclase-activating peptide, and glucagon-like peptide 1.¹⁴

Regarding HCC onset in the context of chronic liver disease (CLD), the efferent nerves of the ANS were shown to be implicated in liver repair and regeneration. One study proposed that after partial hepatectomy, liver regeneration was severely impaired by vagotomy in rats,¹⁵ while two others indicated that adrenergic signaling encouraged regeneration^{16, 17} via hepatocyte growth factor induction¹⁸

and moderation of transforming growth factor $\beta 1$ activity.¹⁹ Hepatic stellate cells express adrenoreceptors, the stimulation of which increases transforming growth factor $\beta 1$ and collagen production, and fosters their orientation toward a profibroblastic phenotype.²⁰ However, unlike vagotomy, sympathetic inhibition by chemicals causes an increase in hepatic oval cells, which are considered to be potential HCC precursors.²⁰ Efferent hepatic vagal activity has anti-inflammatory effects through its action on local IL-1 β and IL-6 secretion.²¹ Hence, an important part of the beneficial association of vagal activity with cancer outcome⁹ may involve this anti-inflammatory pathway. Such findings are summarized in Figure 1.

Heart Rate Variability as an Interesting Marker in CLD

Heart rate variability (HRV) is a systemic marker of vagal activity,²² and high HRV is predictive of survival in pancreatic cancer via its association with reduced inflammation.^{23,24} Concerning the systemic level of vagal nerve activity, HRV significantly and independently predicts cancer prognosis,²² where it is predictive of longer

overall survival or reduced tumor marker levels over time in breast, colon, pancreatic and prostate cancers.^{9,22,24,25} With respect to CLD, HRV significantly predicts mortality and liver transplantation in cirrhotic patients, independently of confounders (eg, disease severity).^{26,27} Finally, HRV frequency domain parameters were significantly and positively correlated with time to death in patients with terminal HCC.²⁸ In line with this finding, from the adrenergic arm perspective, a meta-analysis of 23 randomized trials carried out on >2600 patients with cirrhosis indicates that nonselective sympathetic beta-blockers lead to reduced risk of HCC.²⁹ The hepatic tissue of HCC patients is innervated by both sympathetic and parasympathetic nerves. Though little is known on the role of the ANS in HCC, nerve fiber density seems to be associated with poor prognosis.²⁹ The sympathetic nervous system was speculated to promote hepatocarcinogenesis by modulating inflammation through activation of α_1 -adrenergic receptors of proinflammatory Kupffer cells.³⁰ Such findings including those mentioned in the previous sections are innovative and are especially appealing given the paucity of adequate predictive and curative approaches for HCC.




Figure 1. Pathophysiological functions conveyed both ways by autonomic nervous system (ANS) fibers along the brain–liver axis. GLP1, glucagon-like peptide 1; HOC, hepatic oval cell; HSC, hepatic stellate cell; IL, interleukin; PACAP, pituitary adenylate cyclase-activating peptide; POMC, pro-opiomelanocortin.

Future Directions: Pharmacology, Cell Biology, Animal Models, and Clinical Studies

Despite their potential interest, these new results often rely on correlative studies, and are sometimes conflicting. The precise roles of the sympathetic and parasympathetic branches of the ANS and their causal implication in HCC onset and growth from a biological and mechanistic perspective at the intra-organ level remain unknown. Intriguingly, an acetylcholine-secreting memory phenotype splenic T cell population was shown to regulate inflammation and immune responses,³¹ likely affecting cancer immunosurveillance and immunotherapy. In this context, a number of interesting projects could be proposed. First, *in vitro* experiments could evaluate the influence of cholinergic and adrenergic signaling on the major pathogenic processes (eg, inflammation, fibrogenesis, loss of differentiation and transformation) related to carcinogenesis on each isolated hepatic cell type, including hepatocytes and hepatocyte-like cells, Kupffer cells, endothelial cells and stellate cells, as well as liver mononuclear cells. In particular, hepatic stellate cells in normal liver are arranged in a network of interconnected processes, making contact with sinusoidal endothelial cells and hepatocytes. They are a likely possible portal for mediating effects of the ANS throughout the liver.³² Of note, following liver transplantation, the newly implanted liver is a denervated organ. The implications of such could also be fruitfully considered in clinical investigations in the HCC field, although confounding factors such as immunosuppression need to be taken into consideration. Second, in the event of causal effects observed in the previously mentioned experiments, preclinical *in vivo* models of CLD and HCC could be used to evaluate the sensitivity of HCC onset, evolution, and growth to sympathetic and parasympathetic signaling. Third, retrospective and prospective studies could be implemented at the level of the patient in order to evaluate the prognostic and correlative potential of ANS-derived circulating or resident markers regarding HCC onset and outcome. Finally, pharmacological synergy could be examined between current standard-of-care molecules used for HCC and agonists or antagonists of ANS-related receptors, several of which are present in the tumor local environment and in the liver.¹⁴

Overall, combined with the recently demonstrated anti-inflammatory effects of vagal-stimulating implantable devices in cancer predisposing inflammatory diseases of the GI tract,^{33,34} these findings strengthen the potential interest of liver neurology considerations in future HCC research.

ROMAIN PARENT

UMR5286, CNRS, INSERM U1052, Lyon Cancer Research Center, Lyon, France
Department of Immunology and Virology
University of Lyon, Lyon, France
DevWeCan Laboratories of Excellence Network (Labex)
Lyon, France

YORI GIDRON

SCALAB UMR CNRS 9193, University of Lille
Villeneuve d'Ascq, France

FANNY LEBOSSE

UMR5286, CNRS, INSERM U1052, Cancer Research Centre of Lyon, Lyon, France
Department of Immunology and Virology
University of Lyon, Lyon, France
Hepatogastroenterology Service, Hôpital de la Croix-Rousse, Hospices Civils de Lyon, Lyon, France

THOMAS DECAENS

University of Grenoble-Alpes, Grenoble, France
Department of Hepatogastroenterology, Centre Hospitalier Universitaire Grenoble-Alpes, La Tronche, France
Institute for Advanced Biosciences, CNRS UMR 5309, INSERM U1209
University of Grenoble-Alpes, Grenoble, France

FABIEN ZOULIM

UMR5286, CNRS, INSERM U1052, Cancer Research Centre of Lyon, Lyon, France
Department of Immunology and Virology
University of Lyon, Lyon, France
DevWeCan Laboratories of Excellence Network (Labex), Lyon, France
Hepatogastroenterology Service, Hôpital de la Croix-Rousse, Hospices Civils de Lyon, Lyon, France

References

1. Nault JC, Galle PR, Marquardt JJ. The role of molecular enrichment on future therapies in hepatocellular carcinoma. *J Hepatol* 2018;69:237–247.
2. Llovet JM, Montal R, Sia D, Finn RS. Molecular therapies and precision medicine for hepatocellular carcinoma. *Nat Rev Clin Oncol* 2018;15:599–616.
3. Tomasetti C, Vogelstein B. Cancer etiology. Variation in cancer risk among tissues can be explained by the number of stem cell divisions. *Science* 2015;347:78–81.
4. McCrea ED. The Abdominal Distribution of the Vagus. *J Anat* 1924;59:18–40.
5. Bioulac-Sage P, Lafon ME, Saric J, Balabaud C. Nerves and perisinusoidal cells in human liver. *J Hepatol* 1990;10:105–112.
6. Fukuda Y, Imoto M, Koyama Y, Miyazawa Y, Hayakawa T. Demonstration of noradrenaline-immunoreactive nerve fibres in the liver. *J Int Med Res* 1996;24:466–472.
7. Horn CC, Friedman MI. Separation of hepatic and gastrointestinal signals from the common “hepatic” branch of the vagus. *Am J Physiol Regul Integr Comp Physiol* 2004;287:R120–R126.
8. Buccelletti F, Di Somma S, Iacomini P, Galante A, Pugliese F, Alegiani F, Bertazzoni G, Marsiliani D, Carroccia A, Granato A, Calabrò G, Legramante JM, Zuccalà G, Franceschi F. Assessment of baseline characteristics and risk factors among Emergency

- Department patients presenting with recent onset atrial fibrillation: a retrospective cohort study. *Eur Rev Med Pharmacol Sci* 2013;17(Suppl 1):22–27.
9. Zhou X, Ma Z, Zhang L, Zhou S, Wang J, Wang B, Fu W. Heart rate variability in the prediction of survival in patients with cancer: A systematic review and meta-analysis. *J Psychosom Res* 2016;89:20–25.
 10. Chida Y, Steptoe A. Positive psychological well-being and mortality: a quantitative review of prospective observational studies. *Psychosom Med* 2008;70:741–756.
 11. Nijima A. Afferent discharges from osmoreceptors in the liver of the guinea pig. *Science* 1969;166:1519–1520.
 12. Bennett TD, MacAnespie CL, Rothe CF. Active hepatic capacitance responses to neural and humoral stimuli in dogs. *Am J Physiol* 1982;242:H1000–H1009.
 13. Ueno T, Bioulac-Sage P, Balabaud C, Rosenbaum J. Innervation of the sinusoidal wall: regulation of the sinusoidal diameter. *Anat Rec A Discov Mol Cell Evol Biol* 2004;280:868–873.
 14. Jensen KJ, Alpini G, Glaser S. Hepatic nervous system and neurobiology of the liver. *Compr Physiol* 2013;3:655–665.
 15. Kato H, Shimazu T. Effect of autonomic denervation on DNA synthesis during liver regeneration after partial hepatectomy. *Eur J Biochem* 1983;134:473–478.
 16. Cruise JL, Knechtle SJ, Bollinger RR, Kuhn C, Michalopoulos G. Alpha 1-adrenergic effects and liver regeneration. *Hepatology* 1987;7:1189–1194.
 17. Cruise JL, Michalopoulos G. Norepinephrine and epidermal growth factor: dynamics of their interaction in the stimulation of hepatocyte DNA synthesis. *J Cell Physiol* 1985;125:45–50.
 18. Broten J, Michalopoulos G, Petersen B, Cruise J. Adrenergic stimulation of hepatocyte growth factor expression. *Biochem Biophys Res Commun* 1999;262:76–79.
 19. Houck KA, Cruise JL, Michalopoulos G. Norepinephrine modulates the growth-inhibitory effect of transforming growth factor-beta in primary rat hepatocyte cultures. *J Cell Physiol* 1988;135:551–555.
 20. Oben JA, Roskams T, Yang S, Lin H, Sinelli N, Torbenson M, Smedh U, Moran TH, Li Z, Huang J, Thomas SA, Diehl AM. Hepatic fibrogenesis requires sympathetic neurotransmitters. *Gut* 2004;53:438–445.
 21. Kimura K, Inaba Y, Watanabe H, Matsukawa T, Matsumoto M, Inoue H. Nicotinic alpha-7 acetylcholine receptor deficiency exacerbates hepatic inflammation and fibrosis in a mouse model of non-alcoholic steatohepatitis. *J Diabetes Investig* 2018 Oct 27 [E-pub ahead of print].
 22. Kuo TB, Lai CJ, Huang YT, Yang CC. Regression analysis between heart rate variability and baroreflex-related vagus nerve activity in rats. *J Cardiovasc Electrophysiol* 2005;16:864–869.
 23. Gidron Y, Deschepper R, De Couck M, Thayer JF, Velkeniers B. The vagus nerve can predict and possibly modulate non-communicable chronic diseases: introducing a neuroimmunological paradigm to public health. *J Clin Med* 2018;7:371.
 24. De Couck M, Marechal R, Moorhamers S, Van Laethem JL, Gidron Y. Vagal nerve activity predicts overall survival in metastatic pancreatic cancer, mediated by inflammation. *Cancer Epidemiol* 2016;40:47–51.
 25. De Couck M, Gidron Y. Norms of vagal nerve activity, indexed by heart rate variability, in cancer patients. *Cancer Epidemiol* 2013;37:737–741.
 26. Bhogal AS, De Rui M, Pavanello D, El-Azizi I, Rowshan S, Amodio P, Montagnese S, Mani AR. Which heart rate variability index is an independent predictor of mortality in cirrhosis. *Dig Liver Dis* 2018 Sep 24 [E-pub ahead of print].
 27. Ko FY, Yang AC, Tsai SJ, Zhou Y, Xu LM. Physiologic and laboratory correlates of depression, anxiety, and poor sleep in liver cirrhosis. *BMC Gastroenterol* 2013;13:18.
 28. Chiang JK, Koo M, Kuo TB, Fu CH. Association between cardiovascular autonomic functions and time to death in patients with terminal hepatocellular carcinoma. *J Pain Symptom Manage* 2010;39:673–679.
 29. Zhang L, Wu LL, Huan HB, Chen J, Wen XD, Yang DP, Xia F. Sympathetic and parasympathetic innervation in hepatocellular carcinoma. *Neoplasma* 2017;64:840–846.
 30. Huan HB, Wen XD, Chen XJ, Wu L, Wu LL, Zhang L, Yang DP, Zhang X, Bie P, Qian C, Xia F. Sympathetic nervous system promotes hepatocarcinogenesis by modulating inflammation through activation of alpha1-adrenergic receptors of Kupffer cells. *Brain Behav Immun* 2017;59:118–134.
 31. Rosas-Ballina M, Olofsson PS, Ochani M, Valdés-Ferrer SI, Levine YA, Reardon C, Tusche MW, Pavlov VA, Andersson U, Chavan S, Mak TW, Tracey KJ. Acetylcholine-synthesizing T cells relay neural signals in a vagus nerve circuit. *Science* 2011;334:98–101.
 32. Wake K. Hepatic stellate cells: Three-dimensional structure, localization, heterogeneity and development. *Proc Jpn Acad Ser B Phys Biol Sci* 2006;82:155–164.
 33. Bonaz B, Sinniger V, Pellissier S. Vagus nerve stimulation: a new promising therapeutic tool in inflammatory bowel disease. *J Intern Med* 2017;282:46–63.
 34. Bonaz B, Sinniger V, Pellissier S. Vagus nerve stimulation at the interface of brain-gut interactions. *Cold Spring Harb Perspect Med* 2018 Sep 10 [E-pub ahead of print].

Correspondence

Address correspondence to: Romain Parent, PharmD, PhD, INSERM U1052 (Lyon Cancer Res Center), Department of Immunology and Virology, 151 Cours Albert Thomas, F-69424 Lyon, France. e-mail: romain.parent@inserm.fr; fax: 33-4-72-68-19-71.

Conflicts of interest

This author declares the following: Thomas Decaens has shared interests with AstraZeneca, Bayer, BMS, BTG, Ipsen, and Sirtex (consulting, member of advisory board and/or teaching), as well as Arqule, Genoscience, and Roche (research). The remaining authors disclose no conflicts.

Funding

This work was supported by DevWeCan Laboratories of excellence network (Labex; Grant #ANR LABX-061).

Most current article

© 2019 The Authors. Published by Elsevier Inc. on behalf of the AGA Institute. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2352-345X

<https://doi.org/10.1016/j.jcmgh.2019.03.002>