

# $(\mu 2-\eta 4-N-(2-Butynyl)phthalimide)(hexacarbonyl)dicobalt$

Isabelle Jourdain, Michael Knorr, Tom Charenton, Carsten Strohmann,

Jan-Lukas Kirchhoff, Mohamed Othman

### ► To cite this version:

Isabelle Jourdain, Michael Knorr, Tom Charenton, Carsten Strohmann, Jan-Lukas Kirchhoff, et al.. ( $\mu$ 2- $\eta$ 4-N-(2-Butynyl)phthalimide)(hexacarbonyl)dicobalt. Molbank, 2023, 2023 (1), pp.M1545. 10.3390/M1545. hal-03943146

## HAL Id: hal-03943146 https://hal.science/hal-03943146

Submitted on 19 Jan 2023  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Short Note $(\mu_2 - \eta^4 - N - (2 - Butynyl)phthalimide)(hexacarbonyl)dicobalt$

Isabelle Jourdain <sup>1,\*</sup>, Michael Knorr <sup>1,\*</sup>, Tom Charenton <sup>1</sup>, Carsten Strohmann <sup>2</sup>, Jan-Lukas Kirchhoff <sup>2</sup>, and Mohamed Othman <sup>3</sup>

- <sup>1</sup> Institut UTINAM UMR 6213 CNRS, Université de Franche-Comté, 16 Route de Gray, 25030 Besançon, France
- <sup>2</sup> Anorganische Chemie, Technische Universität Dortmund, Otto-Hahn Straße 6, 44227 Dortmund, Germany
- <sup>3</sup> URCOM, UNILEHAVRE, Normandie University, FR3021, EA 3221, 25 Rue P. Lebon, 76058 Le Havre, France
  - Correspondence: isabelle.jourdain@univ-fcomte.fr (I.J.); michael.knorr@univ-fcomte.fr (M.K.); Tel.: +33-3-8166-6270 (M.K.)

**Abstract**: The reaction of  $[Co_2(CO)_8]$  with an equimolar amount of the internal alkyne *N*-(2-butynyl)phthalimide (1-Phthalimido-2-butyne) **1** in heptane solution yields the title compound  $[Co_2(CO)_6(\mu$ phthalimidoCH<sub>2</sub>C $\equiv$ CMe)] **2**. Compound **2** has been characterized using IR, <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy; the tetrahedrane-type cluster framework has been ascertained using a single-crystal X-ray diffraction study performed at 100 K.

Keywords: cobalt; carbonyl; triphenylphosphite; crystal structure; molecular cluster

#### 1. Introduction

Since the first reports dating from the 1950s on the reaction of dicobalt octacarbonyl with alkynes yielding dicobaltatetrahedranes  $[Co_2(CO)_6(\mu-RC\equiv CR')]$  [1,2], there is nowadays a plethora of articles dealing with this organometallic reaction [3]. The interest in this research is not only driven by merely synthetic aspects, i.e., spectroscopic and structural characterization of these cage-like species, but also by the use of these compounds as precursors for further modifications in material sciences [4] and organic syntheses [5,6]. An important example for the latter domain is the Pauson–Khand reaction, in which  $[Co_2(CO)_8]$  reacts with an alkyne, an alkene, and carbon monoxide to form an  $\alpha,\beta$ -cyclolopentenone [7–11]. In addition to the applications using dicobaltatetrahedranes as synthetic tools, this class of dinuclear organometallic compounds is emerging in bio-organometallic chemistry [12–21]. Some examples of dicobaltatetrahedranes, ligated using terminal functionalized alkynes and displaying biological activity, are depicted in Scheme 1.

Among the organic heterocyclic compounds showing a promising biological and medicinal activity, there are also some representatives based on the phthalimido scaffold [22–24]. In 2005, Gust et al. also reported the reaction of propargyl phtalimide with [ $Co_2(CO)_8$ ] and determined the cytotoxicity and DNA binding efficiency of the resulting complex *N*-(2-propynyl)phthalimide]hexacarbonyldicobalt (see compound **V** of Scheme 1) [21]. In the context of our research on Co–Co carbonyl complexes towards various alkynes producing dicobaltatetrahedranes [25,26], we attempted to synthetize a Co–Co complex ligated using an internal alkyne, namely 1-Phthalimido-2-butyne **1**, for upcoming biological studies. Apart from the purely synthetic aspect, one of the objectives was to compare the impact of the replacement of a terminal alkyne by an internal one (Phtal-C=CH vs. Phtal-C=CMe).



**Citation:** Jourdain, I.; Knorr, M.; Charenton, T.; Strohmann, C.; Kirchhoff, J.-L.; Othman, M. (μ<sub>2</sub>-η<sup>4</sup>-N-(2-Butynyl)phthalimide) (hexacarbonyl)dicobalt. *Molbank* **2023**, 2023, M1545. https://doi.org/ 10.3390/M1545

Academic Editor: Rodrigo Abonia

Received: 8 December 2022 Revised: 26 December 2022 Accepted: 6 January 2023 Published: 11 January 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).



**Scheme 1.** Examples of some dicobaltatetrahedranes displaying biological activity (**I**, **III–V**: antiproliferative activity against cancer cells; **II**: hormonally active compound).

#### 2. Results and Discussion

The title compound **2** was obtained via treatment of  $[Co_2(CO)_8]$  with an equimolar amount of *N*-(2-butynyl)phthalimide **1** in heptane solution at 60 °C as shown in Scheme 2. Upon cooling, small orange-red needle-shaped crystals of air-stable **2** were isolated, and elemental analysis confirmed their composition as  $[Co_2(CO)_6(\mu-\text{phthalimidoCH}_2C\equiv CMe)]$ .



Scheme 2. Synthesis of the title compound 2.

The IR spectrum of this product in cyclohexane, shown in Figure 1, reveals the signals of terminal carbonyl with intense v(CO) vibrations at 2093, 2055, and 2028 cm<sup>-1</sup>. In addition, two further absorptions at 1775 and 1727 cm<sup>-1</sup> are attributed to the carbonyls' stretching modes (symmetric and antisymmetric) of the imide function [27]. These values fit well with those measured in cyclohexane for  $[Co_2(CO)_6(\mu$ -phthalimidoCH<sub>2</sub>C≡CH)] at 2094, 2057, and 2030 as well as 1775 and 1727 cm<sup>-1</sup> (Figure S1). The latter complex has already been described and characterized using IR spectroscopy in solid state as KBr pellets (2095, 2054, 2036, 2016 (Co-CO); 1712 (C=O) cm<sup>-1</sup>) [21]. The infrared spectrum of **2** in the solid state was also recorded in attenuated total reflectance (ATR) mode and is presented in the Supplementary Materials (Figure S2). Despite being a monocrystalline solid, the carbonyl absorptions were not fully resolved (2093, 2052, 2030, and 2008 as well as 1766 and 1708 cm<sup>-1</sup>).



Figure 1. IR spectrum of compound 2 recorded in cyclohexane.

The <sup>1</sup>H-NMR recorded in CDCl<sub>3</sub> reveals a singlet at  $\delta$  2.65 due to the acetylenic methyl group; a further singlet at  $\delta$  5.03 is attributed to the enantiotopic methylene hydrogens. The remaining resonances at  $\delta$ 7.74 and 7.88 are assigned to the aromatic cycle (Figure 2). In the proton-decoupled <sup>13</sup>C-NMR spectrum depicted in Figure 3, only a broadened resonance centered at  $\delta$  199.3 could be observed for the six Co-bound carbonyl groups, suggesting a fluxional behavior of the COs. In contrast, the resonance of the two chemically equivalent imido C=O carbonyl groups at  $\delta$  167.9, the two acetylenic carbons at  $\delta$  93. 2 and 91.8, as well as those of the CH<sub>2</sub> and CH<sub>3</sub> carbons gave rise to well-resolved signals.



**Figure 2.** <sup>1</sup>H-NMR spectrum (400 MHz, CDCl<sub>3</sub>) of compound **2** at 25 °C. The asterisk \* denotes the chloroform resonance.



**Figure 3.**  ${}^{13}C{}^{1}H$ -NMR spectrum (100.62 MHz, CDCl<sub>3</sub>) of compound **2** at 25 °C. The asterisk \* denotes the CDCl<sub>3</sub> resonance.

In addition to the spectroscopic characterization of **2** in solution, the complex was also unambiguously analyzed using X-ray diffraction performed at 100 K. Complex **2** crystallizes in the triclinic space group *P*-1; the asymmetric unit contains two independent molecules with slightly different bond lengths and angles. As shown in Figure 4, in this organometallic species the former alkyne is almost orthogonally  $\mu_2$ - $\eta^4$ -coordinated with respect to the two crystallographically non-equivalent Co1 and Co2 centers, which are linked through a metal–metal bond of 2.4688(14) Å.



**Figure 4.** Molecular structure of one of two independent molecules of **2**. All H atoms are omitted for clarity. Selected bond lengths (Å) and angles (deg): Co1–Co2 2.4688(14), Co1–C2 1.980(6), Co1–C3 1.952(6), Co2–C2 1.984(6), Co2–C3 1.954(7), Co1–C16 1.812(7), Co1–C17 1.784(8), Co1–C18 1.831(7), Co2–C13 1.794(7), Co2–C14 1.812(7), Co2–C15 1.805(7), C2–C3 1.323(9); Co1–Co2–C3 50.75(15), Co2–Co1–C2 50.83(19), Co1–C2–Co2 77.0(2), Co1–C3–Co2 78.4(2).

The resulting tetrahedral skeleton, reminiscent of that of organic tetrahedranes (https: //en.wikipedia.org/wiki/Tetrahedrane, accessed on 26 December 2022), allows for considering this 40-electron species an organometallic cluster of dimetallatetrahedranes, in which two Co(CO)<sub>3</sub> units of the parent cluster [{Co<sub>4</sub>( $\mu_2$ -CO)<sub>3</sub>(CO)<sub>9</sub>] are replaced by isolobal C–R fragments, forming the edges of the cluster core. The former alkyne unit lost its linearity; the torsion angle C1–C2–C3–C4 changed to  $-3.1(18)^\circ$ . The coordination sphere around each Co atom is completed by three terminal carbonyls. The mean Co–C bond length of the four pseudo-equatorial carbonyls is somewhat longer than that of the two pseudo-axial C≡O ligands (1.815(7) vs. 1.789(7) Å). The C2–C3 bond distance of **2** is much shorter than that of a C–C single bond, but considerably elongated with respect to the C≡C bond reported for [phthalimidoCH<sub>2</sub>C≡CCH<sub>2</sub>phthalimido] (CSD refcode ECUFOB) (1.323(9) vs. 1.191(4) Å) [28].

Although there are numerous examples of crystallographically characterized hexacarbonyldicobalt compounds which are  $\mu_2$ - $\eta^2$ , $\eta^2$  capped by an internal alkyne, a survey of the CSD data base revealed only two other examples containing a -CH<sub>2</sub>-C=C-Me motif as is present in **2**, namely (*trans*-5,8,8-trimethyl-*trans*-1,4,5,7-tetrahydro-*cis*-4-(2butynyl)bicyclo(5.1.0)octan-3-one)(hexacarbonyl)dicobalt (**A**) and (pent-3-yn-1-ol)(hexacarbonyl)dicobalt (**B**) (Figure 5) [29,30]. The most relevant averaged bond lengths (comprising molecules 1 and 2) of **2** are listed in Table 1 and compared with those of compounds **A** and **B** (averaged bond lengths).



**Figure 5.** Presentation of the two structurally characterized dicobaltatetrahedranes (A) and (B) incorporating a  $-CH_2-C\equiv C-Me$  motif.

**Table 1.** Comparison of relevant bond lengths (Å) and angles (°) in **2** and the crystallographically characterized  $[Co_2(CO)_6(\mu$ -RCH<sub>2</sub>C $\equiv$ CCH<sub>3</sub>)] tetrahedranes **A** and **B**.

	2	Α	В
Со-Со	2.4679(14)	2.4688(9)	2.4690(9)
C–C <sub>alkyne</sub>	1.330(9)	1.325(6)	1.326(6)
$C_{alkyne} - C_{Me}$	1.481(9)	1.486(6)	1.492(6)
$C_{alkyne} - C_{H2}$	1.499(9)	1.500(7)	1.493(6)
Co–C <sub>alkvne</sub>	1.972(6)	1.974(5)	1.972(4)
Co-C <sub>pseudo-equatorial</sub>	1.815(7)	1.823(5)	1.820(5)
Co-C <sub>pseudo-axial</sub>	1.797(7)	1.785(6)	1.787(5)
CSD reference	This work	DOTKEC [29]	HUKCIA [30]

In the molecular cell of **2** with Z = 4, there are no intermolecular contacts deserving any discussion (Figure 6).



Figure 6. View of the molecular cell of 2 showing the two pairwise arranged independent molecules.

#### 3. Materials and Methods

*N*-(2-butynyl)phthalimide 1 was commercially purchased from Aldrich. <sup>1</sup>H- and <sup>13</sup>C-NMR spectra were recorded on a Brucker AC 400 (Bruker, Wissembourg, France) at 400 and 100.62 MHz, respectively, using CDCl<sub>3</sub> as solvent. Infrared spectra were recorded on a Vertex 70 spectrometer (Bruker, Wissembourg, France) in ATR mode or in solution.

( $\mu_2$ - $\eta^4$ -*N*-(2-butynyl)phthalimide)(hexacarbonyl)dicobalt: Alkyne 1 (199.2 mg, 1 mmol) was added to a stirred solution of Co<sub>2</sub>(CO)<sub>8</sub> (342.0 mg, 1 mmol) in heptane (10 mL). An immediate gas evolution was observed. The reaction mixture was then heated to 60 °C for 5 h. The solution was cooled to room temperature prior to lowering its temperature to 4 °C. Product 2 crystallized as orange-red needles which were collected via filtration. Yield: 78%. Anal. Calc. for C<sub>18</sub>H<sub>9</sub>Co<sub>2</sub>NO<sub>8</sub> (M.W = 485.14 g.mol<sup>-1</sup>): C, 44.56; H, 1.87; N, 2.89%. Detected: C, 44.64; H, 1.92; N, 2.98 %. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) at 298 K: δ 2.65 (s, CH<sub>3</sub>), 5.03 (s, NCH<sub>2</sub>), 7.74 (br, CH Ar), 7.88 (br, CH Ar) ppm. <sup>13</sup>C{<sup>1</sup>H}-NMR (CDCl<sub>3</sub>) at 298 K: δ 20.3 (CH<sub>3</sub>), 39.7 (NCH<sub>2</sub>), 91.8 (Cq), 93.2 (Cq), 123.6 (CH), 132.0 (Cq), 134.4 (CH), 167.9 (C=O), 199.3 (br, CO) ppm.

Since the grown single-crystals of 2 used for the determination of the crystal structure were quite small,  $CuK_{\alpha}$  radiation was employed instead of  $MoK_{\alpha}$  radiation. Moreover, the crystals were twinned. The B-alert found in the checkcif is due to the high electronic residual around the Co–Co bonds, which could not be further refined due to the low resolution at 138.0°.

Crystal data for C<sub>18</sub>H<sub>9</sub>Co<sub>2</sub>NO<sub>8</sub>, M = 485.12 g.mol<sup>-1</sup>, orange-red needles, crystal size 0.377 × 0.194 × 0.15 mm<sup>3</sup>, triclinic, space group *P*-1: *a* = 7.4317(4)Å, *b* = 13.4351(7) Å, *c* = 19.2794(11) Å,  $\alpha$  = 78.462(3°,  $\beta$  = 84.011(3)°,  $\gamma$  = 89.156(3)°, *V* = 1875.73(18) Å<sup>3</sup>, *Z* = 4, D<sub>calc</sub> = 1.718 g/cm<sup>3</sup>, T = 100 K, h = -8 ≤ h ≤ 8, k = -15 ≤ k ≤ 16, I = 0 ≤ 1 ≤ 16, GOF = 1.077, *R*<sub>1</sub> = 0.0855, w*R*2 = 0.2428 for 6962 reflections with I > 2 $\sigma$  (I) and 11958 independent reflections. Largest diff. peak/hole/e Å<sup>-3</sup> 1.35/-1.02. The structure was solved using direct methods and refined using full-matrix least-squares against F<sup>2</sup> (SHELXL, 2015 [31–33]).

Data were collected using graphite-monochromated  $CuK_{\alpha}$  radiation l = 1.54178 Å and were deposited at the Cambridge Crystallographic Data Centre as CCDC 2219351.

(Supplementary Materials). The data can be obtained free of charge from the Cambridge Crystallographic Data Centre via http://www.ccdc.cam.ac.uk/getstructures.

#### 4. Conclusions

We have demonstrated that addition of the internal alkyne *N*-(2-butynyl)phthalimide **1** to  $[Co_2(CO)_8]$  straightforwardly yields the stable complex  $[Co_2(CO)_6(\mu$ -phthalimi-doCH<sub>2</sub>C $\equiv$ CMe)] **2** featuring a dimetallatetrahedrane framework. The investigation of the biological activity of **2** and other related compounds will be the topic of a forthcoming study.

**Supplementary Materials:** The following are available online, CIF file, Check-CIF report, and ATR-IR spectrum.

**Author Contributions:** I.J. and T.C. prepared the compound; C.S. and J.-L.K. collected the X-ray data and solved the structure; I.J., M.K. and M.O. designed the study; I.J. and M.K. analyzed the data and wrote the paper. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Région Bourgogne-Franche-Comté, DeCOmAB project.

Data Availability Statement: The X-ray data are at CCDC as stated in the paper.

Acknowledgments: We thank Stéphanie Beffy for recording some IR and NMR spectra. J.-L.K. acknowledges the receipt of a Kekulé scholarship from the Fonds der Chemischen Industrie (VCI).

Conflicts of Interest: The authors declare no conflict of interest.

#### References

- 1. Greenfield, H.; Sternberg, H.W.; Friedel, R.A.; Wotiz, J.H.; Markby, R.; Wender, I. Acetylenic dicobalt hexacarbonyls. organometallic compounds derived from alkynes and dicobalt octacarbonyl. *J. Am. Chem. Soc.* **1956**, *78*, 120–124. [CrossRef]
- 2. Sly, W.G. The Molecular configuration of dicobalt hexacarbonyl diphenylacetylene. J. Am. Chem. Soc. 1959, 81, 18–20. [CrossRef]
- 3. Barnes, C.E. Comprehensive Organometallic Chemistry II; Cobalt, Rhodium and Iridium; Elsevier: New York, NY, USA, 1995; Volume 8.
- 4. Georgi, C.; Hildebrandt, A.; Waechtler, T.; Schulz, S.E.; Gessner, T.; Lang, H. A cobalt layer deposition study: Dicobaltatetrahedranes as convenient MOCVD precursor systems. *Mat. J. Chem.* **2014**, *2*, 4676–4682. [CrossRef]
- 5. Lockwood, R.F.; Nicholas, K.M. Transition metal-stabilized carbenium ions as synthetic intermediates. I. α-[(alkynyl)dicobalt hexacarbonyl] carbenium ions as propargylating agents. *Tetrahedron Lett.* **1977**, *18*, 4163–4165. [CrossRef]
- Nicholas, K.M. Chemistry and synthetic utility of cobalt-complexed propargyl cations. Acc. Chem. Res. 1987, 20, 207–214. [CrossRef]
- Pauson, P.L.; Khand, I.U. Uses of cobalt-carbonyl acetylene complexes in organic synthesis. Ann. N.Y. Acad. Sci. 1977, 295, 2–14. [CrossRef]
- 8. Hay, A.M.; Kerr, W.J.; Kirk, G.G.; Middlemiss, D. Highly efficient enantioselective Pauson-Khand reactions. *Organometallics* **1995**, 14, 4986–4988. [CrossRef]
- 9. Gibson, S.E.; Stevenazzi, A. The Pauson-Khand reaction: The catalytic age is here! *Angew. Chem. Int. Ed.* **2003**, *4*, 1800–1810. [CrossRef]
- 10. Blanco-Urgoiti, J.; Añorbe, L.; Pérez-Serrano, L.; Domínguez, G.; Pérez-Castells, J. The Pauson–Khand reaction, a powerful synthetic tool for the synthesis of complex molecules. *Chem. Soc. Rev.* **2004**, *33*, 32–42. [CrossRef]
- 11. Werner, H. Peter Ludwig Pauson (1925–2013). Angew. Chem. Int. Ed. 2014, 53, 3309. [CrossRef]
- 12. Ott, I.; Kircher, B.; Dembinski, R.; Gust, R. Alkyne hexacarbonyl dicobalt complexes in medicinal chemistry and drug development. *Expert Opin. Ther. Patents.* 2008, 18, 327–337. [CrossRef]
- Minteanu, C.R.; Suntharalingam, K. Advances in cobalt complexes as anticancer agents. *Dalton Trans.* 2015, 44, 13796–13808. [CrossRef]
- 14. Jung, M.; Kerr, D.E.; Senter, P.D. Bioorganometallic chemistry: Synthesis and antitumor activity of cobalt carbonyl complexes. *Arch. Pharm.* **1997**, *330*, 173–176. [CrossRef] [PubMed]
- 15. Schmidt, K.; Jung, M.; Keilitz, R.; Schnurr, B.; Gust, R. Acetylenehexacarbonyldicobalt complexes, a novel class of antitumor drugs. *Inorg. Chim. Acta.* 2000, 306, 6–16. [CrossRef]
- Ott, I.; Kircher, B.; Gust, R. Investigations on the effects of cobalt-alkyne complexes on leukemia and lymphoma cells: cytotoxicity and cellular uptake. J. Inorg. Biochem. 2004, 98, 485–489. [CrossRef] [PubMed]
- Vessieres, A.; Top, S.; Vaillant, C.; Osella, D.; Mornon, J.P.; Jaouen, G. Estradiols modified by metal carbonyl clusters as suicide substrates for the study of receptor proteins: Application to the estradiol receptor. *Angew. Chem. Int. Ed. Engl.* 1992, *31*, 753–755. [CrossRef]

- Sergeant, C.D.; Ott, I.; Sniady, A.; Meneni, S.; Gust, R.; Rheingold, A.L.; Dembinski, R. Metallo-nucleosides: Synthesis and biological evaluation of hexacarbonyl dicobalt 5-alkynyl-2'-deoxyuridines. *Org. Biomol. Chem.* 2008, *6*, 73–80. [CrossRef]
- Kaczmarek, R.; Korczyński, D.; Królewska-Golińska, K.; Wheeler, K.A.; Chavez, F.A.; Mikus, A.; Dembinski, R. Organometallic Nucleosides: Synthesis and Biological Evaluation of Substituted Dicobalt Hexacarbonyl 2'-Deoxy-5-oxopropynyluridines. *ChemistryOpen* 2018, 7, 237–247. [CrossRef]
- 20. Neukamm, M.A.; Pinto, A.; Metzler-Nolte, N. Synthesis and cytotoxicity of a cobaltcarbonyl-alkyne enkephalin bioconjugate. *Chem. Commun.* **2008**, *2*, 232–234. [CrossRef]
- 21. Ott, I.; Schmidt, K.; Kircher, B.; Schumacher, P.; Wiglenda, T.; Gust, R. Antitumor-active cobalt-alkyne complexes derived from acetylsalicylic acid: Studies on the mode of drug action. *J. Med. Chem.* **2005**, *48*, 622–629. [CrossRef]
- 22. Almeida, M.L.; Oliveira, M.C.V.A.; Pitta, I.R.; Pitta, M.G.R. Advances in Synthesis and Medicinal Applications of Compounds Derived from Phthalimide. *COS* 2020, *17*, 252–270. [CrossRef] [PubMed]
- Sharma, U.; Kumar, P.; Kumar, N.; Singh, B. Recent Advances in the Chemistry of Phthalimide Analogues and their Therapeutic Potential. *Min. Rev. Med. Chem.* 2010, 10, 678–704. [CrossRef] [PubMed]
- 24. Pigeon, P.; Gaschard, M.; Othman, M.; Salmain, M.; Jaouen, G. α-Hydroxylactams as Efficient Entries to Diversely Functionalized Ferrociphenols: Synthesis and Antiproliferative Activity Studies. *Molecules* **2022**, *27*, 4549. [CrossRef]
- Clément, S.; Guyard, L.; Khatyr, A.; Knorr, M.; Rousselin, Y.; Kubicki, M.M.; Mugnier, Y.; Richeter, S.; Gerbier, P.; Strohmann, C. Synthesis, crystallographic and electrochemical study of ethynyl[2.2]paracyclophane-derived cobalt metallatetrahedranes. J. Organomet. Chem. 2012, 699, 56–66. [CrossRef]
- Jourdain, I.; Knorr, M.; Brieger, L.; Strohmann, C. Synthesis of Tris(arylphosphite)-ligated Cobalt(0) Complexes [Co<sub>2</sub>(CO)<sub>6</sub>{P(OAr)<sub>3</sub>}<sub>2</sub>], and their Reactivity towards Phenylacetylene and Diphenylacetylene. *Adv. Chem. Res.* 2020, 2. [CrossRef]
- 27. Hase, Y. The infrared and Raman spectra of phthalimide, N-d-phthalimide and potassium phthalimide. *J. Mol. Struct.* **1978**, *48*, 33–42. [CrossRef]
- Chen, H.; Ai, Z.; Guo, L.; Yao, L.; Li, Y.; Gu, B.; Zhang, Y.; Liu, Y.A.; Tian, B.; Liao, X. Nickel-catalyzed enantioselective domino Heck/Sonogashira coupling for construction of C(sp)-C(sp3) bond-substituted quaternary carbon centers. *Tetrahedron Chem.* 2022, 2, 100021–100031. [CrossRef]
- 29. Saha, M.; Muchmore, S.; van der Helm, D.; Nicholas, K.M. Cobalt-Mediated Cyclopentenone Annulation: An Approach to the Synthesis of Cyclocolorenone. *J. Org. Chem.* **1986**, *51*, 1960–1966. [CrossRef]
- Carniato, F.; Gatti, G.; Gervasio, G.; Marabello, D.; Sappa, E.; Secco, A. Reactions of Co<sub>2</sub>(CO)<sub>8</sub> and of Co<sub>2</sub>(CO)<sub>6</sub>L (L = 3-pentyn-1-ol, 1,4-butyn-diol or 2-methyl-3-butyn-2-ol) with 2(diphenylphosphino)ethyl-triethoxysilane and tris(hydroxymethyl)phosphine for applications to new sol–gel materials. *J. Organomet. Chem.* 2009, 694, 4241–4249. [CrossRef]
- 31. Dolomanov, O.V.; Bourhis, L.J.; Gildea, R.J.; Howard, J.A.; Puschmann, H. OLEX2: A complete structure solution, refinement and analysis program. *J. Appl. Cryst.* 2009, *42*, 339–341. [CrossRef]
- Sheldrick, G. SHELXT-Integrated space-group and crystal-structure determination. Acta Crystallogr. A 2015, 71, 3–8. [CrossRef] [PubMed]
- 33. Sheldrick, G. Crystal structure refinement with SHELXL. Acta Crystallogr. C 2015, 71, 3–8. [CrossRef] [PubMed]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.