



A new species *Zachvatkinibates svanhovdi* sp. nov. (Acari: Oribatida: Punc toribatidae) from Norway with comments on Punc toribatidae in Fennoscandia

Anna Seniczak, Stanislaw Seniczak, Snorre B Hagen, Cornelya F C Klütsch

► To cite this version:

Anna Seniczak, Stanislaw Seniczak, Snorre B Hagen, Cornelya F C Klütsch. A new species *Zachvatkinibates svanhovdi* sp. nov. (Acari: Oribatida: Punc toribatidae) from Norway with comments on Punc toribatidae in Fennoscandia. *Acarologia*, 2023, 63 (1), pp.41-57. 10.24349/skz7-19nt . hal-03943670

HAL Id: hal-03943670

<https://hal.science/hal-03943670>


Submitted on 17 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr



**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2023 (Volume 63): 450 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2021): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France
ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)



Acarologia is under **free license** and distributed under the terms of the Creative Commons-BY

A new species *Zachvatkinibates svanhovdi* sp. nov. (Acari: Oribatida: Punctoribatidae) from Norway with comments on Punctoribatidae in Fennoscandia

Anna Seniczak ^a, Stanisław Seniczak ^b, Snorre B. Hagen^c, Cornelya F. C. Klütsch^c

^a Department of Natural History, University Museum of Bergen, University of Bergen, Bergen, Norway.

^b Department of Evolutionary Biology, Faculty of Biological Sciences, Kazimierz Wielki University, Bydgoszcz, Poland.

^c Norwegian Institute of Bioeconomy Research, NIBIO - Division of Environment and Natural Resources, Svanhovd, Norway.

Original research

ABSTRACT

The adults of the new species *Zachvatkinibates svanhovdi* A. Seniczak & S. Seniczak sp. nov. are described and illustrated from Norway, and this is the first finding of *Zachvatkinibates* Shalbybina, 1973 in Fennoscandia. This species is the most similar to *Z. quadrivertex* (Halbert, 1920), but differs from it mainly by the shape of notogastral setae, posterior tectum of notogaster and lack of postanal porose area *Ap*, which in *Z. quadrivertex* is present. In *Z. svanhovdi*, the prodorsal seta *in* is long, translamella is narrow, notogastral setae are short and distally pliable, notogastral porose areas are usually oval and of medium size, but *Aa* can be larger, especially in males. Dorsal crest on tarsus I is present. The cytochrome oxidase I (COI) barcodes (length: 658 bp) of five specimens of the new species are provided; the maximum variation within the species was 2.41% (p-dist). The morphology and ecology of the new species is compared with other *Zachvatkinibates* species. The knowledge on family Punctoribatidae in Fennoscandia is updated, and *Mycobates carli* (Schweizer, 1922) is first reported from Norway.

Keywords oribatid mites; sexual dimorphism; leg setation; dorsal crest; sex structure; DNA barcoding

Zoobank <http://zoobank.org/9351C829-16CD-4084-A12E-1C98889982CD>

Received 12 August 2022
Accepted 09 December 2022
Published 13 January 2023

Corresponding author
Anna Seniczak :
Anna.Seniczak@uib.no

Academic editor
Baumann, Julia

<https://doi.org/10.24349/skz7-19nt>
ISSN 0044-586X (print)
ISSN 2107-7207 (electronic)


Seniczak A. et al.

Licensed under
Creative Commons CC-BY 4.0

OPEN  ACCESS

Introduction

Zachvatkinibates Shalbybina, 1973 *sensu stricto*, with the type species *Z. nemoralis* Shalbybina, 1973, is a medium sized genus that comprises 18 species, according to Subías (2004, 2022). However, this author included *Minunthozetes selgae* Pérez-Íñigo, 1976 in this genus, which is not consistent with diagnosis of Behan-Pelletier & Eamer (2008). In the later species, the posterior notogastral tectum is complete, whereas in *Zachvatkinibates* it is medially broken.

The diagnosis of the adult of *Zachvatkinibates* was proposed by Shalbybina (1973), but Behan-Pelletier (1988) and Behan-Pelletier & Eamer (2005) consecutively modified it, and in the latter paper it was as follows: notogaster with rounded, convex medial process on anterior tectum and pair of thickened bands bordering medial process, posterior notogastral tectum incomplete medially, indicated by wide margination or narrow slit bordered by overlapping tectal lobes; lenticulus absent, octotaxic system expressing sexually dimorphism in some

How to cite this article Seniczak A. et al. (2023), A new species *Zachvatkinibates svanhovdi* sp. nov. (Acari: Oribatida: Punctoribatidae) from Norway with comments on Punctoribatidae in Fennoscandia. *Acarologia* 63(1): 41-57.
<https://doi.org/10.24349/skz7-19nt>

species, postanal porose area and axillary saccule of subcapitulum present, and mental tectum of subcapitulum absent.

The main aim of this paper is to describe and illustrate the morphology and ecology of *Z. svanhovdi*, and to compare it with congeners. Another aim is to update information on Puncoribatidae in Fennoscandia, and add *Z. svanhovdi* and *Mycobates carli* (Schweizer, 1922) to the list of species in this region.

Material and methods

Study site

The sample was collected at the river outlet (Figure 1) of the Jacob's River in north-eastern Finnmark, Sør-Varanger municipality, Norway. The region is characterized by an oceanic-continental climate with a mean annual temperature of 0.1 °C and a mean annual precipitation of approximately 620 mm (Bjørklund *et al.* 2015). The Jacob's River (Grense Jakobselv; area 237 km²) is a river that runs along the Norwegian-Russian border and discharges into the Varanger Fjord, a bay located in the Barents Sea. The river is part of the Norwegian Protection Plan II3 and hence cannot be used for hydropower purposes. Generally, the vegetation can be characterized as northern boreal in a transition zone between oceanic and continental vegetation sections (Moen 1998, Bjørklund *et al.* 2015) in north-eastern Finnmark. The landscape along the river consists of hills with sparse heath vegetation near the coast, hilly landscapes and plateaus further inland. Along the river, the vegetation is commonly characterized by different vegetation types: closest to the river is often a well-developed sandy beach with key plants like the oyster plant (*Mertensia maritima* L.), followed by a transition zone with oyster plant and sea pea (*Lathyrus japonicus* Willd.), white sandy dunes with sand ryegrass [*Leymus arenarius* (L.) Hochst.] and sea pea, grey sandy dunes with sand ryegrass and red fescue (*Festuca rubra* L.), and dry, species-poor dune grass heath with strong hints of fringed pink (*Dianthus superbus* L.; <https://elvedelta.miljodirektoratet.no/delta-322.htm>, <https://www.biodiversity.no/NiN2.0/T21-C-2>). At the river outlet, where the sample was taken, there is also a semi-limestone beach meadow (Elven & Johansen 1983), which is characterized by freshwater-affected salt weed (*Salicornia* sp.) communities (P.E. Aspholm, NIBIO, Svanhovd; pers. comm.). The sampling location was in an area that is considered brackish water area (<https://elvedelta.miljodirektoratet.no/delta-322.htm>).

Sampling and studying of mites

The adults of *Z. svanhovdi* A. Seniczak *et al.* sp. nov. used in this study were collected by A. Fjellberg on 26.08.2021 from Troms and Finnmark county, Northern Norway, Sør-Varanger community and Gr. Jakobselv locality, at river outlet (69.71258° N, 30.87123° E, 0.5 m a. s. l.). One collective sample was taken on the shore of Jacob's River, in the transect from the river edge to more distant part from it (about 10 m long, Figure 1). This sample was extracted in Berlese funnel for one week. We investigated the density, sex ratio, number of gravid females and carried eggs, and body length and width, based on 30 randomly selected specimens. We measured a total body length (tip of rostrum to posterior edge of notogaster) in lateral aspect and body width (widest part of notogaster) in dorsal aspect, and size of anal and genital openings and setae perpendicularly to their length in µm.

The illustrations are limited to the dorsal, lateral and ventral aspect and some leg segments of adult, and palp and chelicera. Illustrations were prepared from individuals mounted temporarily in lactic acid. In the text and figures we used the following abbreviations: rostral (*ro*), lamellar (*le*), interlamellar (*in*) and exobothridial (*ex*) setae, lamella (*La*), translamella (*Tr*), bothridium (*bo*), bothridial seta (*bs*), tutorium (*Tut*), pteromorph (*Ptm*), notogastral setae (*c₂*, *l*-, *h*-, *p*-series), porose areas (*Aa*, *A1*, *A2*, *A3*), lyrifissures (*ia*, *im*, *ih*, *ip*, *ips*, *iad*), circumpedal carina (*cp*), opisthonotal gland opening (*gla*), subcapitular setae (*a*, *m*, *h*), pedotectum (*Pd*), custodium



Figure 1 General view of the river shore, where *Zachvatkinibates svanhovdi* A. Seniczak et S. Seniczak sp. nov. was collected (photo: Arne Fjellberg).

(*cus*), discidium (*Dis*), genal tooth (*gt*), cheliceral setae (*cha*, *chb*), Trägårdh organ (*Tg*), palp setae (*sup*, *inf*, *l*, *d*, *cm*, *acm*, *lt*, *vt*, *ul*, *su*) and solenidion ω , epimeral setae (*1a-c*, *2a*, *3a-c*, *4a-c*), genital (*g*) and aggenital (*ag*) setae, adanal and anal setae (*ad-*, *an-series*), leg solenidia (σ , φ , ω), famulus (ε) and setae (*bv*, *ev*, *d*, *l*, *ft*, *tc*, *it*, *p*, *u*, *a*, *s*, *pv*, *pl*, *v*). Terminology used follows that of Grandjean (1949, 1953) and Norton and Behan-Pelletier (2009). The species nomenclature follows Subías (2004).

For scanning electron microscopy (SEM), four mites were air-dried and coated with Au/Pd in a Polaron SC502 sputter coater and placed on Al-stubs with double-sided sticky carbontape. Observations and micrographs were made with a ZEISS Supra 55VP scanning electron microscope.

DNA barcoding

Five specimens of *Z. svanhovdi* were used for DNA barcoding. Each specimen was photographed, and the photos are available at Barcode of Life Data System (BOLD, <http://boldsystems.org>). The specimens were subsequently placed in a well containing 50 ml of 90% ethanol in a 96-well microplate and sent to the Canadian Centre for DNA Barcoding (CCDB 2021). Mites were sequenced for the barcode region of the COI gene according to standard protocols at CCDB (<http://www.ccdb.ca>), using either LepF1/LepR1 (Hebert *et al.* 2003) or LCO1490/HCO2198 (Folmer *et al.* 1994) primer pairs. The DNA extracts were placed in archival storage at -80 °C at the Canadian Centre for DNA Barcoding. The sequences are available in GenBank (accession numbers in the section on the Type material). The voucher specimens are stored in the Entomological Collections at the Department of Natural History, University Museum of Bergen, University of Bergen, Bergen, Norway (ZMBN).

Obtained COI sequences were blasted against GenBank to detect and exclude possible contaminations and ensure high quality. Sequence variation within *Z. svanhovdi* specimens was calculated in BOLD using Kimura 2 Parameter distance model, pairwise deletion, and BOLD Aligner (Amino Acid based HMM).

Results

Zachvatkinibates svanhoaudi A. Seniczak et S. Seniczak sp. nov.

Zoobank: A6272084-12BA-4FAC-BA43-7582C1D30CEB

(Figures 2–11)

Diagnosis

Adult of medium size (449–507), with generic characters of *Zachvatkinibates* given by Behan-Pelletier & Eamer (2005). Prodorsal seta *in* long, translamella narrow, notogastral setae short and distally pliable, notogastral porose areas usually oval and of medium size, but *Aa* can be larger, especially in males. Postanal porose area *Ap* absent, dorsal crest on tarsus I present.

Description of adults

Measurements — Holotype (female) length 494, and width 299. Mean length (and range) of females 490.3 ± 7.1 (481–507, n = 15) and males 468.9 ± 12.2 (449–481, n = 15), mean maximum width (and range) of females 299.0 ± 6.2 (293–319) and males 276.7 ± 8.3 (254–293).

Integument — Brown and microporose.




Figure 2 *Zachvatkinibates svanhoaudi* A. Seniczak et S. Seniczak sp. nov., scale bars 50 µm. A – female, dorsal aspect, legs partially drawn; B – porose area *Aa* of another female (enlarged); C – notogaster of male; D – shape of porose area *Aa* of another male.




Figure 3 *Zachvatkinibates svanhovdi* A. Seniczak et S. Seniczak sp. nov., female. A – ventral aspect, legs partially drawn, scale bar 50 µm; B – lateral aspect, legs partially drawn, scale bar 50 µm, mouthparts, right side, scale bars 20 µm; C – chelicera (Trägårdh organ in transparent area); D – palp.

Prodorsum — Sub-triangular, rostrum rounded. Seta *ro* relatively long (47–52), laterally barbed and in lateral position (Figures 2A, 3A, 3B, 5A, 5D, 6A–C, 8C, 8D, 9A, 10A), seta *le* longer (62–66), and seta *in* longest (77–80), both finely barbed; all acuminate; seta *ex* short (17) and smooth. Lamella long (77–110), narrow at base and wider in place of connection with translamella, translamella of medium size (29–31), narrow, lamellar cusp (length 15) rounded. Bothridium with well-developed medial scale, bothridial seta of medium size (43–46), clavate with smooth stalk and barbed head (Figures 2A, 3B, 5A, 5B, 5D, 6A, 6B, 6D, 7B, 7C, 9C).

Notogaster — Longer (females 380–398, males 357–378) than wide (females 293–319, males 254–293) with well-developed movable pteromorphs. Notogastral setae (10 pairs, including *c*₂) short (15–26), and smooth (Figures 2–3B, 4, 5, 6A, 6D, 7A–C, 8A, 9C, 9D). Lyrifissure *ia* posterolateral to seta *c*₂, *im* anterior to seta *h*₃, *ip* anterolateral to seta *p*₁, *ips* anterior to seta *p*₃, *ih* posterior to seta *la*, and opisthonotal gland opening *gla* anterolateral to seta *h*₃ (Figures 2A, 8D). Shape of porose areas (4 pairs) differs between females and males. In females, porose areas oval, *Aa* slightly larger than other porose areas (Figures 2, 3B), in some individuals *Aa* can be elongated. In 15 males investigated, porose areas clearly larger (Figure 2C, 2D) than in females, and in some individuals *Aa* can be elongated, and of crescent shape or divided in two parts. Generally, in both sexual forms shape of other porose areas slightly varies within individuals, and postanal porose area is absent. Posterior tectum of notogaster is broken in middle and rounded ends overlap (Figure 4).

Gnathosoma — Subcapitular setae *h*, *m* and *a* short (18–24) and smooth (Figures 3A, 3B,




Figure 4 *Zachvatkinibates svanhovdi* A. Seniczak et S. Seniczak sp. nov., female, posterior aspect, scale bar 50 µm.

5C, 7B–D, 8C, 8D). Palp short (75), setae relatively short and smooth (Figures 3D, 5D, 9A, 10A), formula of palp setae [trochanter to tarsus (+ solenidion ω)]: 0-2-1-3-9(1). Chelicera longer (96), seta *cha* longer than *chb*, both barbed (Figures 3C, 9A, 10A). Axillary saccule short (12).

Ventral and lateral regions — Most epimeral setae short and smooth (10–13, Figures 3A, 5C, 8B, 8C). Genital setae (six pairs) as short as epimeral setae, *g*₁₋₃ in anterior position. Aggenital setae (one pair) also short, adanal setae longer (three pairs, 22–29) and anal setae longest (two pairs, 29–32), all smooth (Figures 3A, 3B, 4, 5B, 5C, 8A, 8B). Lyrifissure *iad* short, located lateral to anterior part of anal plates. Discidium large, triangular; custodium long (40–42), narrow and pointed; tutorium long (122–126), narrow at base and distally wider and pointed (Figures 3B, 7B–D, 9A, 9B, 10, 11). Pedotectum I large (73 x 30), narrow at base and concave dorsally (Figure 3B), and with longitudinal striae, genal tooth of medium size (30–22). Circumpedal carina well formed. Humerosejugal porose areas *Am* and *Ah* oval.

Legs — Trochanters III and IV and all femora flattened with ventral carina. Porose areas on paraxial side of trochanters III and IV and femora I–IV. Femur I with longitudinal striae, most leg setae barbed (Figures 5, 6, 7, 8B–D, 9A–C). Seta *ft*" on tarsus I relatively long, solenidia ω_1 and ω_2 of similar length, famulus short; solenidia ω_1 and ω_2 and famulus ε located on small dorsal crest. Solenidia ω_1 and ω_2 on tarsus II relatively long and of similar size. Formulae of leg setae [trochanter to tarsus (+ solenidia)]: I – 1-4-3(1)-4(2)-20(2); II – 1-4-3(1)-4(1)-15(2); III – 2-3-1(1)-3(1)-15; IV – 1-2-2-3(1)-12. Tarsi heterotridactylous. Homology of leg setae and solenidia given in Table 1.




Figure 5 *Zachvatkinibates svanhoaudi* A. Seniczak et S. Seniczak **sp. nov.** adult, SEM micrographs. A – dorsal view; B – lateral view; C – ventral view; D – frontolateral view.

Results of DNA barcoding

The barcode based on the cytochrome oxidase I (COI) mitochondrial DNA region (length: 658 bp) of five specimens of the new species is provided under GeneBank accession numbers: OP824977 (SUB2896631 SOITS201-22), OP824978 (SUB2896631 SOITS202-22), OP824979 (SUB2896631 SOITS203-22), OP824980 (SUB2896631 SOITS205-22), OP824981

Table 1 Leg setae (Roman letters) and solenidia (Greek letters) in *Zachvatkinibates svanhoaudi* A. Seniczak et S. Seniczak **sp. nov.**

Legs	Trochanter	Femur	Genu	Tibia	Tarsus
I	v'	$d, bv'', (l)$	$(l), \sigma, v'$	$(l), (v), \varphi_1, \varphi_2$	$l'', (ft), (tc), (it), (pl), (p), v', (pv), (a), s, (u), \varepsilon, \omega_1, \omega_2$
II	v'	$d, bv'', (l)$	$(l), \sigma, v'$	$(l), (v), \varphi$	$(ft), (tc), (it), (p), (pv), s, (a), (u), \omega_1, \omega_2$
III	v', l'	d, ev', l'	l', σ	$(v), l', \varphi$	$(ft), (tc), (it), (p), (pv), s, (a), (u)$
IV	v'	d, ev'	d, l'	$(v), l', \varphi$	$ft'', (tc), (u), (pv), s, (a), (p)$

Note: pairs of setae in parentheses.




Figure 6 *Zachvatkinibates svanhoaudi* A. Seniczak et S. Seniczak **sp. nov.** adult, SEM micrographs. Dorsal view. A – anterior and medial part; B, C – anterior part; D – medial and posterior part.

(SUB2896631 SOITS204-22). The maximum variation within the species was 2.41% (p-dist) that is expected intraspecific variation of COI in oribatids. (e.g., Pfingstl *et al.* 2019, Seniczak *et al.* 2019). These are the first public COI sequences to be reported for the genus *Zachvatkinibates* and therefore, genetic distances to other species could not be calculated.

Distribution, ecology and biology

Zachvatkinibates svanhoaudi was found in Troms and Finnmark county, Northern Norway, in a collective sample, taken in the transect on the shore of the river, so it is not possible to give precisely the species microhabitat. Among the 30 individuals investigated, collected at the end of August, the sex ratio (females to males) was 1:1, and no females were gravid and carried eggs.

Type material

The holotype (female) and four paratypes (2 females and 2 males) are deposited in the University Museum of Bergen, University of Bergen, Bergen, Norway (ZMBN).




Figure 7 *Zachvatkinibates svanhovdi* A. Seniczak et S. Seniczak sp. nov. adult, SEM micrographs. Lateral view. A – posterior and medial part; B – anterior and medial part; C, D – anterior part.

Etymology

The species is named in honour of the biological station NIBIO Svanhovd, which is located in eastern Finnmark in the Pasvik valley, where the project leading to the collection and description of this new species was carried out.

Comparison of morphology of *Zachvatkinibates svanhovdi* with congeners and remarks

Seniczak et al. (2020d) compared selected morphological characters of adults of 15 *Zachvatkinibates* species and stated that the largest among them is *Z. nortoni* Behan-Pelletier et Eamer, 2005, and smallest is *Z. anoporus* (Mahunka, 2006), and the body length of several species overlaps. These authors also compared the length of prodorsal seta *in* and notogastral setae, shape of translamella and some porose areas. To this list *Z. erimo* Shimano et Aoki, 2019 should be added. Based on this list of species, *Z. svanhovdi* A. Seniczak et S. Seniczak sp. nov. is the most similar to *Z. quadrivertex* (Halbert, 1920), but differs from it mainly by the shape of notogastral setae, posterior tectum of notogaster and lack of postanal porose area *Ap*, which in *Z. quadrivertex* is present. In *Z. svanhovdi*, most notogastral setae are distally pliable,




Figure 8 *Zachvatkinibates svanhovdi* A. Seniczak et S. Seniczak **sp. nov.** adult, SEM micrographs. Ventral view. A – posterior part; B – medial part; C, D – anterior part.

whereas in *Z. quadrivertex* they are not. In both species, the posterior tectum of notogaster is broken in the middle, the ends are rounded, but in *Z. svanhovdi* they overlap, whereas in *Z. quadrivertex* they are separated by a certain distance. In *Z. svanhovdi*, lyrifissures *ips* and *ih* are placed at larger distances from seta *p* than in *Z. quadrivertex*. Moreover, in female of *Z. svanhovdi* porose area *Aa* is located between setae *la* and *lm* (vs. anteromedial to seta *la* in *Z. quadrivertex*), and porose areas *A2* and *A3* are as large as *A1*, whereas in *Z. quadrivertex* they are smaller than *A1*. In some males of *Z. svanhovdi*, the porose areas *Aa* are larger than in *Z. quadrivertex*. In the diagnosis for *Zachvatkinibates* by Behan-Pelletier & Eamer (2005), postanal porose area *Ap* is present, but *Z. svanhovdi* has no porose area, so the diagnosis for this genus should be changed to “postanal porose area *Ap* present or absent”.

In *Z. svanhovdi*, sexual dimorphism is observed in the body size and octotaxic system (Table 2). Males are slightly smaller than females, have larger porose areas, and *Aa* can be divided in two parts (vs. in females it is uniform). In six other species of *Zachvatkinibates*, sexual dimorphism is more distinct than in *Z. svanhovdi* (Table 2). Interestingly, the species listed in Table 2 seem to be correlated with littoral habitats of cold climate zones (Behan-Pelletier & Eamer 2010, Behan-Pelletier 2015).




Figure 9 *Zachvatkinibates svanhoedi* A. Seniczak et S. Seniczak **sp. nov.** adult, SEM micrographs. A, B – anterior part; frontolateral view; C – bothridial setae; D – gla opening.

The adult of *Z. svanhoedi* has the dorsal crest on tarsus I, as other species of *Zachvatkinibates* (Shal'dybina & Grishina 1987, Behan-Pelletier 1988, Bayartogtokh & Aoki 1998, Behan-Pelletier & Eamer 2005, Seniczak *et al.* 2020d). In *Minunthozetes* Hull, 1916, the leg crests are better developed than in *Zachvatkinibates* and occur on part of tibiae and tarsi I, II and IV (Seniczak & Seniczak 2018, Seniczak *et al.* 2018) and may suggest close relationship of *Zachvatkinibates* with *Minunthozetes*. In other genera of Puncitoribatidae such as *Mycobates* Hull, 1916 and *Puncitoribates* Berlese, 1908, the leg crests on tibiae and tarsi are absent (Seniczak & Seniczak 2008, Seniczak *et al.* 2015, 2020c). The latter two genera have dorsal projection on tibia II, which in *Zachvatkinibates* and *Minunthozetes* is absent.

Based on the checklists of Oribatida of Norway (Mehl 1979), Sweden (Lundqvist 1987), and Finland (Niemi *et al.* 1997), and other publications listed in Seniczak and Seniczak (2020) and Seniczak *et al.* (2020a,b, 2021), *Z. svanhoedi* represents the first finding of *Zachvatkinibates* in Norway and Fennoscandia. Species of *Zachvatkinibates* are known from Central and Eastern Asia, Northern America, and only few from Europe (Table 4). Their ecology varies a lot. For example, six species inhabit littoral zone and beach debris, other were found in bogs, forests, including forest canopy, and even in the debris in the alpine zone (Table 4).




Figure 10 *Zachvatkinibates svanhoaudi* A. Seniczak et S. Seniczak **sp. nov.** adult, SEM micrographs. Leg I. A – frontolateral view; B – lateral view; C – legs I–III; D – leg III and VI.

Comments on Puncitorbatidae in Fennoscandia and new record of *Mycobates carli*

The family Puncitorbatidae to which *Zachvatkinibates* belongs to is represented in Fennoscandia by four genera and ten nominative species, including *Mycobates carli* (Schweizer, 1922) which is reported here as new to Norway and Scandinavian part of Fennoscandia (Table 3). It was found in alpine tundra in Finse (Vestland, Western Norway, 60°35'14.0"N 7°30'50.2"E, 1228 m a. s. l., 06.08.2022, leg. Anna Seniczak), on forest floor with mosses and lichens. The description of climatic conditions in Finse was given earlier (Seniczak & Seniczak 2021). This species was represented in the sample only by few adults, (density 10 individuals per 500 cm³), and co-occurred with more abundant *M. sarekensis* (Trägårdh, 1910).

Mycobates carli has been mostly considered a southern European (Subías 2004, 2022), European (Weigmann 2006) or Eurosiberian species (Honciuc 1993), known especially from the Central, Southern, Eastern, and South-Eastern parts of Europe. It has been reported from numerous localities in the Alps (Fischer & Schatz 2020, Schatz & Bruckner 2021), the Giant Mountains (Materna 2000, Starý 2006, Miko 2013), Carpathian Mts. (Nae & Băncilă 2017, Skubala & Maslak 2009), and is also known from Caucasus Mts. (Schatz & Bruckner 2021).




Figure 11 *Zachvatkinibates svanhowdi* A. Seniczak et S. Seniczak sp. nov., leg segments of adult (part of femur to tarsus), right side, antiaxial aspect, setae on the opposite side not illustrated are indicated in the legend, scale bar 20 μm . A – leg I, tarsus (pl'); B – leg II; C – part of tibia and tarsus I; D – leg III, tibia (v''); E – leg IV.

Table 2 Selected morphological characters of *Zachvatkinibates* species with distinct dimorphism of notogastral porose areas; ♀ – females, ♂ – males.

Species	Body length (μm)	Seta in reaching rostrum	Shape of Aa	Size of porose areas			Setae ad-, an-series
				A1	A2	A3	
<i>Z. erimo</i> Shimano et Aoki 2019	♀440–449 ♂450	yes	rounded, as large as in male	as large as Aa	as large as A1 A2 and A3 fused	smaller than A2	as long as ag
<i>Z. nortoni</i> Behan-Pelletier et Eamer, 2005	♀525–594 ♂525–574	yes	rounded, as large as in male	smaller than Aa	as large as A1 A1, A2 and A3 fused	as large as A2	as long as ag
<i>Z. quadrivertex</i> (Halbert, 1920)	♀450–500 ♂415–480	yes	oval, smaller than in male elongated	as large as Aa smaller than Aa	smaller than A1 as large as A1	larger than A2 as large as A2	?
<i>Z. shaldybinae</i> Behan-Pelletier et Eamer, 2005	♀499–528 ♂470–508	no	rounded, smaller than in male elongated or divided	smaller than Aa smaller than Aa	as large as A1 smaller than A1	as large as A2 larger than A2	?
<i>Z. schatzi</i> Behan-Pelletier et Eamer, 2005	♀528–554 ♂505–544	no	elongated, smaller than in male	smaller than Aa	smaller than A1 A1, A2 and A3 fused	larger than A2	?
<i>Z. maritimus</i> Shaldybina, 1973	♀473 ♂?	yes	irregular, smaller than in male elongated	smaller than Aa smaller than Aa	smaller than A1 as large as A1	larger than A2 as large as A2	as long as ag
<i>Z. svanhovdi</i> A. et S. Seniczak sp. nov.	♀481–507 ♂449–481	yes	oval, smaller than in male elongated or divided	smaller than Aa smaller than Aa	as large as A1 as large as A1	as large as A2 as large as A2	longer than ag

In the north of Europe it has only been reported from plain tundra of the Kola Peninsula (Leonov & Rakhleeva 2015). It is considered subalpine and alpine (Weigmann 2006, Fischer & Schatz 2007), predominantly lichenicolous (Materna 2000), and muscicolous (Schatz & Bruckner 2021). It has been found in a peatland, 870 m a. s. l. (Stary 2006), at dry limestone plateau, 550 m a. s. l. (Lazarus & Krisper 2014), and in soils polluted with heavy metals in Romania, where it was associated with Cu, As and Mn (Manu et al. 2019). Most often it has been collected from soil or ground vegetation, but has been also reported from the unique microhabitat of the intermediate layer between the base of the soil and the bedrock, called “mesovoid shallow substratum” (Nae & Băncilă 2017) and from decaying wood (Skubała & Maslak 2009).

Acknowledgements

We thank the anonymous reviewers for helpful comments and suggestions that improved the quality of this paper. We also express our gratitude to Dr. Arne Fjellberg for collection of the sample that contained the specimens. Further, we would like to thank Juho Vuolteenaho for

Table 3 Occurrence of Puncitoribatidae in Fennoscandia, N – Norway, S – Sweden, F – Finland; * – new to Norway and Fennoscandia.

Genus	Species	Country
<i>Minunthozetes</i> Hull, 1916	<i>M. pseudofusiger</i> (Schweizer, 1922)	F, N
	<i>M. semirufus</i> (C.L. Koch, 1841)	F, S, N
	<i>M. carli</i> (Schweizer, 1922)*	N
<i>Mycobates</i> Hull, 1916	<i>M. parmeliae</i> (Michael, 1884)	F, S, N
	<i>M. sarekensis</i> (Trägårdh, 1910)	F, S, N
	<i>M. tridactylus</i> Willmann, 1929	N
<i>Puncitoribates</i> Berlese, 1908	<i>P. hexagonus</i> Berlese, 1908	F
	<i>P. punctum</i> (C.L. Koch, 1839)	F, S, N
	<i>P. sellnicki</i> Willmann, 1928	F, S, N
<i>Zachvatkinibates</i> Shaldybina, 1973	<i>Z. svanhovdi</i> A. et S. Seniczak sp. nov.*	N

Table 4 Distribution and ecology of *Zachvatkinibates* species.

Species	Distribution	Habitat	Sexual dimorphism	Reference
<i>Z. anoporus</i> (Mahunka, 2006)	Romania (Poiana Braziilor, Ocna Șugatag, Muntii Piatra, 900 m a. s. l.)	Bog, decaying wood, some <i>Sphagnum</i>	Unknown	Mahunka (2006)
<i>Z. conjunctus</i> Shal'dybina, 1987	Russia (Stolby Nature Reserve, Krasnoyarsky Krai)	Moss cushion at the base of a pine tree	Unknown	Shal'dybina & Grishina (1987)
<i>Z. eoeryi</i> (Mahunka, 1972)	South Europe: Croacia (near Split), Portugal (Lagoon of Faro)	Predominantly marine littoral debris, e.g., algae, detritus	Unknown	Mahunka (1972), Weigmann (2009)
<i>Z. epiphytos</i> Behan-Pelletier, Eamer et Clayton, 2001	Canada (Montane Alternative Silvicultural Systems, Vancouver Island), USA (Willamette National Forest, Oregon)	Epiphytes (mainly lichens) in coniferous forest canopy	No	Behan-Pelletier et al. (2001), Winchester et al. (2008)
<i>Z. erimo</i> Shimano et Aoki, 2019	Japan (Hyakuninhamama, Shono, Erimo City, Horoizumi)	Roots of Poaceae, dead parts of plants, soil on a sandy beach	Yes	Shimano & Aoki (2019)
<i>Z. latilamellatus</i> Bayartogtokh et Aoki, 1998	Mongolia	Siberian larch (<i>Larix sibirica</i> Ledeb.) forest	Unknown	Bayartogtokh & Aoki (1998), Seniczak et al. (2020b)
<i>Z. lobatus</i> (Hammer, 1977)	Pakistan (Lowari Pass, 3400 m a. s. l.)	Thin moss on the soil, grass, <i>Potentilla</i> , <i>Epilobium</i> , <i>Fragaria</i> , <i>Geranium</i> , pine cones	Unknown	Hammer (1977)
<i>Z. maritimus</i> Shal'dybina, 1973	Russia (Primorsky krai), USA (Alaska), Canada (Yukon, Vancouver Island, British Columbia)	Coastal debris, supratidal meadows, coastal and lentic littoral habitats	Yes	Shal'dybina (1973, 1987), Behan-Pelletier (1988), Ryabinin (2015)
<i>Z. mongolicus</i> Bayartogtokh, 2003	Mongolia (Gobi Gurvan Saykhan Nature Reserve, southern Mongolia)	Semidesert, soil under <i>Artemisia santolinifolia</i> (Pamp.) Turcz. ex Krasch and organic debris accumulated	No	Bayartogtokh (2003)
<i>Z. nemoralis</i> Shal'dybina, 1973	Kyrgyzstan and Mongolia	Litter between roots, the alpine region	Unknown	Shal'dybina (1973, 1987)
<i>Z. nortoni</i> Behan-Pelletier et Eamer, 2005	Canada (British Columbia)	Beach debris	Yes	Behan-Pelletier & Eamer (2005)
<i>Z. quadrivertex</i> (Halbert, 1920)	Atlantic coastal regions of Europe, from Denmark to South Portugal, Russia (Islands of White Sea, Far East: Kunashir Island in Kuril Islands)	Lower tidal level of marine saltmarshes, fir forest	Yes	Shal'dybina (1987), Weigmann (2009), Ryabinin (2015)
<i>Z. schatzii</i> Behan-Pelletier et Eamer, 2005	Canada (British Columbia)	Beach debris	Yes	Behan-Pelletier & Eamer (2005)
<i>Z. shal'dybinae</i> Behan-Pelletier et Eamer, 2005	Canada (eastern and northeastern), Greenland	Littoral	Yes	Behan-Pelletier & Eamer (2005), Makarova & Behan-Pelletier (2015)
<i>Z. tetrasklerosis</i> Behan-Pelletier, 1988	USA (California)	Litter of different species of conifers	No	Behan-Pelletier (1988)
<i>Z. volgini</i> Shal'dybina, 1973	Russia (Far East: Shikotan and Yuri islands in Kuril Islands, Sakhalin)	Gull's nest on the coast, quackgrass [<i>Elymus repens</i> (L.) Gould] and brier meadow	Unknown	Shal'dybina (1973, 1987), Ryabinin (2015)

assistance with field work. The study was supported by the grant to NIBIO from the Norwegian Taxonomy Initiative (Grant No. 9-21, 70184244). The DNA sequencing was financed by Norwegian Barcode of Life (NorBOL).

ORCID

Anna Seniczak  <https://orcid.org/0000-0003-0224-5397>

Stanisław Seniczak  <https://orcid.org/0000-0001-7393-5606>

References

- Bayartogtokh B. 2003. Four species of oribatid mites (Acari: Oribatida) from Central and Southern Mongolia, with notes on the genera *Montizetes* and *Zachvatkinibates*. Mong. J. Biol. Sci., 1(1): 3-19. <https://doi.org/10.22353/mjbs.2003.01.01>
- Bayartogtokh B., Aoki J. 1998. Two new species of oribatid mites of the family Mycobatidae (Acari: Oribatei) from Mongolia. Edaphologia, 60: 37-44.
- Behan-Pelletier V.M. 1988. Redefinition of *Zachvatkinibates* (Acari: Mycobatidae), with description of a new species. Can. Entomol., 120: 797-813. <https://doi.org/10.4039/Ent120797-8>
- Behan-Pelletier V.M. 2015. Review of sexual dimorphism in brachypyline oribatid mites. Acarologia, 55(2): 127-146. <https://doi.org/10.1051/acarologia/20152163>
- Behan-Pelletier V.M., Eamer B. 2005. *Zachvatkinibates* (Acari: Oribatida: Mycobatidae) of North America, with descriptions of sexually dimorphic species. Can. Entomol., 137: 631-647. <https://doi.org/10.4039/n05-055>
- Behan-Pelletier V.M., Eamer, B. 2008. Mycobatidae (Acari, Oribatida) of North America. Can. Entomol., 140: 73-110. <https://doi.org/10.4039/n07-027>
- Behan-Pelletier V.M., Eamer B. 2010. The first sexually dimorphic species of *Oribatella* (Acari, Oribatida, Oribatellidae) and a review of sexual dimorphism in the Brachypylina. Zootaxa, 2332: 1-20. <https://doi.org/10.11646/zootaxa.2332.1.1>

- Behan-Pelletier V.M., Eamer B., Clayton M. 2001. Mycobatidae (Acari: Oribatida) of Pacific Northwest canopy habitats. *Can. Entomol.*, 133: 755-776. <https://doi.org/10.4039/Ent133755-6>
- Berlese A. 1908. Elenco di generi e specie nuovi di Acari. *Redia*, 5: 1-15.
- Bjørklund P.K., Rekdal Y., Strand G.-H. 2015. Arealregnskap for utmark. Arealstatistikk for Finnmark. Ressursoversikt fra skog og landskap. 97 pp.
- BOLD. <http://boldsystems.org>
- CCDB 2021. The Canadian Centre for DNA Barcoding website. Available: <https://www.ccdb.ca> (Accessed October 2021).
- Elven R., Johansen V. 1983. Havstrand i Finnmark. Flora, vegetasjon og botaniske verneverdier. Miljøverndepartementet. Rapport T-541.
- Fischer B.M., Schatz H. 2007. Hornmilben (Acari: Oribatida). *Gredleriana*, 7: 414-456.
- Folmer O., Black M., Hoeh W., Lutz, R., Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Marine Biol. Biotechnol.*, 3(5): 294-299.
- Grandjean F. 1949. Formules anales, gastronotiques, génitales et aggénitales du développement numériques des poils chez les Oribates. *Bull. Soc. Zool. Fr.*, 74: 201-225.
- Grandjean F. 1953. Essai de classification des Oribates (Acariens). *Bull. Soc. Zool. Fr.*, 78: 421-446.
- Halbert J.N. 1920. The Acarina of the Seashore. *Proc. R. Ir. Acad.*, 35(B): 105-152.
- Hammer M. 1977. Investigations on the oribatid fauna of North-West Pakistan. *K. dansk. Vidensk. Skr.*, 21(4): 1-108.
- Hebert P.D.N., Cywinska A., Ball S.L., deWaard J.R. 2003. Biological identifications through DNA barcodes. *Proc. Royal Soc. B*, 270: 313-321. <https://doi.org/10.1098/rspb.2002.2218>
- Honciu V. 1993. Researches of oribatid fauna (Acari, Oribatida) in different terrestrial ecosystems in the Danube Delta. *Rev. roum. biol., Sér. biol. anim.*, 38(1): 21-29.
- Hull J.E. 1916. Terrestrial Acari of the Tyne Province, I. Oribatidae. *Trans. Nat. Hist. Soc. Northumbria* (n.s.), 4: 381-410.
- Koch C.L. 1839. Deutschlands Crustaceen, Myriapoden und Arachnidien. Regensburg: Bd. 28-30.
- Koch C.L. 1841. Deutschlands Crustaceen, Myriapoden und Arachnidien. Regensburg: Bd. 31-32.
- Lazarus S., Krisper, G. 2014. Diversity of the oribatid mite fauna (Acari, Oribatida) in two dry meadows in Styria (Austria). *Soil Org.*, 86(2): 117-124.
- Leonov V.D., Rakhee A.A. 2015. New information on similarity and difference between highland and plain tundras of the Kola Peninsula based on oribatid mite data (Acari: Oribatida). *Evraziat. Entomol. Z.*, 14(5): 489-499.
- Lundqvist L. 1987. Bibliografi och checklist över Sveriges oribatider (Acari: Oribatei). *Entomol. Tidskr.*, 108(1): 3-12.
- Mahunka S. 1972. *Punctoribates eoeryi* sp. n., eine neue Milbenart vom Meerstrand aus Jugoslawien (Acari, Oribatei). *Ann. Hist. Nat. Mus. Nat. Hung.*, 64: 355-357.
- Mahunka S. 2006. Oribatids from Maramures (Romania, Transylvania) (Acari: Oribatida). *Stud. Univ. Vasile Goldis Arad Ser. Stiint. Vietii*, 17: 59-75.
- Makarova O., Behan-Pelletier V. 2015. Oribatida (=Cryptostigmata, Beetle mites). In: Böcher J., Kristensen N.P., Pape T., Vilhelmsen L. (Eds.): *The Greenland Entomofauna. An Identification Manual of Insects, Spiders and Their Allies*. Brill, Leiden - Boston. p. 802-845.
- Manu M., Honciuc V., Neagoe A., Băncilă R.I., Iordache V., Onete M. 2019. Soil mite communities (Acari: Mesostigmata, Oribatida) as bioindicators for environmental conditions from polluted soils. *Sci. Rep.*, 9: 20250. <https://doi.org/10.1038/s41598-019-56700-8>
- Materna J. 2000. Oribatid communities (Acari: Oribatida) inhabiting saxicolous mosses and lichens in the Krkonoše Mts. (Czech Republic). *Pedobiologia*, 44: 40-62. [https://doi.org/10.1078/S0031-4056\(04\)70027-X](https://doi.org/10.1078/S0031-4056(04)70027-X)
- Mehl R. 1979. Checklist of Norwegian ticks and mites (Acari). *Fauna Norv. B.*, 26(1): 31-45.
- Michael A.D. 1884. British Oribatidae. Vol I. Ray Society, London: 1-336.
- Miko L. 2013. History of oribatid studies (Acarina, Oribatida) in the Krkonoše National Park (the Giant Mountains, Czech Republic), with a revised checklist of all known species of the Giant Mountains. *Opera Corcontica*, 50/S: 143-164.
- Moen A. 1998. Nasjonalatlas for Norge: Vegetasjon. Statens kartverk, Hønefoss.
- Nae I., Băncilă R.I. 2017. Mesovoid shallow substratum as a biodiversity hotspot for conservation. *Acarologia*, 57(4): 855-868. <https://doi.org/10.24349/acarologia/20174202>
- Niemi R., Karppinen E., Uusitalo M. 1997. Catalogue of the Oribatida (Acari) of Finland. *Acta Zool. Fenn.*, 207(1): 1-39.
- Norton R.A., Behan-Pelletier, V.M. 2009. Suborder Oribatida. Chapter 15. In: Krantz G.W., Walter D.E. (Eds.) *A Manual of Acarology*, Texas Tech University Press, Lubbock. p. 430-564.
- Pérez-Íñigo C. 1976. Acaros oribátidos de la Isla de Tenerife. (Acari, Oribatei). II Contribución. *EOS-Rev. Esp. Entomol.*, 51: 85-141.
- Pfingstl T., Lienhard A., Baumann J., Koblmüller S. 2021. A taxonomist's nightmare - Cryptic diversity in Caribbean intertidal arthropods (Arachnida, Acari, Oribatida). *Mol. Phylogenetic Evol.*, 163: 107240. <https://doi.org/10.1016/j.ympev.2021.107240>
- Ryabinin N.A. 2015. Oribatid mites (Acari, Oribatida) in soils of the Russian Far East. *Zootaxa*, 3914(3): 201-244. <https://doi.org/10.11646/zootaxa.3914.3.1>
- Schatz H. 2020. Hornmilben (Acari, Oribatida) von Vorarlberg - Allgemeine Verbreitung und Habitat-bindung. *Inatura - Forschung online*, 76: 9.
- Schatz H., Bruckner A. 2021. Hornmilben (Acari: Oribatida) aus dem Wildnisgebiet Dürrenstein und dem Urwald Rothwald (Österreich). *Silva Fera*, 8: 42-62.

- Schweizer J. 1922. Beiträge zur Kenntnis der terrestrischen Milbenfauna der Schweiz. Verh. Naturforsch. Ges. Basel., 23: 23-112. <https://doi.org/10.5962/bhl.part.3644>
- Seniczak A., Seniczak S. 2020. Diversity of oribatid mites (Acari: Oribatida) in the Svalbard archipelago: a historical overview, Zootaxa, 4834(1): 41-65. <https://doi.org/10.11646/zootaxa.4834.1.3>
- Seniczak A., Seniczak S. 2021. Morphological ontogeny and ecology of *Ceratozetes parvulus* (Acari: Oribatida: Ceratozetidae). Zootaxa, 5086(1): 111-134. <https://doi.org/10.11646/zootaxa.5086.1.9>
- Seniczak A., Seniczak S., Iturrondebeitia J.C., Solhoy T., Flatberg K.I. 2020a. Diverse *Sphagnum* mosses support rich moss mite communities (Acari, Oribatida) in mires of western Norway. Wetlands, 40, 1339-1351. <https://doi.org/10.1007/s13157-019-01236-w>
- Seniczak A., Seniczak S., Jordal B.H. 2019. Integrated taxonomy approach: molecular data and ontogeny studies clarify systematic status of *Chamobates borealis* (Acari, Oribatida). Syst. Appl. Acarol., 24(12): 2409-2426. <https://doi.org/10.11158/saa.24.12.9>
- Seniczak A., Seniczak S., Schwarzfeld M.D., Coulson S.J., Gwiazdowicz D.J. 2020b. Diversity and distribution of mites (Acari: Ixodida, Mesostigmata, Trombidiformes, Sarcoptiformes) in the Svalbard Archipelago. Diversity, 12(9): 323. <https://doi.org/10.3390/d12090323>
- Seniczak A., Seniczak S., Starý J., Kaczmarek S., Jordal B.H., Kowalski J., Roth S., Djursvoll P., Bolger T. 2021. High diversity of mites (Acari: Oribatida, Mesostigmata) supports the high conservation value of a broadleaf forest in eastern Norway. Forests, 12(8): 1098. <https://doi.org/10.3390/f12081098>
- Seniczak S., Seniczak A. 2008. Morphology of three European species of the genus *Puncitoribates* Berlese, 1908 (Acari: Oribatida: Mycobatidae). Ann. Zool., 58(3): 473-485. <https://doi.org/10.3161/000345408X364328>
- Seniczak S., Seniczak A. 2018. Morphological ontogeny of *Minunthozetes semirufus* (Acari, Oribatida, Puncitoribatidae). Zootaxa, 4540(1): 73-92. <https://doi.org/10.11646/zootaxa.4540.1.8>
- Seniczak S., Ivan O., Marquardt T., Seniczak, A. 2020c. Morphological ontogeny of *Puncitoribates astrachanicus* (Acari: Oribatida: Puncitoribatidae), and comments on *Puncitoribates* Berlese. Zootaxa, 4900(1): 043-061. <https://doi.org/10.11646/zootaxa.4900.1.6>
- Seniczak S., Seniczak A., Coulson S.J. 2015. Morphology, distribution and biology of *Mycobates sarekensis* (Acari: Oribatida: Puncitoribatidae). Int. J. Acarol., 41(8): 663-675. <https://doi.org/10.1080/01647954.2015.1102968>
- Seniczak S., Seniczak A., Kaczmarek S., Marquardt, T. 2018. Morphological ontogeny of *Minunthozetes pseudofusiger* (Acari, Oribatida, Puncitoribatidae) and comments on *Minunthozetes* Hull. Syst. Appl. Acarol., 23(6): 1155-1168. <https://doi.org/10.11158/saa.23.6.11>
- Seniczak S., Seniczak A., Kaczmarek S., Marquardt T., Sarsenova B. 2020d. Morphological ontogeny of *Zachvatkinibates latilamellatus* (Acari: Oribatida: Puncitoribatidae), and comments on *Zachvatkinibates* Shaldybina. Zootaxa, 4857(1): 97-116. <https://doi.org/10.11646/zootaxa.4857.1.6>
- Shaldybina E.S. 1973. New species of oribatid mites of the subfamily Minunthozetinae (Oribatei, Mycobatidae) from the territory of the Soviet Union. Zool. Zhurnal, 52(5): 689-699. (In Russian)
- Shaldybina E.S., Grishina L.G. 1987. Three new species of mites of subfamily Minunthozetinae (Oribatei, Mycobatidae). Insects, Mites and Helminths, Novosibirsk. p. 63-70. (In Russian)
- Shimano S., Aoki J. 2019. A new species of Japanese oribatid mite, *Zachvatkinibates erimo* sp. nov., showing sexual dimorphism (Acariformes: Oribatida: Puncitoribatidae). Zootaxa, 4647(1): 362-367. <https://doi.org/10.11646/zootaxa.4647.1.22>
- Skubala P., Maslak M. 2009. Succession of oribatid fauna (Acari, Oribatida) in fallen spruce trees: Deadwood promotes species and functional diversity. In: Sabelis M.W., Bruin J. (Eds.), Trends in Acarology. p. 123-128. https://doi.org/10.1007/978-90-481-9837-5_19
- Subías L.S. 2004. Listado sistemático, sinónímico y biogeográfico de los Ácaros Oribátidos (Acariformes, Oribatida) del mundo (1758-2002). Graellsia, 60 (número extraordinario): 3-305. <https://doi.org/10.3989/graelessia.2004.v60.iExtra.218>
- Subias, L.S. 2022. Listado sistemático, sinónímico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del mundo (excepto fósiles). Monografías electrónicas SEA, 12: 1-539. <http://www.sea-entomologia.org>
- Starý J. 2006. Contribution to the knowledge of the oribatid mite fauna (Acari, Oribatida) of peat bogs in Bohemian Forest. Silva Gabreta, 12: 35-47.
- Trägårdh I. 1910. Acariden aus dem Sarekgebirge. Naturw. Untersuch. des Sarekgebirges, 4: 375-586.
- Weigmann G. 2006. Hornmilben (Oribatida). Die Tierwelt Deutschlands, 76. Teil. Goecke & Evers, Keltern, 520 pp.
- Weigmann G. 2009. Oribatid mites (Acari: Oribatida) from the coastal region of Portugal. II. The genera *Zachvatkinibates* and *Puncitoribates* (Mycobatidae). Soil Org., 81(1): 85-105.
- Willmann C. 1928. Die Oribatidenfauna nordwestdeutscher und einiger süddeutscher Moore. Abh. Naturwiss. Ver. Bremen, 27(1): 143-176.
- Willmann C. 1929. Neue Oribatiden II. Zoologischer Anzeiger, 80(1-2): 43-46.
- Winchester N.N., Lindo Z., Behan-Pelletier V.M. 2008. Oribatid mite communities in the canopy of montane *Abies amabilis* and *Tsuga heterophylla* trees on Vancouver Island, British Columbia. Environ. Entomol., 37(2): 464-471. <https://doi.org/10.1093/ee/37.2.464>