



HAL
open science

New flavonoids from *Portulaca oleracea* L. and their activities

Xu Yang, Wenjie Zhang, Xixiang Ying, Didier Stien

► **To cite this version:**

Xu Yang, Wenjie Zhang, Xixiang Ying, Didier Stien. New flavonoids from *Portulaca oleracea* L. and their activities. *Fitoterapia*, 2018, 127, pp.257-262. 10.1016/j.fitote.2018.02.032 . hal-03945739

HAL Id: hal-03945739

<https://hal.science/hal-03945739v1>

Submitted on 18 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New flavonoids from *Portulaca oleracea* L. and their activities

Xu Yang^a, WenjieZhang^a, XixiangYing^a, DidierStien^b

^aSchool of Pharmacy, Liaoning University of Traditional Chinese Medicine, Dalian, PR China

^bSorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire de Biodiversité et Biotechnologies Microbiennes (LBBM), France

Abstract

Three new compounds, identified as (3S)-5-hydroxy-3-(2-hydroxybenzyl)-7-methoxychroman-4-one, oleracone C (**1**), 5-hydroxy-3-(2-hydroxybenzyl)-7-methoxy-4H-chromen-4-one, oleracone D (**2**), and 1-(2-hydroxy-4,6-dimethoxyphenyl)-3-(2-hydroxyphenyl)propan-1-one, oleracone E (**4**), together with one new natural product, 5,7-dimethoxy-4-O-2'-cycloflavan (**3**) already known from synthesis, and two known compounds, (2S)-5,2'-dihydroxy-7-methoxyflavanone (**5**) and 2',4'-dihydroxy-4,6'-dimethoxychalcone (**6**) were isolated from the *Portulaca oleracea* L. for the first time. Their structures were elucidated using spectroscopic methods, including one- and two-dimensional nuclear magnetic resonance, high-resolution electrospray ionization time-of-flight mass spectrometry and circular dichroism spectrometry. Oleracone C (**1**), D (**2**) and E (**4**) presented scavenging activities in 1,1-diphenyl-2-picryl-hydrazyl (DPPH) radical quenching assay, with IC₅₀ values of 23.27, 11.73, 13.17 μM, respectively and anticholinesterase activities with IC₅₀ values ranging between 59.55 and 78.85 μM.

Keywords

Portulaca oleracea L.

Flavonoids

Isolation

Antioxidant activity

Anticholinesterase activity

Abbreviations

DPPH, 1,1-diphenyl-2-picrylhydrazyl

BHA, butylated hydroxyl anisole

ATCI, acetylthiocholine iodide

AChE, acetylcholinesterase

DTNB, 5,5-dithiobis-2-nitrobenzoic acid

PBS, phosphate buffer saline

AD, Alzheimer disease

CAM, crassulacean acid metabolism

1. Introduction

Portulaca oleracea L. a herbaceous succulent annual plant, of which stems and leaves are edible with a slightly acidic and salty taste [1], belonging to the family of Portulacaceae, is widely distributed in tropical and subtropical areas [2]. *P. oleracea*, cold in nature and acid in flavor [3], is a nutritious vegetable crop, rich in essential ω -3 fatty acids, α -linolenic acid, α -tocopherol, ascorbic acid, linoleic acid, β -carotene, glutathione [4], and highest dietary minerals, such as K, Ca, Mg, P, Fe and Zn [5], which are essential for normal human healthy growth. As a result, the vigorous, invasive, yet edible plant *P. oleracea* is widely used in folk medicine or for other applications [6]. In view of its popularity, more and more pharmacological effects have been studied, including anti-inflammatory [7], analgesic [8], antibacterial [9], antioxidant [10], and neuroprotective [11] ones. More recently, *P. oleracea* being a crassulacean acid metabolism (CAM) photosynthetic plant [12], many scholars also concentrated on its constituents, such as phenolic acids [13], terpenes [14], coumarins [15], flavonoids [16] and alkaloids [17], in which flavonoids can be produced via its photosynthesis [18] that can also effectively prevent oxidative damage to the plant cells to enhance its antioxidant constituents [19]. In the study, three new flavonoids including oleracone C (**1**), oleracone D (**2**) and oleracone E (**3**) were found, especially, the absolute configuration of oleracone C was elucidated, (2*R*, 4*R*)-5,7-dimethoxy-4-*O*-2'-cycloflavan being the new natural product whose absolute configuration was also determined for the first time (Fig. 1). Flavonoids, with high antioxidation was related to many pharmacological effects, such as cardiovascular disease, senile dementia, cancer, etc. [20]. At the same time, flavonoids as acetylcholinesterase (AChE) inhibitor, can promote the production of acetylcholine effectively [21]. Therefore, the antioxidant and anticholinesterase activities of three new compounds isolated from *P. oleracea*, including oleracone C (**1**), D (**2**) and E (**3**) were studied.




Fig. 1. Chemical structures of compounds isolated from *P. oleracea* L.

2. Experimental

2.1. Chemical and reagents

1,1-Diphenyl-2-picrylhydrazyl radical (DPPH^{*}) (purity \geq 99%) was purchased from Sigma Co. (USA). Butylated hydroxyanisole (BHA) (purity \geq 98%) was purchased from Shanghai Xiangrui Biological Technology Co., Ltd. (Shanghai, China). Methanol and formic acid are all of HPLC grade provided by Damao Chemical Reagent Plant (Tianjin, China, purity \geq 99.9%). All other reagents are of analytical grade purchased from Jinfeng Chemical Factory (Tianjin, China), and the water is WAHAHA purified water (Shenyang, China). Acetylthiocholine iodide (ATCI) (purity \geq 99%) and acetylcholinesterase (AChE) (vitality \geq 200 units/mg protein) were purchased from Dalian Meilun Biotechnology Co., Ltd. (Dalian, China). Eserine (purity \geq 98%) was purchased from Shanghai Hanxiang Biotechnology Co., Ltd. (Shanghai, China). 5,5-dithiobis-2-nitrobenzoic acid (DTNB) (purity \geq 99%) was purchased from Shanghai Jinshui Biotechnology Co., Ltd. (Shanghai, China). Na₂HPO₄ and NaH₂PO₄ were purchased from Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China) (purity \geq 99.0%).

2.2. Plant material

The whole herbs of *P. oleracea* were collected in Shijiazhuang (Hebei, China) in June 2014, and identified by Prof. Xixiang Ying. Voucher specimens (No. 20140312) were deposited at School of Pharmacy, Liaoning University of Traditional Chinese Medicine.

2.3. General experimental procedures

In the separation process, column chromatography (CC) included silica gel (100–220 and 200–300 mesh, Qingdao Marine Chemical Co., Qingdao, China), polyamide resin (80–100 mesh, Taizhou Luqiao Sijia Biochemical Plastic Factory, Zhejiang, China) and ODS (20–40 μ m, GE Healthcare, Marlborough, MA). TLC was performed on silica gel GF₂₅₄ (Qingdao Marine Chemical Co., Qingdao, China).

2.4. Equipment

The UV spectrum and data of absorbance were determined by using a HITACHI U-3010 spectrophotometer (Hitachi Ltd., Tokyo, Japan). The IR spectrum was obtained by using an IR200 spectrophotometer (Thermo Electron Corporation, Waltham, MA). The Circular dichroism spectra were obtained using J-810 Circular Dichroism spectrometer (JASCO Corporation, Japan). Optical rotations were measured with Autopol I automatic polarimeter (Rudolph Research Analytical, Hackettstown, NJ). The NMR spectra were recorded using an AVANCE 500 MHz instrument (Bruker Corporation, Switzerland). All compounds were dissolved in DMSO-*d*₆ or CDCl₃. Relative molecular mass were recorded by using a 6520 quadrupole-time-of-flight mass spectrometer (Agilent, Palo Alto, CA). Purity was detected on a Nexera X2 UHPLC LC-30A system (Shimadzu, Kyoto, Japan), using a Kromasil C18 column (150 mm \times 4.6 mm, 5 μ m, Dalian Johnson Separation Science and Technology Corporation). The oven temperature was maintained at 40 °C. Water containing 0.1% formic acid served as solvent system A, and methanol served as solvent system B. The flow rate was 1.0 mL/min, and injection volume was 5 μ L. 96-well microplate reader (HBS-1096A) was purchased from Nanjing Detie Experimental Equipment Co., Ltd. (Nanjing, China).

2.5. Extraction and isolation

The dried whole herb of *P. oleracea* (150 kg) were extracted twice with a 10-fold amount 50% ethanol for 2 h each time, and then the ethanol extract was concentrated under reduced pressure reflux, obtaining the extract (15 kg), which was latter absorbed on a 100–200 mesh silica-gel column chromatography (82 \times 59 cm, approximately, 118.5 kg) with a 2-

fold amount ethyl acetate added aqueous ammonia (100:1, v/v) 3 times as the isocratic elution, affording the fraction evaporated (510 g). Then, the fraction, via the a 80–100 mesh polyamide resin column chromatography (120 × 8 cm, approximately, 2.5 kg), was eluted with water, 30%, 50%, 70% and pure ethanol to obtain 5 fractions (4 L each). The fraction of 50% extracting was condensed under the reflux to obtain the *P. oleracea* extract of 100 g, which was subjected to a 200–300 mesh silica-gel column (120 × 8 cm, approximately, 2.5 kg) chromatography with ethyl acetate, ethyl acetate and methanol (5:1, 2:1, 1:2, v/v) as the gradient eluent in proper order, obtaining 4 fractions (Frs. 1–4, 500 mL each). Then the fractions were spotted on a thin-layer chromatography (TLC) plate and sprayed with ferric chloride reagent. Frs. 1 (62 g) turned cyan and afterwards passed through the 20–40 μm ODS (25 × 3 cm, approx. 150 g, Ultimate XB-C18) under medium pressure, with a gradient of methanol/water (84:16, 93:7, 97:3, v/v), acquiring 3 fractions (G1-G3, 200 mL each). Then the fractions were repeatedly spotted on a thin-layer chromatography (TLC) plate and sprayed with ferric chloride reagent. G1 (15 g) turned cyan and then was prepared by with ultra-high-performance liquid chromatography (UHPLC), using MeOH - 0.1% formic acid different ratio as the mobile phase, with flow rate of 1.0 mL/min, and resulted in compound **1** (40 mg, purity of > 99% with UHPLC, MeOH - 0.1% formic acid, 61:39, v/v), compound **2** (30 mg, purity of >98% with UHPLC, MeOH - 0.1% formic acid, 58:42, v/v), compound **3** (3 mg, purity of >98% with UHPLC, MeOH - 0.1% formic acid, 52:48, v/v), compound **4** (10 mg, purity of >98% with UHPLC, MeOH - 0.1% formic acid, 57:43, v/v), compound **5** (4 mg, purity of >97% with UHPLC, MeOH - 0.1% formic acid, 52:48, v/v), compound **6** (3 mg, purity of >96% with UHPLC, MeOH - 0.1% formic acid, 56:44, v/v).

2.5.1. Oleracone C (**1**)

Yellowish-brown powder; $[\alpha]_D^{20} + 0.8$ (c 0.1, MeOH); UV (CH₃OH) λ_{\max} 288, 214 nm; IR (KBr) ν_{\max} 3425, 2928, 1650, 1603, 1506, 1456, 1303, 1231, 1193, 1162 cm⁻¹. ¹H and ¹³C NMR data (Table 1. and Table S1. in Supplementary material); HR-ESI-MS m/z 299.0930 [M – H]⁻ (calcd. for C₁₇H₁₅O₅, 299.0925).

2.5.2. Oleracone D (**2**)

Yellowish powder; UV (CH₃OH) λ_{\max} 286, 254 nm; IR (KBr) ν_{\max} 3428, 2925, 2853, 1659, 1634, 1596, 1384 cm⁻¹. ¹H and ¹³C NMR data (Table 1. and Table S2. in Supplementary material); HR-ES-IMS m/z 299.0911 [M + H]⁺ (calcd. for C₁₇H₁₅O₅, 299.0914).

2.5.3. 5,7-Dimethoxy-4-O-2'-cycloflavan (**3**)

Yellowish powder; $[\alpha]_D^{20} - 14$ (c 0.01, MeOH); ¹H and ¹³C NMR data (Table 1. and Table S3. in Supplementary material); HR-ESI-MS m/z 285.1118 [M + H]⁺ (calcd. for C₁₇H₁₇O₄, 285.1121).

2.5.4. Oleracone E (**4**)

Yellowish powder; UV (CH₃OH) λ_{\max} 290, 212 nm; IR (KBr) ν_{\max} 3446, 2925, 2862 1653, 1559, 1540, 1506, 1459 cm⁻¹. ¹H and ¹³C NMR data (Table 1. and Table S4. in Supplementary material); HR-ESI-MS m/z 303.1229 [M + H]⁺ (calcd. for C₁₇H₁₉O₅, 303.1227).

2.6. Assay of scavenging DPPH radicals

An assay of DPPH radical scavenging activity was performed by the modified method described formerly [22]. For each compound, a dilution into five different concentrations in methanol was prepared and the methanol was used to zero the spectrophotometer. A

methanolic solution of the sample (1 mL) was mixed with a 126.80 μM DPPH (1 mL) dissolved in methanol. The reaction mixture was allowed to stand for 10 min at room temperature, measured at 517 nm, and then followed at 30 min. The DPPH solution (1 mL) added to methanol (1 mL) were measured as the blank control. At the same time, the sample solution (1 mL), with methanol (1 mL) was operated as the control group. BHA (butylated hydroxyanisole) was tested as a positive control as the same series of concentrations as other samples. Each sample was measured three times in parallel, and the percentage of DPPH radical scavenging was calculated in the following way [23]:

$$\text{DPPH radical scavenging effect (\%)} = (1 - (A_{\text{sample}} - A_{\text{control}}) / A_{\text{blank}}) \times 100\%$$

where, A_{control} is the absorbance of the control group, A_{sample} is the absorbance of the test compound, and A_{blank} is the absorbance of the blank control. The antioxidant capacity was evaluated by the value of IC_{50} , and values are expressed as the means \pm SD for $n = 3$.

2.7. Assay of anticholinesterase activity

Anticholinesterase activity was determined against AChE, using the modified method according to Ellman [24]. The 20 μL sample solutions in methanol diluted into five different concentrations and 15 μL AChE solution in buffer of PBS (pH 8.0, containing 0.1 mol/L Na_2HPO_4 and NaH_2PO_4) were mixed in wells a 96-well microplate and incubated for 10 min at 37 $^\circ\text{C}$, and 10 μL DTNB in buffer was added. Then 10 μL ATCI was added to initiate the reaction and the absorbance was monitored spectrophotometrically at 405 nm. Eserine, as the same concentration as samples, was used as the positive control. Instead of samples, methanol was included in the blank control. Each sample was measured three times in parallel, and the percentage of inhibition of AChE was calculated using the following formula [25]: $\text{Anticholinesterase activity\%} = (A_{\text{blank}} - A_{\text{sample}}) / A_{\text{blank}} \times 100\%$ where, A_{sample} is the absorbance of the test compound, and A_{blank} is the absorbance of the blank control. The anticholinesterase activity was evaluated by the value of IC_{50} , and values are expressed as the means \pm SD for $n = 3$.

3. Results and discussion

Compound **1**, $[\alpha]_{\text{D}}^{20} + 0.8$ (c 0.1, MeOH), was obtained as yellowish-brown powder turning cyan upon spraying ferric chloride. Its molecular formula was deduced as $\text{C}_{17}\text{H}_{16}\text{O}_5$ based on a molecular ion peak at m/z 299.0930 $[\text{M}-\text{H}]^-$ in the HR-ESI-MS spectrum (calcd. for $\text{C}_{17}\text{H}_{15}\text{O}_5$, 299.0925). The degree of unsaturation of compound **1** is 10. The ^1H NMR spectrum exhibited six aromatic protons at δ_{H} 6.06 (1H, d, $J = 2.3$ Hz), 6.07 (1H, d, $J = 2.3$ Hz), 6.73 (1H, td, $J = 7.4, 1.2$ Hz), 6.82 (1H, dd, $J = 8.4, 1.2$ Hz), 7.05 (1H, brtd, $J = 7.4, 1.6$ Hz), and 7.06 (1H, brd, $J = 7.4$ Hz), a methoxy group at δ_{H} 3.79 (3H, s), and two hydroxyl groups at δ_{H} 9.59 (1H, brs) and 12.12 (1H, brs) (Table 1). According to the ^{13}C NMR and DEPT 135 spectrum, 17 carbon signals were detected, including 7 quaternary carbons with one carbonyl at δ_{C} 198.7, 7 methines, two methylenes (δ_{C} 27.2, 69.4), and one methoxy group (δ_{C} 56.0).

The signals at δ_{H} 6.06/ δ_{C} 93.6 and δ_{H} 6.07/ δ_{C} 94.8 are typical of a tetrasubstituted aromatic A-ring of a flavonoid. Both protons correlate with the methoxyl at δ_{H} 3.79 in NOESY, indicating that the methoxy substituent is in C-7 (at δ_{C} 167.4). The relative position of these

two protons was difficult to ascertain. Both correlate with carbons at δ_C 102.2 (C-4a) and 167.4 (C-7). Eventually, it was found that proton at δ_H 6.06 correlates with carbon at δ_C 162.9 (C-8a) while proton at δ_H 6.07 correlates with carbon at δ_C 163.6 (C-5). These correlations allowed us to assign δ_H 6.06/ δ_C 93.6 at position 8 and δ_H 6.07/ δ_C 94.8 at position 6.

Compound **1** also contained a CH₂-CH-CH₂ system identified by the sequence of COSY correlations H-2a, b/H-3/H-9a, b and coupling constants in the system. Both protons H-2 correlate with C-8a at δ_C 162.9 through oxygen 1, and also correlate to carbonyl moiety at δ_C 198.7 which had to be linked to C-3, forming the C-ring of an isoflavonoid. However, in the isoflavonoids, an aromatic ring is usually linked to C-3. Here, C-9 is inserted between C-3 and an aromatic ring as shown by correlations of protons H-9 with nearby carbons C-1', C-6' (and C-2' for H-9a). Carbons C-1' to C-6' describe a disubstituted phenyl moiety as confirmed by the coupling figures and COSY correlations of protons H-3' to H-6', along with respective HMBC correlations (Fig. 2). It should be mentioned that H-6' also correlates to C-9 further confirming the C-9-C-1' link. The OH signals were attributed based on NOE correlation H-3'/HO-C2'. Finally, the absolute configuration at C-3 was assigned as *S* on the basis of the positive Cotton effect at 288 nm [26,27]. (Fig. S41–42, Supporting information).

Overall, compound **1** was identified as the (3*S*)-5-hydroxy-3-(2-hydroxybenzyl)-7-methoxychroman-4-one (or (3*S*)-5,2'-dihydroxy-7-methoxyhomoisoflavanone) and was given the trivial name oleracone C.

data of compounds 1–4 in dmsd- <i>d</i> ₆ .		2		3		Pos
	δ_C	δ_H (<i>J</i> in Hz)	δ_C	δ_H (<i>J</i> in Hz)	δ_H (<i>J</i> in Hz)	
1)	154.9	8.05, s	66.6	5.38, brq (2.5)		1
5)	121.0		26.0	a:2.11, dt (14.1, 2.8) b:2.18, dt (14.1, 2.8)		2
3)	181.8		61.2	5.56, brq (1.9)		3
	105.1		103.0			1'
	161.3		158.7			2'
		12.74, brs				2''-C
	97.9	6.38, d (1.9)	55.6	3.79, s		3'
	165.1		92.0	5.96, d (2.3)		4'
	56.0	3.84, s	161.4			5'
	92.3	6.59, d (1.9)	55.1	3.65, s		6'
	157.7		91.3	6.09, d (2.3)		1''
	24.4	3.61, s	154.4			2''
3)						2''-C
3)	124.5		121.4			3''
	155.2		153.3			4''
	115.0	9.65, brs				4''-C
	127.4	6.82, dd (7.9, 1.1)	116.6	6.78, brd (8.2)		5''
	118.8	7.03, brtd (7.6, 1.6)	130.3	7.19, ddd (8.2, 7.3, 1.7)		6''
	129.8	6.70, brtd (7.4, 1.1)	119.9	6.87, brtd (7.6, 1.6)		6''-C
		7.08, dd (7.4, 1.6)	131.0	7.35, dd (7.7, 1.7)		




Fig. 2. ^1H - ^1H COSY and HMBC correlations for compounds 1-4.

Compound **2** was obtained as yellowish powder, which turned cyan after spraying ferric chloride. Its molecular formula was inferred as $\text{C}_{17}\text{H}_{14}\text{O}_5$ based on a molecular ion peak at m/z 299.0911 $[\text{M} + \text{H}]^+$ in positive-ion HR-ESI-MS (calcd. for $\text{C}_{17}\text{H}_{15}\text{O}_5$, 299.0914). The degree of unsaturation of compound **2** is 11. The ^1H NMR spectrum contains six aromatic proton signals including the two protons at δ_{H} 6.38 (1H, d, $J = 1.9$ Hz) and 6.59 (1H, d, $J = 1.9$ Hz) attributable to H-6 and H-8 although they appear more downfield than in compound **1**. Other aromatic signals are at δ_{H} 6.82 (1H, dd, $J = 7.9, 1.1$ Hz), 6.70 (1H, brtd, $J = 7.4, 1.1$ Hz), 7.03 (1H, brtd, $J = 7.6, 1.6$ Hz), and 7.08 (1H, dd, $J = 7.4, 1.6$ Hz). These correspond to the sequence of protons H-3' to H-6', with chemical shifts almost identical to those of the same protons in **1**. An olefinic proton at δ_{H} 8.05 (1H, s), a methylene proton at δ_{H} 3.61 (2H, s), a methoxy group at δ_{H} 3.84 (3H, s), and two hydroxyl groups at δ_{H} 9.66 (1H, s) and 12.76 (1H, s) were also seen. Seventeen carbons were detected in the ^{13}C NMR and DEPT 135 spectra. These include one carbonyl (δ_{C} 181.8) and 13 other sp^2 carbons, one methylene (δ_{C} 24.4), one methine (δ_{C} 154.9), and one methoxy group (δ_{C} 56.0) (Table 1). Compound **2** was clearly related to **1** with 1 more double bond. The HMBC correlations from H-9 (δ_{H} 3.61) to C-1' (δ_{C} 124.5), C-2' (δ_{C} 155.2), C-6' (δ_{C} 129.8), C-2 (δ_{C} 154.9), C-3 (δ_{C} 121.0), C-4 (δ_{C} 181.8), and from H-2 (δ_{H} 8.05) to C-3 (δ_{C} 121.0), C-4 (δ_{C} 181.8) revealed direct connection of C-3 with C-4, and C-1' with C-9. HMBC correlations from the CH_3 at δ_{H} 3.84 (3H, s) to C-7 (δ_{C} 165.1) indicated that the methoxy group should be linked to C-7, like in **1** (Fig. 2). As a result, compound **2** was identified as the 5-hydroxy-3-(2-hydroxybenzyl)-7-methoxy-4*H*-chromen-4-one (or 5,2'-dihydroxy-7-methoxy-homoisoflavone) and was given the trivial name oleracone D (**2**).

Compounds **3**, $[\alpha]_{\text{D}}^{20} -14$ (c 0.01, MeOH), was obtained as yellowish powder. Its molecular formula was identified as $\text{C}_{17}\text{H}_{16}\text{O}_4$ based on a molecular ion peak at m/z 285.1118 $[\text{M} + \text{H}]^+$ in its positive-ion HR-ESI-MS spectrum (calcd. for $\text{C}_{17}\text{H}_{17}\text{O}_4$, 285.1121). The degree of unsaturation of the compound **3** is 10. The ^1H NMR spectrum contained six aromatic proton signals at δ_{H} 5.96 (1H, d, $J = 2.3$ Hz), 6.09 (1H, d, $J = 2.3$ Hz), 6.78 (1H, brd, $J = 8.2$ Hz), 6.87 (1H, brtd, $J = 7.6, 1.1$ Hz), 7.19 (1H, ddd, $J = 8.2, 7.3, 1.7$ Hz) and 7.35 (1H, dd, $J = 7.7, 1.7$ Hz), one methylene with non-equivalent protons at δ_{H} 2.11 (1H, dt, $J = 14.1, 2.8$ Hz) and 2.18 (1H, dt, $J = 14.1, 2.8$ Hz), two methine protons at δ_{H} 5.38 (1H, brq, $J = 2.5$ Hz), 5.56 (1H, brq, $J = 1.9$ Hz), and two methoxy groups at δ_{H} 3.65 (3H, s) and 3.79 (3H, s). In the ^{13}C NMR and DEPT 135 spectrum, there were 17 carbon signals, including 12 sp^2 carbons, one methylene carbon (δ_{C} 26.0), two methine carbons (δ_{C} 61.2, 66.6), two methoxy groups (δ_{C} 55.1, 55.6) (Table 1). The COSY and HMBC data, along with comparison with compounds **1** and **2** provide evidence that this compound also contain the same di- and tetrasubstituted aromatic rings (B- and A-ring, respectively, Fig. 2). The methoxy substituents were located

based of HMBC correlations between the methoxy groups at δ_{H} 3.79 and 3.66 with C-5 (δ_{C} 158.7) and C-7 (δ_{C} 161.4), respectively.

A CH–CH₂–CH sequence was also identified based on ¹H–¹H couplings and COSY experiment. Unfortunately, it was not possible to detect any HMBC correlations for these protons. However, the methines chemical shifts at δ_{H} 5.38 (H-2) and 5.56 (H-4) indicate that both CH are deshielded presumably by an oxygen and an aromatic ring. Interestingly, H-2 and H-4 are coupled to each other ($J \approx 1.9$ – 2.5 Hz). This is specific to systems constrained in a W-shaped conformation, typical of the 4-*O*-2'-cycloflavan skeleton. In the closely related 3'-hydroxy-5,7-dimethoxy-4-*O*-2'-cycloflavan isolated from *Iris tenuifolia*, H-2 and H-4 are reported at δ_{H} 5.34 and 5.62 in dms-*d*₆, respectively [28]. These values are very similar to those of compound **3**. Overall, it can be concluded that **3** is the 5,7-dimethoxy-4-*O*-2'-cycloflavan. This compound has been obtained in the past by synthesis but its analytical data have not been described [29,30]. It is isolated for the first time as a natural product. The circular dichroism spectrum of **3** was compared with the one reported for the (2*R*,4*R*)-4'-hydroxy-3',5'-methyl-6,7-methylenedioxy-4-*O*-2'-cycloflavan [31]. This compound is the only 4-*O*-2'-cycloflavan, not hydroxylated in C-3 as in the cyanomaclurin series, for which the CD spectrum has been reported. Compound **3** presented 3 positive maxima at 217, 242 and 281 nm (Fig. S41–42, Supporting information), while the reported CD spectrum had negative Cotton effect at these wavelengths. This indicated that the absolute configuration of compound **3** is 2*S* and 4*S*, as represented in Fig. 1.

Compound **4** was obtained as yellowish powder, which became cyan after spraying Ferric chloride. Its molecular formula was identified as C₁₇H₁₈O₅ based on a molecular ion peak at m/z 303.1229 [M + H]⁺ in its positive-ion HR-ESI-MS spectrum (calcd. for C₁₇H₁₉O₅, 303.1227). The degree of unsaturation of compound **4** is 9. The ¹H NMR spectrum exhibited six aromatic proton signals at δ_{H} 6.08 (1H, d, $J = 2.3$ Hz), 6.09 (1H, d, $J = 2.3$ Hz), 6.68 (1H, td, $J = 7.4, 1.0$ Hz), 6.78 (1H, dd, $J = 8.0, 1.0$ Hz), 6.98 (1H, brtd, $J = 7.6, 1.6$ Hz) and 7.05 (1H, dd, $J = 7.4, 1.6$ Hz), two methylene protons at δ_{H} 2.80 (2H, brt, $J = 7.7$ Hz), 3.18 (2H, brt, $J = 7.7$ Hz), two methoxy groups at δ_{H} 3.79 (3H, s), 3.81 (3H, s), and two hydroxyl groups at δ_{H} 9.57 (1H, s) and 13.59 (1H, s). The ¹³C NMR and DEPT 135 spectrum indicated that 17 carbon signals, including one carbonyl (δ_{C} 204.8) and 12 other sp² carbons, two methylene carbons (δ_{C} 25.1, 43.5), and two methoxy groups (δ_{C} 55.6, 56.0) (Table 1). Comparison with compounds **1**–**3** provided straightforward identification of the tetrasubstituted A-ring and the *ortho*-disubstituted B-ring. However in this case, it was eventually found that **4** was a dihydrochalcone, with both aromatic rings linked by a propane-1-one core. This linkage was identified based on the H-2–H-3 coupling and the HMBC correlations from H-3 (δ_{H} 2.80) to C-1' (δ_{C} 127.3), C-2' (δ_{C} 155.3), C-6' (δ_{C} 129.7), C-2 (δ_{C} 43.5), C-1 (δ_{C} 204.8), and from H-2 (δ_{H} 3.19) to C-1 (δ_{C} 204.8), C-3 (δ_{C} 25.1), and C-1' (δ_{C} 127.3). The HMBC correlations revealed that the methoxy groups (δ_{H} 3.79) was coupled to the C-4'' (δ_{C} 165.4), and another methoxy groups (δ_{H} 3.82) was coupled to the C-6'' (δ_{C} 162.3), which implied that two methoxy groups are affiliated with C-4'' and C-6'', respectively (Fig. 2). As a result, compound **4** was determined as the 1-(2-hydroxy-4,6-dimethoxyphenyl)-3-(2-hydroxyphenyl)propan-1-one and was given the trivial name oleracone E.

Compounds **5** and **6** had been described before and were identified as the (2*S*)-5,2'-dihydroxy-7-methoxyflavanone and 2',4'-dihydroxy-4,6'-dimethoxychalcone respectively, according to the NMR data and comparison with the literature [32,33].

Previously, the antioxidant activity of extracts was assayed by using the DPPH radical scavenging method [34], which is now applied to evaluate the oxidation resistance of the three new compounds. Phenolic compounds are good hydrogen donors and are usually effective DPPH radical quenchers [35]. Indeed, oleracones C (**1**), D (**2**) and E (**4**) were found more effective than reference compound butylated hydroxyanisole (BHA) (Fig. S43–44, Supporting information). Their antioxidant potential was dose-dependent with IC₅₀ values of 11.7, 13.2 and 23.3 μM, respectively (Table 2). The biological activity profiles were somewhat similar at 30 min, with IC₅₀ values of 12.0, 13.7 and 23.5 μM, respectively (Table 2). Oleracone D (**2**) remained the most effective. Overall, the compared radical scavenging activities on DPPH were oleracone D (**2**)>E (**4**)>C (**1**)>BHA.

Acetylcholinesterase (AChE) inhibitors, inhibiting the production of choline, play an significant role in the treatment of Alzheimer disease (AD) [36]. As a result, antioxidant activities of phenolic compounds can be implicated in the neuroprotective effect at the same time [37], of which flavonoids are responsible to be the multipotent agents to fight against AD [38]. Consequently, the anticholinesterase activities of three new compounds, oleracone C (**1**), D (**2**) and E (**3**), were determined. All of three new compounds had different degree of inhibitory activities against the AChE (Fig. S45, Supporting Information). Table 2 indicated that although the anticholinesterase activity of compound **2** was lower than that of eserine at the same concentration, it had the highest anticholinesterase activity with IC₅₀, the lowest was that of compound **3**.

Table 2

IC₅₀ (μM) for antioxidant and acetylcholinesterase activities of compounds **1**, **2** and **4** (n = 3).

Compounds and standard inhibitors	DPPH IC ₅₀ (μM)		AChE IC ₅₀ (μM)
	10 min	30 min	
1	23.27 ± 0.02	23.53 ± 0.03	68.87 ± 0.09
2	11.73 ± 0.02	12.02 ± 0.02	59.56 ± 0.50
4	13.17 ± 0.01	13.72 ± 0.07	78.85 ± 0.16
Eserine	–	–	31.13 ± 0.02
BHA	59.14 ± 0.04	59.15 ± 0.11	–

Eserine and BHA as positive control, and values are expressed as the means ± SD for n = 3.

Conflict of interest statement

The authors declare no competing financial interest.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant 81573546).

References

[1]

Y.Y. Lim, E.P.L. Quah

Antioxidant properties of different cultivars of *Portulaca oleracea*

Food Chem., 103 (2007), pp. 734-740

[2]

Y.X. Zhou, H.L. Xin, K. Rahman, S.J. Wang, C. Peng, H. Zhang

***Portulaca oleracea* L.: a review of phytochemistry and pharmacological effects**

Biomed. Res. Int., 2015 (2015), p. 925631

[3]

X. Lei, J. Li, B. Liu, N. Zhang, H. Liu

Separation and identification of four new compounds with antibacterial activity from *Portulaca oleracea* L

Molecules, 20 (2015), pp. 16375-16387

[4]

U.R. Palaniswamy, R.J. Mcavoy, B.B. Bible

Stage of harvest and polyunsaturated essential fatty acid concentrations in purslane (*Portulaca oleraceae*) leaves

J. Agric. Food Chem., 49 (2001), pp. 3490-3493

[5]

M.K. Uddin, A.S. Juraimi, M.S. Hossain, M.A. Nahar, M.E. Ali, M.M. Rahman

Purslane weed (*Portulaca oleracea*): a prospective plant source of nutrition, omega-3 fatty acid, and antioxidant attributes

Sci. World J., 2014 (2014), pp. 1-6

[6]

D. Liu, S. Tao, X. Lan

Two antioxidant alkaloids from *Portulaca oleracea* L

Helv. Chim. Acta., 94 (2011), pp. 497-501

[7]

Y. Meng, Z. Ying, Z. Xiang, D. Hao, W. Zhang, Y. Zheng, Y. Gao, X. Ying

The anti-inflammation and pharmacokinetics of a novel alkaloid from *Portulaca oleracea* L

J. Pharm. Pharmacol., 68 (2016), pp. 397-405

[8]

K. Chan, M.W. Islam, M. Kamil, R. Radhakrishnan, M.N.M. Zakaria, M. Habibullah, A. Attas

The analgesic and anti-inflammatory effects of *Portulaca oleracea* L. subsp. sativa (Haw.) Celak

J. Ethnopharmacol., 73 (2000), pp. 445-451

[9]

X.J. Zhang, Y.B. Ji, Z.Y. Qu, J.C. Xia, L. Wang

Experimental studies on antibiotic functions of *Portulaca oleracea* L. in vitro

Chinese J. Microecol., 14 (2002), pp. 277-280

[10]

S. Sirithon, S. Maitree

Microchemical components and antioxidant activity of different morphological parts of Thai wild purslane (*Portulaca oleracea*)

Weed Sci., 58 (2010), pp. 182-188

[11]

A.E.A. Moneim, M.A. Dkhil, S. Al-Quraishy

The potential role of *Portulaca oleracea* as a neuroprotective agent in rotenone-induced neurotoxicity and apoptosis in the brain of rats

Pestic. Biochem. Physiol., 105 (2013), pp. 203-212

[12]

R.A. Kennedy

Characteristics of Crassulacean acid metabolism in the succulent C₄ dicot, *Portulaca oleracea* L

Plant Physiol., 65 (1980), pp. 193-197

[13]

N. Erkan

Antioxidant activity and phenolic compounds of fractions from *Portulaca oleracea* L

Food Chem., 133 (2012), pp. 775-781

[14]

E.S. Elkhayat, S.R. Ibrahim, M.A. Aziz

Portulene, a new diterpene from *Portulaca oleracea* L

J. Asian Nat. Prod. Res., 10 (2008), pp. 1039-1043

[15]

A.N. Rasheed, F.U. Afifi, M. Shaedah, M.O. Taha

Investigation of the active constituents of *Portulaca oleraceae* L. (Portulacaceae) growing in Jordan

Pak. J. Pharm. Sci., 17 (2004), pp. 37-45

[16]

X. Xu, L. Yu, G. Chen

Determination of flavonoids in *Portulaca oleracea* L. by capillary electrophoresis with electrochemical detection

J. Pharm. Biomed. Anal., 41 (2006), pp. 493-499

[17]

C. Li, Y. Meng, Z. Ying, N. Xu, D. Hao, M. Gao, W. Zhang, L. Xu, Y. Gao, X. Ying

Three novel alkaloids from *Portulaca oleracea* L. and their anti-inflammatory effects

J. Agric. Food Chem., 64 (2016), pp. 5837-5844

[18]

T.A. Akhtar, H.A. Lees, M.A. Lampi, D. Enstone, R.A. Brain, B.M. Greenberg

Photosynthetic redox imbalance influences flavonoid biosynthesis in *Lemna gibba*

Plant Cell Environ., 33 (2010), pp. 1205-1219

[19]

B. Demmigadams, W.W. Adams

Antioxidants in photosynthesis and human nutrition

Science, 298 (2002), pp. 2149-2153

[20]

S.J. Stohs

The role of free radicals in toxicity and disease

J. Basic Clin. Physiol. Pharmacol., 6 (1995), pp. 205-228

[21]

C. Bensouici, A. Kabouche, A. Karioti, M. Öztürk, M.E. Duru, A.R. Bilia, Z. Kabouche

Compounds from Sedum caeruleum with antioxidant, anticholinesterase, and antibacterial activities

Pharm. Biol., 54 (2016), pp. 174-179

[22]

J.K. Liu, L. Hu, Z.J. Dong, Q. Hu

DPPH radical scavenging activity of ten natural p-terphenyl derivatives obtained from three edible mushrooms indigenous to China

Chem. Biodivers., 1 (2004), pp. 601-605

[23]

W. Xu, F. Zhang, Y. Luo, L. Ma, X. Kou, K. Huang

Antioxidant activity of a water-soluble polysaccharide purified from Pteridium aquilinum

Carbohydr. Res., 344 (2009), pp. 217-222

[24]

G.L. Ellman, K.D. Courtney, A.V. Jr, R.M. Featherstone

A new and rapid colorimetric determination of acetylcholinesterase activity

Biochem. Pharmacol., 7 (1961), pp. 88-90

[25]

T. Dirmenci, B. Culhaoglu, F. Bahadori, S.D. Hatipoglu, M. Boga, M. Ozturk, U. Kolak, G.

Topcu

Investigation of anticholinesterase activity of a series of salvia extracts and the constituents of Salvia staminea

Nat. Products J., 3 (2013), pp. 3-9

[26]

M. Adinolfi, G. Barone, M.M. Corsaro, L. Mangoni, R. Lanzetta, M. Parrilli

Absolute configuration of homoisoflavanones from muscari species

Tetrahedron, 44 (1988), pp. 6565-6574

[27]

Y. Dai, L. Harinantenaina, P.J. Brodie, M. Goetz, Y. Shen, K. Tendyke, D.G. Kingston
Antiproliferative Homoisoflavonoids and Bufatrienolides from *Urginea depressa*
J. Nat. Prod., 76 (2013), pp. 865-872

[28]

Y.M. Cui, H. Wang, Q.R. Liu, M. Han, Y. Lu, C.Q. Zhao
Flavans from *Iris tenuifolia* and their effects on β -amyloid aggregation and neural stem cells proliferation in vitro
Bioorg. Med. Chem. Lett., 21 (2011), pp. 4400-4403

[29]

P.M. Nair, P.C. Parthasarathy, P.V. Radhakrishnan, K. Venkataraman
Stereochemistry of cyanomaclurin
Tetrahedron Lett., 7 (1966), pp. 5357-5361

[30]

G.D. Bhatia, S.K. Mukerjee, T.R. Seshadri
Synthesis of cyanomaclurin—I: synthesis of some model compounds
Tetrahedron, 22 (1966), pp. 139-143

[31]

W. Peng, C. Yang, R. Zhan, Y. Chen
Two new flavans from the trunk and leaves of *Horsfieldia glabra*
Nat. Prod. Res., 30 (2016), pp. 2350-2355

[32]

V. Seidel, F. Bailleul, P.G. Waterman
(Rel)-1 β ,2 α -di-(2,4-dihydroxy-6-methoxybenzoyl)-3 β , 4 α -di-(4-methoxyphenyl)-cyclobutane and other flavonoids from the aerial parts of *Goniothalamus gardneri* and *Goniothalamus thwaitesii*
Phytochemistry, 55 (2000), pp. 439-446

[33]

B. Jayaprakasam, A.G. Damu, D. Gunasekar, A. Blond, B. Bodo
ChemInform abstract: dihydroechioidinin, a flavanone from *Andrographis echiodes*
Phytochemistry, 52 (1999), pp. 935-937

[34]

W. Brandwilliams, M.E. Cuvelier, C. Berset
Use of a free radical method to evaluate antioxidant activity
LWT Food Sci. Technol., 28 (1995), pp. 25-30

[35]

G. Cao, E. Sofic, R.L. Prior
Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships
Free Radic. Biol. Med., 22 (1997), pp. 749-760

[36]

E.K. Akkol, I.E. Orhan, E. Yeşilada

Anticholinesterase and antioxidant effects of the ethanol extract, ethanol fractions and isolated flavonoids from *Cistus laurifolius* L. leaves

Food Chem., 131 (2012), pp. 626-631

[37]

C. Ramassamy

Emerging role of polyphenolic compounds in the treatment of neurodegenerative diseases: a review of their intracellular targets

Eur. J. Pharmacol., 545 (2006), pp. 51-64

[38]

H.F. Ji, H.Y. Zhang

Theoretical evaluation of flavonoids as multipotent agents to combat Alzheimer's disease

J Mol Struc.Theochem., 767 (2006), pp. 3-9