



HAL
open science

Reduced bacterial mortality and enhanced viral productivity during sinking in the ocean

Wei Wei, X Chen, Markus G Weinbauer, Nianzhi Jiao, R Zhang

► **To cite this version:**

Wei Wei, X Chen, Markus G Weinbauer, Nianzhi Jiao, R Zhang. Reduced bacterial mortality and enhanced viral productivity during sinking in the ocean. *The International Society of Microbiological Ecology Journal*, 2022, 10.1038/s41396-022-01224-9 . hal-03946294

HAL Id: hal-03946294

<https://hal.science/hal-03946294>

Submitted on 23 Mar 2023




HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ARTICLE OPEN



Reduced bacterial mortality and enhanced viral productivity during sinking in the ocean

Wei Wei ^{1,2,3}, Xiaowei Chen^{2,4}, Markus G. Weinbauer⁵, Nianzhi Jiao ^{2,4}✉ and Rui Zhang ^{2,4,6}✉

© The Author(s) 2022

Particle sinking is an important process in the ocean, influencing the biogeochemical cycle and driving the long-term preservation of carbon into the deep sea via the biological pump. However, as an important component of marine ecosystems, the role of viruses during sinking is still poorly understood. Therefore, we performed a series of transplantation experiments in the South China Sea to simulate environmental changes during sinking and investigate their effects on viral eco-dynamics and life strategy. Our study demonstrated increased viral production but decreased virus-mediated bacterial mortality after transplantation. A larger burst size and switch from the lysogenic to lytic strategy were shown to contribute to enhanced viral productivity. We provide experimental evidence that surface viral ecological characteristics changed dramatically after transplantation into deep-sea waters, indicating a potential importance of viruses during vertical sinking in the ocean. This effect probably provides positive feedback on the efficiency of the biological pump.

The ISME Journal (2022) 16:1668–1675; <https://doi.org/10.1038/s41396-022-01224-9>

INTRODUCTION

The vertical sinking of particles is one of the most important mechanisms driving oceanic dynamics throughout physical, chemical, and biological processes on a global scale. It is a major channel to transport and redistribute fixed carbon through photosynthesis from the sea surface to the deep ocean [1]. As a key component of the “biological pump” (BP) in the ocean, the particulate organic matter (POM) produced in the photic zone aggregates by a complex interplay among faecal pellets, carcasses of zooplankton, phytoplankton cells, and prokaryotic cells [2, 3] and regulates the global climate by long-term sequestration of atmospheric CO₂ [4]. For instance, *ca.* 1–40% of the photosynthetically fixed CO₂ flows into the dark zone of the ocean in the form of sinking POM [5]. Moreover, the sinking POM is continuously degraded and releases the dissolved organic matter (DOM) and nutrients, which are used by microbial communities and made available for recycling into the microbial loop [6]. As the main microbial communities on POM, colonizing bacteria have been revealed to harbor distinct features regarding diversity, structure, and dynamics, which are significantly different from those of the surrounding free-living bacterioplankton [7–9]. They are considered an important transformer in driving the disaggregation and remineralization of sinking POM to regulate the efficiency of the BP [10–13], and play a substantial role in global biogeochemical cycling [14, 15].

Over the past three decades, viruses have been found to be the most abundant “life forms” in global oceans, playing an important ecological role [16, 17]. They infected bacteria and are responsible

for high bacterial mortality [18, 19]. Thus, the “viral shunt” could enhance the transformation of POM (e.g., host cells) into DOM (e.g., nucleic acid and protein) [20] and subsequently impact biogeochemical cycling in the global oceans [6, 17, 21, 22]. During sinking, viruses can be transplanted into the deep ocean directly through adsorption on POM or indirectly through infected cell hosts (e.g., viral particles in the infected bacterial cells) [23, 24]. However, up to present, the bacterium-virus interaction and the role of viruses in these processes are unclear. A recent study found that the transferred viral particles still retain infectivity for years in the deep-ocean environment [25], indicating that the vertical transportation of these viruses may continuously contribute to the activity of the deep-ocean viral population [26, 27]. Thus, we hypothesize that such direct and indirect vertical transportation in the ocean could impact the virus-host interaction, and subsequently influence the ecological role of microbes in the sinking process, as well as the efficiency of the BP.

To verify our hypothesis and better understand how the vertical transportation affects virus-host interactions by altering viral activity, dynamics and life strategy, we performed 18 transplantation experiments at 6 stations in the South China Sea (SCS) (Fig. 1). Our study for the first time reveals the effect of vertical transplantation on viral eco-dynamics and life strategies, such as the lytic and lysogenic viral production rate (VP_R), viral decay rate (VD_R), burst size (BS), virus-mediated mortality (VMM) and fraction of lysogenic cells (FLC), and highlights the importance of viruses in vertical sinking in the ocean, which may influence the efficiency of the BP.

¹College of the Environment and Ecology, Xiamen University, Xiamen 361102, PR China. ²State Key Laboratory of Marine Environmental Science, Fujian Key Laboratory of Marine Carbon Sequestration, Xiamen University, Xiamen 361102, PR China. ³School of Environmental Ecology and Biological Engineering, Wuhan Institute of Technology, Wuhan 430205, PR China. ⁴College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, PR China. ⁵Sorbonne Universités, UPMC, Université Paris 06, CNRS, Laboratoire d’Océanographie de Villefranche (LOV), Villefranche-sur-Mer 06230, France. ⁶Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai 519080, PR China. ✉email: jiao@xmu.edu.cn; ruizhang@xmu.edu.cn

Received: 28 April 2021 Revised: 14 January 2022 Accepted: 14 March 2022

Published online: 1 April 2022




Fig. 1 Map of sampling stations and the design of transplantation experiments in the South China Sea. The seawater samples were collected from the total 19 layers of the six stations (a, b, generated by Ocean Data View 4 software, <https://odv.awi.de/>). The transplantation experiments of surface bacterial/viral concentrate (5 m depth) mixed with virus-free seawater of each deeper layer were performed at the six stations (c, d). The schematic diagram of the transplantation experiments is shown (d) and detailed in the text. Abbreviations represent tangential flow filtration (TFF), incubation with mitomycin C (+MC), incubation without mitomycin C (–MC), viral production (VP) and viral decay (VD).

MATERIALS AND METHODS

Study area, sampling, and environmental parameters

Seawater samples were obtained at six stations along the Pearl River Estuary to the pelagic area of the SCS during a research cruise from August to September 2014 (Fig. 1a, b). The sampling covered three typical pelagic zones: upper epipelagic (0–75 m), lower epipelagic (75–200 m), and mesopelagic ocean (200–1000 m). At each station, water samples were collected with a CTD (SBE9/11 plus, Sea-Bird Electronics, Inc., USA) rosette sampler equipped with 12 Niskin bottles (12 l each) and then prefiltered with 20-µm mesh filters to remove zooplankton. Although this operation removed particles larger than 20 µm, previous investigations showed that these large POM aggregates occupy relatively small fraction (ca. 0.1%) in total particles in surface ocean [28, 29]. The prefiltered water was prepared for the determinations of picoplankton and virioplankton abundances and the following viral dynamics and life strategy experiment. The CTD was also equipped with sensors for measuring in situ data of depth, temperature, and salinity in each water layer of the stations. In addition, samples (500 ml) for nutrient analysis were stored at -20°C , and then the concentrations of $\text{NO}_3^- + \text{NO}_2^-$, NO_2^- , SiO_3^{2-} and PO_4^{3-} were determined by an Auto Analysis III, AA3 instrument (Bran-Luebbe, Germany) in the laboratory.

Picoplankton and virioplankton abundance

The subsamples (1.98 ml of seawater) for picoplankton and virioplankton abundance measurement were fixed with glutaraldehyde (20 µl, Sangon) at a final concentration of 0.5%. After incubation at 4°C for 15 min in the dark, they were flash-frozen in liquid nitrogen and then stored at -80°C until analysis [30]. The abundances of picoeukaryotes, *Synechococcus*, *Prochlorococcus*, heterotrophic bacteria, and viruses were measured by using a flow cytometer (FCM, Epics Altra II, Beckman Coulter, USA) equipped with an air-cooled laser with a wavelength of 488 nm (Sapphire, Coherent, USA).

Specifically, the samples for virus counting were diluted in Tris–EDTA buffer (pH: 8; Sigma), stained with SYBR Green I (0.5×10^{-4} dilution of the commercial stock solution; Molecular Probes), incubated for 10 min at 80°C in a thermostat water bath (DKB-501A, Shanghai Jinghong, China), and then cooled to room temperature for FCM analysis [31, 32]. Heterotrophic bacterial counting was performed on samples that were incubated with SYBR Green I (1×10^{-4} dilution of the commercial stock solution) for 15 min in the dark. In addition, the abundances of picoeukaryotes, *Synechococcus*, and *Prochlorococcus* were directly determined by FCM without staining [33]. For all these samples, a suspension of yellow-green fluorescent beads (1 µm in diameter; Molecular Probes) was added as an internal standard, and all data were analysed with FCS Express V3 software (De Novo Software, <http://www.denovosoftware.com/>). The typical scatter plots of each microbial group were shown in Fig. S1 and additional details about the FCM analysis and data treatment can be found in the previous reports [34, 35].

Lytic and lysogenic viral production

Lytic and lysogenic VP was measured through the dilution approach [36] for samples from the different depth layers and transplant experiments. Fifty milliliters of the bacterial concentrate obtained through tangential flow filtration (TFF) with a 0.22-µm polyvinylidene difluoride (PVDF) cartridge (Labscale, Millipore) from 600 ml of in situ seawater was mixed with 250 ml of virus-free filtrate produced from the same water sample obtained through TFF with a 30 kDa molecular weight polysulfone cartridge (Labscale, Millipore). This operation made viral abundance decline to ca. 10–20% of the initial abundance, and the bacterial concentration was similar to the in situ status [37]. Finally, the experiments measuring lytic and lysogenic VP were performed in 50 ml aseptic tubes (duplicate) with and without adding $1 \mu\text{g ml}^{-1}$ mitomycin C (final concentration; Roche) [20, 38], which were then incubated in dry bath incubators (MK-20, Hangzhou

Allsheng, China) at in situ temperature in the dark. Subsamples (1 ml) were collected to enumerate the viral abundance at 0, 3, 6, 9, and 12 h of incubation. The lytic and lysogenic VP (viruses $\text{ml}^{-1} \text{h}^{-1}$) was estimated through viral accumulation in each 12 h incubation with the VIPCAL online program (<http://www.univie.ac.at/nuhag-php/vipcal/>) [39]. Furthermore, the VP_R was calculated as $\text{VP}_R (\% \text{h}^{-1}) = \frac{\text{VP} (\text{viruses ml}^{-1} \text{h}^{-1})}{\text{VA}_0} \times 100\%$, where VA_0 represents the viral abundance at 0 h in the VP experiment. The BS was defined as $\text{BS} = \frac{\text{VP} (\text{viruses ml}^{-1} \text{h}^{-1})}{\Delta \text{BA} (\text{cells ml}^{-1} \text{h}^{-1})}$, where ΔBA represents the amount of bacterial death determined from the reduction in bacterial abundance per hour during the incubation of the VP experiments, and is calculated by using the equation $\Delta \text{BA} = \left(\frac{\text{BA}_{\text{max } n} - \text{BA}_{\text{min } 1}}{t_{\text{min } 1} - t_{\text{max } 1}} + \dots + \frac{\text{BA}_{\text{max } n} - \text{BA}_{\text{min } n}}{t_{\text{min } n} - t_{\text{max } n}} \right) / n$. This formula for BS calculation is an improved method according to prior studies [40, 41], where $\text{BA}_{\text{max } n}$ and $\text{BA}_{\text{min } n}$ represent the maximum and minimum bacterial abundance at the n^{th} peak of the polyline with bacterial abundance versus time (i.e., 0, 3, 6, 9, and 12 h) in the VP experiment, and $t_{\text{max } n}$ and $t_{\text{min } n}$ are the corresponding times. Since grazers were destroyed or inactivated by tangential flow filtration [37], the bacterial mortality in the incubation system is mainly due to viral lysis in our study. Therefore, the VMM was calculated by using the equation $\text{VMM} = \frac{\Delta \text{BA}}{\text{BA}_0} \times 12 \text{h}$, where BA_0 represents the 0 h bacterial abundance in the incubation of the VP experiment. In addition, FLC, i.e., the percent of lysogenic cells, was estimated by using the formula reported in the VIPCAL online program (<http://www.univie.ac.at/nuhag-php/vipcal/>) [39].

Viral decay

The VD_R was measured through the reported approach [42]. Sample water was filtered through a 0.22- μm PVDF cartridge (Labscale, Millipore) in 50 ml aseptic tubes (duplicate) and incubated in dry bath incubators (MK-20, Hangzhou Allsheng, China) at in situ temperature in the dark. Subsamples (1 ml) were collected at 0, 3, 6, 9, and 12 h of incubation, and the viral abundances were determined at each time point by using FCM. We calculated the VD_R by fitting a linear regression to the decline in ln-transformed viral abundance versus time in the incubation (see Fig. S4 in Wei et al. [41] as an example). Therefore, the slope of the line multiplied by 100 was the decay rate and was expressed as a percentage per hour [42].

Experimental setup of transplantation

The bacterial/viral concentrate obtained from the 5 m depth was mixed with virus-free seawater of each deeper layer at the six stations (Fig. 1c), for example, in Station J4 with four experiments (Fig. 1d): (1) 5 m bacterial concentrate + 5 m virus-free seawater with/without mitomycin C (in situ lysogenic/lytic VP); (2) 5 m viral concentrate + 5 m virus-free seawater (in situ VD); (3) 5 m bacterial concentrate + 90 m virus-free seawater with/without mitomycin C (transplanted lysogenic/lytic VP); and (4) 5 m viral concentrate + 90 m virus-free seawater (transplanted VD). Each sample was added into a 50 ml aseptic tube (duplicate) and incubated in a dry bath incubator for 12 h with the in situ temperature of each layer and in the dark to measure the (lytic and lysogenic) VP and VD as described above. Similar to the dilution approach for measuring VP_R [36], the mixture of viral concentrate (50 ml) with each deeper layer of virus-free water (250 ml) was used to determine the VD_R in the transplant experiments.

Statistical analysis

Using the SPSS (19.0) software package (SPSS Inc., Chicago, IL, USA), Pearson's correlation analysis was applied to assess the degree of correlation among the biotic or abiotic parameters investigated, and Student's *t*-test (paired design) was performed to assess the significant difference in the (lytic and lysogenic) VP_R , VD_R , BS, FLC and VMM between surface water and transplantation treatments. A *p* value < 0.05 was used to indicate statistical significance. In addition, distance-based multivariate analysis for a linear model using forward selection (DISTLM-forward) was used to test the relationships between viral and environmental parameters in Primer 6 software with the PERMANOVA + package (Primer-E, Plymouth, United Kingdom).

RESULTS AND DISCUSSION

In situ environmental and microbial parameters

The in situ environmental variables at the six stations of the SCS with a total of 19 layers were investigated, covering the typical regions of the SCS, from coastal to offshore sea. The temperature ranged from 4.4 to 29.8 °C and dropped sharply with increasing

depth (Fig. S2 and Table S1). The salinity content tended to increase rapidly from the surface of the coastal sea to the deep layer of the offshore sea, varying from 29.6 to 34.6 (Fig. S2 and Table S1). Similarly, the concentrations of $\text{NO}_3^- + \text{NO}_2^-$, SiO_3^{2-} and PO_4^{3-} in the SCS displayed a trend showing that the deep water contained higher nutrient versus surface water, with mean values of $12.10 \pm 16.32 \mu\text{mol l}^{-1}$, $22.38 \pm 34.23 \mu\text{mol l}^{-1}$ and $1.26 \pm 2.15 \mu\text{mol l}^{-1}$, respectively (Fig. S3 and Table S1).

The autotrophic microbes decrease in abundance with depth at three stations at open sea. Specifically, the *Synechococcus*, *Prochlorococcus* and eukaryotic abundances varied from $3.31 \pm 0.25 \times 10^5 \text{ cells ml}^{-1}$ to $1.04 \pm 0.09 \times 10^4 \text{ cells ml}^{-1}$, $1.08 \pm 0.03 \times 10^5 \text{ cells ml}^{-1}$ to $9.10 \pm 4.30 \times 10^1 \text{ cells ml}^{-1}$ and $6.54 \pm 0.69 \times 10^3 \text{ cells ml}^{-1}$ to $1.50 \pm 3.60 \times 10^1 \text{ cells ml}^{-1}$, respectively (Table S1). The heterotrophic bacterial and viral abundances were $7.23 \pm 5.44 \times 10^5 \text{ cells ml}^{-1}$ and $6.41 \pm 4.24 \times 10^6 \text{ viruses ml}^{-1}$ on average, respectively, and tended to increase with depth at Stations A9 and J2 and decrease with depth at the other four stations. Horizontally, the heterotrophic bacterial abundance decreased from the coastal ($1.77 \pm 0.07 \times 10^6 \text{ cells ml}^{-1}$) to the offshore area ($7.21 \pm 0.02 \times 10^5 \text{ cells ml}^{-1}$) at the surface, while the viral abundance showed a maximum of $1.01 \pm 0.01 \times 10^7 \text{ viruses ml}^{-1}$ at Station J4.

Vertically, the lytic VP_R tended to increase with depth at five of the six stations, and the lysogenic VP_R increased with depth at Stations J2, K3, and SEATS but tended to decrease with depth at Stations A9, J4, and I1. Horizontally, the lytic VP_R tended to increase from the coast to the offshore area at the surface, while the lysogenic VP_R did not show a clear trend. The VD_R showed a downward trend with depth at each station and a weak rising trend from the coastal to the offshore area at the surface, with a range from $0.58 \pm 0.10\% \text{ h}^{-1}$ (30 m at Station A9) to $4.14 \pm 0.50\% \text{ h}^{-1}$ (5 m at Station I1). Overall, the distribution of microbiological parameters was consistent with previous studies in the SCS and other marine systems [31, 32, 42–48].

The virus-to-bacterium ratio (VBR) is an index to represent the relationship between viruses and bacteria and usually reflects the balance of VP and VD [49, 50]. In our investigation, the VBR ranged from 4.78 to 23.97 and was higher in the deeper layers of water than in the surface water (9.45 ± 1.35 vs. 9.15 ± 1.10 on average). However, there was no significant correlation between VBR and lytic VP_R , lysogenic VP_R and VD_R in our investigation (Pearson's correlation analysis, data not shown). We further found that only 57% of the total variation in the VBR was explained by viral abundance, PO_4^{3-} concentration and *Synechococcus* abundance (DISTLM analysis, Table S2). These suggest that additional environmental processes may strongly regulate the distribution of the VBR. In addition to viral particles originating from the in situ deep ocean, they can be continuously transported from surface water through several potential mechanisms. First, viral particles are transported into the deep ocean through adsorption on POM [23]. Second, infected cells were inevitably adsorbed by POM to transfer into the deep ocean, indirectly resulting in the sinking of numerous viral particles [24, 51]. Additionally, internal waves, mesoscale eddies, and other processes promote the mixing of water masses [52–54], driving the potentially vertical mixing of bacterioplankton and virioplankton. Therefore, the relatively high VBR in the deep ocean may be explained by viruses sinking from surface water [50, 55, 56]. Other factors influencing VBR are discussed further down.

Vertical transplantation affecting viral production and viral decay rates

The lytic VP_R of the transplantation treatments (i.e., the surface bacterial concentrate transplanted into virus-free seawater of each deeper layer) at each station was significantly higher than that of the surface water ($n = 12$, $p < 0.05$), at $9.21 \pm 4.92\% \text{ h}^{-1}$ vs. $3.81 \pm 1.66\% \text{ h}^{-1}$ on average (Fig. 2a, b). This increasing trend was most obvious at Station I1, where the lytic VP_R increased with ca. an




Fig. 2 Viral activity in the transplantation experiments. **a** Lytic and lysogenic viral production and viral decay in control and transplantation experiments performed in six stations. Error bars indicate the standard errors calculated from duplicate sample measurements. **b** Comparison of lytic and lysogenic viral production and viral decay in transplantation experiments with control treatments. Error bars indicate the standard deviations calculated from all measurements in control or in transplantation experiments. ns, no significant difference; * $p < 0.05$; ** $p < 0.01$.

order of magnitude from the surface water ($2.47 \pm 2.02\% \text{ h}^{-1}$) to the deepest layer transplantation treatments ($24.78 \pm 6.43\% \text{ h}^{-1}$), and the lowest increase was observed at Station SEATS, which exceeded 17%. This result is unexpected since the viral abundance and production showed a relatively low level in the deep ocean in our study (Table S1) and other deep ocean environments, which could be explained by the low temperature, high pressure and limited input of exogenous organic carbon [46, 47, 56].

Three possible reasons could result in an increase in lytic VP_R in the transplantation treatments: (1) increased host infection rate, (2) increased BS, and (3) switch from the lysogenic to lytic strategy. In the VP experiments, the production rate was estimated by the reduction and reoccurrence assay [36]. This dilution approach resulted in *ca.* an order of magnitude reduction in viral abundance, thus greatly decreasing the virus-host contact rate (i.e., blocking new infections) [37]. Therefore, the production of progeny viruses was likely not due to an enhanced host infection rate; rather it was due to the lysis of already infected cells. In fact, the VMM was lower in the transplantation treatments than in the surface water ($n = 12$, $p < 0.01$; Fig. 3a). This observation is most likely due to a decrease in temperature, which slows down intracellular viral replication in bacterial cells thus causing a delay of the latent period and a decline in VMM [57]. Interestingly, the BS was 3.89-fold higher in the transplantation treatments than in the surface water (Fig. 3b), which was consistent with the increase of BS in natural environment with depth [20]. An increase in BS implied that the host had increased metabolic activity during transplantation, which provided additional substances and energy for virus replication [51]. The composition of the viral particles is relatively rich in the elements of N and P [22], so viral replication in infected cells often is limited by the scarcity of P and N in oligotrophic surface oceans [58]. Inhibition of viral production was found during the incubation of both isolates and natural communities under the condition of nutrient limitation [59, 60]. Another study showed that the addition of inorganic nutrients increased the production rate of viral community by 14–52% in

dark incubation [61]. Motegi and colleagues also suggested that addition of inorganic P and N can enhance prokaryotic growth and viral production in the surface ocean [58, 60]. Indeed, in 5 of 6 stations, the concentrations of $\text{NO}_3^- + \text{NO}_2^-$, SiO_3^{2-} and PO_4^{3-} clearly increased with depth (Fig. S3). Therefore, when the surface bacteria were transplanted into the deep ocean, the bacterial activity could have been increased by surrounding sufficient nutrients, which in turn could facilitate the synthesis of progeny viruses in infected cells and increased VP [62]. In addition, our data showed that the increased BS offset the decline in VP due to reduced VMM and caused a general increase in lytic VP_R .

The shift from lysogeny to lysis could have been an alternative or additional reason for increased lytic VP_R in the deeper-layer transplanted incubation. Lysogeny is a viral life strategy in which the viral genome integrates into the host's genome to form prophages and initiate a symbiotic relationship with its host until the lytic cycle is induced [63]. Previous studies found that a high rate of lysogenic bacteria survived in conditions of low prokaryote abundance, activity, and production [49, 64]. In general, the lysogenic strategy has been reported to be dependent on changes in temperature, salinity and nutrient concentrations [49, 65–69], and environmental variation or shock was considered an inducing agent. For example, salt stress was found to shock the stability of bacteriophage $\lambda^{\text{imm}434}$ lysogens [70]. PO_4^{3-} was considered as an important factor in the lysis-lysogeny switch by previous studies. For example, Tuomi et al. suggested that the increase in inorganic P concentration may induce the lysis of some lysogenic bacterial populations in dark incubation, resulting in the release of viral particles [71]. Therefore, when surface host cells were transplanted into deeper seawater, they experienced changes in environmental factors, e.g., lower temperature, higher salinity, and higher concentrations of inorganic nutrients. The lytic VP_R can be increased by the switch from viral lysogenic to lytic strategies, which release progeny virus particles. Indeed, the FLC was significantly lower in the transplantation treatments than in surface water, $1.31 \pm 0.32\%$ vs. $3.30 \pm 0.53\%$ ($n = 12$, $p < 0.05$;




Fig. 3 Viral impacts on bacterioplankton in the transplantation experiments. **a** Virus-mediated bacterial mortality. **b** Burst size of lytic viral production. **c** Fraction of lysogenic bacterial cells. **d** Burst size of lysogenic viral production. Error bars indicate the standard deviations calculated from all measurements in control or in transplantation experiments. BS burst size, VP viral production; ns, no significant difference; * $p < 0.05$; ** $p < 0.01$.

Fig. 3c), suggesting that environmental changes due to transplantation can trigger the switch from lysogeny to lysis and cause lysis afterward to release more viral particles. Furthermore, the lysogenic VP_R was significantly higher in transplantation treatments than in surface water ($n = 12$, $p < 0.05$), with a range of increase from 28.0 to 180.7% (Fig. 2a, b), which was also probably due to the increased BS when the lysogenic strategy switched to the lytic strategy in the transplantation treatments ($n = 12$, $p < 0.05$; Fig. 3d). Altogether, the results suggested that vertical

sinking may enhance both lytic and lysogenic VP_R via the promotion of BS, probably due to additional nutrient supply, and trigger the switch from the lysogenic to lytic strategy, probably due to environmental changes.

In contrast to lytic and lysogenic VP_R, the VD_R in the surface water was significantly higher than that in the transplantation treatments ($n = 12$, $p < 0.01$) and tended to decrease with the depth of transplantation (Fig. 2a, b). Comparing the data between surface water and deepest layer transplantation treatments, the




Fig. 4 Schematic overview depicting the impact of sinking on viral eco-dynamics and life strategy. When the surface bacterial/viral communities sank into the deep ocean, both lytic and lysogenic VP_R significantly increased, which was mainly due to the increased BS , and triggered the switch from the lysogenic to lytic strategy, which was confirmed by the decline in the FLC . However, the VMM and VD_R decreased obviously, probably due to cooling inhibition.

strongest reduction was 59.1% at Station K3 and that the weakest reduction was 28.6% at Station J2. These results suggested that the VD_R was inhibited by sinking into the deep ocean, which may be mainly caused by the low temperature and low activity of extracellular enzymes [32, 42, 72]. Low temperature contributes to the stability of membrane lipids and the protein capsids of viral particles, e.g., it protects the biomolecular elasticity and molecular structure of proteins, thus reducing the degradation of viral particles [73]. In addition, when the surface viroplankton were transplanted into the deeper seawater, fewer extracellular enzymes due to less biomass produced and low temperature in the transplanted incubation could have decreased their activity [47, 74, 75], resulting in a reduction in VD_R .

Biogeochemical significance

Both bacteria colonized on POM and surrounding free-living bacteria are considered important drivers causing rapid degradation of the sinking POM [15, 76, 77]. Our study shows that viral particles continued to be produced by the lysis of infected bacteria in vertical transplantations, probably resulting in new infection and lysis events. Together with the low VD_R in the deep ocean, these viruses indirectly transmitted from surface water are an underlying mechanism contributing to the high VBR , a common phenomenon observed in deep-ocean environments, such as the North Atlantic Ocean [55], South Atlantic Ocean [78], and Pacific Ocean [79]. In

addition, viral lysis occurring during particle sinking will regulate bacterial diversity and community structure and, subsequently, impact the biological utilization and transformation of both particular and dissolved organic matters. This indirect influence of viruses on quantity and quality of organic matters can be revealed by monitoring bacterial and viral community with molecular ecological techniques, distinguishing group-specific responses with cell sorting techniques [80, 81], and detecting taxon-specific lysis with polony and ipolony techniques [82].

Viruses have been found to be beneficial for increasing the abundance, size, and stability of aggregates [83]. Attributed to lysis products and cell residues that acted as biological glue, causing the particles to stick together [65, 84], the release of cell contents by viral lysis could enhance aggregate formation [65, 85]. For example, an investigation suggested that aggregate formation and vertical carbon fluxes were enhanced by the Coccolithovirus infections of coccolithophore, *Emiliania huxleyi*, blooms in the North Atlantic [86]. These indicate that viral infection can “shuttle” surface organic particles into deep waters, and may promote the efficiency of BP in the ocean [87]. In this study, for the first time, we revealed that environmental changes caused by vertical transplantation decreased the VMM , but significantly increased lytic and lysogenic VP by increasing BS and triggering the switch from the lysogenic to lytic strategy to release additional viral particles into the surrounding water (Fig. 4). This could enhance viral productivity, potentially resulting in additional infection and lysis events afterwards. During sinking, such an effect would not only reduce the bacterial consumption of POM by viral lysis but also promote the formation and aggregation of POM by lysis products. This could provide a double-positive feedback for the efficiency of the BP hence increasing the downward flux of organic carbon. However, it is worth noting that the POM-aggregation can also lead to viral decay via irreversible binding, which could be a significant mechanism causing the removal of viruses in the ocean [18, 43, 73]. Hewson and Fuhrman reported that between 20 and 90% of natural marine viruses can be absorbed by POM, dependent on their concentration, size, and source [88]. Therefore, the interactions among particles, bacteria and viruses are complex and further investigations are required to evaluate the net effect of viruses on the efficiency of BP in the ocean.

REFERENCES

- Volk T, Hoffert MI. Ocean carbon pumps: Analysis of relative strengths and efficiencies in ocean-driven atmospheric CO_2 changes. In: Sundquist ET, Broecker WS. (eds). The carbon cycle and atmospheric CO_2 : Natural variations archean to present. American Geophysical Union, Geophysical Monograph, Washington, DC: 1985. p. 32:99–110.
- Scharek R, Tupas LM, Karl DM. Diatom fluxes to the deep sea in the oligotrophic North Pacific gyre at Station ALOHA. *Mar Ecol-Prog Ser.* 1999;182:55–67.
- Simon M, Grossart H, Schweitzer B, Ploug H. Microbial ecology of organic aggregates in aquatic ecosystems. *Aquat Micro Ecol.* 2002;28:175–211.
- Siegenthaler U, Sarmiento JL. Atmospheric carbon dioxide and the ocean. *Nature.* 1993;365:119–25.
- Ducklow H, Steinberg DK. Upper ocean carbon export and the biological pump. *Oceanography.* 2001;14:50–58.
- Jiao N, Herndl GJ, Hansell DA, Benner R, Kattner G, Wilhelm SW, et al. Microbial production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean. *Nat Rev Microbiol.* 2010;8:593–9.
- DeLong EF, Franks DG, Alldredge AL. Phylogenetic diversity of aggregate-attached vs. free-living marine bacterial assemblages. *Limnol Oceanogr.* 1993;38:924–34.
- Allen AE, Allen LZ, McCrow JP. Lineage specific gene family enrichment at the microscale in marine systems. *Curr Opin Microbiol.* 2013;16:605–17.
- D’Ambrosio L, Zierovogel K, MacGregor B, Teske A, Arnosti C. Composition and enzymatic function of particle-associated and free-living bacteria: a coastal/off-shore comparison. *ISME J.* 2014;8:2167–79.
- Martin JH, Knauer GA, Karl DM, Broenkow WW. VERTEX: carbon cycling in the northeast Pacific. *Deep-Sea Res Part I-Oceanogr Res Pap.* 1987;34:267–85.

11. Buesseler KO. The decoupling of production and particulate export in the surface ocean. *Glob Biogeochem Cycle*. 1998;12:297–310.
12. Schlitzer R. Applying the adjoint method for biogeochemical modeling: export of particulate organic matter in the world ocean. In: Kasibhata P, editor. *Inverse Methods in Global biogeochemical Cycles*. Washington, DC: American Geophysical Union; 2000. p. 114:107–24.
13. Steinberg DK, Van Mooy BAS, Buesseler KO, Boyd PW, Kobari T, Karl DM. Bacterial vs. zooplankton control of sinking particle flux in the ocean's twilight zone. *Limnol Oceanogr*. 2008;53:1327–38.
14. Cho BC, Azam F. Major role of bacteria in biogeochemical fluxes in the ocean's interior. *Nature*. 1988;332:441–3.
15. Herndl GJ, Reinthaler T. Microbial control of the dark end of the biological pump. *Nat Geosci*. 2013;6:718–24.
16. Bergh Ø, Borsheim KY, Bratbak G, Heldal M. High abundance of viruses found in aquatic environments. *Nature*. 1989;340:467–8.
17. Suttle CA. Viruses in the sea. *Nature*. 2005;437:356–61.
18. Zhang R, Wei W, Cai L. The fate and biogeochemical cycling of viral elements. *Nat Rev Microbiol*. 2014;12:850–1.
19. Middelboe M, Lyck PG. Regeneration of dissolved organic matter by viral lysis in marine microbial communities. *Aquat Micro Ecol*. 2002;27:187–94.
20. Weinbauer MG, Brettar I, Hofle MG. Lysogeny and virus-induced mortality of bacterioplankton in surface, deep, and anoxic marine waters. *Limnol Oceanogr*. 2003;48:1457–65.
21. Fuhrman JA. Marine viruses and their biogeochemical and ecological effects. *Nature*. 1999;399:541–8.
22. Jover LF, Effler TC, Buchan A, Wilhelm SW, Weitz JS. The elemental composition of virus particles: implications for marine biogeochemical cycles. *Nat Rev Microbiol*. 2014;12:519–28.
23. Bongiorno L, Magagnini M, Armeni M, Noble R, Danovaro R. Viral production, decay rates, and life strategies along a trophic gradient in the North Adriatic Sea. *Appl Environ Microbiol*. 2005;71:6644–50.
24. Weinbauer MG, Bettarel Y, Cattaneo R, Luef B, Maier C, Motegi C, et al. Viral ecology of organic and inorganic particles in aquatic systems: avenues for further research. *Aquat Micro Ecol*. 2009;57:321–41.
25. Tian Y, Cai L, Xu Y, Luo T, Zhao Z, Wang Q, et al. Stability and infectivity of allochthonous viruses in deep sea: A long-term high pressure simulation experiment. *Deep-Sea Res Part I-Oceanogr Res Pap*. 2020;161:103302.
26. Lara E, Vaqué D, Sà EL, Boras JA, Gomes A, Borrull E, et al. Unveiling the role and life strategies of viruses from the surface to the dark ocean. *Sci Adv*. 2017;3:e1602565.
27. Zhang R, Li Y, Yan W, Wang Y, Cai L, Luo T, et al. Viral control of biomass and diversity of bacterioplankton in the deep sea. *Commun Biol*. 2020;3:256.
28. Woźniak SB, Stramski D, Stramska M, Reynolds RA, Wright VM, Miksic EY, et al. Optical variability of seawater in relation to particle concentration, composition, and size distribution in the nearshore marine environment at Imperial Beach, California. *J Geophys Res*. 2010;115:C08027.
29. White AE, Letelier RM, Whitmire AL, Barone B, Bidigare RR, Church MJ, et al. Phenology of particle size distributions and primary productivity in the North Pacific subtropical gyre (Station ALOHA). *J Geophys Res-Oceans*. 2015;120:7381–99.
30. Vault D, Courties C, Partensky F. A simple method to preserve oceanic phytoplankton for flow cytometric analyses. *Cytom Part A*. 1989;10:629–35.
31. Chen X, Liu H, Weinbauer M, Chen B, Jiao N. Viral dynamics in the surface water of the western South China Sea in summer 2007. *Aquat Micro Ecol*. 2011;63:145–60.
32. Wei W, Zhang R, Peng L, Liang Y, Jiao N. Effects of temperature and photosynthetically active radiation on virioplankton decay in the western Pacific Ocean. *Sci Rep*. 2018;8:1525–34.
33. Marie D, Partensky F, Vault D, Brussaard C. Numeration of phytoplankton, bacteria and viruses in marine samples. *Curr Protoc Cytom*. 1999;11:1–15.
34. Marie D, Brussaard CPD, Thyrhaug R, Bratbak G, Vault D. Enumeration of marine viruses in culture and natural samples by flow cytometry. *Appl Environ Microbiol*. 1999;65:45–52.
35. Brussaard CP. Optimization of procedures for counting viruses by flow cytometry. *Appl Environ Microbiol*. 2004;70:1506–13.
36. Wilhelm SW, Brigden SM, Suttle CA. A dilution technique for the direct measurement of viral production: a comparison in stratified and tidally mixed coastal waters. *Micro Ecol*. 2002;43:168–73.
37. Weinbauer MG, Rowe JM, Wilhelm SW. Determining rates of virus production in aquatic systems by the virus reduction approach. In: Wilhelm SW, Weinbauer MG, Suttle CA. (eds) *Manual of Aquatic Viral Ecology*. American Society of Limnology and Oceanography Inc., Waco, TX: 2010. p. 1–8.
38. Chen X, Wei W, Wang J, Li H, Sun J, Ma R, et al. Tide driven microbial dynamics through virus-host interactions in the estuarine ecosystem. *Water Res*. 2019;160:118–29.
39. Luef B, Luef F, Peduzzi P. Online program 'vipcal' for calculating lytic viral production and lysogenic cells based on a viral reduction approach. *Environ Microbiol Rep*. 2009;1:78–85.
40. Winget DM, Helton RR, Williamson KE, Bench SR, Williamson SJ. Repeating patterns of virioplankton production within an estuarine ecosystem. *Proc Natl Acad Sci USA*. 2011;108:11506–11.
41. Wei W, Wang N, Cai L, Zhang C, Jiao N, Zhang R. Impacts of freshwater and seawater mixing on the production and decay of virioplankton in a subtropical estuary. *Micro Ecol*. 2019;78:843–54.
42. Noble RT, Fuhrman JA. Virus decay and its causes in coastal waters. *Appl Environ Microbiol*. 1997;63:77–83.
43. Suttle CA, Chen F. Mechanisms and rates of decay of marine viruses in seawater. *Appl Environ Microbiol*. 1992;58:3721–9.
44. Rowe JM, Saxton MA, Cottrell MT, DeBruyn JM, Berg GM, Kirchman DL, et al. Constraints on viral production in the Sargasso Sea and North Atlantic. *Aquat Micro Ecol*. 2008;52:233–44.
45. Evans C, Pearce I, Brussaard CP. Viral-mediated lysis of microbes and carbon release in the sub-Antarctic and Polar Frontal zones of the Australian Southern Ocean. *Environ Microbiol*. 2009;11:2924–34.
46. De Corte D, Sintès E, Winter C, Yokokawa T, Reinthaler T, Herndl GJ. Links between viral and prokaryotic communities throughout the water column in the (sub)tropical Atlantic Ocean. *ISME J*. 2010;4:1431–42.
47. Li Y, Lou T, Sun J, Cai L, Liang Y, Jiao N, et al. Lytic viral infection of bacterioplankton in deep waters of the western Pacific Ocean. *Biogeosciences*. 2014;11:2531–42.
48. Liang Y, Zhang Y, Zhang Y, Luo T, Rivkin R, Jiao N. Distributions and relationships of virio- and picoplankton in the epi-, meso- and bathypelagic zones of the Western Pacific Ocean. *FEMS Microbiol Ecol*. 2017;93:fiw238.
49. Wommack KE, Colwell RR. Virioplankton: viruses in aquatic ecosystems. *Microbiol Mol Biol Rev*. 2000;64:69–114.
50. Parikka KJ, Le Romancer M, Wauters N, Jacquet S. Deciphering the virus-to-prokaryote ratio (VPR): insights into virus-host relationships in a variety of ecosystems. *Biol Rev*. 2016;92:1081–1100.
51. Parada V, Herndl GJ, Weinbauer MG. Viral burst size of heterotrophic prokaryotes in aquatic systems. *J Mar Biol Assoc UK*. 2006;86:613–21.
52. Yuan D. A numerical study of the South China Sea deep circulation and its relation to the Luzon Strait transport. *Acta Oceano Sin*. 2002;21:187–202.
53. Tian J, Yang Q, Zhao W. Enhanced diapycnal mixing in the South China Sea. *J Phys Oceanogr*. 2009;39:3191–203.
54. Alford MH, Lien R, Simmons H, Klymak J, Ramp S, Yang YJ, et al. Speed and evolution of nonlinear internal waves transiting the South China Sea. *J Phys Oceanogr*. 2010;40:1338–55.
55. Parada V, Sintès E, Van Aken HM, Weinbauer MG, Herndl GJ. Viral abundance, decay, and diversity in the meso- and bathypelagic waters of the north atlantic. *Appl Environ Microbiol*. 2007;73:4429–38.
56. De Corte D, Sintès E, Yokokawa T, Reinthaler T, Herndl GJ. Links between viruses and prokaryotes throughout the water column along a North Atlantic latitudinal transect. *ISME J*. 2012;6:1566–77.
57. Zachary A. An ecological study of bacteriophages of *Vibrio natriegens*. *Appl Environ Microbiol*. 1978;24:321–4.
58. Motegi C, Nagata T. Enhancement of viral production by addition of nitrogen or nitrogen plus carbon in subtropical surface waters of the South Pacific. *Aquat Micro Ecol*. 2007;48:27.
59. Bratbak G, Egge JK, Heldal M. Viral mortality of the marine alga *Emiliania huxleyi* (Haptophyceae) and termination of algal blooms. *Mar Ecol-Prog Ser*. 1993;93:39–48.
60. Motegi C, Kaiser K, Benner R, Weinbauer MG. Effect of P-limitation on prokaryotic and viral production in surface waters of the Northwestern Mediterranean Sea. *J Plankton Res*. 2015;37:16–20.
61. Hewson I, O'Neil JM, Fuhrman JA, Dennison WC. Virus-like particle distribution and abundance in sediments and overlying waters along eutrophication gradients in two subtropical estuaries. *Limnol Oceanogr*. 2001;46:1734–46.
62. Wilson WH, Mann NH. Lysogenic and lytic viral production in marine microbial communities. *Aquat Micro Ecol*. 1997;13:95–100.
63. Paul JH. Prophages in marine bacteria: dangerous molecular time bombs or the key to survival in the sea? *ISME J*. 2008;2:579–89.
64. Chibani-Chennoufi S, Bruttin A, Dillmann ML, Brussow H. Phage-host interaction: an ecological perspective. *J Bacteriol*. 2004;186:3677–86.
65. Weinbauer MG. Ecology of prokaryotic viruses. *FEMS Microbiol Rev*. 2004;28:127–81.
66. Williamson SJ, Paul JH. Nutrient stimulation of lytic phage production in bacterial populations of the Gulf of Mexico. *Aquat Micro Ecol*. 2004;36:9–17.
67. Williamson SJ, Paul JH. Environmental factors that influence the transition from lysogenic to lytic existence in the ϕ HSIC/Listonella pelagia marine phage-host system. *Micro Ecol*. 2006;52:217–25.

68. Cissoko M, Desnues A, Bouvy M, Sime-Ngando T, Verling E, Bettarel Y. Effects of freshwater and seawater mixing on viro- and bacterioplankton in a tropical estuary. *Freshw Biol.* 2008;53:1154–62.
69. Bettarel Y, Bouvier T, Agis M, Bouvier C, Van Chu T, Combe M, et al. Viral distribution and life strategies in the Bach Dang Estuary, Vietnam. *Micro Ecol.* 2011;62:143–54.
70. Shkilnyj P, Koudelka GB. Effect of salt shock on stability of λ^{imm434} lysogens. *J Bacteriol.* 2007;189:3115–23.
71. Tuomi P, Fagerbakke KM, Bratbak G, Heldal M. Nutritional enrichment of a microbial community: the effects on activity, elemental composition, community structure and virus production. *FEMS Microbiol Ecol.* 1995;16:23–134.
72. Dell'Anno A, Corinaldesi C, Danovaro R. Virus decomposition provides an important contribution to benthic deep-sea ecosystem functioning. *Proc Natl Acad Sci USA.* 2015;112:E2014–E2019.
73. Mojica KD, Brussaard CP. Factors affecting virus dynamics and microbial host-virus interactions in marine environments. *FEMS Microbiol Ecol.* 2014;89:495–515.
74. Zweifel UL. Factors controlling accumulation of labile dissolved organic carbon in the Gulf of Riga. *Estuar Coast Shelf Sci.* 1999;48:357–70.
75. Pomeroy LR, Wiebe WJ. Temperature and substrates as interactive limiting factors for marine heterotrophic bacteria. *Aquat Micro Ecol.* 2001;23:187–204.
76. Ploug H, Grossart H, Azam F, Jørgensen BB. Photosynthesis, respiration, and carbon turnover in sinking marine snow from surface waters of Southern California Bight: implications for the carbon cycle in the ocean. *Mar Ecol-Prog Ser.* 1999;179:1–11.
77. Azam F, Malfatti F. Microbial structuring of marine ecosystems. *Nature.* 2007;5:782–91.
78. De Corte D, Sintes E, Yokokawa T, Lekunberri I, Herndl GJ. Large-scale distribution of microbial and viral populations in the South Atlantic Ocean. *Environ Microbiol Rep.* 2016;8:305–15.
79. Yang YH, Yokokawa T, Motegi C, Nagata T. Large-scale distribution of viruses in deep waters of the Pacific and Southern Oceans. *Aquat Micro Ecol.* 2014;71:193–202.
80. Labonté JM, Swan BK, Poulos B, Luo H, Koren S, Hallam SJ, et al. Single-cell genomics-based analysis of virus-host interactions in marine surface bacterioplankton. *ISME J.* 2015;9:2386–99.
81. Martínez-Hernández F, Fornas Ò, Lluésma Gomez M, García-Heredia I, Maestre-Carballa L, López-Pérez M, et al. Single-cell genomics uncover Pelagibacter as the putative host of the extremely abundant uncultured 37-F6 viral population in the ocean. *ISME J.* 2019;13:232–6.
82. Mruwat N, Carlson MCG, Goldin S, Ribalet F, Kirzner S, Hulata Y, et al. A single-cell polony method reveals low levels of infected *Prochlorococcus* in oligotrophic waters despite high cyanophage abundances. *ISME J.* 2021;15:41–54.
83. Peduzzi P, Weinbauer M. Effect of concentrating the virus-rich 2–200 nm size fraction of seawater on the formation of algal flocs (marine snow). *Limnol Oceanogr.* 1993;38:1562–5.
84. Uitz J, Stramski D, Baudoux A, Reynolds RA, Wright VM, Dubranna J, et al. Variations in the optical properties of a particle suspension associated with viral infection of marine bacteria. *Limnol Oceanogr.* 2010;55:2317–30.
85. Sullivan MB, Weitz JS, Wilhelm SW. Viral ecology comes of age. *Environ Microbiol Rep.* 2017;9:33–35.
86. Laber CP, Hunter JE, Carvalho F, Collins JR, Hunter EJ, Schieler BM, et al. Coccolithovirus facilitation of carbon export in the North Atlantic. *Nat Microbiol.* 2018;3:537–47.
87. Kranzler CF, Brzezinski MA, Cohen NR, Lampe RH, Maniscalco M, Till CP, et al. Impaired viral infection and reduced mortality of diatoms in iron-limited oceanic regions. *Nat Geosci.* 2021;4:231–7.
88. Hewson I, Fuhrman JA. Viriobenthos production and virioplankton sorptive scavenging by suspended sediment particles in coastal and pelagic waters. *Micro Ecol.* 2003;46:337–47.

ACKNOWLEDGEMENTS

The authors wish to thank Prof. Farooq Azam of University of California San Diego for his constructive reviews to improve the manuscript and Prof. Jun Sun of Tianjin University of Science and Technology for providing some environmental data. This work was supported by the National Natural Science Foundation of China (42188102, 91951209, 41906085, 41861144018) and the China Postdoctoral Science Foundation (2019M662236, 2020T130066ZX). Markus Weinbauer was supported by the international collaborative project on “Marine Biogeochemistry and Ecotoxicology” funded by the Program of Introducing Talents of Discipline to Universities (BP0719030).

AUTHOR CONTRIBUTIONS

WW, NJ and RZ conceived and designed the study. WW and XC carried out the experiments. WW, XC and RZ performed the data analysis and figure drawing. MGW and RZ provided critical biological interpretations of the data. NJ and RZ supervised this work. WW wrote the original draft and all authors reviewed and edited the draft manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41396-022-01224-9>.

Correspondence and requests for materials should be addressed to Nianzhi Jiao or Rui Zhang.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022