



HAL
open science

Long-term air pollution exposure and Parkinson's disease mortality in a large pooled European cohort: An ELAPSE study

Tom Cole-Hunter, Jiawei Zhang, Rina So, E. Samoli, Shuo Liu, Jie Chen, Maciej Strak, Kathrin Wolf, Gudrun Weinmayr, Sophia Rodopoulou, et al.

► To cite this version:

Tom Cole-Hunter, Jiawei Zhang, Rina So, E. Samoli, Shuo Liu, et al.. Long-term air pollution exposure and Parkinson's disease mortality in a large pooled European cohort: An ELAPSE study. *Environment International*, 2023, 171, 10.1016/j.envint.2022.107667 . hal-03949602

HAL Id: hal-03949602

<https://hal.science/hal-03949602v1>

Submitted on 10 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Contents lists available at ScienceDirect

Environment International

journal homepage: www.elsevier.com/locate/envint

Full length article



Long-term air pollution exposure and Parkinson's disease mortality in a large pooled European cohort: An ELAPSE study

Thomas Cole-Hunter^a, Jiawei Zhang^a, Rina So^a, Evangelia Samoli^b, Shuo Liu^a, Jie Chen^c, Maciej Strak^{c,d}, Kathrin Wolf^e, Gudrun Weinmayr^f, Sophia Rodopoulou^b, Elizabeth Remfry^g, Kees de Hoogh^{h,i}, Tom Bellander^{j,k}, Jørgen Brandt^{l,m}, Hans Concinaⁿ, Emanuel Zitt^{n,o}, Daniela Fecht^p, Francesco Forastiere^{q,r}, John Gulliver^{p,s}, Barbara Hoffmann^t, Ulla A. Hvidtfeldt^u, Karl-Heinz Jöckel^v, Laust H. Mortensen^{w,x}, Matthias Ketzel^{l,y}, Diego Yacamán Méndez^{z,ai}, Karin Leander^j, Petter Ljungman^{j,aa}, Elodie Faure^{ab}, Pei-Chen Lee^{ab,ac}, Alexis Elbaz^{ab}, Patrik K.E. Magnusson^{ad}, Gabriele Nagel^f, Göran Pershagen^{j,k}, Annette Peters^{e,ae}, Debora Rizzuto^{af,ag}, Roel C.H. Vermeulen^{c,ah}, Sara Schramm^v, Massimo Stafoggia^{j,q}, Klea Katsouyanni^{b,r}, Bert Brunekreef^c, Gerard Hoek^c, Youn-Hee Lim^a, Zorana J. Andersen^{a,*}

^a Section of Environmental Health, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

^b Department of Hygiene, Epidemiology and Medical Statistics, Medical School, National and Kapodistrian University of Athens, Athens, Greece

^c Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands

^d National Institute for Public Health and the Environment, Bilthoven, the Netherlands

^e Institute of Epidemiology, Helmholtz Zentrum München, Neuherberg, Germany

^f Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany

^g Wolfson Institute of Population Health, Queen Mary University of London, United Kingdom

^h Swiss Tropical and Public Health Institute, Basel, Switzerland

ⁱ University of Basel, Basel, Switzerland

^j Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

^k Centre for Occupational and Environmental Medicine, Region Stockholm, Stockholm, Sweden

^l Department of Environmental Science, Aarhus University, Roskilde, Denmark

^m iClimate, interdisciplinary Centre for Climate Change, Aarhus University, Roskilde, Denmark

ⁿ Agency for Preventive and Social Medicine (aks), Bregenz, Austria

^o Department of Internal Medicine 3, LKH Feldkirch, Feldkirch, Austria

^p MRC Centre for Environment and Health, School of Public Health, Imperial College London, London, United Kingdom

^q Department of Epidemiology, Lazio Region Health Service / ASL Roma 1, Rome, Italy

^r MRC Centre for Environment and Health, Environmental Research Group, School of Public Health, Imperial College London, London, United Kingdom

^s Centre for Environmental Health and Sustainability & School of Geography, Geology and the Environment, University of Leicester, Leicester, United Kingdom

^t Institute for Occupational, Social and Environmental Medicine, Centre for Health and Society, Medical Faculty, Heinrich Heine University Düsseldorf, Germany

^u Danish Cancer Society Research Center, Copenhagen, Denmark

^v Institute for Medical Informatics, Biometry and Epidemiology, Medical Faculty, University of Duisburg-Essen, Essen, Germany

^w Section of Epidemiology, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

^x Statistics Denmark, Copenhagen, Denmark

^y Global Centre for Clean Air Research (GCARE), University of Surrey, Guildford GU2 7XH, United Kingdom

^z Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

^{aa} Department of Cardiology, Danderyd University Hospital, Stockholm, Sweden

^{ab} University Paris-Saclay, UVSQ, Inserm, Gustave Roussy, "Exposome and Heredity" team, CESP UMR1018, 94805 Villejuif, France

^{ac} Department of Public Health, National Cheng Kung University, Tainan, Taiwan

^{ad} Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

^{ae} Ludwig Maximilians Universität München, München, Germany

^{af} Department of Neurobiology, Care Sciences, and Society, Karolinska Institutet and Stockholm University, Stockholm, Sweden

^{ag} Stockholm Gerontology Research Center, Stockholm, Sweden

^{ah} Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands

^{ai} Centre for Epidemiology and Community Medicine, Region Stockholm, Stockholm, Sweden

* Corresponding author.

E-mail address: zorana.andersen@sund.ku.dk (Z.J. Andersen).

<https://doi.org/10.1016/j.envint.2022.107667>

Received 29 August 2022; Received in revised form 22 November 2022; Accepted 27 November 2022

Available online 30 November 2022

0160-4120/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

ARTICLE INFO

Handling Editor: Adrian Covaci

Keywords:

Air pollution
Adults
Parkinson's Disease
Long-term exposure
Low-level exposure
Pooled-cohort study

ABSTRACT

Background: The link between exposure to ambient air pollution and mortality from cardiorespiratory diseases is well established, while evidence on neurodegenerative disorders including Parkinson's Disease (PD) remains limited.

Objective: We examined the association between long-term exposure to ambient air pollution and PD mortality in seven European cohorts.

Methods: Within the project 'Effects of Low-Level Air Pollution: A Study in Europe' (ELAPSE), we pooled data from seven cohorts among six European countries. Annual mean residential concentrations of fine particulate matter (PM_{2.5}), nitrogen dioxide (NO₂), black carbon (BC), and ozone (O₃), as well as 8 PM_{2.5} components (copper, iron, potassium, nickel, sulphur, silicon, vanadium, zinc), for 2010 were estimated using Europe-wide hybrid land use regression models. PD mortality was defined as underlying cause of death being either PD, secondary Parkinsonism, or dementia in PD. We applied Cox proportional hazard models to investigate the associations between air pollution and PD mortality, adjusting for potential confounders.

Results: Of 271,720 cohort participants, 381 died from PD during 19.7 years of follow-up. In single-pollutant analyses, we observed positive associations between PD mortality and PM_{2.5} (hazard ratio per 5 µg/m³: 1.25; 95% confidence interval: 1.01–1.55), NO₂ (1.13; 0.95–1.34 per 10 µg/m³), and BC (1.12; 0.94–1.34 per 0.5 × 10⁻⁵ m⁻¹), and a negative association with O₃ (0.74; 0.58–0.94 per 10 µg/m³). Associations of PM_{2.5}, NO₂, and BC with PD mortality were linear without apparent lower thresholds. In two-pollutant models, associations with PM_{2.5} remained robust when adjusted for NO₂ (1.24; 0.95–1.62) or BC (1.28; 0.96–1.71), whereas associations with NO₂ or BC attenuated to null. O₃ associations remained negative, but no longer statistically significant in models with PM_{2.5}. We detected suggestive positive associations with the potassium component of PM_{2.5}.

Conclusion: Long-term exposure to PM_{2.5}, at levels well below current EU air pollution limit values, may contribute to PD mortality.

1. Introduction

The association between exposure to air pollution and cardiovascular and respiratory diseases is well documented (WHO, 2021). Recently, attention has been drawn to the potential negative impact of air pollution on disorders that affect the brain and central nervous system, including neurodegenerative diseases such as dementia, Alzheimer's Disease, and Parkinson's Disease (PD) (Kim et al., 2020; Raggi and Leonardi, 2020). PD is the second most prevalent neurodegenerative disorder, affecting an estimated six million individuals globally, preceded in prevalence only by Alzheimer's Disease (Dorsey et al., 2018). Known risk factors for PD include age, being male, smoking, high consumption of dairy, exposure to pesticides, and genetic predisposition (Ascherio and Schwarzschild, 2016).

It has been suggested that both gaseous pollutants and fine particles can directly or indirectly damage the brain by crossing the blood–brain barrier, causing oxidative stress, neuroinflammation and abnormal aggregation of proteins, damaging the olfactory bulb and frontal cortex, leading to the development of PD (Ascherio and Schwarzschild, 2016). Despite PD being a progressive disease that leads to death in older age, epidemiological evidence on long-term exposure to air pollution and PD consists mainly of studies on the incidence of (rather than mortality from) PD. These studies on incidence present mixed findings, with some studies reporting associations with particulate matter of diameter < 10 µm (PM₁₀), particulate matter of diameter < 2.5 µm (PM_{2.5}), nitrogen dioxide (NO₂) or ozone (O₃) (Chen et al., 2017; Kirrane et al., 2015; Rhew et al., 2021;16(7):e0253253.; Ritz et al., 2016; Shi et al., 2020; Shin et al., 2018; Yu et al., 2021; Yuchi et al., 2020), while other studies find no associations (Cerza et al., 2018; Lee et al., 2016; Liu et al., 2016; Palacios et al., 2017; Palacios et al., 2014; Toro et al., 2019) (see supplemental Table S1). Only two studies examined long-term exposure to air pollution with respect to PD mortality: one only examining O₃, detecting significant positive associations (Zhao et al., 2021), and another examining PM_{2.5} and finding suggestive positive associations, which were weaker than those detected with incidence (compared to mortality) of PD in the same study (Rhew et al., 2021). Two studies went further to examine the association of specific components of PM_{2.5} and PD. Nunez and colleagues examined black carbon (BC), organic matter (OM), nitrate, sulfate, sea salt, and soil PM_{2.5} components, detecting

associations of nitrate and OM with PD hospitalization (Nunez et al., 2021). Palacios and colleagues suggested an association of mercury with PD incidence (Palacios et al., 2014). Therefore, further study is warranted to elucidate the long-term effects of specific air pollution components, and strengthen the evidence base for potential effects at low pollutant concentrations, on mortality from neurodegenerative diseases including PD among large cohorts, such as through the 'Effects of Low-Level Air Pollution: A Study in Europe' (ELAPSE) project. ELAPSE has recently been able to show that long-term exposure to low levels of air pollution was associated with premature mortality (Stafoggia et al., 2022; Strak et al., 2021), as well as incidence of cardiovascular diseases (Wolf et al., 2021), adult-onset asthma (Liu et al., 2021), chronic obstructive pulmonary disease (COPD) (Liu et al., 2021), lung cancer (Hvidtfeldt et al., 2021) and liver cancer (So et al., 2021), as well as mortality from dementia, psychiatric disorders, and suicide (Andersen et al., 2022). Here, we aimed to examine the association of long-term exposure to PM_{2.5}, NO₂, BC, and O₃, as well as eight PM_{2.5} components, with mortality from PD among seven cohorts of the ELAPSE project.

2. Materials and methods

2.1. Study population

Within the ELAPSE project, we analysed pooled data from seven cohorts (and their sub-cohorts) among six European countries (see supplemental Figure S1), including information on potential confounders:

- Cardiovascular Effects of Air Pollution and Noise in Stockholm (CEANS) cohort in Sweden, which combined four sub-cohorts: Stockholm Diabetes Prevention Program (SDPP) (Eriksson et al., 2008); Stockholm Cohort of 60-year-olds (SIXTY) (Wändell et al., 2007); Stockholm Screening Across the Lifespan Twin study (SALT) (Magnusson et al., 2013); Swedish National Study on Aging and Care in Kungsholmen (SNAC-K) (Lagergren et al., 2004);
- Danish Nurse Cohort (DNC) in Denmark (Hundrup et al., 2012), including two cohort recruitment rounds in 1993 and 1999;

- (c) Etude Epidémiologique auprès de femmes de la Mutuelle Générale de l'Education Nationale (E3N) in France (Clavel-Chapelon, 2015);
- (d) European Prospective Investigation into Cancer and Nutrition-Netherlands (EPIC-NL) cohort in the Netherlands, which included two sub-cohorts: Monitoring Project on Risk Factors and Chronic Diseases in the Netherlands (Morgen) and Prospect (Beulens et al., 2010);
- (e) Heinz Nixdorf Recall study (HNR) in Germany (Schmermund et al., 2002);
- (f) Cooperative Health Research in the Region of Augsburg (KORA) in Germany (Holle et al., 2005), combining two sub-cohorts from baseline rounds in 1994–1995 (S3) and 1999–2001 (S4); and;
- (g) Vorarlberg Health Monitoring and Prevention Programme (VHM&PP) in Austria (Ulmer et al., 2007).

The cohorts were recruited in the 1990s or early 2000s, from one or several large cities and their surrounding towns, except for the two nationwide cohorts E3N and DNC. Detailed information of each cohort has been described previously (Chen et al., 2021). All cohorts were approved by the medical ethics committees in their respective countries.

2.2. Air pollution exposure assessment

The method for assessment of air pollution exposure via modelling has previously been described in detail (Hvidtfeldt et al., 2021; Liu et al., 2021; de Hoogh et al., 2018). Briefly, annual mean concentrations of PM_{2.5}, NO₂, BC, and O₃ (limited to the maximum running 8-hour average in the boreal warm season, April–September) were estimated for 2010 at participants' baseline residential addresses utilising standardized Europe-wide hybrid land use regression (LUR) models (de Hoogh et al., 2018). These models were developed based on air pollution concentration data from routine and research study monitors, satellites, estimates from chemical transport models, and land use and road variables as predictors.

PM_{2.5}, NO₂, BC, and O₃ LUR models used a 100 m × 100 m (grid) spatial scale that performed satisfactorily in fivefold hold-out validation, explaining 66%, 58%, 51 and 60%, respectively, of measured spatial variation. BC was measured by the reflectance of PM_{2.5} filters from 2009 and 2010, expressed in absorbance units.

In addition, we extrapolated pollutant concentrations for each year from baseline to the end of follow-up (Hvidtfeldt et al., 2021; Liu et al., 2021), and incorporated dynamic residential address history during follow-up. The extrapolation method was applied by utilising estimated monthly average concentrations, at a spatial resolution of 26 km × 26 km, from the Danish Eulerian Hemispheric Model (DEHM) since 1990 (Brandt et al., 2012). The extrapolation was performed after checking the agreement between ground measurements versus DEHM predictions for the four pollutants, as detailed previously to show little impact by temporal misalignment (Chen et al., 2021). The DEHM provided predicted modelling exposure data for a complete database to perform harmonious extrapolation for the four pollutants; ground measurements for PM_{2.5} components were not available back in time. Pollutant concentrations were extrapolated for cohorts with available residential history information, using both a difference method and a ratio method with 2010 as the reference year (Stafoggia et al., 2022; Strak et al., 2021).

We estimated exposure to eight components of PM_{2.5} for 2010 at the participants' baseline residential addresses using Europe-wide LUR models based on the standardized ESCAPE project monitoring data, with model details published previously (Chen et al., 2020). The eight components were selected to represent major pollution sources: copper (Cu), iron (Fe), and zinc (Zn) for non-tailpipe traffic emissions; sulfur (S) for long-range transport of secondary inorganic aerosols; nickel (Ni) and vanadium (V) for mixed oil burning/industry; silicon (Si) for earth

crustal material; and, potassium (K) for biomass burning. We included large-scale satellite-model and chemical transport-model estimates of components to represent background concentrations, in addition to land-use, road, population, and industrial point source data to model local spatial variability. We applied two algorithms, supervised linear regression (SLR) (de Hoogh et al., 2018), and random forest (RF) (Chen et al., 2019), to develop models for the eight components. The models explained a moderate-to-large fraction of the measured concentration variation at the European scale (ranging from 41% to 90% across components).

2.3. Mortality outcome definition

We defined PD mortality from mortality registries, based solely on the underlying cause of death (as contributing causes of death were not available) using the International Classification of Diseases (ICD)-9: 332 (PD); and, ICD-10: G20 (PD), G21 (secondary Parkinsonism), G22 (Parkinsonism in diseases classified elsewhere), and F02.3 (dementia in PD).

2.4. Statistical analysis

We performed Cox proportional hazards models with age as the underlying timescale (Thiébaud and Bénichou, 2004) to examine associations between long-term exposure to air pollution and PD, following the general analytical framework of ELAPSE (Samoli et al., 2021). Censoring occurred at the time of event of interest, death from other causes, emigration, loss to follow-up, or the end of follow-up (ranging from 2011 to 2015 depending on sub-cohort), whichever came first. The start of follow-up was the year of enrolment in individual cohorts which ranged from 1985 to 2001 (supplemental Table S2). Air pollution exposure was included in models as a linear term. We examined associations using three models including *a priori* defined individual and area-level covariates, all assessed at the cohort baseline. Model 1 included age (time axis), sex (strata), sub-cohort (strata), and calendar year of baseline (year of enrolment). Model 2 was additionally adjusted for smoking status (never, former, current), smoking duration (years) for current smokers, smoking intensity (linear and squared term: cigarettes/day) for current smokers, body mass index (BMI, categories: <18.5, 18.5–24.9, 25.0–29.9, and ≥ 30 kg/m²), marital status (married/cohabiting, divorced/separated, single, widowed), and employment status (employed/self-employed, other). Model 3 (main model) was further adjusted for area-level mean annual (2001) income as a proxy for socio-economic status (SES). Only participants with complete exposure and covariate information for Model 3 were included in the analyses to ensure comparability among model results.

For explorative purposes, we performed a subset analysis with Model 3 by excluding participants of exposure levels above certain pre-defined values (PM_{2.5}: 25, 20, 15 µg/m³; NO₂: 40, 30, 20 µg/m³; BC: 3, 2.5, 2, 1.5 × 10⁻⁵ m⁻¹; O₃: 120, 100 µg/m³), based partially on existing EU and US limit values and previous (2005) WHO guidelines. To assess the shape of concentration–response functions (associations from Model 3), we modelled pollutants as natural cubic splines with two degrees of freedom and tested for deviation of linearity by comparing with linear models using a likelihood ratio test. Further, for explorative purposes, we assessed a potential effect modification on these associations by age (<65 or ≥ 65 years old), sex (female or male), overweight status (BMI ≥ 25 kg/m² or not), smoking status (current, former, or never smoker), and employment status, by including an interaction term into Model 3 tested by the Wald test. In addition, we performed two-pollutant models based on Model 3 to investigate the contribution of individual pollutants.

We performed several sensitivity analyses to check the robustness of our associations. We compared the results of year 2010 exposure in Model 3 with results of back-extrapolated baseline year exposures and time-varying annual exposures. Time-varying (exposure) analyses were

performed for cohorts with available information on residential address history (CEANS, EPIC-NL, and VHM&PP), with 1-year strata of calendar time to account for time trends in air pollution and mortality. Finally, we compared effect estimates in Model 3 using different datasets excluding one cohort at a time.

The results are presented as hazard ratios (HR) and 95% confidence intervals (CI) for pollutant unit increases of $5 \mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$, $10 \mu\text{g}/\text{m}^3$ for NO_2 , $0.5 \times 10^{-5}\text{m}^{-1}$ for BC, $10 \mu\text{g}/\text{m}^3$ for O_3 , and IQR for $\text{PM}_{2.5}$ composition. All statistical analyses were performed in software R (version 3.4.0).

3. Results

3.1. Population description

Our pooled cohort included 324,728 participants. We excluded

53,008 of those participants with covariate data missing for Model 3. Of 271,720 remaining participants in the final analyses, 381 died from PD (380 from PD and 1 from secondary Parkinsonism) during a mean follow-up time of 19.7 years (Table 1) and a mean age at end of follow-up of 66.9 years (supplemental Table S2). The large VHM&PP cohort contributed with the majority (284, or ~ 75%) of the deaths. Baseline characteristics of participants varied widely across sub-cohorts (Table 1), supporting the use of strata for sub-cohorts to adjust for differences in baseline hazard. Mean age of participants at baseline was 47.1 years, ranging from 42.1 in VHM&PP to 72.9 years in CEANS-SNACK. The majority of participants (69%) were female, as three cohorts/sub-cohorts were female-only by design (DNC, E3N, and EPIC-NL-Prospect). The proportion of current smokers ranged from 13% in E3N to 37% in DNC-1993. Almost half of our participants (41%) were overweight, with the lowest proportion (21%) in E3N and the highest (74%) in HNR (Table 1).

Table 1
Baseline demographic characteristics of participants by the pooled cohort and sub-cohorts.

Cohort/ sub- cohort	N	Deaths, N	Follow- up time, years	Age, years	Female (%)	Current smokers (%)	Smoking duration, years*	Smoking intensity, n/day*	Over weight (%) ϕ	Married/ cohabiting (%)	Employed (%)	Mean income, euro \ddagger
Pooled Cohort	271,720	381	19.7	47.1 \pm 14.0	69	22	21.9 \pm 12.5	14.7 \pm 8.9	41	72	68	20.1 \pm 6.2
CEANS	20,702	12	13	56.3 \pm 11.4	58	22	33.6 \pm 11.0	13.1 \pm 7.7	51	72	69	25.3 \pm 5.6
SDPP	7,727	0	15.9	47.1 \pm 4.9	61	26	27.9 \pm 8.6	13.5 \pm 7.4	52	84	91	24.3 \pm 4.2
SIXTY	3,969	3	15.5	60.0 \pm 0.0	52	21	36.3 \pm 9.9	13.4 \pm 7.6	65	74	68	24.7 \pm 6.9
SALT	6,176	1	10.4	57.8 \pm 10.6	55	21	37.9 \pm 9.3	12.7 \pm 8.0	40	68	64	25.3 \pm 6.6
SNACK	2,830	8	7.4	72.9 \pm 10.4	62	14	43.3 \pm 13.6	11.7 \pm 8.2	53	46	23	28.7 \pm 2.2
DNC	25,171	27	17.3	53.5 \pm 8.3	100	35	30.4 \pm 9.5	13.7 \pm 8.0	29	70	78	19.1 \pm 2.5
1993	17,043	27	18.7	56.2 \pm 8.4	100	37	31.6 \pm 9.9	13.9 \pm 8.2	28	68	70	19.2 \pm 2.6
1999	8,128	0	14.4	47.9 \pm 4.2	100	29	27.1 \pm 7.1	13.3 \pm 7.3	30	76	95	19.0 \pm 2.4
E3N	39,006	29	16.7	53.0 \pm 6.8	100	13	28.6 \pm 7.6	11.4 \pm 9.2	21	83	68	11.2 \pm 3.0
EPIC-NL	32,872	17	16.7	49.5 \pm 11.9	75	29	28.9 \pm 11.2	15.0 \pm 8.7	52	70	61	12.6 \pm 1.6
Morgen	18,302	5	16.8	42.9 \pm 11.2	55	35	24.8 \pm 10.6	15.7 \pm 8.6	50	65	69	12.2 \pm 1.6
Prospect	14,570	12	16.4	57.7 \pm 6.1	100	23	36.8 \pm 7.6	13.7 \pm 8.7	55	77	51	13.1 \pm 1.4
HNR	4,733	8	12	59.7 \pm 7.8	50	24	34.5 \pm 9.4	18.6 \pm 12.0	74	75	40	25.2 \pm 8.2
KORA	4,853	4	14.3	49.4 \pm 13.9	51	21	24.7 \pm 11.8	16.1 \pm 9.5	68	80	57	37.3 \pm 6.0
S3	2,572	2	15.6	49.4 \pm 13.9	51	20	25.2 \pm 12.1	16.5 \pm 9.5	67	80	55	36.7 \pm 4.4
S4	2,281	2	12.9	49.3 \pm 13.8	51	23	24.3 \pm 11.6	15.7 \pm 9.5	69	79	59	38.0 \pm 7.3
VHM&PP	144,383	284	23.1	42.1 \pm 15.0	56	20	13.4 \pm 8.3	15.6 \pm 8.9	43	69	70	22.9 \pm 1.7

Results of participants' characteristics at baseline are presented as Mean \pm SD, Number, or Percentage.

*: Smoking duration and smoking intensity are only for current smokers. We set these variables to zero for never and former smokers.

ϕ : BMI $\geq 25 \text{ kg}/\text{m}^2$ indicates overweight according to the World Health Organization (WHO) categories.

\ddagger : Area-level mean year income in euros \times 1,000 in the year 2001. The spatial scale of an area varied from neighborhoods and city districts (CEANS, E3N, EPIC-NL, and HNR) to municipalities (DNC, KORA, and VHM&PP).

Definition of abbreviation: BMI, body mass index; SD, standard deviation.

3.2. Air pollution description

Fig. 1 represents the distribution of air pollution levels in 2010 by pooled cohort and sub-cohorts. Exposure distributions varied between cohorts with the lowest concentrations of PM_{2.5} and BC in the Nordic cohorts (CEANS and DNC). Almost all participants were exposed to annual PM_{2.5} levels below the EU limit value (25 µg/m³) but above the US limit value (12 µg/m³) and the new WHO guideline (5 µg/m³). Similarly, almost all participants were exposed to annual NO₂ levels below the EU limit value (40 µg/m³) but above the new WHO guideline (10 µg/m³) (Fig. 1). We also observed varying exposure levels across cohorts and sub-cohorts for the baseline year exposure (supplemental Figure S2). Comparing with 2010 exposure levels, the concentrations of PM_{2.5} were much higher at baseline, with smaller differences observed for the other pollutants. Pearson correlations between NO₂ and BC were moderate to high in sub-cohorts (0.67–0.93) except for in CEANS-SNACK (0.43; Table S3). PM_{2.5} was moderately positively correlated with exposure to BC and NO₂ in most sub-cohorts, and (warm season) O₃ was moderately negatively correlated with other pollutants in most sub-cohorts (Table S3).

3.3. Main analyses

In our fully adjusted model (Model 3), significant positive associations were observed between PM_{2.5} exposure and PD mortality (HR: 1.25; 95% CI: 1.01–1.55), while suggestive positive associations were evident for NO₂ (1.13; 0.95–1.34) and BC (1.12; 0.94–1.34) (Table 2). We observed a negative association for O₃ (0.74; 0.58–0.94). Adjustment for individual covariates minimally affected the air pollution effect estimates (Model 2 versus Model 1). Adjustment for area-level income mildly decreased the effect estimates (Model 3 versus Model 2).

In two-pollutant models, the association with PM_{2.5} was reasonably robust when including NO₂ or BC although with attenuation towards the null for either pollutant, and the association with O₃ was mostly unchanged (Table 2). Two-pollutant results as contributions of NO₂ and BC individually are difficult to interpret because of the high correlations between NO₂ and BC in some sub-cohorts (Table S3).

We observed linear associations of PM_{2.5}, NO₂, and BC with PD mortality, with no evidence of a lower threshold below which air pollution was not associated with PD mortality (Fig. 2, Table S4). At the extremes of concentration distribution, the uncertainty of the shape of




Fig. 1. Distribution of annual average concentrations of air pollution for the year 2010 by pooled cohort and sub-cohorts (N = 271,720). The bold lines in the middle of the box indicate the median values (the 50th percentile). The lower and upper hinges correspond to the 25th and 75th percentiles. The lower and upper whiskers extend to the 5th and 95th percentiles. Red dashed lines represent different limited values in EU, U.S., and WHO guidelines (2021 version). For PM_{2.5}, they indicate the annual average limited/guideline values of WHO (5 µg/m³), U.S. (12 µg/m³), and EU (25 µg/m³). For NO₂, they indicate the annual average limited/guideline values of WHO (10 µg/m³) and EU (40 µg/m³). For O₃, it indicates the 8-hour mean, peak season guideline value of WHO (60 µg/m³). O₃ was in the boreal warm season from April 1st through September 30th. Definition of abbreviation: PM_{2.5}, particulate matters with aerodynamic diameters of less than 2.5 µm; NO₂, nitrogen dioxide; BC, black carbon (measured by the reflectance of PM_{2.5} filters from 2009 and 2010, expressed in absorbance units); O₃, ozone. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Table 2

Associations between long-term air pollution exposure and Parkinson's disease mortality (N = 271,720).

	Model 1	Model 2	Model 3	Model 3 + NO ₂	Model 3 + PM _{2.5}	Model 3 + BC	Model 3 + O ₃
Parkinson's disease (381deaths)							
NO ₂	1.19 (1.01, 1.40)	1.18 (1.01, 1.39)	1.13 (0.95, 1.34)	/	1.01 (0.81, 1.26)	1.11 (0.74, 1.68)	0.93 (0.72, 1.20)
PM _{2.5}	1.30 (1.05, 1.60)	1.30 (1.05, 1.60)	1.25 (1.01, 1.55)	1.24 (0.95, 1.62)	/	1.28 (0.96, 1.71)	1.11 (0.85, 1.45)
BC	1.18 (1.00, 1.40)	1.17 (0.99, 1.39)	1.12 (0.94, 1.34)	1.02 (0.67, 1.55)	0.97 (0.76, 1.25)	/	0.91 (0.71, 1.18)
O ₃	0.70 (0.56, 0.89)	0.70 (0.56, 0.89)	0.74 (0.58, 0.94)	0.68 (0.48, 0.98)	0.79 (0.59, 1.07)	0.67 (0.47, 0.96)	/

Results are presented as hazard ratio and 95% confidence interval [HR (95%CI)] for the following increases: 10 µg/m³ for NO₂, 5 µg/m³ for PM_{2.5}, 0.5 × 10⁻⁵m⁻¹ for BC and 10 µg/m³ for O₃. BC was measured by the reflectance of PM_{2.5} filters from 2009 and 2010, expressed in absorbance units.

Model 1: adjusted for age (time axis), sex (strata), sub-cohort (strata), and calendar year of baseline;

Model 2: additionally adjusted for smoking (status, duration, intensity, and intensity²), BMI (category), marital status, and employment status;

Model 3: further adjusted for area-level mean year income.




Fig. 2. Concentration-response curves for the associations between long-term exposure to air pollution and Parkinson's disease mortality. Natural cubic splines with two degrees of freedom were fit for air pollutants based on Model 3, where the hazard ratios equal to one were for minimum pollutant exposures. Solid black (horizontal) lined-curves indicate hazard ratio values and dashed black lines (with grey shading) indicate their 95% confidence intervals. Solid black (vertical) lines indicate the 5th and 95th percentiles of air pollutants' concentrations. Dashed red (vertical) lines represent existing EU, US, and WHO guideline limit values. The histograms show the distributions of exposures in 2010. X-axes are truncated at 60 µg/m³ for NO₂ and 3 × 10⁻⁵m⁻¹ for BC. Definition of abbreviation: PM_{2.5}, particulate matter with aerodynamic diameter of less than 2.5 µm; NO₂, nitrogen dioxide; BC, black carbon (measured by the reflectance of PM_{2.5} filters from 2009 and 2010, expressed in absorbance units); O₃, ozone. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

the relationship was large (Fig. 2). The likelihood ratio test (results not shown) indicated no significant deviations from a linear relationship. The majority of participants (74% to 100%) in the two Nordic cohorts (CEANS, DNC) were exposed to PM_{2.5} levels mostly below 15 µg/m³ (Fig. 1). This disparity in exposure level across sub-cohorts was less prevalent for NO₂ and O₃ but not for BC (Fig. 1).

We observed that the association between PD mortality with PM_{2.5} was statistically stronger in participants who were normal or under weight, as compared to overweight participants (*p*-value for interaction = 0.05). We also note stronger association in males, although no

significant effect modification was observed (Table S5).

3.4. Sensitivity analyses

A number of sensitivity analyses showed robustness in associations for all pollutants. Associations with PM_{2.5}, NO₂ and BC were (borderline) significant when using either of the back-extrapolated (ratio or difference method) baseline year exposures (Table S6). The HRs and confidence intervals for all three pollutants were substantially smaller because of the higher exposure levels and variability of baseline

exposure compared to 2010 exposure (Figure S2). For time-varying exposures, available in a subset of three cohorts (N = 132,952) with available data (CEANS, EPIC-NL, and VHM&PP), HRs were generally very similar compared to those with the 2010 exposure (Table S7). Further, the associations with PD mortality were largely robust to excluding one sub-cohort at a time, except for NO₂ association when excluding DNC. Associations for PM_{2.5}, NO₂ and BC became very imprecise and non-significant when excluding the large VHM&PP which contributed the majority of PD deaths (Table S8).

PM_{2.5} component data were available in a subset of 271,003 participants. Exposure to the eight PM_{2.5} components estimated by the two different models, SLR and RF, and their correlation with PM_{2.5} mass and NO₂ can be seen in Supplementary Material (Tables S9–S12). In single SLR models, Cu, Fe, K, and S were positively associated with PD mortality, and associations were attenuated in two-pollutant models with PM_{2.5} and NO₂ (Fig. 3). Single-pollutant RF models showed significant associations between K and PD mortality, and no associations for other components (Fig. 3).

4. Discussion

In the pooled analysis of 271,720 adults from seven European cohorts, we found that long-term exposure to PM_{2.5}, NO₂, and BC were associated with PD mortality, with the strongest and most robust associations for PM_{2.5}. We observed that associations persisted at low-levels of pollutant concentration, well below current EU limit values. This study of a large European dataset on Europeans from seven countries contributes to the evidence base on air pollution and PD risk, albeit that this evidence base remains mixed. Air pollution is a complex mix of PM and gases, with some components able to access the brain through the circulatory or olfactory system. While further research into the relevant biological pathways is required, several hypotheses exist including that air pollution components enter the brain and disrupt proteostasis, causing injury to mitochondria and inducing inflammation.

Of 14 studies investigating the association of long-term exposure to PM_{2.5} with PD development or mortality, the majority reported positive associations, in agreement with our findings (Table S1). Seven of these studies were included in a recent review and meta-analysis of PM_{2.5} and PD incidence, reporting a relative risk (RR) estimate of 1.08 (95% CI: 0.98–1.19) per 10 µg/m³ increase (Han et al., 2020), which is weaker than our HR estimate of 1.25 (95% CI: 1.01–1.55) per 5 µg/m³. A similar

comparison can be made for NO₂, for which Han and colleagues reported a RR of 1.03 (95% CI: 0.99–1.07) per 10 µg/m³ compared to our HR of 1.13 (95% CI: 0.95–1.34) per 10 µg/m³. We found generally stronger associations than those reported in the literature, which are in line with generally stronger associations reported in the ELAPSE project with overall mortality (Stafoggia et al., 2022; Strak et al., 2021) compared to those in the literature and recent meta-analyses (Chen and Hoek, 2020). We observed a clear linear PM_{2.5} exposure–response function with PD mortality, with a linear function for NO₂ and BC exposure especially at lower levels, suggesting that there are no lower threshold levels below which air pollution is not harmful (Fig. 2), consistent with observations made for other health outcomes in ELAPSE and other studies (Stafoggia et al., 2022; Strak et al., 2021).

Our results on long-term exposure to NO₂ and PD mortality are in accordance with current evidence of a suggestive positive association, however, not to O₃, as reported by a 2019 meta-analysis by Kasdagli and colleagues (Kasdagli et al., 2019). They found a significant positive association with O₃, while we found a negative association with O₃ in single pollutant models. In two-pollutant models with PM_{2.5}, associations with O₃ remained negative but no longer statistically significant. Our negative association with O₃ may be due to its negative correlation with PM_{2.5}, BC and NO₂, the small exposure contrasts within each sub-cohort (Fig. 1), or generally low levels of O₃ exposure in our study. The inverse relationship between O₃ and PD mortality could also be explained by the moderate-to-strong negative correlation of O₃ with NO₂ in our pooled cohort, and strong negative correlations of O₃ with NO₂ and BC in some of our larger populated sub-cohorts (e.g., VHM&PP) (Table S3). Alternatively, this negative association between O₃ and PD mortality could be explained by bias related to the study design and lack of power due to low numbers of cases. In our study, we exploit exposure contrasts within relatively small study areas compared to other studies, making it thus less informative for O₃ which tends to vary on a large spatial scale (Brunekreef et al., 2021). Moreover, one previous study has shown that ambient ozone concentration is not a suitable surrogate for individual exposure assessment and may lead to large misclassification of exposure (Niu et al., 2018). A large US study has previously suggested the existence of a threshold at 56 ppb (~110 µg/m³) for the effect of warm-season O₃ on all-cause mortality (Jerrett et al., 2009). Of five studies on O₃ and PD incidence, three (Shin et al., 2018; Cerza et al., 2018; Zhao et al., 2021) have detected positive associations, whereas two have found none (Chen et al., 2017; Lee et al., 2016). We present the




Fig. 3. Associations of PM_{2.5} components with Parkinson's disease mortality in single-pollutant and two-pollutant models in SLR and RF analyses. (N = 271,003; Parkinson's mortality = 381). Results are presented as hazard ratio and 95% confidence interval [HR (95%CI)] for the following increments: 5 µg/m³ for PM_{2.5}, 10 µg/m³ for NO₂, and IQR increments for each PM_{2.5} component. The main model was adjusted for sub-cohort identification, age, sex, year of enrollment, smoking (status, duration, intensity, and intensity²), BMI categories, marital status, employment status, and 2001 neighborhood-level mean income. In two-pollutant models, PM_{2.5} and NO₂ exposures were estimated using LUR. Definition of abbreviation: BMI, body mass index; CI, confidence interval; Cu, copper; Fe, iron; HR, hazard ratio; K, potassium; Ni, nickel; PM_{2.5}, fine particulate matter; S, sulfur; Si, silicon; V, vanadium; Zn, zinc.

novel result of a positive association between long-term exposure to BC and PD mortality, with a HR of 1.12 (0.94, 1.34) per $0.5 \times 10^{-5} \text{m}^{-1}$, which is in contrast to the only other study on BC and incidence of PD reporting an inverse association (RR: 0.96; 95%CI: 0.94–0.99) (Yang et al., 2018). These varied findings could be partly explained by a lack of specificity regarding particle component exposure levels across studies, with components expected to vary in presence and level across study sites due to varying air pollution sources contributing to the same component (Nunez et al., 2021).

In our particle component analysis, we detected a suggestive association between the PM_{2.5} component of K and mortality from PD (Fig. 3). This association, however, was attenuated when adjusting for either PM_{2.5} mass or NO₂. The particle component results shed some light on the potential local contributing (combustion or otherwise) sources. K has been seen as a tracer of biomass burning, which in Europe may be in the form of residential wood fire places or municipal water boilers. K has also been seen in roadside samples of Poland as an element of salt sprinkles used by local authorities to de-ice road surfaces during winter, which may suspend as airborne particles outside of winter (Skorbiłowicz and Skorbiłowicz, 2019). An analysis from the US suggested that 80% of the variability of K can be explained by factors associated with soil dust, traffic and biomass burning, however more accurately indicating traffic when considered alongside Fe (for which we saw a less suggestive association) (Pachon et al., 2013). Our findings differ from those of Nunez and colleagues, who detected associations of traffic-related nitrate and OM with PD hospitalization (Nunez et al., 2021), as well as of Palacios and colleagues, who found the relevance of mercury for PD incidence (Palacios et al., 2014). OM is a complex mixture of compounds including the potentially neurotoxic polycyclic aromatic hydrocarbons, which are known to evoke an inflammatory response and contribute to degenerative disease-like syndromes in animal studies (Nunez et al., 2021). While our study was not able to analyse this specific PM component, it is moderately-to-highly correlated with PM_{2.5} and may explain those associated findings. Based on the literature, more studies are needed to determine which PM component is most biologically relevant for neuropathology related to PD (Nunez et al., 2021).

We observed significant effect modification for the association of PM_{2.5} with PD mortality by BMI, with the strongest associations in those who were not overweight (BMI < 25 kg/m², as defined by WHO), which may be indicative of PD-related weight loss with progressive disease stages due to malnutrition (van der Marck et al., 2012). We hypothesise that those individuals with more progressed disease and that are closer to death would have a lower body weight, also as a function of age-related weight loss – suggested separately by a stronger association of PM_{2.5} with PD mortality among individuals ≥ 65 years of age.

A strength of our study is the pooling of data from seven European cohorts within the ELAPSE framework, which allows for a larger sample size to investigate mortality from PD. Another strength is that we had harmonized exposure data based on the Europe-wide hybrid LUR models of a fine spatial scale. Furthermore, we had detailed and harmonised information available on potential confounders including sample characteristics at the individual and area level. While we did have information on an individual's history of smoking, we did not have information on history of traumatic brain injury, another major personal risk factor for PD, which is somewhat controlled for in our models by controlling for sex: males show increased likelihood to suffer this injury due to their greater exposure to events such as road traffic accidents or contact sports; as well as more likely to smoke, as another major personal risk factor (Rocca, 2018).

The main limitation of our study is related to the definition of PD based on mortality data. Numerous studies show that PD is not well ascertained in death certificates and it underestimates the true burden of PD (Shi and Counsell, 2021; Hobson and Meara, 2018). Furthermore, many PD patients die from competing illnesses, and it is a weakness that we could not identify those, as we did not have information on deaths

from PD as a contributing cause. Although the sample size achieved by pooling seven European cohorts was large, we had limited statistical power due to a small number of deaths due to PD. Another limitation is that exposure data were based on the year 2010, and applied at the baseline of the cohorts recruited in the 1990s and early 2000s. A previous study reported stable spatial distribution of NO₂ over 10 years in the Netherlands prior to 2011 (Eeftens et al., 2011). Furthermore, predictions from our year 2010 model were highly correlated ($R^2 > 76\%$) with year 2000 and 2005 models for NO₂ and O₃, and the year 2013 model for PM_{2.5} (de Hoogh et al., 2018), indicating a limited impact of temporal misalignment by exposures based on the year 2010. Furthermore, our sensitivity analyses showed robust associations when using either back-extrapolated baseline year exposure (Table S6) or time-varying annual exposure in three cohorts with available information on address history (Table S7). Another limitation is inherent exposure misclassification due to lack of information on time spent outdoors and commuting to work (as personal exposure from outdoor sources), other sources of air pollution indoors, as well as outdoor and indoor sources at work. Finally, we did not have environmental noise exposure information for the VHM&PP cohort, which contains the majority of our cases, to treat as a potential confounder for the effect of air pollution on PD mortality.

5. Conclusions

In conclusion, we found that long-term exposure to PM_{2.5}, NO₂ and BC were associated with the risk of dying from PD, with PM_{2.5} found to be the most relevant pollutant for this risk. We also found that associations persisted at low levels of pollutant concentration, well below current EU air quality standards. This study based on a large European population adds strong novel evidence in support of an association between air pollution and PD.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

This study was supported by the Health Effects Institute (HEI) (#4954-RFA14-3/16-5-3) and the Novo Nordisk Foundation Challenge Programme [NNF17OC0027812]. The HEI is an organization jointly funded by the United States Environmental Protection Agency (EPA) (Assistance Award No. R-82811201) and certain motor vehicle and engine manufacturers. The contents of this article do not necessarily reflect the views of HEI, or its sponsors, nor do they necessarily reflect the views and policies of the EPA or motor vehicle and engine manufacturers. While HEI has reviewed and approved the study design, it was not involved in data collection and analysis, decision to publish, or preparation of the manuscript. We give thanks to all participants in the pooled cohort studies and the respective study teams of the ELAPSE project for their hard work and effort. Accordingly, we especially thank Marjan Tewis for conducting data management tasks when creating the pooled cohort database. In addition, we specifically thank the National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands, for their contribution to the ELAPSE project. SALT and TwinGene are sub-studies of The Swedish Twin Registry (STR), which is managed by Karolinska Institutet and receives additional funding through the Swedish Research Council (No. 2017-00641). The KORA study was initiated and financed by the Helmholtz Zentrum München –

German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research (BMBF) and by the State of Bavaria. The Novo Nordisk Foundation, the Swedish Research Council, the German Federal Ministry of Education and Research, and the State of Bavaria were not involved in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2022.107667>.

References

- Andersen, Z.J., Zhang, J., Jørgensen, J.T., Samoli, E., Liu, S., Chen, J., et al., 2022. Long-term exposure to air pollution and mortality from dementia, psychiatric disorders, and suicide in a large pooled European cohort: ELAPSE study. *Environ. Int.* 170.
- Ascherio, A., Schwarzschild, M.A., 2016. The epidemiology of Parkinson's disease: risk factors and prevention. *Lancet Neurol.* 15 (12), 1257–1272.
- Beulens, J.W., Monninkhof, E.M., Verschuren, W.M., van der Schouw, Y.T., Smit, J., Ocke, M.C., et al., 2010. Cohort profile: the EPIC-NL study. *Int. J. Epidemiol.* 39 (5), 1170–1178.
- Brandt, J., Silver, J.D., Frohn, L.M., Geels, C., Gross, A., Hansen, A.B., et al., 2012. An integrated model study for Europe and North America using the Danish Eulerian Hemispheric Model with focus on intercontinental transport of air pollution. *Atmos. Environ.* 53, 156–176.
- Brunekreef, B., Strak, M., Chen, J., Andersen, Z.J., Atkinson, R., Bauwelinck, M., et al., 2021. Mortality and Morbidity Effects of Long-Term Exposure to Low-Level PM_{2.5}, BC, NO₂, and O₃: An Analysis of European Cohorts in the ELAPSE Project. *Res Rep Health Eff Inst.* 208, 1–127.
- Cerza, F., Renzi, M., Agabiti, N., Marino, C., Gariazzo, C., Davoli, M., et al., 2018. Residential exposure to air pollution and incidence of Parkinson's disease in a large metropolitan cohort. *Environ. Epidemiol.* 2 (3).
- Chen, J., de Hoogh, K., Gulliver, J., Hoffmann, B., Hertel, O., Ketzel, M., et al., 2019. A comparison of linear regression, regularization, and machine learning algorithms to develop Europe-wide spatial models of fine particles and nitrogen dioxide. *Environ. Int.* 130, 104934.
- Chen, J., Hoek, G., 2020. Long-term exposure to PM and all-cause and cause-specific mortality: A systematic review and meta-analysis. *Environ. Int.* 143, 105974.
- Chen, J., de Hoogh, K., Gulliver, J., Hoffmann, B., Hertel, O., Ketzel, M., et al., 2020. Development of Europe-Wide Models for Particle Elemental Composition Using Supervised Linear Regression and Random Forest. *Environ. Sci. Technol.* 54 (24), 15698–15709.
- Chen, C.Y., Hung, H.J., Chang, K.H., Hsu, C.Y., Muo, C.H., Tsai, C.H., et al., 2017. Long-term exposure to air pollution and the incidence of Parkinson's disease: A nested case-control study. *PLoS One* 12 (8), e0182834.
- Chen, J., Rodopoulou, S., de Hoogh, K., Strak, M., Andersen, Z.J., Atkinson, R., et al., 2021. Long-Term Exposure to Fine Particle Elemental Components and Natural and Cause-Specific Mortality—a Pooled Analysis of Eight European Cohorts within the ELAPSE Project. *Environ. Health Perspect.* 129 (4), 47009.
- Clavel-Chapelon, F., 2015. Cohort Profile: The French E3N Cohort Study. *Int. J. Epidemiol.* 44 (3), 801–809.
- de Hoogh, K., Chen, J., Gulliver, J., Hoffmann, B., Hertel, O., Ketzel, M., et al., 2018. Spatial PM_{2.5}, NO₂, O₃ and BC models for Western Europe - Evaluation of spatiotemporal stability. *Environ. Int.* 120, 81–92.
- de Hoogh, K., Chen, J., Gulliver, J., Hoffmann, B., Hertel, O., Ketzel, M., et al., 2018. Spatial PM_{2.5}, NO₂, O₃ and BC models for Western Europe - Evaluation of spatiotemporal stability. *Environ. Int.* 120, 81–92.
- Dorsey, E.R., Elbaz, A., Nichols, E., Abbasi, N., Abd-Allah, F., Abdelalim, A., et al., 2018. Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* 17 (11), 939–953.
- Eeftens, M., Beelen, R., Fischer, P., Brunekreef, B., Meliefste, K., Hoek, G., 2011. Stability of measured and modelled spatial contrasts in NO₂ over time. *Occup. Environ. Med.* 68 (10), 765.
- Eriksson, A.K., Ekblom, A., Granath, F., Hilding, A., Eféndic, S., Ostenson, C.G., 2008. Psychological distress and risk of pre-diabetes and Type 2 diabetes in a prospective study of Swedish middle-aged men and women. *Diabet. Med. : J. British Diabet. Assoc.* 25 (7), 834–842.
- Han, C., Lu, Y., Cheng, H., Wang, C., Chan, P., 2020. The impact of long-term exposure to ambient air pollution and second-hand smoke on the onset of Parkinson disease: a review and meta-analysis. *Publ. Health* 179, 100–110.
- Hobson, P., Meara, J., 2018. Mortality and quality of death certification in a cohort of patients with Parkinson's disease and matched controls in North Wales, UK at 18 years: a community-based cohort study. *BMJ Open* 8 (2), e018969.
- Holle, R., Hoppich, M., Löwel, H., Wichmann, H.E., 2005. KORA—a research platform for population based health research. *Gesundheitswesen.* 67 (Suppl 1), S19–S25.
- Hundrup, Y.A., Simonsen, M.K., Jørgensen, T., Obel, E.B., 2012. Cohort profile: the Danish nurse cohort. *Int. J. Epidemiol.* 41 (5), 1241–1247.
- Hvidtfeldt, U.A., Severi, G., Andersen, Z.J., Atkinson, R., Bauwelinck, M., Bellander, T., et al., 2021. Long-term low-level ambient air pollution exposure and risk of lung cancer - A pooled analysis of 7 European cohorts. *Environ. Int.* 146, 106249.
- Jerrett, M., Burnett, R.T., Pope 3rd, C.A., Ito, K., Thurston, G., Krewski, D., et al., 2009. Long-term ozone exposure and mortality. *N. Engl. J. Med.* 360 (11), 1085–1095.
- Kasdagli, M.I., Katsouyanni, K., Dimakopoulou, K., Samoli, E., 2019. Air pollution and Parkinson's disease: A systematic review and meta-analysis up to 2018. *Int. J. Hyg. Environ. Health.* 222 (3), 402–409.
- Kim, H., Kim, W.H., Kim, Y.Y., Park, H.Y., 2020. Air Pollution and Central Nervous System Disease: A Review of the Impact of Fine Particulate Matter on Neurological Disorders. *Front. Publ. Health.* 8, 575330.
- Kirrane, E.F., Bowman, C., Davis, J.A., Hoppin, J.A., Blair, A., Chen, H., et al., 2015. Associations of Ozone and PM_{2.5} Concentrations With Parkinson's Disease Among Participants in the Agricultural Health Study. *J. Occup. Environ. Med.* 57 (5), 509–517.
- Lagergren, M., Fratiglioni, L., Hallberg, I.R., Berglund, J., Elmståhl, S., Hagberg, B., et al., 2004. A longitudinal study integrating population, care and social services data. The Swedish National study on Aging and Care (SNAC). *Aging Clin. Exp. Res.* 16 (2), 158–168.
- Lee, P.C., Liu, L.L., Sun, Y., Chen, Y.A., Liu, C.C., Li, C.Y., et al., 2016. Traffic-related air pollution increased the risk of Parkinson's disease in Taiwan: A nationwide study. *Environ. Int.* 96, 75–81.
- Liu, S., Jørgensen, J.T., Ljungman, P., Pershagen, G., Bellander, T., Leander, K., et al., 2021. Long-term exposure to low-level air pollution and incidence of chronic obstructive pulmonary disease: The ELAPSE project. *Environ. Int.* 146, 106267.
- Liu, S., Jørgensen, J.T., Ljungman, P., Pershagen, G., Bellander, T., Leander, K., et al., 2021. Long-term exposure to low-level air pollution and incidence of asthma: the ELAPSE project. *Eur. Respir. J.* 57 (6).
- Liu, S., Jørgensen, J.T., Ljungman, P., Pershagen, G., Bellander, T., Leander, K., et al., 2021. Long-term exposure to low-level air pollution and incidence of chronic obstructive pulmonary disease: The ELAPSE project. *Environ. Int.* 146, 106267.
- Liu, R., Young, M.T., Chen, J.C., Kaufman, J.D., Chen, H., 2016. Ambient Air Pollution Exposures and Risk of Parkinson Disease. *Environ. Health Perspect.* 124 (11), 1759–1765.
- Magnusson, P.K., Almqvist, C., Rahman, I., Ganna, A., Viktorin, A., Walum, H., et al., 2013. The Swedish Twin Registry: establishment of a biobank and other recent developments. *Twin Res. Hum. Genet.: Offic. J. Int. Soc. Twin Stud.* 16 (1), 317–329.
- Niu, Y., Cai, J., Xia, Y., Yu, H., Chen, R., Lin, Z., et al., 2018. Estimation of personal ozone exposure using ambient concentrations and influencing factors. *Environ. Int.* 117, 237–242.
- Nunez, Y., Boehme, A.K., Li, M., Goldsmith, J., Weisskopf, M.G., Re, D.B., et al., 2021. Parkinson's disease aggravation in association with fine particle components in New York State. *Environ. Res.* 201, 111554.
- Pachon, J.E., Weber, R.J., Zhang, X., Mulholland, J.A., Russell, A.G., 2013. Revising the use of potassium (K) in the source apportionment of PM_{2.5}. *Atmospheric. Pollut. Res.* 4 (1), 14–21.
- Palacios, N., Fitzgerald, K.C., Hart, J.E., Weisskopf, M.G., Schwarzschild, M.A., Ascherio, A., et al., 2014. Particulate matter and risk of Parkinson disease in a large prospective study of women. *Environ. Health.* 13, 80.
- Palacios, N., Fitzgerald, K.C., Hart, J.E., Weisskopf, M., Schwarzschild, M.A., Ascherio, A., et al., 2017. Air Pollution and Risk of Parkinson's Disease in a Large Prospective Study of Men. *Environ. Health Perspect.* 125 (8), 087011.
- Raggi, A., Leonardi, M., 2020. Burden of brain disorders in Europe in 2017 and comparison with other non-communicable disease groups. *J. Neurol. Neurosurg. Psychiatry* 91 (1), 104–105.
- Rhew, S.H., Kravchenko, J., Lyerly, H.K., 2021. Exposure to low-dose ambient fine particulate matter PM_{2.5} and Alzheimer's disease, non-Alzheimer's dementia, and Parkinson's disease in North Carolina. *PLoS One* 16 (7), e0253253.
- Ritz, B., Lee, P.C., Hansen, J., Lassen, C.F., Ketzel, M., Sorensen, M., et al., 2016. Traffic-Related Air Pollution and Parkinson's Disease in Denmark: A Case-Control Study. *Environ. Health Perspect.* 124 (3), 351–356.
- Rocca, W.A., 2018. The burden of Parkinson's disease: a worldwide perspective. *Lancet Neurol.* 17 (11), 928–929.
- Samoli, E., Rodopoulou, S., Hvidtfeldt, U.A., Wolf, K., Stafoggia, M., Brunekreef, B., et al., 2021. Modeling multi-level survival data in multi-center epidemiological cohort studies: Applications from the ELAPSE project. *Environ. Int.* 147, 106371.
- Schmermund, A., Möhlenkamp, S., Stang, A., Grönemeyer, D., Seibel, R., Hirche, H., et al., 2002. Assessment of clinically silent atherosclerotic disease and established and novel risk factors for predicting myocardial infarction and cardiac death in healthy middle-aged subjects: rationale and design of the Heinz Nixdorf RECALL Study. Risk Factors, Evaluation of Coronary Calcium and Lifestyle. *Am. Heart J.* 144 (2), 212–218.
- Shi, H., Counsell, C., 2021. Accuracy of death certificates for recording parkinsonian syndromes and associated dementia. *J. Neurol.* 268 (1), 140–146.
- Shi, L., Wu, X., Danesh Yazdi, M., Braun, D., Abu Awad, Y., Wei, Y., et al., 2020. Long-term effects of PM_{2.5} on neurological disorders in the American Medicare population: a longitudinal cohort study. *Lancet Planet Health* 4 (12), e557–e65.
- Shin, S., Burnett, R.T., Kwong, J.C., Hystad, P., van Donkelaar, A., Brook, J.R., et al., 2018. Effects of ambient air pollution on incident Parkinson's disease in Ontario, 2001 to 2013: a population-based cohort study. *Int. J. Epidemiol.* 47 (6), 2038–2048.
- Skorbiłowicz, M., Skorbiłowicz, E., 2019. Content of calcium, magnesium, sodium and potassium in the street dust from the area of Białystok (Poland). *J. Ecol. Eng.* 20 (10), 125–131.

- So, R., Chen, J., Mehta, A.J., Liu, S., Strak, M., Wolf, K., et al., 2021. Long-term exposure to air pollution and liver cancer incidence in six European cohorts. *Int. J. Cancer*. 149 (11), 1887–1897.
- Stafoggia, M., Oftedal, B., Chen, J., Rodopoulou, S., Renzi, M., Atkinson, R.W., et al., 2022. Long-term exposure to low ambient air pollution concentrations and mortality among 28 million people: results from seven large European cohorts within the ELAPSE project. *Lancet Planet. Health*. 6 (1), e9–e18.
- Strak, M., Weinmayr, G., Rodopoulou, S., Chen, J., de Hoogh, K., Andersen, Z.J., et al., 2021. Long term exposure to low level air pollution and mortality in eight European cohorts within the ELAPSE project: pooled analysis. *BMJ* 374, n1904.
- Thiébaud, A.C., Bénichou, J., 2004. Choice of time-scale in Cox's model analysis of epidemiologic cohort data: a simulation study. *Stat. Med.* 23 (24), 3803–3820.
- Toro, R., Downward, G.S., van der Mark, M., Brouwer, M., Huss, A., Peters, S., et al., 2019. Parkinson's disease and long-term exposure to outdoor air pollution: A matched case-control study in the Netherlands. *Environ. Int.* 129, 28–34.
- Ulmer, H., Kelleher, C.C., Fitz-Simon, N., Diem, G., Concin, H., 2007. Secular trends in cardiovascular risk factors: an age-period cohort analysis of 698,954 health examinations in 181,350 Austrian men and women. *J. Intern. Med.* 261 (6), 566–576.
- van der Marck, M.A., Dicke, H.C., Uc, E.Y., Kentin, Z.H., Borm, G.F., Bloem, B.R., et al., 2012. Body mass index in Parkinson's disease: a meta-analysis. *Parkinsonism Relat. Disord.* 18 (3), 263–267.
- Wändell, P.E., Wajngot, A., de Faire, U., Hellénus, M.L., 2007. Increased prevalence of diabetes among immigrants from non-European countries in 60-year-old men and women in Sweden. *Diabetes Metab.* 33 (1), 30–36.
- WHO, 2021. WHO global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization.
- Wolf, K., Hoffmann, B., Andersen, Z.J., Atkinson, R.W., Bauwelinck, M., Bellander, T., et al., 2021. Long-term exposure to low-level ambient air pollution and incidence of stroke and coronary heart disease: a pooled analysis of six European cohorts within the ELAPSE project. *Lancet Planet. Health*. 5 (9), e620–e632.
- Yang, Y., Tang, R., Qiu, H., Lai, P.C., Wong, P., Thach, T.Q., et al., 2018. Long term exposure to air pollution and mortality in an elderly cohort in Hong Kong. *Environ. Int.* 117, 99–106.
- Yu, Z., Wei, F., Zhang, X., Wu, M., Lin, H., Shui, L., et al., 2021. Air pollution, surrounding green, road proximity and Parkinson's disease: A prospective cohort study. *Environ. Res.* 197, 111170.
- Yuchi, W., Sbihi, H., Davies, H., Tamburic, L., Brauer, M., 2020. Road proximity, air pollution, noise, green space and neurologic disease incidence: a population-based cohort study. *Environ. Health*. 19 (1), 8.
- Zhao, N., Pinault, L., Toyib, O., Vanos, J., Tjepkema, M., Cakmak, S., 2021. Long-term ozone exposure and mortality from neurological diseases in Canada. *Environ. Int.* 157, 106817.