



## High-flow nasal oxygen for suspension laryngoscopy: a multicenter open-label study

Mary Saad, Aline Albi-Feldzer, Rabah Taouachi, Isabelle Wagner, Marc Fischler, Pierre Alexandre Squara, Morgan Le Guen

### ► To cite this version:

Mary Saad, Aline Albi-Feldzer, Rabah Taouachi, Isabelle Wagner, Marc Fischler, et al.. High-flow nasal oxygen for suspension laryngoscopy: a multicenter open-label study. *Journal of International Medical Research*, 2022, 50 (12), 10.1177/03000605221140685 . hal-03950154

**HAL Id: hal-03950154**

**<https://hal.science/hal-03950154>**

Submitted on 6 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



# High-flow nasal oxygen for suspension laryngoscopy: a multicenter open-label study

Journal of International Medical Research

2022, Vol. 50(12) 1–11

© The Author(s) 2022


Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/03000605221140685

journals.sagepub.com/home/imr



Mary Saad<sup>1</sup> , Aline Albi-Feldzer<sup>1</sup>,  
Rabah Taouachi<sup>2</sup>, Isabelle Wagner<sup>3,4</sup>,  
Marc Fischler<sup>4,5</sup>, Pierre Squara<sup>6</sup> and  
Morgan Le Guen<sup>4,5</sup>

## Abstract

In this study, we aimed to assess the efficacy of high-flow nasal oxygen (HFNO) to maintain blood peripheral oxygen saturation (SpO<sub>2</sub>) in patients undergoing suspension laryngoscopy under general anesthesia. Adult patients were included in this bicenter study. After face-mask oxygenation, HFNO at a flow rate of 70 L min<sup>-1</sup> and fraction of inspired oxygen 100% was initiated at loss of consciousness. At the end of HFNO, blood gas analysis was performed. Of the 29 included patients, five (17.2%; 95% confidence interval [CI]: 7.6–34.5) presented SpO<sub>2</sub> <95% during the first 15 minutes of the procedure and eight patients (27.6%; 95% CI: 14.7–45.7) presented SpO<sub>2</sub> <95% throughout the procedure. Six patients (20.7%; 95% CI: 9.8–38.4) required rescue jet ventilation. Median apnea time before SpO<sub>2</sub> <95% was 13.5 (interquartile range [IQR]: 10–17.7) minutes. Arterial carbon dioxide tension at the end of the procedure or at the time of study discontinuation was 9.73 (IQR: 8.8–10.9) kPa and was higher than 8 kPa in 88.9% (95% CI: 71.9–96.1) of patients. HFNO was associated with a relatively high incidence of suboptimal oxygen saturation and hypercapnia during suspension laryngoscopy under general anesthesia and may not be considered the reference technique.

## Keywords

General anesthesia, adult, apnea, airway management, larynx, surgery

Date received: 16 March 2022; accepted: 4 November 2022

<sup>1</sup>Institut Curie, PSL Research University, Department of Anesthesia, EPST, Saint Cloud, France

<sup>2</sup>Institut Curie, PSL Research University, Department of ENT, EPST, Saint Cloud, France

<sup>3</sup>Hôpital Foch, Department of ENT, Suresnes, France

<sup>4</sup>University Versailles Saint-Quentin en Yvelines, Versailles, France

<sup>5</sup>Hôpital Foch, Department of Anesthesia, Suresnes, France

<sup>6</sup>Research Unit, Clinique Ambroise Paré, Neuilly-sur-Seine, France

### Corresponding author:

Mary Saad, Institut Curie, 35 rue Dailly, Paris, Île-de-France 75248, France.

Email: mary.saad@curie.fr



Creative Commons NonCommercial-NoDerivs CC BY-NC-ND: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial-NoDerivs 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial use, reproduction and distribution of the work as published without adaptation or alteration, without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

## Introduction

Anesthetic management of suspension laryngoscopy must meet several requirements: ensuring sufficiently deep anesthesia to avoid memory of the procedure, guaranteeing satisfactory gas exchange, and causing the least possible disturbance to the operative vision.<sup>1</sup> No technique can fully meet all three elements. A recent French national survey on pan-endoscopy revealed that in 2018, the top two techniques were orotracheal intubation (62%) and general anesthesia with spontaneous ventilation (37%). Other techniques therefore have anecdotal use.<sup>2</sup> However, it is likely that the use of high-flow nasal oxygen (HFNO) will increase for this indication, particularly because HFNO is indicated for the treatment for hypoxic patients with COVID-19.<sup>3</sup> HFNO consists of transnasal high-flow insufflation of heated, humidified, oxygen (O<sub>2</sub>)-enriched air. The procedure has several advantages for conscious patients compared with the other available noninvasive ventilation techniques, particularly enhanced comfort and tolerance and reduced work of breathing.<sup>4-6</sup> Patel and Nouraei reported the use of OptiFlow™ (Fisher and Paykel Healthcare Limited, Auckland, New Zealand), a commercial device delivering HFNO in anesthetized and paralyzed patients.<sup>7</sup> Their technique, called transnasal humidified rapid-insufflation ventilatory exchange (THRIVE), ensured a prolonged median apnea time of 14 minutes with an interquartile range of 9 and 19 minutes and extreme values of 5 and 65 minutes, with a rate of increase in end-tidal carbon dioxide of 0.15 kPa min<sup>-1</sup>.

The aim of this open-label prospective two-center study was to assess the efficacy of HFNO to maintain blood oxygen saturation in apneic patients undergoing suspension laryngoscopy under general anesthesia.

## Methods

This prospective observational, multi-center cohort study was conducted in two French hospitals (Institut Curie, Saint-Cloud and Foch Hospital, Suresnes). The study was initiated after receiving Ethics Committee approval (East Regional Research Ethics Committee, Chairperson JP Quenot, ID RCB 2018-A00396-49 on 12 March 2018) and registration at ClinicalTrials.gov (NCT03472118 on 21 March 2018). Written informed consent was obtained from patients prior to their participation in the study. This report follows the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for observational cohort studies.<sup>8</sup>

It was decided that the sponsor of the study (i.e., the data monitoring committee) would be kept informed of the occurrence of adverse events and would make the decision whether to continue the study.

### Study population

We included adult patients with American Society of Anesthesiologists physical status class I–III scheduled for short diagnostic and/or therapeutic suspension laryngoscopy (except laser surgery and cavoscopy). Exclusion criteria were pregnancy, epilepsy, arrhythmia, pulmonary hypertension, obstructive laryngeal tumors, lung disease with basal SpO<sub>2</sub> <95%, and obesity (body mass index >30 kg/m<sup>2</sup>).

### Study design

Patients were provided with information about the study and gave their written informed consent at a pre-anesthetic visit. Patients did not receive any premedication. An intravenous line was placed on arrival in the operating room. Standard monitoring included electrocardiogram, noninvasive

arterial blood pressure monitoring, pulse oximetry (SpO<sub>2</sub>), capnography, bispectral index, and train-of-four monitoring. Patients were pre-oxygenated with 100% O<sub>2</sub> using a tight-fitting face mask with end-tidal gas monitoring for 3 minutes or longer, if necessary, until an end-tidal O<sub>2</sub> concentration  $\geq 90\%$  was achieved. Induction of anesthesia was performed with target controlled infusion (Alaris<sup>TM</sup> PK syringe pump, Cardinal Health, Rolle, Switzerland) of remifentanyl (Ultiva<sup>®</sup>, GlaxoSmithKline AB, Solna, Sweden) and propofol (Propofol Lipuro<sup>®</sup>, B. Braun Melsungen AG, Melsungen, Germany) followed by neuromuscular blockade with 0.6 mg kg<sup>-1</sup> rocuronium (Esmeron<sup>®</sup>, MSD, Haarlem, Netherlands). Anesthesia was maintained with target concentration infusions of propofol and remifentanyl with bispectral index objectives between 40 and 60, and mean arterial pressure within 20% of baseline, with reinjection of rocuronium to maintain a null response to train-of-four monitoring. HFNO was started after the onset of apnea using a high-flow nasal cannula and the simplified Optiflow<sup>®</sup> device (Fisher and Paykel Healthcare Ltd., Auckland, New Zealand) with a flow rate of 70 L min<sup>-1</sup> and fraction of inspired oxygen (FiO<sub>2</sub>) 100%. A transglottic jet ventilation catheter was inserted into the trachea through the mouth. HFNO was stopped and replaced by high-frequency jet ventilation when SpO<sub>2</sub> fell to below 92% or when the procedure lasted more than 30 minutes or when the patient presented arrhythmia. Jaw thrust was applied from onset of apnea until the start of surgery and then from the end of surgery until awakening. Neuromuscular blockade was reversed using prostigmine or sugammadex (Bridion<sup>®</sup>, MSD, Hertfordshire, United Kingdom), depending on the train-of-four monitoring results. HFNO was stopped as

soon as the patient regained spontaneous breathing. At the end of surgery, or earlier when SpO<sub>2</sub> was less than 92%, an arterial blood sample was drawn for blood gas analysis. Surgeons were asked to evaluate their satisfaction with the ventilation technique using an 11-point numeric rating scale, with 0 indicating "completely dissatisfied" and 10 indicating "completely satisfied." Patients were transferred to the post-anesthesia care unit (PACU) for routine care. All patients were routinely placed on supplemental O<sub>2</sub> (3 L min<sup>-1</sup>) for the first 30 minutes of their PACU stay. On the first postoperative day, ambulatory patients were contacted by phone and any post-discharge complications were noted. Hospitalized patients were also screened for complications on the first postoperative day.

## Outcomes

The primary outcome of the study was the incidence of peripheral oxygen saturation (SpO<sub>2</sub>) below 95% during the first 15 minutes of apnea. Secondary endpoints included the incidence of SpO<sub>2</sub> below 95% throughout the entire procedure (maximum 30 minutes), the incidence of SpO<sub>2</sub> below 92% throughout the procedure, the time between onset of apnea and onset of SpO<sub>2</sub> <95%, the incidence of hypercapnia (defined as arterial carbon dioxide tension >8 kPa or 60 mmHg) at the end of the procedure (or at the end of the experiment, whichever came first), blood pH at the end of the procedure (or at the end of the experiment, whichever came first), time-course of arterial blood pressure, heart rate, intra-operative and post-operative complications, and the surgeon's satisfaction.

## Statistical analysis

Because this study was purely descriptive, no power analysis was performed; we


planned to include 100 patients. An interim analysis was planned after inclusion of the first 30 patients. Results are expressed as percentage (95% confidence interval, CI) and median (interquartile range, IQR). Data were analyzed using IBM SPSS Statistics, version 22 (IBM Corp., Armonk, NY, USA).

Results

The study was prematurely terminated for safety concerns owing to the development of severe hypercapnia in a number of

patients (inclusion of the first patient on July 5th, 2018 and inclusion of the last one on January 14th, 2019). At the time of study discontinuation, 30 patients had been included and 29 had completed the study protocol (17 at Foch Hospital and 12 at Institut Curie) (see Figure 1). Baseline patient demographic characteristics are presented in Table 1. The median age of participants was 56 years and the male-to-female ratio was 62% to 38%.

Median total operating time was 12 [10.00–22.00] minutes (Table 2). Five



**Figure 1.** Flow diagram.  
SpO2, peripheral oxygen saturation.

patients (17.2%; 95% CI: 7.6–34.5) presented blood oxygen saturation below 95% during the first 15 minutes of the procedure and eight patients (27.6%; 95% CI: 14.7–45.7) presented blood oxygen saturation below 95% throughout surgery. Six patients (20.7%; 95% CI: 9.8–38.4) required rescue jet ventilation for blood oxygen saturation below 92%. Median apnea time before SpO<sub>2</sub> <95% was 13.5 (IQR: 10.0–17.7) minutes with a minimum of 2 minutes and a maximum of 49 minutes.

**Table 1.** Patient characteristics.

	N = 29
Age (years), median [IQR]	56 [49–66]
Sex (men:women)	18:11 (62:38)
BMI (kg m <sup>-2</sup> ), median [IQR]	24 [21–26]
Comorbidities, n (%)	
Smoking	19 (66)
Diabetes	1 (3)
Hypertension	4 (14)
Ischemic heart disease	1 (3)
Pulmonary disease	2 (7)
Procedure, n (%)	
Diagnostic suspension laryngoscopy	15 (52)
Vocal cord lesion resection	8 (28)
Vocal cord fat injection	3 (10)
Vocal cord heterologous substance injection	3 (10)

BMI, body mass index; IQR, interquartile range.

The evolution of SpO<sub>2</sub> in all patients is shown in Figure 2, with all normoxic patients (SpO<sub>2</sub> ≥92%) in the left panel and hypoxic patients in the right panel. Median arterial partial pressure of carbon dioxide (PaCO<sub>2</sub>) at the end of the procedure, or at the time of study discontinuation, was 9.73 (IQR: 8.8–10.9) kPa and was higher than 8 kPa (60 mmHg) in 88.9% (95% CI: 71.9–96.1) of patients. One patient had a PaCO<sub>2</sub> of 13.45 kPa. The lowest reported pH was 7.11.

No patients experienced cardiac arrhythmia. The mean time from sugammadex injection and discontinuation of propofol infusion to the patients opening their eyes was 4 minutes (95% CI: 3.0–5.5). No seizures were reported. All patients remained hemodynamically stable. One patient experienced intra-operative epistaxis. No other intra-operative complications or postoperative complications were reported.

The median surgeon's satisfaction score with laryngeal visibility was 10 (IQR: 10–10).


## Discussion

In this case series, we observed that patients undergoing suspension laryngoscopy and receiving high-flow apneic nasal oxygenation could occasionally develop early oxygen desaturation and severe hypercapnia.

**Table 2.** Outcome results.

	N = 29
Total surgery time (min), median [IQR]	12 [10.00–22.00]
SpO <sub>2</sub> <95% in first 15 min of procedure, % (95% CI)	17.2 (7.6–34.5)
SpO <sub>2</sub> <95% before procedure end, % (95% CI)	27.6 (14.7–45.7)
SpO <sub>2</sub> <92% before procedure end, % (95% CI)	20.7 (9.8–38.4)
Apnea time without desaturation (SpO <sub>2</sub> <95%) (min), median [IQR]	13.5 [10.17.7]
PaCO <sub>2</sub> at end of procedure/study discontinuation (kPa), median [IQR]	9.73 [8.8–10.9]
PaCO <sub>2</sub> >8 kPa at end of procedure/study discontinuation, % (95% CI)	88.9 (71.9–96.1)
Surgeon satisfaction score, median [IQR]	10 [10–10]

SpO<sub>2</sub>, peripheral oxygen saturation; PaCO<sub>2</sub>, arterial partial pressure of carbon dioxide; CI, confidence interval; IQR, interquartile range.



**Figure 2.** Individual evolutions of pulse oximetry. Left panel: evolution of pulse oximetry for each patient whose SpO<sub>2</sub> value was  $\geq 92\%$ . Right panel: evolution of pulse oximetry for each of the 6 patients with SpO<sub>2</sub>  $< 92\%$ . SpO<sub>2</sub>, peripheral oxygen saturation.

All anesthetic techniques used for suspension laryngoscopy and for other surgical laryngeal procedures have advantages and disadvantages.<sup>1</sup> Briefly, total intra-venous anesthesia and maintenance of spontaneous ventilation are widely used but the balance between sufficiently deep anesthesia and satisfactory gas exchange may be difficult or impossible to achieve.<sup>9</sup> In contrast, tracheal intubation avoids most ventilatory problems at the cost of discomfort for the operator, even if a small-diameter intubation probe is used. High frequency jet ventilation can be used with i) a laryngoscope with a specific channel for jet ventilation (i.e., supraglottic jet ventilation),<sup>10</sup> ii) a tracheal catheter placed through the mouth (i.e., subglottic jet ventilation),<sup>11</sup> or iii) a tracheal catheter placed percutaneously through the cricothyroid membrane.<sup>12</sup> However, high frequency jet ventilation remains a marginal technique owing to the requirement for a specific ventilator and the risk of a specific complication, notably pneumothorax.<sup>13</sup> Therefore, HFNO appears to be the ideal

anesthetic technique for laryngoscopy suspension because it is non-invasive and does not interfere with operative vision to ensure satisfactory gas exchange.

Oxygen flow delivered by conventional nasal cannula is generally limited to  $10 \text{ L min}^{-1}$  whereas HFNO uses oxygen flows between 50 and  $120 \text{ L min}^{-1}$  administered via specially designed nasal cannulas. These high O<sub>2</sub> flows mean that little ambient air is drawn into the cannula, allowing for a high FiO<sub>2</sub> (95% to 100%).

HFNO now has a prominent role in intensive care as an alternative to noninvasive ventilation owing to several advantages, particularly clearance of the anatomical dead space,<sup>14</sup> improved CO<sub>2</sub> elimination,<sup>15</sup> creation of positive end-expiratory pressure roughly proportional to flow (1 cm H<sub>2</sub>O for every  $10 \text{ L min}^{-1}$  of flow) when the patient keeps their mouth closed,<sup>16,17</sup> reduced work of breathing, and attenuation of inspiratory resistance.<sup>18</sup> These characteristics of HFNO result in improvement of patient comfort as a



result of humidification and heating of the delivered mixture.<sup>19</sup>

HFNO is also used in anesthesia, especially since the publication by Patel and Nouraei reporting prolonged median desaturation-free apnea time in paralyzed patients under general anesthesia, leading to the introduction of THRIVE.<sup>7</sup> HFNO has also been tested for preoxygenation prior to induction of general anesthesia;<sup>20</sup> continuous oxygenation during intubation, especially when rapid sequence intubation is indicated;<sup>21</sup> oxygenation during fiberoptic intubation;<sup>22</sup> oxygenation during intravenous anesthesia, especially in the presence of airway obstruction;<sup>17</sup> and postoperative O<sub>2</sub> therapy,<sup>23</sup> particularly after cardiac<sup>24</sup> and cardiothoracic<sup>25</sup> procedures.

THRIVE could be the best technique for diagnostic and surgical suspension laryngoscopy because it permits complete visualization of the glottis. Evaluation of the safety of THRIVE consists of evaluation of the risk of oxygen desaturation and hypercapnia during prolonged apnea under general anesthesia with muscle paralysis. In 2015, Patel and Nouraei<sup>7</sup> reported a prolonged apneic window for up to 65 minutes with HFNO, with a median apnea time of 14 minutes (IQR: 9–19) with no desaturation below 90% and an apnea time ranging from 5 minutes to 65 minutes. Other authors have reported similar apnea times: 19 (IQR: 15–24) minutes in a series of patients undergoing laryngeal or tracheal surgery performed under apneic conditions<sup>26</sup> and 18 (IQR: 10–27) minutes in a group of patients undergoing surgery for subglottic stenosis.<sup>27</sup> Of particular interest is a study by Gustafsson et al. on patients undergoing brief laryngeal surgery who were oxygenated using THRIVE with 100% O<sub>2</sub> at 40–70 L min<sup>-1</sup>. Blood gas analyses were performed every 5 minutes during the procedure. The authors defined the apnea time as the interval between the onset of apnea

and the end of the procedure, or earlier in the following situations: apnea time exceeding 40 minutes, PaCO<sub>2</sub> >11 kPa, pH <7.15, SpO<sub>2</sub> <90%, or the presence of malignant arrhythmia. Gustafsson et al. reported a mean apnea time of 22.5 ± 4.5 minutes,<sup>28</sup> which would appear to be suitable for suspension laryngoscopy performed as a diagnostic procedure or to treat pharyngolaryngeal lesions as the procedure duration is only slightly longer than reported apnea times. Contrary to these data showing the safety of HFNO for these indications, we report disquieting results in our series: oxygen desaturation occurred before completion of surgery in 20.6% of cases, and we decided to introduce rescue high-frequency jet ventilation when SpO<sub>2</sub> fell below 92% to avoid subsequent rapid desaturation because such SpO<sub>2</sub> values are close to the steep slope of the oxyhemoglobin dissociation curve. Moreover, we think that the study of HFNO safety should not stop at the analysis of mean or median apnea times; examination of individual data shows that a relatively high percentage of patients experience oxygen desaturation, whether defined as SpO<sub>2</sub> <95%, <92%, or <90%.<sup>7,26,28</sup>

Our protocol differs from that reported in previous publications in terms of choice of preoxygenation technique. Previous publications have reported preoxygenation using HFNO with different flows and durations: 70 L min<sup>-1</sup> for 10 minutes,<sup>7</sup> 40 L min<sup>-1</sup> for 3 minutes,<sup>28</sup> and 80 L min<sup>-1</sup> for 3 minutes.<sup>26</sup> In our study, we used face mask preoxygenation to achieve an end-tidal O<sub>2</sub> ≥90%, which constitutes the gold standard for preoxygenation and is the only way to document lung denitrogenation, a necessary condition for ventilatory flow motion, the physiological basis of apneic oxygenation.<sup>29</sup>

We observed one case of early desaturation (2 minutes after the onset of apnea) despite preoxygenation performed



according to the protocol and application of jaw thrust as soon as apnea was obtained. A possible explanation for this case of early desaturation could be unidentified obstruction of the nasal cavity. Another hypothesis is this was owing to the presence of unidentified tracheal or bronchial collapse. To date, no published studies have identified a patient population at risk of early oxygen desaturation during apneic HFNO. In the absence of easily identifiable risk factors for HFNO failure, practitioners should always have an alternate airway management plan if they decide to use HFNO during apnea.

Several mechanisms have been proposed to explain CO<sub>2</sub> elimination during HFNO: rapid clearance of nasal cavities,<sup>14</sup> positive airway pressure generation,<sup>30,31</sup> reduced O<sub>2</sub> dilution by nitrogen<sup>32</sup> and, more recently, interaction between entrained and highly turbulent supraglottic flow vortices created by HFNO and cardiogenic oscillations.<sup>33,34</sup> Laviola et al. proposed a combination of cardiac oscillations with a high degree of gas mixing in the anatomical dead space and high-frequency glottic pressure oscillations as a mechanism of CO<sub>2</sub> clearance with HFNO.<sup>34</sup> Increased blood carbon dioxide levels during THRIVE has been reported, with lower values in a laboratory model (0.15 kPa min<sup>-1</sup>)<sup>33</sup> and in adults under general anesthesia (from 0.17<sup>27</sup> to 0.24<sup>28</sup> kPa min<sup>-1</sup>) than in our study (0.3 kPa min<sup>-1</sup> assuming a preoperative median value of 5 kPa). Caution must be urged when capnography is used; it has been shown that the increase in PaCO<sub>2</sub> is twice that measured using end-tidal CO<sub>2</sub>.<sup>28</sup> However, no difference has been reported between transcutaneous CO<sub>2</sub> monitoring and PaCO<sub>2</sub>.<sup>28</sup>

As expected, surgeons reported a high level of satisfaction with THRIVE owing to the absence of interference between the ventilation technique and the surgical procedure. However, the main drawback was

that THRIVE could not be used for procedures requiring the use of a laser owing to the risk of combustion. The use of laser is also contraindicated when using jet ventilation where a FiO<sub>2</sub> of 50% is used in laser surgery, although the FiO<sub>2</sub> actually delivered is lower owing to air entrainment.<sup>35</sup> However, this dilution cannot be evaluated in practice, which has led many practitioners to prefer the use of positive pressure ventilation via endotracheal tubes specifically designed for laser surgery.

This study has several limitations. This was a case series without a control group, which considerably limits the conclusions to be drawn. Furthermore, our study had to be prematurely terminated owing to patient safety concerns. Regular review of study data by the data monitoring committee revealed a high incidence of sometimes severe hypercapnia; the data monitoring committee therefore recommended termination of the study. We used an SpO<sub>2</sub> monitor that cannot differentiate oxyhemoglobin and carboxyhemoglobin; this device may therefore have underestimated hypoxemia when patients had smoked several hours before anesthesia.<sup>36</sup> We used face mask preoxygenation in this study whereas all previous studies used HFNO preoxygenation before apneic oxygenation; this makes it difficult to compare our results with previously reported data and suggests that differences in the preoxygenation technique may be at least partly responsible for differences in the reported results. Further studies are needed to determine the optimal preoxygenation technique before apneic HFNO. Our study population also did not include any obese patients or patients with obstructive airway tumors, precluding generalization of our results to those clinical settings.

## Conclusions

In contrast to several studies showing the safety of HFNO, our case series showed

less favorable results. We observed a relatively high incidence of suboptimal oxygen saturation and occasional early desaturation during suspension laryngoscopy under general anesthesia. It is possible that the safety of HFNO can be improved if this technique is also used to provide pre-oxygenation. Our results indicate caution is needed if HFNO is chosen, and a reliable rescue technique must be readily available.

### Acknowledgements

The authors would like to thank Cecile Naudin for her help in the management of the study. The authors would also like to thank Polly Gobin for her linguistic assistance.

### Author contributions

Conceptualization: M.S., M.F., M.L-G.

Methodology: P.S.

Validation: M.S., M.F., P.S.

Formal analysis: P.S.

Investigation: M.S., A.A-F., R.T., I.W., M.L-G.

Data curation: M.S., A.A-F., R.T., I.W., M.F., M.L-G.

Visualization: M.F.

Supervision: M.F.

Project administration, Funding acquisition, Resources: P.S.

Writing—original draft preparation: M.S.

Writing—review and editing: M.F.

All authors have read and agreed to the published version of the manuscript.

### Data availability statement

The dataset generated during the current study is available from the corresponding author on reasonable request.

### Declaration of conflicting interests


The authors declare that there is no conflict of interest.

### Funding

The authors disclosed receipt (pending publication) of the following financial support for the research, authorship, and/or publication of this article: This research was funded by Clinique

Ambroise Paré, Neuilly sur Seine, France. Paykel and Fisher provided material but did not have any involvement with the study design and conduct nor analysis of the data.

### ORCID iD

Mary Saad  <https://orcid.org/0000-0002-2918-1330>

### References

1. Jaquet Y, Monnier P, Van Melle G, et al. Complications of different ventilation strategies in endoscopic laryngeal surgery: a 10-year review. *Anesthesiology* 2006; 104: 52–59.
2. Habrial P and Lasocki S. [National survey on ENT panendoscopy] 2018 [Available from: <https://sfar.org/enquete-nationale-sur-les-panendoscopies-en-ori/>. Accessed on May 16, 2022.
3. Bonnet N, Martin O, Boubaya M, et al. High flow nasal oxygen therapy to avoid invasive mechanical ventilation in SARS-CoV-2 pneumonia: a retrospective study. *Ann Intensive Care* 2021; 11: 37.
4. Spoletini G, Alotaibi M, Blasi F, et al. Heated humidified high-flow nasal oxygen in adults: Mechanisms of action and clinical implications. *Chest* 2015; 148: 253–261.
5. Beng Leong L, Wei Ming N and Wei Feng L. High flow nasal cannula oxygen versus noninvasive ventilation in adult acute respiratory failure: a systematic review of randomized-controlled trials. *Eur J Emerg Med* 2019; 26: 9–18.
6. Guitton C, Ehrmann S, Volteau C, et al. Nasal high-flow preoxygenation for endotracheal intubation in the critically ill patient: a randomized clinical trial. *Intensive Care Med* 2019; 45: 447–458.
7. Patel A and Nouraei SA. Transnasal Humidified Rapid-Insufflation Ventilatory Exchange (THRIVE): a physiological method of increasing apnoea time in patients with difficult airways. *Anaesthesia* 2015; 70: 323–329.
8. Von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for

- reporting observational studies. *Int J Surg* 2014; 12: 1495–1499.
9. Habrial P, Léger M, Costerousse F, et al. Spontaneous Breathing for Panendoscopy? Retrospective Cohort and Results of a French Practice Survey. *OTO Open* 2022; 6: 2473974x211065015.
  10. Bacher A, Pichler K and Aloy A. Supraglottic combined frequency jet ventilation versus subglottic monofrequent jet ventilation in patients undergoing microlaryngeal surgery. *Anesth Analg* 2000; 90: 460–465.
  11. Hunsaker DH. Anesthesia for microlaryngeal surgery: the case for subglottic jet ventilation. *Laryngoscope* 1994; 104: 1–30.
  12. Bourgain JL, Desruennes E, Fischler M, et al. Transtracheal high frequency jet ventilation for endoscopic airway surgery: a multicentre study. *Br J Anaesth* 2001; 87: 870–875.
  13. Cook TM and Alexander R. Major complications during anaesthesia for elective laryngeal surgery in the UK: a national survey of the use of high-pressure source ventilation. *Br J Anaesth* 2008; 101: 266–272.
  14. Moller W, Celik G, Feng S, et al. Nasal high flow clears anatomical dead space in upper airway models. *J Appl Physiol (1985)* 2015; 118: 1525–1532.
  15. Sivieri EM, Foglia EE and Abbasi S. Carbon dioxide washout during high flow nasal cannula versus nasal CPAP support: An in vitro study. *Pediatr Pulmonol* 2017; 52: 792–798.
  16. Parke RL and McGuinness SP. Pressures delivered by nasal high flow oxygen during all phases of the respiratory cycle. *Respir Care* 2013; 58: 1621–1624.
  17. Booth AWG, Vidhani K, Lee PK, et al. Spontaneous Respiration using Intravenous anaesthesia and Hi-flow nasal oxygen (STRIVE Hi) maintains oxygenation and airway patency during management of the obstructed airway: an observational study. *Br J Anaesth* 2017; 118: 444–451.
  18. Dysart K, Miller TL, Wolfson MR, et al. Research in high flow therapy: mechanisms of action. *Respir Med* 2009; 103: 1400–1405.
  19. Chikata Y, Izawa M, Okuda N, et al. Humidification performance of two high-flow nasal cannula devices: a bench study. *Respir Care* 2014; 59: 1186–1190.
  20. Simon M, Wachs C, Braune S, et al. High-flow nasal cannula versus bag-valve-mask for preoxygenation before intubation in subjects with hypoxemic respiratory failure. *Respir Care* 2016; 61: 1160–1167.
  21. Mir F, Patel A, Iqbal R, et al. A randomised controlled trial comparing transnasal humidified rapid insufflation ventilatory exchange (THRIVE) pre-oxygenation with facemask pre-oxygenation in patients undergoing rapid sequence induction of anaesthesia. *Anaesthesia* 2017; 72: 439–443.
  22. Badiger S, John M, Fearnley RA, et al. Optimizing oxygenation and intubation conditions during awake fibre-optic intubation using a high-flow nasal oxygen-delivery system. *Br J Anaesth* 2015; 115: 629–632.
  23. Lu Z, Chang W, Meng S, et al. The effect of high-flow nasal oxygen therapy on post-operative pulmonary complications and hospital length of stay in postoperative patients: A systematic review and meta-analysis. *J Intensive Care Med* 2020; 35: 1129–1140.
  24. Zochios V, Collier T, Blandszun G, et al. The effect of high-flow nasal oxygen on hospital length of stay in cardiac surgical patients at high risk for respiratory complications: a randomised controlled trial. *Anaesthesia* 2018; 73: 1478–1488.
  25. Wu X, Cao W, Zhang B, et al. Effect of high-flow nasal cannula oxygen therapy vs conventional oxygen therapy on adult post-cardiothoracic operation: A meta-analysis. *Medicine (Baltimore)* 2018; 97: e12783.
  26. Lyons C and Callaghan M. Apnoeic oxygenation with high-flow nasal oxygen for laryngeal surgery: a case series. *Anaesthesia* 2017; 72: 1379–1387.
  27. To K, Harding F, Scott M, et al. The use of Transnasal Humidified Rapid-Insufflation Ventilatory Exchange in 17 cases of subglottic stenosis. *Clin Otolaryngol* 2017; 42: 1407–1410.
  28. Gustafsson IM, Lodenius Å, Tunelli J, et al. Apnoeic oxygenation in adults under general anaesthesia using Transnasal Humidified Rapid-Insufflation Ventilatory Exchange

- (THRIVE) – a physiological study. *Br J Anaesth* 2017; 118: 610–617.
29. Bartlett RG Jr, Brubach HF and Specht H. Demonstration of a ventilatory mass flow during ventilation and apnea in man. *J Appl Physiol* 1959; 14: 97–101.
  30. Ritchie JE, Williams AB, Gerard C, et al. Evaluation of a humidified nasal high-flow oxygen system, using oxygraphy, capnography and measurement of upper airway pressures. *Anaesth Intensive Care* 2011; 39: 1103–1110.
  31. Parke RL, Bloch A and McGuinness SP. Effect of Very-High-Flow Nasal Therapy on Airway Pressure and End-Expiratory Lung Impedance in Healthy Volunteers. *Respir Care* 2015; 60: 1397–1403.
  32. Ward JJ. High-flow oxygen administration by nasal cannula for adult and perinatal patients. *Respir Care* 2013; 58: 98–122.
  33. Hermez LA, Spence CJ, Payton MJ, et al. A physiological study to determine the mechanism of carbon dioxide clearance during apnoea when using transnasal humidified rapid insufflation ventilatory exchange (THRIVE). *Anaesthesia* 2019; 74: 441–449.
  34. Laviola M, Das A, Chikhani M, et al. Computer simulation clarifies mechanisms of carbon dioxide clearance during apnoea. *Br J Anaesth* 2019; 122: 395–401.
  35. Seigneur F, Fischler M, Bourreli B, et al. Air entrainment during high-frequency jet-ventilation. Simulation of a bronchoscopy with a lung model. *Bull Eur Physiopathol Respir* 1986; 22: 341–347.
  36. Barker SJ and Tremper KK. The effect of carbon monoxide inhalation on pulse oximetry and transcutaneous PO<sub>2</sub>. *Anesthesiology* 1987; 66: 677–679.