



## Estimation of physical conditions in PDRs Bayesian approach with spatial regularization

Pierre Palud, Franck Le Petit, Pierre Chainais, Emeric E. Bron,  
Pierre-Antoine Thouvenin

### ► To cite this version:

Pierre Palud, Franck Le Petit, Pierre Chainais, Emeric E. Bron, Pierre-Antoine Thouvenin. Estimation of physical conditions in PDRs Bayesian approach with spatial regularization. Multi-line Diagnostics of the Interstellar Medium, Apr 2022, Nice, France. . hal-03953044

HAL Id: hal-03953044

<https://hal.science/hal-03953044v1>

Submitted on 23 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Pierre Palud<sup>1,2</sup>, F. Le Petit<sup>1</sup>, P. Chainais<sup>2</sup>,  
E. Bron<sup>1</sup>, P.-A. Thouvenin<sup>2</sup>, ORIONB consortium  
<sup>1</sup>LERMA, Obs. de Paris, PSL Research Univ., CNRS, Meudon  
<sup>2</sup>Univ. Lille, CNRS, Centrale Lille, UMR 9189 – CRIStAL, France

### Aims and contributions

New large hyper-spectral surveys in radio-astronomy: game changer for study of star formation, feedback mechanisms and chemistry of interstellar gas.  
 ⇒ now possible to observe full molecular clouds (10 pc size) at dense core scale (<0.1 pc) spatial resolution. For instance, the IRAM-30m Large Program «Orion B» [1] provides a  $10^6$  pixel observation map, covering  $\sim 250$  pc<sup>2</sup> of the Orion B cloud with emission of dozens of molecules.

**Goal** To estimate maps of thermal pressure  $P_{th}$ , UV radiation intensity  $G_0$ , visual extinction  $A_V$ , scaling factor  $\kappa$  and the associated uncertainties.


**Challenge** Variability of the SNR: the brightest regions can be well constrained, but the regions with low SNR lead to degenerate solutions.

**Approach**

- ★ Mixture of both additive and multiplicative noises + limited detectability ⇒ more realistic observation model
- ★ Spatial regularization ⇒ exploitation of the information contained in neighbouring pixels.
- ★ Bayesian approach ⇒ access to accurate credibility intervals.

### Problem formulation

→ Meudon PDR code [2]: solves radiative transfer, chemistry, thermal balance on stationary 1D plane-parallel model.



→ Assumption: physics in PDR = interpolation of a grid of simulations of Meudon PDR code. Defines function  $f$  s.t. for one pixel  $n$ :


$$f : x_n = (\kappa, P_{th}, G_0, A_V) \mapsto \underbrace{(y_{n,\ell})_{\ell=1}^L}_{\text{physical parameters (to estimate)}} \quad \underbrace{(y_{n,\ell})_{\ell=1}^L}_{\text{integrated intensities (observed)}}$$

→ observation model involves detectability limit  $\omega$ , gaussian noise  $\varepsilon_{n,\ell}^{(a)}$  and lognormal noise  $\varepsilon_{n,\ell}^{(m)}$

$$y_{n,\ell} = \begin{cases} \omega & \text{if } \varepsilon_{n,\ell}^{(m)} f_\ell(x_n) + \varepsilon_{n,\ell}^{(a)} \leq \omega \\ \varepsilon_{n,\ell}^{(m)} f_\ell(x_n) + \varepsilon_{n,\ell}^{(a)} & \text{else} \end{cases}$$

⇒ defines a **likelihood** function  $\pi(y | x)$ .

### Application on toy dataset (100 pixels)



### Bayesian approach

→ State-of-the-art in millimeter/IR astronomy estimations: Maximum Likelihood Estimates (MLE) ⇒ very sensitive to noise

→ For robust estimators: spatial regularization prior  $\pi(x)$  (norm of image gradient or laplacian)

→ Posterior probability density function:


$$\pi(x | y) \propto \pi(y | x) \pi(x)$$

→ To derive estimators and credibility intervals from posterior distribution: **sample** from it with Monte Carlo Markov Chain (MCMC).

→ Our sampler mixes two kernels: one identifies local minima, one efficiently explores them.

→ Our punctual estimator: MMSE (Minimum Mean Squared Error) = mean of posterior.

### NGC 7023 (1 pixel)



→ Estimation from 10 lines of CO and 2 lines of CI.

→ In contrast with [3],  $G_0$  and  $A_V$  estimated along with  $\kappa$  and  $P_{th}$ .

→ Two regions of acceptable solutions identified.

Summary for main mode

	MLE [3]	MMSE	2.5%	97.5%
$\kappa$	0.7	4.8	2.5	8.6
$P_{th} (\times 10^8)$	1	1.1	1.0	1.3
$G_0 (\times 10^3)$	2.6	1.6	1.0	2.6
$A_V$	10	3.1	1.6	30.5

### References

- [1] Pety et al., The anatomy of the Orion B Giant Molecular Cloud: A local template for studies of nearby galaxies, 2016
- [2] Le Petit et al., A model for atomic and molecular interstellar gas the Meudon PDR code, 2009
- [3] Joblin et al., Structure of photodissociation fronts in star-forming regions revealed by Herschel observations of high- $J$  CO emission lines, 2018

### Conclusions

**Achieved:**

- ✓ Spatial regularization ⇒ more robust estimations of maps of parameters,
- ✓ Bayesian approach ⇒ more complete description than punctual estimators.

**Future Work:**

- Application to other environments than PDR (shocks, dark clouds, etc.),
- Code acceleration to scale to  $\sim 10^6$  pixel observations (e.g., Orion B IRAM Large Programm).