



**HAL**  
open science

# Goal-oriented error estimation based on equilibrated flux reconstruction for the approximation of the harmonic formulations in eddy current problems

Emmanuel Creusé, Serge Nicaise, Zuqi Tang

## ► To cite this version:

Emmanuel Creusé, Serge Nicaise, Zuqi Tang. Goal-oriented error estimation based on equilibrated flux reconstruction for the approximation of the harmonic formulations in eddy current problems. 2023. hal-03954215v1

**HAL Id: hal-03954215**

**<https://hal.science/hal-03954215v1>**

Preprint submitted on 24 Jan 2023 (v1), last revised 30 Jan 2024 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Goal-oriented error estimation based on equilibrated flux reconstruction for the approximation of the harmonic formulations in eddy current problems

Emmanuel Creusé\*, Serge Nicaise

Univ. Polytechnique Hauts-de-France, INSA Hauts-de-France,

CERAMATHS - Laboratoire de Matériaux Céramiques et de Mathématiques,

F-59313 Valenciennes, France

and

Zuqi Tang

Univ. Lille, Arts et Métiers Institute of Technology, Centrale Lille, Junia,

ULR 2697-L2EP, F-59000 Lille, France

January 24, 2023

## Abstract

In this work, we propose an *a posteriori* goal-oriented error estimator for the harmonic ( $\mathbf{A} - \varphi$ ) formulation arising in the modeling of eddy current problems, approximated by non-conforming finite element methods. It is based on the resolution of a dual problem associated with the initial one. For each of these two problems, a guaranteed equilibrated estimator is developed using some flux reconstructions. These fluxes also allow to obtain a goal-oriented error estimator that is fully computable and can be split in a principal part and a reminder one. Our theoretical results are illustrated by numerical experiments.

**Keywords** : Maxwell equations, potential formulations, goal-oriented a posteriori estimators, finite element method.

## 1 Introduction

The finite element method is widely used to solve a large variety of electromagnetic problems, and many papers have been devoted to this topic for the last decades. More particularly, in the context of low-frequency electromagnetics allowing a quasi-static approximation, some specific models are usually introduced. Among them, the so-called ( $\mathbf{A} - \varphi$ ) formulation consists in computing a magnetic vector potential  $\mathbf{A}$  as well as an electric scalar potential  $\varphi$ , allowing to obtain approximated values of the magnetic flux density  $\mathbf{B}$  as well as of the electric field  $\mathbf{E}$  arising in the Maxwell equations. From there, an important question to address is to ensure the good quality of the numerical solutions obtained. Consequently, some *a posteriori* error estimators have been developed in order to provide a global upper bound of the numerical error, as well as some local lower bounds, very useful to drive a mesh-refinement strategy. We refer to [12, 26, 5] for residual estimators and to [13, 10] for equilibrated ones, allowing in this second case to obtain a sharp upper bound of the error without any unknown multiplicative constant. All these estimators have been tested in several configurations, from academic to more industrial ones (see e.g. [27, 28, 30]). In all the papers quoted above, the error to control is defined globally, and

---

\*Corresponding author: emmanuel.creuse@uphf.fr

corresponds to the value of a global energy, directly linked to the bilinear form arising in the variational formulation of the model.

Nevertheless, in many applications, engineers are interested in some local physical quantities, that are called quantities of interest (QOI). To stay in the field of electromagnetic problems, let us mention for example the computation of the flux through a coil [20] or the magnetic flux density intensity at a given point of an electromagnetic device [17, 21]. Some specific estimators, called "goal-oriented estimators", have consequently to be developed in order to derive an upper bound for this kind of errors. In the case of magnetostatic problems, we refer e.g. to [29] where a dual formulation of the model is used.

In this paper, we derive a goal-oriented *a posteriori* estimator for the harmonic  $\mathbf{A} - \varphi$  formulation, to control an error defined by a linear form operating on the magnetic vector potential  $\mathbf{A}$ . Using some flux reconstructions of the primal and dual formulations, the upper bound is obtained without any multiplicative constant. Moreover, this fully computable estimator can be split in a principal part and a reminder one, the last one being in some cases of higher order and can be most of the time disregarded. This result can be seen as an extension of [11], which is devoted to classical diffusion problems. Moreover, this new goal-oriented estimator is based on some equilibrated error estimators developed for the primal and the dual formulations of the problem. They are based on potential reconstructions of the vector and scalar potentials, and have already been derived in a previous work for conforming finite element approximations [10]. Let us note that the present work also allows to generalize them to the case of non-conforming approximations.

The paper is organized as follows. Section 2 gives the continuous formulation of the eddy-current problem in the  $\mathbf{A} - \varphi$  formulation. Moreover, some regularity results are derived on the solution (see Lemma 2.2 to 2.4), which have their own interest but will be useful in the interpretation of our numerical results. Section 3 provides a guaranteed equilibrated error estimator for non-conforming approximations of the problem, leading to Theorem 3.1 ensuring the guaranteed reliability of the estimation. Then, section 4 details the goal-oriented functional as well as the associated adjoint problem and its discrete approximation. In section 5, the error representation of the error is given in Theorem 5.1, and Theorem 5.2 allows to prove that the remainder term can be controlled by the product of the equilibrated estimators devoted to the direct and dual problems. Finally, the obtained theoretical results are illustrated by some numerical experiments in section 6.

## 2 The continuous formulation

Let us consider a bounded simply connected polyhedral domain  $D \subset \mathbb{R}^3$  with a Lipschitz and connected boundary  $\Gamma = \partial D$ .  $D$  is composed of two subdomains: the conducting domain  $D_c$  and the non-conducting domain  $D_{nc} = D \setminus \bar{D}_c$ . Let us remark that  $D_c$  is supposed bounded, simply connected with a Lipschitz connected boundary  $\partial D_c$  and strictly included into  $D$ , in the sense that  $\bar{D}_c \subset D$ . The source domain  $D_s$  where the divergence free current density  $\mathbf{J}_s$  is imposed is usually included into  $D_{nc}$ , but our mathematical analysis does not require this assumption. The eddy current problem is given by:

Find the electric field  $\mathbf{E}$  and the magnetic field  $\mathbf{H}$  solution of

$$\begin{cases} \operatorname{curl} \mathbf{E} &= -j\omega \mathbf{B} & \text{in } D, \\ \operatorname{curl} \mathbf{H} &= \mathbf{J}_s + \mathbf{J}_e & \text{in } D, \\ \operatorname{div} \mathbf{B} &= 0 & \text{in } D, \end{cases} \quad (1)$$

where  $j^2 = -1$  is the unit imaginary number,  $\omega$  the pulsation, and the magnetic flux  $\mathbf{B}$  and the eddy current  $\mathbf{J}_e$  are given by the constitutive laws

$$\mathbf{B} = \mu \mathbf{H} \text{ in } D \text{ and } \mathbf{J}_e = \sigma \mathbf{E} \text{ in } D_c, \quad (2)$$

where  $\mu$  denotes the magnetic permeability and  $\sigma$  the electric conductivity. Here we suppose that  $\mu, \sigma \in L^\infty(D)$  and that there exists positive real numbers  $\mu_0, \sigma_0$  such that

$$\begin{aligned} \mu &\geq \mu_0 && \text{a.e. in } D \\ \sigma &\geq \sigma_0 && \text{a.e. in } D_c, \\ \sigma &= 0 && \text{a.e. in } D_{nc}. \end{aligned}$$

Note that the divergence free property of  $\mathbf{J}_s$ , the fact that  $\partial D_c \subset D$ , and the second equation of (1) implies that

$$\operatorname{div} \mathbf{J}_e = 0 \text{ in } D_c, \quad (3)$$

as well as

$$\mathbf{J}_e \cdot \mathbf{n} = 0 \text{ on } \partial D_c, \quad (4)$$

where  $\mathbf{n}$  stands for the unit outward normal to  $D$  or  $D_c$  depending on the context.

System (1)-(2) is completed with the following boundary conditions on  $\Gamma$

$$\mathbf{B} \cdot \mathbf{n} = 0 \text{ on } \Gamma. \quad (5)$$

Before stating the  $\mathbf{A} - \varphi$  formulation of this problem, let us introduce some notations used throughout the paper. On a given domain  $\mathcal{D}$ , the  $L^2(\mathcal{D})$ -norm is denoted by  $\|\cdot\|_{\mathcal{D}}$ , and the corresponding  $L^2(\mathcal{D})$ -inner product by  $(\cdot, \cdot)_{\mathcal{D}}$ . In the case of  $\mathcal{D} = D$ , the index  $D$  is dropped.  $H_0^1(\mathcal{D})$  is the subspace of  $H^1(\mathcal{D})$  with vanishing trace on  $\partial \mathcal{D}$  and

$$H_0(\operatorname{curl}, \mathcal{D}) = \left\{ \mathbf{F} \in L^2(\mathcal{D})^3 : \operatorname{curl} \mathbf{F} \in L^2(\mathcal{D})^3, \mathbf{F} \times \mathbf{n} = 0 \text{ on } \partial \mathcal{D} \right\}.$$

Finally, in order to ensure later the uniqueness of the fields, let us introduce the gauge spaces:

$$\begin{aligned} \tilde{X}(\mathcal{D}) &= \left\{ \mathbf{F} \in H_0(\operatorname{curl}, \mathcal{D}) : (\mathbf{F}, \nabla \xi)_{\mathcal{D}} = 0, \forall \xi \in H_0^1(\mathcal{D}) \right\}, \\ \tilde{H}^1(\mathcal{D}) &= \left\{ f \in H^1(\mathcal{D}) : (f, 1)_{\mathcal{D}} = 0 \right\}. \end{aligned}$$

The harmonic  $\mathbf{A} - \varphi$  formulation is based on the introduction of a magnetic vector potential  $\mathbf{A}$  in  $D$  and an electric scalar potential  $\varphi$  in  $D_c$  such that:

$$\mathbf{B} = \operatorname{curl} \mathbf{A} \text{ in } D \text{ and } \mathbf{E} = -j\omega \mathbf{A} - \nabla \varphi \text{ in } D_c. \quad (7)$$

From system (1), the harmonic  $\mathbf{A} - \varphi$  formulation reads:

$$\operatorname{curl} (\mu^{-1} \operatorname{curl} \mathbf{A}) + \sigma (j\omega \mathbf{A} + \nabla \varphi) = \mathbf{J}_s \text{ in } D, \quad (8a)$$

$$\operatorname{div} (\sigma (j\omega \mathbf{A} + \nabla \varphi)) = 0 \text{ in } D_c, \quad (8b)$$

with the boundary conditions, derived from (4)-(5), given by

$$\mathbf{A} \times \mathbf{n} = 0 \text{ on } \Gamma \text{ and } \sigma (j\omega \mathbf{A} + \nabla \varphi) \cdot \mathbf{n} = 0 \text{ on } \partial D_c. \quad (9)$$

The Coulomb gauge on  $\mathbf{A}$ , namely  $\operatorname{div} \mathbf{A} = 0$ , and the zero mean of the potential  $\varphi$  in  $D_c$  ensure the uniqueness of these potentials. Since  $\varphi$  does not make sense in  $D_{nc}$ , we fix an arbitrary extension of  $\varphi$  in the whole domain  $D$ , this choice does not impact the problem since  $\sigma \equiv 0$  in  $D_{nc}$ . The corresponding weak formulation is given by:

Find  $(\mathbf{A}, \varphi) \in \widetilde{X}(D) \times \widetilde{H}^1(D_c)$  such that

$$B((\mathbf{A}, \varphi), (\mathbf{A}', \varphi')) = (\mathbf{J}_s, \mathbf{A}'), \quad \forall (\mathbf{A}', \varphi') \in \widetilde{X}(D) \times \widetilde{H}^1(D_c). \quad (10)$$

where

$$\begin{aligned} B((\mathbf{A}, \varphi), (\mathbf{A}', \varphi')) &= (\mu^{-1} \operatorname{curl} \mathbf{A}, \operatorname{curl} \mathbf{A}')_D \\ &+ j\omega^{-1} (\sigma(j\omega \mathbf{A} + \nabla \varphi), (j\omega \mathbf{A}' + \nabla \varphi'))_{D_c}, \quad \forall (\mathbf{A}, \varphi), (\mathbf{A}', \varphi') \in \widetilde{X}(D) \times \widetilde{H}^1(D_c). \end{aligned}$$

Lemma 2.1 of [12] ensures the existence and uniqueness of the weak solution  $(\mathbf{A}, \varphi)$  of this problem since it was shown there that

$$\|(\mathbf{A}', \varphi')\|_B := |B((\mathbf{A}', \varphi'), (\mathbf{A}', \varphi'))|^{\frac{1}{2}}, \quad \forall (\mathbf{A}', \varphi') \in \widetilde{X}(D) \times \widetilde{H}^1(D_c),$$

is a norm on  $\widetilde{X}(D) \times \widetilde{H}^1(D_c)$  equivalent to the natural one

$$\|(\mathbf{A}, \varphi)\|_V = \left( \|\mathbf{A}'\|_D^2 + \|\mu^{-1/2} \operatorname{curl} \mathbf{A}'\|_D^2 + |\varphi'|_{1, D_c}^2 \right)^{\frac{1}{2}}, \quad \forall (\mathbf{A}', \varphi') \in \widetilde{X}(D) \times \widetilde{H}^1(D_c).$$

Recall that from the definition of  $B$ , we have

$$|B((\mathbf{A}', \varphi'), (\mathbf{A}', \varphi'))|^{\frac{1}{2}} = \left( \left\| \mu^{-1/2} \operatorname{curl} \mathbf{A}' \right\|_D^2 + \left\| \omega^{-1/2} \sigma^{1/2} (j\omega \mathbf{A}' + \nabla \varphi') \right\|_{D_c}^2 \right)^{1/2},$$

for all  $(\mathbf{A}', \varphi') \in \widetilde{X}(D) \times \widetilde{H}^1(D_c)$ .

As  $\mathbf{J}_s$  is divergence free, it was further shown in Lemma 2.3 of [12] that (10) remains valid for non divergence free fields  $\mathbf{A}'$ , namely

$$B((\mathbf{A}, \varphi), (\mathbf{A}', \varphi')) = (\mathbf{J}_s, \mathbf{A}'), \quad \forall (\mathbf{A}', \varphi') \in H_0(\operatorname{curl}, D) \times \widetilde{H}^1(D_c). \quad (11)$$

As usual, the convergence of numerical schemes are related to regularity results of the solution  $(\mathbf{A}, \varphi)$  of (10). But up to now, such results are not available in the literature. Let us then give some of them in some particular cases. Before stating them, we introduce the following subspace of  $H_0(\operatorname{curl}, D)$  (see [1, 6]) defined that

$$X_N(D) = \{\mathbf{A}' \in H_0(\operatorname{curl}, D) : \operatorname{div} \mathbf{A}' \in L^2(D)\},$$

that is a Hilbert space with its natural inner product. With this definition, we can formulate the following results.

**Lemma 2.1** *If  $(\mathbf{A}, \varphi) \in \widetilde{X}(D) \times \widetilde{H}^1(D_c)$  is solution of (10), then*

1.  $\mathbf{A}$  belongs to  $X_N(D)$  and is solution of the (regularized) Maxwell problem

$$\int_D (\mu^{-1} \operatorname{curl} \mathbf{A} \cdot \operatorname{curl} \overline{\mathbf{A}'} + \operatorname{div} \mathbf{A} \operatorname{div} \overline{\mathbf{A}'}) = \int_D \mathbf{f} \cdot \overline{\mathbf{A}'}, \quad \forall \mathbf{A}' \in X_N(D). \quad (12)$$

where for any extension  $\tilde{\varphi} \in H^1(D)$  of  $\varphi$  to  $D$ ,

$$\mathbf{f} = \mathbf{J}_s + \sigma(j\omega \mathbf{A} + \nabla \tilde{\varphi}) \quad (13)$$

belongs to  $L^2(D)^3$  and is divergence free,

2.  $\varphi$  belongs to  $\widetilde{H}^1(D_c)$  and is solution of the non-homogeneous Neumann problem

$$\int_{D_c} \sigma \nabla \varphi \cdot \nabla \overline{\varphi'} = -j\omega \int_{D_c} \sigma \mathbf{A} \cdot \nabla \overline{\varphi'}, \quad \forall \varphi' \in H^1(D_c). \quad (14)$$

**Proof.** The second property is a direct consequence of (10) by taking  $\mathbf{A}' = \mathbf{0}$  (noticing that the obtained identity remains valid for any  $\varphi' \in H^1(D_c)$ ). For the first property, we notice that (11) with  $\varphi' = 0$  yields

$$\int_D \mu^{-1} \operatorname{curl} \mathbf{A} \cdot \operatorname{curl} \overline{\mathbf{A}'} - \int_{D_c} \sigma(j\omega \mathbf{A} + \nabla \varphi) \cdot \overline{\mathbf{A}'} = (\mathbf{J}_s, \mathbf{A}'), \quad \forall \mathbf{A}' \in H_0(\operatorname{curl}, D).$$

As  $\mathbf{A}$  is divergence free, it clearly belongs to  $X_N(D)$ , and as  $X_N(D)$  is a subset of  $H_0(\operatorname{curl}, D)$ , the previous identity directly implies

$$\int_D (\mu^{-1} \operatorname{curl} \mathbf{A} \cdot \operatorname{curl} \overline{\mathbf{A}'} + \operatorname{div} \mathbf{A} \operatorname{div} \overline{\mathbf{A}'} - \int_{D_c} \sigma(j\omega \mathbf{A} + \nabla \varphi) \cdot \overline{\mathbf{A}'} = (\mathbf{J}_s, \mathbf{A}'), \quad \forall \mathbf{A}' \in X_N(D).$$

Since  $\sigma = 0$  on  $D \setminus D_c$ , the second term of this left-hand side can be written as

$$\int_{D_c} \sigma(j\omega \mathbf{A} + \nabla \varphi) \cdot \overline{\mathbf{A}'} = \int_D \sigma(j\omega \mathbf{A} + \nabla \varphi) \cdot \overline{\mathbf{A}'}.$$

These two identity directly lead to (12). The divergence free property of  $\mathbf{f}$  comes from the divergence free of  $\mathbf{J}_s$  and from (14). ■

According to this Lemma, the regularity of  $\mathbf{A}$  is related to the regularity of the Maxwell problem (12), while the regularity of  $\varphi$  is related to the regularity of the Neumann problem (14). Let us start with the regularity of  $\mathbf{A}$ .

**Lemma 2.2** *Let  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  be the unique solution of (10) with a divergence free field  $\mathbf{J}_s \in L^2(D)$ . Then the next results hold:*

1. *If  $D$  has a  $C^{1,1}$  boundary and  $\mu \in C^{0,1}(D)$ , then  $\mathbf{A} \in H^2(D)^3$ .*
2. *If  $D$  is a convex polyhedron and  $\mu = 1$ , then  $\mathbf{A} \in H^{1+\varepsilon}(D)^3$ , for some  $\varepsilon \in (0, 1]$  that depends on the interior angles along the edges of  $D$  and of the corner singularities of the Neumann problem in  $D$ .*
3. *If  $D$  is a parallelepiped and  $\mu = 1$ , then  $\mathbf{A} \in H^2(D)^3$ .*

**Proof.** Point 1 follows from the fact that the system associated with (12) is an elliptic system of order 2 (see [9, §4.5]), hence the  $H^2$  regularity of  $\mathbf{A}$  follows from a standard shift theorem as  $\mathbf{f}$  is in  $L^2(D)^3$ .

Points 2 and 3 follow from [6, §4.4.2 (b) and (c)]. ■

For  $\mu$  piecewise smooth and/or  $D$  a non convex polyhedron, some regularity results for (12) are also available, see for instance [7].

Let us go on with the regularity of  $\varphi$ .

**Lemma 2.3** *Let  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  be the unique solution of (10) with a divergence free field  $\mathbf{J}_s \in L^2(D)^3$ . Then the next results hold:*

1. *If  $\mathbf{A} \in H^2(D)^3$ ,  $D_c$  has a  $C^{2,1}$  boundary, and  $\sigma \in C^{1,1}(D_c)$ , then  $\varphi \in H^3(D_c)$ .*
2. *If  $\mathbf{A} \in H^1(D)^3$ ,  $D_c$  has a  $C^{1,1}$  boundary and  $\sigma \in C^{0,1}(D_c)$ , then  $\varphi \in H^2(D_c)$ .*
3. *If  $\mathbf{A} \in H^1(D)^3$ ,  $D_c$  is a convex polyhedron and  $\sigma = 1$ , then  $\varphi \in H^{1+s}(D_c)$ , for all  $s \in (0, 1)$ .*

**Proof.** As the strong formulation of (14) is (see (8b) and (9))

$$\begin{cases} \operatorname{div}(\sigma \nabla \varphi) &= -j\omega \operatorname{div}(\sigma \mathbf{A}) \text{ in } D_c \\ \sigma \nabla \varphi \cdot \mathbf{n} &= -j\omega \sigma \mathbf{A} \cdot \mathbf{n} \text{ on } \partial D_c, \end{cases} \quad (15)$$

both points follow from an appropriate shift theorem for the non-homogeneous Neumann problem in  $D_c$  and trace theorems.

For point 1, the regularities on  $\mathbf{A}$  and  $\sigma$  guarantee that  $\sigma \mathbf{A} \in H^2(D_c)^3$ , hence  $\operatorname{div}(\sigma \mathbf{A}) \in H^1(D_c)$ , and its trace  $\sigma \mathbf{A} \in H^{3/2}(\partial D_c)^3$ . From the regularity of the boundary of  $D_c$ , its normal trace  $\mathbf{n}$  belongs to  $C^{1,1}(\partial D_c)$  and therefore  $\mathbf{n} \in W^{2,p}(\partial D_c)$ , for all  $p > 1$ . By Theorem 1.4.4.2

from [18] (multiplication in Sobolev spaces), we deduce that  $\sigma \mathbf{A} \cdot \mathbf{n} \in H^{3/2}(\partial D_c)$ . By a standard shift theorem for (15), we deduce that  $\varphi \in H^3(D_c)$ .

Point 2 is obtained similarly, but here we only get  $\sigma \mathbf{A} \in H^1(D_c)^3$ , hence  $\operatorname{div}(\sigma \mathbf{A}) \in L^2(D_v)$ ,  $\sigma \mathbf{A} \in H^{1/2}(\partial D_c)^3$ . and  $\mathbf{n}$  belongs to  $C^{0,1}(\partial D_c)$  (and therefore  $\mathbf{n} \in W^{1,p}(\partial D_c)$ , for all  $p > 1$ ). Again by Theorem 1.4.4.2 from [18], we deduce that  $\sigma \mathbf{A} \cdot \mathbf{n} \in H^{1/2}(\partial D_c)$ . By a standard shift theorem for (15), we deduce that  $\varphi \in H^2(D_c)$ .

For the third point, the problem comes from the poor regularity of the normal trace along  $\partial D_c$ . In that case, we can only say that  $\mathbf{n}$  is piecewise smooth, and therefore deduce that  $\sigma \mathbf{A} \cdot \mathbf{n} \in H^{1/2}(F)$ , for all faces  $F$  of  $\partial D_c$ . Owing to Corollary 1.4.4.5 of [18], we obtain that  $\sigma \mathbf{A} \cdot \mathbf{n} \in H^{s-1/2}(\partial D_c)$ , for all  $s \in [0, 1)$ . The conclusion then follows from Theorem 23.3 of [14].

■

Finally a bootstrapping argument allows to obtain improved regularities for  $\mathbf{A}$ . For shortness, we restrict ourselves to some particular cases.

**Lemma 2.4** *Let  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  be the unique solution of (10) with a divergence free field  $\mathbf{J}_s \in H^{1/2}(D)^3$ . Assume that  $D_c$  has a  $C^{1,1}$  boundary and  $\sigma \in C^{0,1}(D_c)$  or  $D_c$  is a convex polyhedron and  $\sigma = 1$ . Then if  $D$  has a  $C^{2,1}$  boundary and  $\mu \in C^{1,1}(D)$  or  $D$  is a parallelepiped and  $\mu = 1$ ,  $\mathbf{A} \in H^{2+s}(D)^3$ , for all  $s \in (0, 1/2)$ .*

**Proof.** The assumptions on  $D$  and  $\mu$  allow to apply Lemma 2.2, and obtain  $\mathbf{A} \in H^2(D)^3$ . In a second step, due to our assumption on  $D_c$  and  $\sigma$ , we can apply Lemma 2.3 (point 2 or 3) and get  $\varphi \in H^{1+s}(D_c)$ , for all  $s \in (0, 1)$ . This regularity and the regularity of  $\mathbf{A}$  yield

$$\sigma(j\omega \mathbf{A} + \nabla \varphi) \in H^s(D_c), \forall s \in (0, 1).$$

And again by Corollary 1.4.4.5 of [18], we deduce that

$$\sigma(j\omega \mathbf{A} + \nabla \tilde{\varphi}) \in H^s(D), \forall s \in (0, 1/2).$$

Using the definition (13) of  $\mathbf{f}$  and the assumption on  $\mathbf{J}_s$ , we conclude that  $\mathbf{f}$  belongs to  $H^s(D)$ , for all  $s \in (0, 1/2)$ .

Coming back to problem (12), we find  $\mathbf{A} \in H^{2+s}(D)^3$ , for all  $s \in (0, 1/2)$ , by a standard shift theorem if  $D$  has a  $C^{2,1}$  boundary and  $\mu \in C^{1,1}(D)$  or by [6, §4.4.2 (c)] if  $D$  is a parallelepiped and  $\mu = 1$ . ■

### 3 A guaranteed equilibrated error estimator for non conforming approximations of the $\mathbf{A} - \varphi$ formulation

In this section, we obtain a guaranteed equilibrated error estimator for nonconforming approximations of (10), extending the results from [10, §3.2 and §4] to nonconforming and higher order approximations with general equilibrated fluxes.

More precisely, to discretize problems (10), we suppose given a partition  $\mathcal{T}$  of  $D$  into polyhedral elements  $T$  that covers exactly  $D$ . Each element  $T$  of  $\mathcal{T}$  is assumed to belong either to  $\bar{D}_c$  or to  $\overline{D_{nc}}$ , furthermore we denote by  $h_T$  its diameter. For simplicity, we set  $\mathcal{T}_c = \{T \in \mathcal{T} : T \subset \bar{D}_c\}$ . On such a mesh we introduce the so-called broken Sobolev space

$$\begin{aligned} H^1(\mathcal{T}) &= \{v \in L^2(D) \mid v|_T \in H^1(T), \forall T \in \mathcal{T}\}, \\ H^1(\mathcal{T}_c) &= \{v \in L^2(D_c) \mid v|_T \in H^1(T), \forall T \in \mathcal{T}_c\}. \end{aligned}$$

As in [15, 22, 11], in order to analyse simultaneously different approximation schemes, the problem is approximated in a finite dimensional subspace  $V_h$  of  $H^1(\mathcal{T})^3 \times H^1(\mathcal{T}_c)$ . In other words, we suppose given an approximation  $(\mathbf{A}_h, \varphi_h) \in V_h$  of the solution  $(\mathbf{A}, \varphi)$  of (10).

For the sake of simplicity, for  $\mathbf{A}'_h \in H^1(\mathcal{T})^3$  (resp.  $\varphi'_h \in H^1(\mathcal{T}_c)$ ), we denote by  $\text{curl}_h \mathbf{A}'_h$  (resp. its  $\nabla_h \varphi'_h$ ) its broken rotation (resp. its broken gradient), namely

$$\begin{aligned}\text{curl}_h \mathbf{A}'_h &= \text{curl} \mathbf{A}'_h && \text{on } T, \forall T \in \mathcal{T}, \\ \nabla_h \varphi'_h &= \nabla \varphi'_h && \text{on } T, \forall T \in \mathcal{T}_c.\end{aligned}$$

We now introduce the discrete counterparts of (7) by setting

$$\mathbf{B}_h = \text{curl}_h \mathbf{A}_h \text{ and } \mathbf{E}_h = -(j\omega \mathbf{A}_h + \nabla_h \varphi_h). \quad (16)$$

We further assume that some flux reconstructions  $\mathbf{H}_h$  and  $\mathbf{J}_{e,h}$  are available (using  $(\mathbf{A}_h, \varphi_h)$  and the datum  $\mathbf{J}_s$ ) that belong respectively to  $H(\text{curl}, D)$  and  $H(\text{div}, D_c)$  and satisfy the following conservation properties (compare with [15, identity (18)] and [10, Lemma 4.1])

$$(\text{curl} \mathbf{H}_h - \tilde{\mathbf{J}}_{e,h} - \mathbf{J}_s, \mathbf{e})_T = 0, \forall T \in \mathcal{T}, \mathbf{e} \in \mathbb{C}^3, \quad (17)$$

$$\text{div} \mathbf{J}_{e,h} = 0 \text{ in } D_c, \quad (18)$$

$$\mathbf{J}_{e,h} \cdot \mathbf{n} = 0 \text{ on } \partial D_c. \quad (19)$$

For further uses, we denote by  $\tilde{\mathbf{J}}_{e,h}$  the extension of  $\mathbf{J}_{e,h}$  by zero outside  $D_c$ , that remains divergence free.

Note that  $\mathbf{H}_h$  represents an approximation of  $\mu^{-1} \text{curl} \mathbf{A}$ , while  $\mathbf{J}_{e,h}$  is an approximation of  $\sigma \mathbf{E}$ .

Remark that (17) is a minimal assumption to guarantee that  $\mathbf{H}_h$  is a correct approximation of the continuous flux  $\mu^{-1} \text{curl} \mathbf{A}$ , while (18) and (19) is the exact counterpart of (3) and (4), therefore  $\mathbf{J}_{e,h}$  is a correct approximation of  $\mathbf{J}_e = \sigma \mathbf{E}$ .

For further uses, we also recall the following Poincaré inequality

$$\|u - \mathcal{M}_T u\|_T^2 \leq c_{P,T} h_T^2 |u|_{1,T}^2, \forall u \in H^1(T), T \in \mathcal{T}, \quad (20)$$

where  $\mathcal{M}_T u = |T|^{-1} \int_T u(x)$  is the mean of  $u$  on  $T$ , and  $c_{P,T}$  is a positive constant depending only on  $T$ . If  $T$  is convex, then  $c_{P,T} \leq \frac{1}{\pi^2}$  (hence for practical uses, in (20), we can replace  $c_{P,T}$  by  $\frac{1}{\pi^2}$ ), see [2, 25]. For nonconvex  $T$ , some estimations of  $c_{P,T}$  can be found in [3, Lemma 10.2] or [16, §2].

Besides the above estimate, we also need the following consequence of [8, Thm 3.4] (see also [4, p. 45]). Namely there exists a positive constant  $C_L$  such that for all  $\mathbf{A}' \in \tilde{X}(D)$ , there exist a unique function  $\Phi \in H_0^1(D)$  and a unique vector field  $\Psi \in H_0(\text{curl}, D) \cap H^1(D)^3$  such that

$$\mathbf{A}' = \nabla \Phi + \Psi, \quad (21)$$

with

$$|\Phi|_{1,D} + |\Psi|_{1,D} \leq C_L \|\text{curl} \mathbf{A}'\|_D. \quad (22)$$

Note that  $C_L = 1$  if  $D$  is convex, due to [6, Theorem 1.1] and [8, Theorem 2.1 and Lemma 2.2] since in that case  $\Phi = 0$  and  $\Psi = \mathbf{A}'$ .

Given a potential reconstruction  $(\mathbf{S}_h, \psi_h) \in H_0(\text{curl}, D) \times \widetilde{H}^1(D_c)$  of  $(\mathbf{A}_h, \varphi_h)$ , we now define the global estimator  $\eta$  defined as

$$\eta = 2\eta_{NC} + \eta_{\text{flux}} + \eta_{\mathcal{O}}, \quad (23)$$

where the nonconforming estimator  $\eta_{NC}$  is defined by

$$\begin{aligned}\eta_{NC} &= \left( \left\| \mu^{-1/2} \text{curl}_h(\mathbf{A}_h - \mathbf{S}_h) \right\|^2 \right. \\ &\quad \left. + \left\| \omega^{-1/2} \sigma^{1/2} (j\omega(\mathbf{A}_h - \mathbf{S}_h) + \nabla_h(\varphi_h - \psi_h)) \right\|_{D_c}^2 \right)^{1/2},\end{aligned} \quad (24)$$



the flux estimator  $\eta_{\text{flux}}$  is defined by (see [10, (15)])

$$\eta_{\text{flux}} = (\eta_{\text{magn}}^2 + \eta_{\text{elec}}^2)^{1/2}, \quad (25)$$

where  $\eta_{\text{magn}}$  and  $\eta_{\text{elec}}$  are defined by

$$\eta_{\text{magn}} = \left\| \mu^{1/2}(\mathbf{H}_h - \mu^{-1}\mathbf{B}_h) \right\|_D, \quad \eta_{\text{elec}} = \left\| (\omega\sigma)^{-1/2}(\mathbf{J}_{e,h} - \sigma\mathbf{E}_h) \right\|_{D_c}, \quad (26)$$

and finally the oscillation estimator  $\eta_{\mathcal{O}}$  is defined by

$$\eta_{\mathcal{O}} = C_L \mu^{\frac{1}{2}\max} \left( \sum_{T \in \mathcal{T}} c_{P,T} h_T^2 \|\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}\|_T^2 \right)^{\frac{1}{2}}, \quad (27)$$

where

$$\mu_{\max} = \sup_{x \in D} \mu(x).$$

We are ready to prove the following upper error bound (compare with [10, Theorem 4.2]) of the energy norm of the  $\mathbf{A} - \varphi$  error  $\epsilon_{A,\varphi}$ , given by:

$$\epsilon_{A,\varphi} = \left( \left\| \mu^{-1/2} \text{curl}_h \epsilon_A \right\|^2 + \left\| \omega^{-1/2} \sigma^{1/2} (j\omega\epsilon_A + \nabla_h \epsilon_\varphi) \right\|_{D_c}^2 \right)^{1/2}, \quad (28)$$

where

$$\epsilon_A = \mathbf{A} - \mathbf{A}_h \quad \text{and} \quad \epsilon_\varphi = \varphi - \varphi_h.$$

**Theorem 3.1** *Let us suppose that  $\mathbf{J}_s \in (L^2(D))^3$ , that  $\mathbf{H}_h \in H(\text{curl}, D)$ , and that  $\mathbf{J}_{e,h} \in H(\text{div}, D_c)$  satisfy (17)-(19). Then the following upper bound holds:*

$$\epsilon_{A,\varphi} \leq \eta. \quad (29)$$

**Proof.** Introduce

$$\epsilon_S = \mathbf{A} - \mathbf{S}_h \quad \text{and} \quad \epsilon_\psi = \varphi - \psi_h.$$

as well as

$$\begin{aligned} \epsilon_{S,\psi} &= |B((\epsilon_S, \epsilon_\psi), (\epsilon_S, \epsilon_\psi))|^{\frac{1}{2}} \\ &= \left( \left\| \mu^{-1/2} \text{curl} \epsilon_S \right\|^2 + \left\| \omega^{-1/2} \sigma^{1/2} (j\omega\epsilon_S + \nabla \epsilon_\psi) \right\|_{D_c}^2 \right)^{1/2}. \end{aligned} \quad (30)$$

By the triangular inequality, we directly deduce that

$$\epsilon_{A,\varphi} \leq \epsilon_{S,\psi} + \eta_{NC}. \quad (31)$$

Hence it remains to estimate  $\epsilon_{S,\psi}$ .

From the definition of  $B$ , we have

$$\begin{aligned} B((\epsilon_S, \epsilon_\psi), (\epsilon_S, \epsilon_\psi)) &= \int_D \mu^{-1} \text{curl}(\mathbf{A} - \mathbf{S}_h) \cdot \text{curl} \overline{\epsilon_S} \\ &\quad + \int_{D_c} \frac{j\sigma}{\omega} (j\omega(\mathbf{A} - \mathbf{S}_h) + \nabla(\varphi - \psi_h)) \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)}. \end{aligned}$$

By defining (compare with (16))

$$\mathbf{B}_h^S = \text{curl} \mathbf{S}_h \quad \text{and} \quad \mathbf{E}_h^S = -(j\omega \mathbf{S}_h + \nabla_h \psi_h), \quad (32)$$

and adding the quantities  $\pm \int_D \mathbf{H}_h \cdot \text{curl} \overline{\epsilon_S} \pm \frac{j}{\omega} \int_{D_c} \mathbf{J}_{e,h} \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)}$ , the above identity becomes

$$\begin{aligned} B((\epsilon_S, \epsilon_\psi), (\epsilon_S, \epsilon_\psi)) &= B((\mathbf{A}, \varphi), (\epsilon_S, \epsilon_\psi)) \\ &+ \int_D (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \cdot \text{curl} \overline{\epsilon_S} \\ &+ \frac{j}{\omega} \int_{D_c} (\sigma \mathbf{E}_h^S - \mathbf{J}_{e,h}) \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)} + \frac{j}{\omega} \int_{D_c} \mathbf{J}_{e,h} \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)} \\ &- \int_D \mathbf{H}_h \cdot \text{curl} \overline{\epsilon_S}. \end{aligned}$$

Using the weak formulation (10) valid for test-functions in  $H_0(\text{curl}, D) \times \widetilde{H}^1(D_c)$  see [12, Lemma 2.2]) and applying Green's formula to the last term of this right-hand side, we find term

$$\begin{aligned} B((\epsilon_S, \epsilon_\psi), (\epsilon_S, \epsilon_\psi)) &= \int_D \mathbf{J}_s \cdot \overline{\epsilon_S} - \int_D \text{curl} \mathbf{H}_h \cdot \overline{\epsilon_S} \\ &+ \int_D (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \cdot \text{curl} \overline{\epsilon_S} + \frac{j}{\omega} \int_{D_c} (\sigma \mathbf{E}_h^S - \mathbf{J}_{e,h}) \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)} \\ &+ \frac{j}{\omega} \int_{D_c} \mathbf{J}_{e,h} \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)}. \end{aligned}$$

Keeping unchanged the third and fourth terms of this right-hand side and rearranging the other terms, we find

$$\begin{aligned} B((\epsilon_S, \epsilon_\psi), (\epsilon_S, \epsilon_\psi)) &= \int_D (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \cdot \text{curl} \overline{\epsilon_S} \\ &+ \frac{j}{\omega} \int_{D_c} (\sigma \mathbf{E}_h^S - \mathbf{J}_{e,h}) \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)} \\ &+ \int_D (\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \overline{\epsilon_S} \\ &+ \frac{j}{\omega} \int_{D_c} \mathbf{J}_{e,h} \cdot \nabla \overline{\epsilon_\psi}. \end{aligned}$$

Using the assumptions (18)-(19), the last term vanishes, therefore by the identity (30), and the triangular inequality, we arrive at

$$\begin{aligned} \epsilon_{S,\psi}^2 &\leq \left| \int_D (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \cdot \text{curl} \overline{\epsilon_S} \right| \\ &+ \frac{1}{\omega} \left| \int_{D_c} (\sigma \mathbf{E}_h^S - \mathbf{J}_{e,h}) \cdot \overline{(j\omega\epsilon_S + \nabla\epsilon_\psi)} \right| \\ &+ \left| \int_D (\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \overline{\epsilon_S} \right|. \end{aligned} \tag{33}$$

Let us estimate each term of the right hand-side of this estimate. But we first transform (and then estimate) the third term of this right-hand side. For that purpose, we use the Helmholtz decomposition of [23, Lemma 4.5]

$$H_0(\text{curl}, D) = \nabla H_0^1(D) \oplus \widetilde{X}(D), \tag{34}$$

so that

$$\epsilon_S = \nabla \phi + \epsilon_\perp, \tag{35}$$

with  $\phi \in H_0^1(D)$  and  $\epsilon_\perp \in \widetilde{X}(D)$  and

$$\|\epsilon_S\|_D^2 = \|\nabla \phi\|_D^2 + \|\epsilon_\perp\|_D^2. \tag{36}$$

This decomposition, Green's formula and the divergence free property of  $\tilde{\mathbf{J}}_{e,h}$  allow to get

$$\int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\epsilon}_S = \int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\epsilon}_\perp. \quad (37)$$

Now as  $\epsilon_\perp \in \tilde{X}(D)$ , there exist a unique function  $\Phi \in H_0^1(D)$  and a unique vector field  $\Psi \in H_0(\text{curl}, D) \cap H^1(D)^3$  such that (see (21) and (22))

$$\epsilon_\perp = \nabla\Phi + \Psi, \quad (38)$$

with

$$|\Phi|_{1,D} + |\Psi|_{1,D} \leq C_L \|\text{curl}\epsilon_\perp\|.$$

Furthermore since  $\text{curl}\epsilon_S = \text{curl}\epsilon_\perp$ , the previous estimate implies

$$\begin{aligned} |\Phi|_{1,D} + |\Psi|_{1,D} &\leq C_L \|\text{curl}\epsilon_S\| \\ &\leq C_L \mu_{\max}^{\frac{1}{2}} \|\mu^{-1/2} \text{curl}\epsilon_S\|. \end{aligned}$$

Therefore by the definition of  $\epsilon_{S,\psi}$ , we get

$$|\Phi|_{1,D} + |\Psi|_{1,D} \leq C_L \mu_{\max}^{\frac{1}{2}} \epsilon_{S,\psi}. \quad (39)$$

Using (38) into (37) and using Green's formula and the divergence free property of  $\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}$ , we obtain

$$\int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\epsilon}_S = \int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\Psi}.$$

By the property (17), we obtain

$$\int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\epsilon}_S = \int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot (\bar{\Phi} - \mathcal{M}_h \bar{\Psi}),$$

where  $\mathcal{M}_h \bar{\Psi} \in H^1(\mathcal{T})^3$  is defined by

$$(\mathcal{M}_h \bar{\Psi})|_T = \mathcal{M}_T(\bar{\Psi}|_T), \forall T \in \mathcal{T}.$$

By (continuous and discrete) Cauchy-Schwarz's inequality and Poincaré's inequality (20), we deduce that

$$\int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\epsilon}_S \leq \left( \sum_{T \in \mathcal{T}} c_{P,T} h_T^2 \|\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}\|_T^2 \right)^{\frac{1}{2}} |\Psi|_{1,D}.$$

Using the estimate (39), we arrive at

$$\left| \int_D (\mathbf{J}_s - \text{curl}\mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}) \cdot \bar{\epsilon}_S \right| \leq \eta_{\mathcal{O}} \epsilon_{S,\psi}. \quad (40)$$

A simple consequence of (continuous and discrete) Cauchy-Schwarz's inequality allows to estimate the first two terms of the right-hand side of (33) as follows

$$\begin{aligned} &\left| \int_D (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \cdot \text{curl}\bar{\epsilon}_S \right| + \frac{1}{\omega} \left| \int_{D_c} (\sigma \mathbf{E}_h^S - \mathbf{J}_{e,h}) \cdot (\overline{j\omega\epsilon_S + \nabla\epsilon_\psi}) \right| \\ &\leq \left\| \mu^{1/2} (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \right\|_D \left\| \mu^{-1/2} \text{curl}\epsilon_S \right\|_D \\ &\quad + \left\| (\omega\sigma)^{-1/2} (\mathbf{J}_{e,h} - \sigma \mathbf{E}_h^S) \right\|_{D_c} \left\| \sigma^{1/2} \omega^{-1/2} (j\omega\epsilon_S + \nabla\epsilon_\psi) \right\|_{D_c} \\ &\leq \left( \left\| \mu^{1/2} (\mathbf{H}_h - \mu^{-1} \mathbf{B}_h^S) \right\|_D^2 + \left\| (\omega\sigma)^{-1/2} (\mathbf{J}_{e,h} - \sigma \mathbf{E}_h^S) \right\|_{D_c}^2 \right)^{\frac{1}{2}} \epsilon_{S,\psi}. \end{aligned} \quad (41)$$

Let us now transform the first factor of this right-hand side in order to display the estimator  $\eta_{flux}$ . Indeed inserting  $\pm \mathbf{E}_h$  and  $\pm \mathbf{B}_h$  and using the discrete Cauchy-Schwarz's inequality, and using the definition (25) of  $\eta_{flux}$ , we have

$$\begin{aligned} & \left( \left\| \mu^{1/2}(\mathbf{H}_h - \mu^{-1}\mathbf{B}_h^S) \right\|_D^2 + \left\| (\omega\sigma)^{-1/2}(\mathbf{J}_{e,h} - \sigma\mathbf{E}_h^S) \right\|_{D_c}^2 \right)^{\frac{1}{2}} \\ & \leq \eta_{flux} \\ & + \left( \left\| \mu^{-1/2}(\mathbf{B}_h - \mathbf{B}_h^S) \right\|_D^2 + \left\| \omega^{-1/2}\sigma^{1/2}(\mathbf{E}_h - \mathbf{E}_h^S) \right\|_{D_c}^2 \right)^{\frac{1}{2}}, \end{aligned}$$

The definition of  $\mathbf{E}_h$ ,  $\mathbf{E}_h^S$ ,  $\mathbf{B}_h$  and  $\mathbf{B}_h^S$  directly give

$$\mathbf{B}_h - \mathbf{B}_h^S = \text{curl}_h(\mathbf{A}_h - \mathbf{S}_h), \text{ and } \mathbf{E}_h - \mathbf{E}_h^S = j\omega(\mathbf{A}_h - \mathbf{S}_h) + \nabla(\varphi_h - \psi_h),$$

which leads to

$$\left( \left\| \mu^{-1/2}(\mathbf{B}_h - \mathbf{B}_h^S) \right\|_D^2 + \left\| \omega^{-1/2}\sigma^{1/2}(\mathbf{E}_h - \mathbf{E}_h^S) \right\|_{D_c}^2 \right)^{\frac{1}{2}} = \eta_{NC}.$$

Therefore we have found that

$$\left( \left\| \mu^{1/2}(\mathbf{H}_h - \mu^{-1}\mathbf{B}_h^S) \right\|_D^2 + \left\| (\omega\sigma)^{-1/2}(\mathbf{J}_{e,h} - \sigma\mathbf{E}_h^S) \right\|_{D_c}^2 \right)^{\frac{1}{2}} \leq \eta_{flux} + \eta_{NC}.$$

This estimate in (41) leads to

$$\begin{aligned} & \left| \int_D (\mathbf{H}_h - \mu^{-1}\mathbf{B}_h^S) \cdot \text{curl} \bar{\epsilon}_S \right| + \frac{1}{\omega} \left| \int_{D_c} (\sigma\mathbf{E}_h^S - \mathbf{J}_{e,h}) \cdot (j\omega\epsilon_S + \nabla\epsilon_\psi) \right| \\ & \leq (\eta_{flux} + \eta_{NC})\epsilon_{S,\psi}. \end{aligned}$$

Using this last estimate and (40) in (33), we arrive at

$$\epsilon_{S,\psi} \leq \eta_{flux} + \eta_{NC} + \eta_{\mathcal{O}}.$$

With the help of (31), we conclude that (29) holds (using the definition (23) of  $\eta$ ). ■

## 4 The goal oriented functional and the adjoint problem

We here consider the output functional that represents the physical quantity of interest given by

$$Q(\mathbf{A}) = \int_D \mathbf{q} \cdot \text{curl} \bar{\mathbf{A}} \, dx, \forall \mathbf{A} \in H(\text{curl}, D), \quad (42)$$

where  $\mathbf{q} \in L^2(D)^3$  is a given function.

**Remark 4.1** *In many engineering applications, engineers are interested in the computation of the flux through a coil where the divergence free current  $\mathbf{J}_s$  is imposed. In that case, it is equivalent to replace the function  $\mathbf{q}$  by  $\mathbf{H}_s$  with  $\text{curl} \mathbf{H}_s = \mathbf{J}_s$ , see [20, identity (7)]. It corresponds to the numerical test proposed in subsections 6.1 and 6.2. In other applications, engineers can be interested in the computation of the magnetic flux density at a given point of an electromagnetic device, see [17, 21]. It corresponds to the numerical test proposed in subsection 6.3.*

Accordingly by setting

$$\begin{aligned} B^*((\mathbf{A}, \varphi), (\mathbf{A}', \varphi')) &= \overline{B((\mathbf{A}', \varphi'), (\mathbf{A}, \varphi))} \\ &= (\mu^{-1} \operatorname{curl} \mathbf{A}, \operatorname{curl} \mathbf{A}')_D \\ &\quad - j\omega^{-1} (\sigma(j\omega \mathbf{A} + \nabla \varphi), (j\omega \mathbf{A}' + \nabla \varphi'))_{D_c}, \quad \forall (\mathbf{A}, \varphi), (\mathbf{A}', \varphi') \in \tilde{X}(D) \times \widetilde{H}^1(D_c), \end{aligned}$$

the associated dual problem consists in looking for  $(\mathbf{A}^*, \varphi^*) \in \tilde{X}(D) \times \widetilde{H}^1(D_c)$  solution of the adjoint problem

$$B^*((\mathbf{A}^*, \varphi^*), (\mathbf{A}', \varphi')) = Q(\mathbf{A}'), \quad \forall (\mathbf{A}', \varphi') \in \tilde{X}(D) \times \widetilde{H}^1(D_c), \quad (43)$$

that also has a unique solution. Note that the Helmholtz decomposition (34) implies that (43) remains valid for any  $(\mathbf{A}', \varphi') \in H_0(\operatorname{curl}, D) \times \widetilde{H}^1(D_c)$ . The strong formulation of (43) is clearly

$$\begin{aligned} \operatorname{curl}(\mu^{-1} \operatorname{curl} \mathbf{A}^*) - \sigma(j\omega \mathbf{A}^* + \nabla \varphi^*) &= \operatorname{curl} \mathbf{q} \text{ in } D, \\ \operatorname{div}(\sigma(j\omega \mathbf{A}^* + \nabla \varphi^*)) &= 0 \text{ in } D_c. \end{aligned}$$

Let us notice that Lemmas 2.2, 2.3 and 2.4 remain valid for the adjoint problem.

Similarly to the primal problem (see the beginning of section 3), the dual one is approximated in a finite dimensional subspace  $V_h^*$  of  $H^1(\mathcal{T})^3 \times H^1(\mathcal{T}_c)$ , that may be different from  $V_h$ . In other words, we suppose given an approximation  $(\mathbf{A}_h, \varphi_h) \in V_h$  of the solution  $(\mathbf{A}, \varphi)$  of (10) and  $(\mathbf{A}_h^*, \varphi_h^*) \in V_h^*$  of the solution  $(\mathbf{A}^*, \varphi^*)$  of (43). As already specified, we assume that some flux reconstructions  $\mathbf{H}_h$  and  $\mathbf{J}_{e,h}$  are available (using  $(\mathbf{A}_h, \varphi_h)$  and the datum  $\mathbf{J}_s$ ) that belong respectively to  $H(\operatorname{curl}, D)$  and  $H(\operatorname{div}, D_c)$  and satisfy the conservation properties (17)-(19). In the same manner, by assuming that  $\mathbf{q} \in H(\operatorname{curl}, D)$ , we suppose that some flux reconstructions  $\mathbf{H}_h^*$  and  $\mathbf{J}_{e,h}^*$  are available (using  $(\mathbf{A}_h^*, \varphi_h^*)$  and the datum  $\operatorname{curl} \mathbf{q}$ ) that belongs respectively to  $H(\operatorname{curl}, D)$  and  $H(\operatorname{div}, D_c)$  satisfy the following conservation properties:

$$(\operatorname{curl} \mathbf{H}_h^* + \tilde{\mathbf{J}}_{e,h}^* - \operatorname{curl} \mathbf{q}, \mathbf{e})_T = 0, \quad \forall T \in \mathcal{T}, \mathbf{e} \in \mathbb{C}^3, \quad (44)$$

$$\operatorname{div} \mathbf{J}_{e,h}^* = 0 \text{ in } D_c, \quad (45)$$

$$\mathbf{J}_{e,h}^* \cdot \mathbf{n} = 0 \text{ on } \partial D_c. \quad (46)$$

Recall that  $\mathbf{H}_h$  represents an approximation of  $\mu^{-1} \operatorname{curl} \mathbf{A}$ ,  $\mathbf{J}_{e,h}$  an approximation of  $\sigma \mathbf{E}$ . Similarly, note that  $\mathbf{H}_h^*$  represents an approximation of  $\mu^{-1} \operatorname{curl} \mathbf{A}^*$  and  $\mathbf{J}_{e,h}^*$  an approximation of  $\sigma \mathbf{E}^* = -\sigma(j\omega \mathbf{A}^* + \nabla \varphi^*)$ .

Let us now state a guaranteed error estimate for the dual problem. For that purpose, we suppose given a potential reconstruction  $(\mathbf{S}_h^*, \psi_h^*) \in H_0(\operatorname{curl}, D) \times \widetilde{H}^1(D_c)$  of  $(\mathbf{A}_h^*, \varphi_h^*)$ . Then we define the global estimator  $\eta^*$  as follows

$$\eta^* = 2\eta_{NC}^* + \eta_{\text{flux}}^* + \eta_{\mathcal{O}}^*, \quad (47)$$

where the nonconforming estimator  $\eta_{NC}^*$  is defined by

$$\begin{aligned} \eta_{NC}^* &= \left( \left\| \mu^{-1/2} \operatorname{curl}_h (\mathbf{A}_h^* - \mathbf{S}_h^*) \right\|^2 \right. \\ &\quad \left. + \left\| \omega^{-1/2} \sigma^{1/2} (j\omega (\mathbf{A}_h^* - \mathbf{S}_h^*) + \nabla_h (\varphi_h^* - \psi_h^*)) \right\|_{D_c}^2 \right)^{1/2}, \end{aligned} \quad (48)$$

the flux estimator  $\eta_{\text{flux}}^*$  is defined by

$$\eta_{\text{flux}}^* = ((\eta_{\text{magn}}^*)^2 + (\eta_{\text{elec}}^*)^2)^{1/2}, \quad (49)$$

where  $\eta_{\text{magn}}^*$  and  $\eta_{\text{elec}}^*$  are defined by

$$\eta_{\text{magn}}^* = \left\| \mu^{1/2}(\mathbf{H}_h^* - \mu^{-1}\mathbf{B}_h^*) \right\|_D, \quad \eta_{\text{elec}}^* = \left\| (\omega\sigma)^{-1/2}(\mathbf{J}_{e,h}^* - \sigma\mathbf{E}_h^*) \right\|_{D_c}, \quad (50)$$

and finally the oscillation estimator  $\eta_{\mathcal{O}}^*$  is defined by

$$\eta_{\mathcal{O}}^* = C_L \mu^{\frac{1}{2} \max} \left( \sum_{T \in \mathcal{T}} c_{P,T} h_T^2 \|\text{curl } \mathbf{H}_h^* + \tilde{\mathbf{J}}_{e,h}^* - \text{curl } \mathbf{q}\|_T^2 \right)^{\frac{1}{2}}. \quad (51)$$

The same arguments that the ones used in the proof of Theorem 3.1 allow to prove the following guaranteed error estimate on the error

$$\epsilon_{A^*, \varphi^*} = \left( \left\| \mu^{-1/2} \text{curl}_h \epsilon_{A^*} \right\|^2 + \left\| \omega^{-1/2} \sigma^{1/2} (j\omega \epsilon_{A^*} + \nabla_h \epsilon_{\varphi^*}) \right\|_{D_c}^2 \right)^{1/2}, \quad (52)$$

where

$$\epsilon_{A^*} = \mathbf{A}^* - \mathbf{A}_h^* \quad \text{and} \quad \epsilon_{\varphi^*} = \varphi^* - \varphi_h^*.$$

**Corollary 4.2** *Under the assumptions from this section, we have*

$$\epsilon_{A^*, \varphi^*} \leq \eta^*. \quad (53)$$

## 5 The error representation

With the help of the equilibrated fluxes of the primal and dual problems, in the spirit of [24] (see also [22, 11]), we here show that the error on the quantity of interest can be expressed into a fully computable expression and a remainder that will be estimated by a fully computable quantity (but is usually of higher order and can then be disregarded).

**Theorem 5.1** *Let  $(\mathbf{S}_h, \psi_h) \in H_0(\text{curl}, D) \times \widetilde{H}^1(D_c)$  be a potential reconstruction of  $(\mathbf{A}_h, \varphi_h)$ , then the error on the quantity of interest defined by*

$$\mathcal{E} = \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \text{curl} \overline{(\mathbf{A} - \mathbf{A}_h)} dx,$$

admits the splitting

$$\mathcal{E} = \eta_{\text{QOI}} + \mathcal{R}, \quad (54)$$

where the estimator  $\eta_{\text{QOI}}$  is given by

$$\begin{aligned} \eta_{\text{QOI}} &= \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \text{curl} \overline{(\mathbf{S}_h - \mathbf{A}_h)} dx \\ &+ \int_D \mathbf{S}_h^* \cdot \overline{(\mathbf{J}_s - \text{curl } \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx \\ &- j\omega^{-1} \int_{D_c} \sigma^{-1} \mathbf{J}_{e,h}^* \cdot \overline{(\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})} dx \\ &- \int_D \mathbf{H}_h^* \cdot (\text{curl } \overline{\mathbf{S}_h} - \mu \overline{\mathbf{H}_h}) dx \end{aligned} \quad (55)$$

while the remainder term  $\mathcal{R}$  is defined by

$$\begin{aligned} \mathcal{R} &= \int_D (\mathbf{A}^* - \mathbf{S}_h^*) \cdot \overline{(\mathbf{J}_s - \text{curl } \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx \\ &+ j\omega^{-1} \int_{D_c} (\sigma^{-1} \mathbf{J}_{e,h}^* - \mathbf{E}^*) \cdot \overline{(\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})} dx \\ &- \int_D (\mu^{-1} \text{curl } \mathbf{A}^* - \mathbf{H}_h^*) \cdot (\text{curl } \overline{\mathbf{S}_h} - \mu \overline{\mathbf{H}_h}) dx \end{aligned} \quad (56)$$

**Proof.** First we notice that

$$(\mathbf{J}_s, \mathbf{A}^*)_D = B((\mathbf{A}, \varphi), (\mathbf{A}^*, \varphi^*)) = \overline{B^*((\mathbf{A}^*, \varphi^*), (\mathbf{A}, \varphi))} = \overline{Q(\mathbf{A})}.$$

Hence by the definition of the error, we have

$$\mathcal{E} = \overline{(\mathbf{J}_s, \mathbf{A}^*)_D} - \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \text{curl } \overline{\mathbf{A}_h} dx.$$

Introducing artificially  $\mathbf{S}_h$  and using (43) with  $(\mathbf{A}', \varphi') = (\mathbf{S}_h, \psi_h)$ , this is equivalent to

$$\begin{aligned} \mathcal{E} &= \overline{(\mathbf{J}_s, \mathbf{A}^*)_D} - \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \text{curl } \overline{(\mathbf{A}_h - \mathbf{S}_h)} dx \\ &\quad - (\mu^{-1} \text{curl } \mathbf{A}^*, \text{curl } \mathbf{S}_h)_D \\ &\quad + j\omega^{-1} (\sigma(j\omega \mathbf{A}^* + \nabla \varphi^*), (j\omega \mathbf{S}_h + \nabla \psi_h))_{D_c}. \end{aligned} \tag{57}$$

Using (16), and adding and subtracting the terms

$$\int_D \text{curl } \mathbf{A}^* \cdot \overline{\mathbf{H}_h} dx, \text{ and } j\omega^{-1} \int_{D_c} \nabla \varphi^* \cdot \overline{\mathbf{J}_{e,h}} dx,$$

we find

$$\begin{aligned} \mathcal{E} &= \overline{(\mathbf{J}_s, \mathbf{A}^*)_D} - \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \text{curl } \overline{(\mathbf{A}_h - \mathbf{S}_h)} dx \\ &\quad - \int_D \text{curl } \mathbf{A}^* \cdot (\mu^{-1} \text{curl } \overline{\mathbf{S}_h} - \overline{\mathbf{H}_h}) dx \\ &\quad - \int_D \text{curl } \mathbf{A}^* \cdot \overline{\mathbf{H}_h} dx \\ &\quad - \int_{D_c} \sigma \mathbf{A}^* \cdot \overline{(j\omega \mathbf{S}_h + \nabla \psi_h)} dx \\ &\quad + j\omega^{-1} \int_{D_c} \nabla \varphi^* \cdot \overline{(\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})} dx \\ &\quad - j\omega^{-1} \int_{D_c} \nabla \varphi^* \cdot \overline{\mathbf{J}_{e,h}} dx. \end{aligned}$$

Using Green's formula in the fourth and seventh terms of this right-hand side and using (18)-(19), we find

$$\begin{aligned} \mathcal{E} &= \overline{(\mathbf{J}_s, \mathbf{A}^*)_D} - \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \text{curl } \overline{(\mathbf{A}_h - \mathbf{S}_h)} dx \\ &\quad - \int_D \text{curl } \mathbf{A}^* \cdot (\mu^{-1} \text{curl } \overline{\mathbf{S}_h} - \overline{\mathbf{H}_h}) dx \\ &\quad - \int_D \mathbf{A}^* \cdot \text{curl } \overline{\mathbf{H}_h} dx \\ &\quad - \int_{D_c} \sigma \mathbf{A}^* \cdot \overline{(j\omega \mathbf{S}_h + \nabla \psi_h)} dx \\ &\quad + j\omega^{-1} \int_{D_c} \nabla \varphi^* \cdot \overline{(\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})} dx. \end{aligned}$$

Adding and subtracting the term  $\int_{D_c} \mathbf{A}^* \cdot \overline{\mathbf{J}_{e,h}} dx$  and rearranging the terms of this right-hand side, we find equivalently

$$\begin{aligned} \mathcal{E} &= - \sum_{T \in \mathcal{T}} \int_T \mathbf{q} \cdot \overline{\text{curl}(\mathbf{A}_h - \mathbf{S}_h)} dx \\ &+ \int_D \mathbf{A}^* \cdot \overline{(\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx \\ &- j\omega^{-1} \int_{D_c} \mathbf{E}^* \cdot \overline{(\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})} dx \\ &- \int_D \text{curl} \mathbf{A}^* \cdot (\mu^{-1} \text{curl} \overline{\mathbf{S}_h} - \overline{\mathbf{H}_h}) dx, \end{aligned}$$

where we recall that  $\tilde{\mathbf{J}}_{e,h}$  means the extension by zero of  $\mathbf{J}_{e,h}$  outside  $D_c$  and that  $\mathbf{E}^* = -(j\omega \mathbf{A}^* + \nabla \varphi^*)$ .

Writing  $\mathbf{A}^* = \mathbf{S}_h^* + \mathbf{A}^* - \mathbf{S}_h^*$ ,  $\mathbf{E}^* = \sigma^{-1} \mathbf{J}_{e,h}^* + \mathbf{E}^* - \sigma^{-1} \mathbf{J}_{e,h}^*$  and  $\mu^{-1} \text{curl} \mathbf{A}^* = \mathbf{H}_h^* + \mu^{-1} \text{curl} \mathbf{A}^* - \mathbf{H}_h^*$ , we arrive at (54). ■

Let us now show that the remainder can be explicitly estimated using the error estimators for  $(\mathbf{A}, \varphi)$  and  $(\mathbf{A}^*, \varphi^*)$  obtained in section 3.

**Theorem 5.2** *With  $\eta$  (resp.  $\eta^*$ ) defined before, we have*

$$|\mathcal{R}| \leq 10\eta\eta^*. \quad (58)$$

**Proof.** We estimate each term of  $\mathcal{R}$  separately. For the first term,

$$\mathcal{R}_1 = \int_D (\mathbf{A}^* - \mathbf{S}_h^*) \cdot \overline{(\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx,$$

as in the proof of Theorem 3.1 we use the Helmholtz decomposition

$$\mathbf{A}^* - \mathbf{S}_h^* = \nabla \phi^* + \epsilon_{\perp}^*, \quad (59)$$

with  $\phi^* \in H_0^1(D)$  and  $\epsilon_{\perp}^* \in \tilde{X}(D)$  and

$$\|\mathbf{A}^* - \mathbf{S}_h^*\|_D^2 = \|\nabla \phi^*\|_D^2 + \|\epsilon_{\perp}^*\|_D^2. \quad (60)$$

As  $\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}$  is divergence free, we then get

$$\mathcal{R}_1 = \int_D \epsilon_{\perp}^* \cdot \overline{(\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx.$$

Again as in the proof of Theorem 3.1, as  $\epsilon_{\perp}^* \in \tilde{X}(D)$ , there exist a unique function  $\Phi^* \in H_0^1(D)$  and a unique vector field  $\Psi^* \in H_0(\text{curl}, D) \cap H^1(D)^3$  such that (see (21) and (22))

$$\epsilon_{\perp}^* = \nabla \Phi^* + \Psi^*,$$

with

$$|\Phi^*|_{1,D} + |\Psi^*|_{1,D} \leq C_L \|\text{curl} \epsilon_{\perp}^*\|_D \leq C_L \mu_{\max}^{\frac{1}{2}} \|\mu^{-1/2} \text{curl}(\mathbf{A}^* - \mathbf{S}_h^*)\|, \quad (61)$$

since  $\text{curl}(\mathbf{A}^* - \mathbf{S}_h^*) = \text{curl} \epsilon_{\perp}^*$ . As  $\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h}$  is divergence free and using the property (17), we then get

$$\begin{aligned} \mathcal{R}_1 &= \int_D \Psi^* \cdot \overline{(\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx \\ &= \int_D (\Psi^* - \mathcal{M}_h \Psi^*) \cdot \overline{(\mathbf{J}_s - \text{curl} \mathbf{H}_h + \tilde{\mathbf{J}}_{e,h})} dx. \end{aligned}$$



Cauchy-Schwarz's inequality and Poincaré's inequality (20) yield

$$|\mathcal{R}_1| \leq \eta_{\mathcal{O}} \|\mu^{-1/2} \operatorname{curl}(\mathbf{A}^* - \mathbf{S}_h^*)\|, \quad (62)$$

where we have used the estimate (61) and (27). Coming back to  $\mathbf{A}^* - \mathbf{A}_h^*$ , we get

$$\begin{aligned} |\mathcal{R}_1| &\leq \eta_{\mathcal{O}} (\|\mu^{-1/2} \operatorname{curl}_h(\mathbf{A}^* - \mathbf{A}_h^*)\| + \|\mu^{-1/2} \operatorname{curl}_h(\mathbf{A}_h^* - \mathbf{S}_h^*)\|) \\ &\leq \eta_{\mathcal{O}} (\epsilon_{A^*, \varphi^*} + \eta_{NC}^*), \end{aligned}$$

where we recall that  $\epsilon_{A^*, \varphi^*}$  is defined by (52). Owing to the estimate (53), we get finally

$$|\mathcal{R}_1| \leq 2 \eta_{\mathcal{O}} \eta^*. \quad (63)$$

For the second term

$$\mathcal{R}_2 = j\omega^{-1} \int_{D_c} (\sigma^{-1} \mathbf{J}_{e,h}^* - \mathbf{E}^*) \cdot \overline{(\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})} dx,$$

we use Cauchy-Schwarz's inequality, to get

$$|\mathcal{R}_2| \leq \|(\omega\sigma)^{-1/2} (\mathbf{J}_{e,h}^* - \sigma \mathbf{E}^*)\|_{D_c} \|(\omega\sigma)^{-1/2} (\sigma(j\omega \mathbf{S}_h + \nabla \psi_h) + \mathbf{J}_{e,h})\|_{D_c}.$$

Adding  $\pm(\omega\sigma)^{-1/2} \mathbf{E}_h^*$  and  $\pm(\omega\sigma)^{-1/2} \mathbf{E}_h$ , and using the triangular inequality, we get

$$\begin{aligned} |\mathcal{R}_2| &\leq \left( \|(\omega\sigma)^{-1/2} (\mathbf{J}_{e,h}^* - \sigma \mathbf{E}_h^*)\|_{D_c} + \|\omega^{-1/2} \sigma^{1/2} (\mathbf{E}_h^* - \mathbf{E}^*)\|_{D_c} \right) \times \\ &\quad \left( \|(\omega\sigma)^{-1/2} (-\sigma \mathbf{E}_h + \mathbf{J}_{e,h})\|_{D_c} + \|\omega^{-1/2} \sigma^{1/2} (j\omega \mathbf{S}_h + \nabla \psi_h + \mathbf{E}_h)\|_{D_c} \right). \end{aligned}$$

Reminding (7), (16), and the definitions of  $\eta_{\text{flux}}$ ,  $\eta_{NC}$ ,  $\epsilon_{A, \varphi}$ , and  $\epsilon_{A^*, \varphi^*}$ , we find

$$|\mathcal{R}_2| \leq (\eta_{\text{flux}}^* + \epsilon_{A^*, \varphi^*}) (\eta_{\text{flux}} + \eta_{NC} + \epsilon_{A, \varphi}).$$

By (29) and (53), we get finally

$$|\mathcal{R}_2| \leq (\eta_{\text{flux}} + \eta) (\eta_{\text{flux}}^* + \eta^*) \leq 4\eta\eta^* \quad (64)$$

Again for the third term

$$\mathcal{R}_3 = - \int_D (\mu^{-1} \operatorname{curl} \mathbf{A}^* - \mathbf{H}_h) \cdot (\operatorname{curl} \overline{\mathbf{S}_h} - \mu \overline{\mathbf{H}_h}) dx,$$

we use Cauchy-Schwarz's inequality to obtain

$$|\mathcal{R}_3| \leq \|\mu^{1/2} (\mu^{-1} \operatorname{curl} \mathbf{A}^* - \mathbf{H}_h)\|_D \|\mu^{1/2} (\mu^{-1} \operatorname{curl} \mathbf{S}_h - \mathbf{H}_h)\|_D.$$

Introducing  $\operatorname{curl}_h \mathbf{A}_h^*$  and  $\operatorname{curl}_h \mathbf{A}_h$ , the triangular inequality yields

$$\begin{aligned} |\mathcal{R}_3| &\leq \left( \|\mu^{1/2} (\mu^{-1} \operatorname{curl} \mathbf{A}_h^* - \mathbf{H}_h)\|_D + \|\mu^{-1/2} \operatorname{curl}_h(\mathbf{A}^* - \mathbf{A}_h^*)\|_D \right) \times \\ &\quad \left( \|\mu^{1/2} (\mu^{-1} \operatorname{curl} \mathbf{A}_h - \mathbf{H}_h)\|_D + \|\mu^{-1/2} \operatorname{curl}_h(\mathbf{S}_h - \mathbf{A}_h)\|_D \right). \end{aligned}$$

As before, this means that

$$|\mathcal{R}_3| \leq (\eta_{\text{flux}} + \eta_{NC} + \epsilon_{A, \varphi}) (\eta_{\text{flux}}^* + \eta_{NC}^*) \leq 4\eta\eta^*. \quad (65)$$

The estimates (63) to (65) lead to (58). ■

Before going on, if  $(\mathbf{A}_h, \varphi_h)$  is a conforming approximation of  $(\mathbf{A}, \varphi)$ , let us give an estimate on the error of quantity of interest that will be used for our numerical examples but has also its own interest.

**Lemma 5.3** Let  $\mathbf{V}_h$  be a finite dimensional subspace of  $\tilde{X}(D) \times \tilde{H}^1(D_c)$ . Let  $(\mathbf{A}_h, \varphi_h) \in \mathbf{V}_h$  be the Galerkin approximation of the solution  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  of (10), namely the unique solution to

$$B((\mathbf{A}_h, \varphi_h), (\mathbf{A}'_h, \varphi'_h)) = (\mathbf{J}_s, \mathbf{A}'_h), \quad \forall (\mathbf{A}'_h, \varphi'_h) \in \mathbf{V}_h. \quad (66)$$

Then the error on the quantity of interest is equal to

$$\mathcal{E} = B^*((\mathbf{A}^* - \mathbf{A}'_h, \varphi^* - \varphi'_h), (\mathbf{A} - \mathbf{A}_h, \varphi - \varphi_h)), \quad \forall (\mathbf{A}'_h, \varphi'_h) \in \mathbf{V}_h. \quad (67)$$

**Proof.** By its definition and our assumption on  $\mathbf{V}_h$ , we have

$$\mathcal{E} = \int_D q \cdot \text{curl}(\mathbf{A} - \mathbf{A}_h).$$

Therefore by the definition of the adjoint problem, we have

$$\mathcal{E} = B^*((\mathbf{A}^*, \varphi^*), (\mathbf{A} - \mathbf{A}_h, \varphi - \varphi_h)). \quad (68)$$

As Lemma 2.3 of [12] proved that

$$B((\mathbf{A} - \mathbf{A}_h, \varphi - \varphi_h), (\mathbf{A}'_h, \varphi'_h)) = 0, \quad \forall (\mathbf{A}'_h, \varphi'_h) \in \mathbf{V}_h,$$

by the definition of  $B^*$ , we get equivalently

$$B^*((\mathbf{A}'_h, \varphi'_h), (\mathbf{A} - \mathbf{A}_h, \varphi - \varphi_h)), \quad \forall (\mathbf{A}'_h, \varphi'_h) \in \mathbf{V}_h.$$

Subtracting this identity to (68), we obtain (67). ■

**Corollary 5.4** Suppose that  $D$  and  $D_c$  are polyhedra and define (see [12])

$$\mathbf{V}_h = \{(\mathbf{A}'_h, \varphi'_h) \in \tilde{X}(D) \times \tilde{H}^1(D_c) : A'_{h|T} \in \mathcal{ND}_0(T), \text{ and } \varphi'_{h|T} \in \mathbb{P}_1(T), \forall T \in \mathcal{T}\}, \quad (69)$$

where  $\mathcal{ND}_0(T)$  is the low-order Nédélec element defined by

$$\mathcal{ND}_0(T) = \{p \in \mathbb{P}_1(T)^3 : \exists a, b \in \mathbb{C}^3 \text{ s.t. } p(x) = a + b \times x, \forall x \in T\}.$$

Let  $(\mathbf{A}_h, \varphi_h) \in \mathbf{V}_h$  be the Galerkin approximation of the solution  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  of (10). Suppose further that the solution  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  of (10) belongs to  $H^2(D)^3 \times H^2(D_c)$  and that the solution  $(\mathbf{A}^*, \varphi^*) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  of (43) belongs to  $H^2(D)^3 \times H^{1+s}(D_c)$ , for some  $s \in (0, 1]$ . If  $\mathcal{T}$  is a regular triangulation, then

$$|\mathcal{E}| \leq Ch^{1+s}, \quad (70)$$

for a positive constant  $C$  that does not depend on  $h$ .

**Proof.** By the identity (67) with  $\mathbf{A}'_h = I_{ND}\mathbf{A}^*$  and  $\varphi'_h = I_L\varphi^*$ , where  $I_{ND}$  (resp.  $I_L$ ) is the (low-order) Nédélec (resp.  $\mathbb{P}_1$  Lagrange) interpolant, we have

$$|\mathcal{E}| \leq C_1(\|\mathbf{A}^* - I_{ND}\mathbf{A}^*\|_{H(\text{curl}, D)} + |\varphi^* - I_L\varphi^*|_{H^1(D_c)})(\|\mathbf{A} - \mathbf{A}_h\|_{H(\text{curl}, D)} + |\varphi - \varphi_h|_{H^1(D_c)}),$$

for some positive constant  $C_1$  independent of  $h$ . As there exists a positive constant  $C_2$  independent of  $h$  such that

$$\|\mathbf{A} - \mathbf{A}_h\|_{H(\text{curl}, D)} + |\varphi - \varphi_h|_{H^1(D_c)} \leq C_2(\|\mathbf{A} - I_{ND}\mathbf{A}\|_{H(\text{curl}, D)} + |\varphi - I_L\varphi|_{H^1(D_c)}),$$

we conclude that

$$\begin{aligned} |\mathcal{E}| &\leq C_1 \max\{C_2, 1\}(\|\mathbf{A}^* - I_{ND}\mathbf{A}^*\|_{H(\text{curl}, D)} + |\varphi^* - I_L\varphi^*|_{H^1(D_c)}) \\ &\quad (\|\mathbf{A} - I_{ND}\mathbf{A}\|_{H(\text{curl}, D)} + |\varphi^* - I_L\varphi|_{H^1(D_c)}). \end{aligned}$$

We conclude by standard interpolation error estimates for the Nédélec and Lagrange elements. ■




Figure 1: Domains configuration.

## 6 Numerical validation

In this section we propose a numerical test inspired by [12] (see section 5.1), in order to underline and confirm our theoretical predictions, using the FreeFem++ software [19]. The data are built in order to have in hand an exact solution, allowing to compare the estimator to the exact error. The domains are defined by  $D = [-2, 5] \times [-2, 2] \times [-2, 2]$ ,  $D_s = [-1, 1]^3$  and  $D_c = [2, 4] \times [-1, 1] \times [-1, 1]$  (see Figure 1). We set  $\mu \equiv 1$  in  $D$ ,  $\sigma \equiv 1$  in  $D_c$  and  $\omega = 2\pi$ . Then, the exact solution is given by  $\varphi \equiv 0$  and

$$\mathbf{A} = \text{curl} \begin{pmatrix} f \\ 0 \\ 0 \end{pmatrix} \text{ with } f(x, y, z) = \begin{cases} (x^2 - 1)^4 (y^2 - 1)^4 (z^2 - 1)^4 & \text{in } D_s, \\ 0 & \text{in } D \setminus D_s. \end{cases}$$

The value of  $\mathbf{J}_s$  is computed accordingly using (8a), and is given by

$$\mathbf{J}_s = \text{curlcurl}\mathbf{A},$$

as the support of  $\mathbf{A}$  is equal to  $D_s$  (that is disjoint to  $D_c$ ).

First of all, we compute  $(\mathbf{A}_h, \varphi_h) \in \mathbf{V}_h$ , with  $\mathbf{V}_h$  defined by (69), as the Galerkin approximation of the solution  $(\mathbf{A}, \varphi) \in \tilde{X}(D) \times \tilde{H}^1(D_c)$  of (10). The estimator  $\eta$  defined by (23), with  $\mathbf{S}_h = \mathbf{A}_h$  and  $\psi_h = \varphi_h$ , is computed in the same manner as in [10]. As mentioned in remark 4.1, we choose

$$\mathbf{q} = \mathbf{H}_s \tag{71}$$

with  $\text{curl}\mathbf{H}_s = \mathbf{J}_s$ , and we are interested in the value of  $\mathcal{E}$  given by:

$$\mathcal{E} = \int_D \mathbf{H}_s \cdot \text{curl} \overline{(\mathbf{A} - \mathbf{A}_h)} dx = \int_D \mathbf{J}_s \cdot \overline{(\mathbf{A} - \mathbf{A}_h)} dx.$$

Consequently in order to validate the identity (54) and the estimate (58), it remains to compute a numerical approximation  $(\mathbf{A}_h^*, \varphi_h^*)$  of the solution  $(\mathbf{A}^*, \varphi^*)$  of the adjoint problem (43), and the estimator  $\eta^*$ . Let us first notice that for this example, we directly check that  $(\mathbf{A}^*, \varphi^*) = (\mathbf{A}, 0)$ . Here we take  $(\mathbf{A}_h^*, \varphi_h^*) \in \mathbf{V}_h^*$  as the Galerkin approximation of  $(\mathbf{A}^*, \varphi^*)$ , with  $\mathbf{V}_h^*$  defined as (69), but with piecewise Nédélec elements of higher degree (namely polynomial of degree two in each tetrahedron) for  $\mathbf{A}_h^*$ , and with piecewise polynomial of degree two for  $\varphi_h^* \in H^1(D_c)$ . The estimator  $\eta^*$  defined by (47), with  $\mathbf{S}_h^* = \mathbf{A}_h^*$  and  $\psi_h^* = \varphi_h^*$ , is computed in the same manner as in [10]. Finally, the estimator  $\eta_{\text{QOI}}$  is computed by (55).

Tests are performed using two sets of meshes as described below.




Figure 2: Mesh 4, global refinement : 12 482 elements in  $D_s$ , 12 974 elements in  $D_c$  and 153 178 elements in  $D \setminus (D_s \cup D_c)$ .


Mesh	Nb of elements	Nb dof for $(\mathbf{A}_h, \varphi_h)$	Nb dof for $(\mathbf{A}_h^*, \varphi_h^*)$
1	5 935	8 097	41 843
2	49 642	63 031	334 722
3	110 430	137 324	733 305
4	178 634	222 824	1 189 235
5	944 475	1 144 479	6 170 254

Table 1: The meshes used for the simulations, global refinement.


## 6.1 Results with globally refined meshes

The first set consists in some uniformly refined meshes. Figure 2 displays the fourth mesh of the series (Mesh 4), and Table 1 indicates for each mesh the number of tetrahedra and the number of degrees of freedom associated to each formulation (primal or dual one). Recall that since the adjoint problem is computed with finite elements of higher degrees, the number of degrees of freedom is more important on the same mesh.

To begin with, we plot in Figure 4 the values of  $\mathcal{E}$ ,  $\eta_{\text{QOI}}$  and  $10\eta\eta^*$  as a function of the number of elements  $N$ , in a log-log scale. More precisely, the real parts are displayed in Figure 4(a) and the imaginary ones in Figure 4(b). First, it can be seen in Figure 4(a) that  $\Re(\mathcal{E})$  goes towards zero as  $O(N^{-2/3}) = O(h^2)$ . This is a consequence of Corollary 5.4 (with  $s = 1$ ), since in this example  $\mathbf{A} = \mathbf{A}^*$  are in  $H^3(D)^3$ , while  $\varphi = \varphi^* = 0$ . There it can be seen that the value of  $10\eta\eta^*$  converges faster towards zero than the real part  $\Re(\mathcal{E})$  and becomes smaller than  $\Re(\mathcal{E})$  when the mesh is sufficiently refined. Again from the regularity of  $\mathbf{A}$  and  $\mathbf{A}^*$ , the error  $\mathbf{A} - \mathbf{A}_h$  (resp.  $\mathbf{A}^* - \mathbf{A}_h^*$ ) will be of order  $h$  (resp.  $h^2$ ), and therefore we may expect a convergence in  $h$  for  $\eta$  and in  $h^2$  for  $\eta^*$ . From Theorem 5.2 and estimate (58), it shows that the reminder  $\mathcal{R}$  is a superconvergent term. Moreover,  $\Re(\eta_{\text{QOI}})$  has exactly the same behaviour as  $\Re(\mathcal{E})$ , that once again from Theorem 5.1 and estimate (54) shows that the reminder  $\mathcal{R}$  can be neglected. Concerning the imaginary parts, it can be seen in Figure 4(b) that the values are very small, so that they are not significant, even if the behaviour of the imaginary part  $\Im(\mathcal{E})$  and  $\Im(\eta_{\text{QOI}})$  are once again exactly the same (and  $\Im(\mathcal{E})$  seems to tend towards zero faster than  $O(N^{-2/3})$ ). Then we plot in Figure 5 the values of  $\Re(\mathcal{E})/\Re(\eta_{\text{QOI}})$  (Figure 5(a)) and  $\Im(\mathcal{E})/\Im(\eta_{\text{QOI}})$  (Figure 5(b)), defined as the real effectivity index and the imaginary effectivity index. We can see that they are both exactly equal to one whatever the mesh in consideration. It illustrates the fact that the estimator  $\eta_{\text{QOI}}$  gives a very accurate evaluation of the error  $\mathcal{E}$ .




(a)  $\Re(\mathcal{E})$ ,  $\Re(\eta_{QOI})$  and  $10\eta^*$




(b)  $\Im(\mathcal{E})$  and  $\Im(\eta_{QOI})$

Figure 3: Error and Estimator, global refinement,  $\mathbf{q} = \mathbf{H}_s$ .




(a)  $\Re(\mathcal{E})$ ,  $\Re(\eta_{QOI})$  and  $10\eta^*$




(b)  $\Im(\mathcal{E})$  and  $\Im(\eta_{QOI})$

Figure 4: Error and Estimator, global refinement,  $\mathbf{q} = \mathbf{H}_s$ .



(a)  $\Re(\mathcal{E})/\Re(\eta_{QOI})$



(b)  $\Im(\mathcal{E})/\Im(\eta_{QOI})$

Figure 5: Ratios of error over estimator, real and imaginary parts, global refinement,  $\mathbf{q} = \mathbf{H}_s$ .




Figure 6: Mesh 3, local refinement : 243 232 elements in  $D_s$ , 5 794 elements in  $D_c$  and 103 136 elements in  $D \setminus (D_s \cup D_c)$ .

Mesh	Nb of elements	Nb dof for $(\mathbf{A}_h, \varphi_h)$	Nb dof for $(\mathbf{A}_h^*, \varphi_h^*)$
1	86 335	108 740	577 696
2	121 005	149 374	797 644
3	352 162	420 454	2 264 432
4	2 128 618	2 500 710	13 530 768

Table 2: The meshes used for the simulations, local refinement.

## 6.2 Results with locally refined meshes

The second set of meshes consists in some locally refined meshes. Here, the meshes are more accurate in  $D_s$ , corresponding to the support of  $\mathbf{J}_s$ . Figure 6 displays the third mesh of the series (Mesh 3), and Table 2 indicates for each mesh the number of tetrahedra and the number of degrees of freedom associated to each formulation (primal or dual one). The same tests are performed, and Figures 7 and 8 display the same results as Figures 4 and 5 on this second set of meshes. The conclusions are very similar, and we can see moreover that the value of  $10\eta\eta^*$  goes towards zero faster than in the globally refined case, what can be explained by the fact that the support of  $\mathbf{q} = \mathbf{H}_s$ , namely  $D_s$ , is here better refined.

## 6.3 Results with another value of $\mathbf{q}$

A last test is now proposed. It consists in the same as in subsection 6.1, but instead of defining  $\mathbf{q}$  by (71), we choose:

$$\mathbf{q} = \begin{pmatrix} \rho_s \\ 0 \\ 0 \end{pmatrix}, \quad (72)$$

with

$$\rho_s(x, y, z) = e^{-\frac{(x-3)^2 + y^2 + z^2}{\log(10)^4}}, \forall (x, y, z) \in D.$$

The error we are interested in is given by (42). Figures 9 and 10 display the same results as Figures 4 and 5 for this new error definition. The difference with the previous example relies on the limited regularity of  $\varphi^*$ . Indeed Lemma 2.2 (point 3) guarantees  $\mathbf{A}^* \in H^2(D)^3$  and Lemma 2.3 (point 3) guarantees  $\varphi^* \in H^{1+s}(D_c)$ , for all  $s \in (0, 1)$ . This means that  $\eta^*$  could converge to 0 only in  $h^s$ , for all  $s \in (0, 1)$ . This is indeed the case since it can be seen that the quantity  $10\eta\eta^*$  does no more converge faster towards zero than the error  $\mathcal{E}$ , and remains significantly higher. Nevertheless, the error  $\mathcal{E}$  and the estimator  $\eta_{\text{QOI}}$  remain nearly the same, and converge




Figure 7: Error and Estimator, local refinement,  $\mathbf{q} = \mathbf{H}_s$ .




Figure 8: Ratios of error over estimator, real and imaginary parts, local refinement,  $\mathbf{q} = \mathbf{H}_s$ .




Figure 9: Error and Estimator, global refinement,  $\mathbf{q}$  given by (72).




Figure 10: Ratios of error over estimator, real and imaginary parts, global refinement,  $\mathbf{q}$  given by (72).

towards zero in  $h^2$  (consequence of Corollary 5.4, with  $s$  as close as we want to 1). It shows that in that case, the estimation (58) given by Theorem 5.2 strongly overestimates the value of  $\mathcal{R}$ . Nevertheless and despite this phenomenon, the estimator  $\eta_{\text{QOI}}$  provides once again a very accurate evaluation of the error  $\mathcal{E}$ .

## References

- [1] C. Amrouche, C. Bernardi, M. Dauge, and V. Girault. Vector potentials in three-dimensional non-smooth domains. *Math. Methods Appl. Sci.*, 21(9):823–864, 1998.
- [2] M. Bebendorf. A note on the Poincaré inequality for convex domains. *Z. Anal. Anwendungen*, 22(4):751–756, 2003.
- [3] C. Carstensen and S. A. Funken. Constants in Clément-interpolation error and residual based a posteriori error estimates in finite element methods. *East-West J. Numer. Math.*, 8(3):153–175, 2000.
- [4] T. Chaumont-Frelet, A. Ern, and M. Vohralík. Stable broken  $\mathbf{H}(\mathbf{curl})$  polynomial extensions and  $p$ -robust a posteriori error estimates by broken patchwise equilibration for the curl-curl problem. *Math. Comp.*, 91(333):37–74, 2021.
- [5] C. Chen, E. Creusé, S. Nicaise, and Z. Tang. Residual-based a posteriori estimators for the potential formulations of electrostatic and time-harmonic eddy current problems with voltage or current excitation. *Internat. J. Numer. Methods Engrg.*, 107(5):377–394, 2016.
- [6] M. Costabel and M. Dauge. Singularities of electromagnetic fields in polyhedral domains. *Arch. Rational Mech. Anal.*, 151:221–276, 2000.
- [7] M. Costabel, M. Dauge, and S. Nicaise. Singularities of Maxwell interface problems. *RAIRO Modél. Math. Anal. Numér.*, 33:627–649, 1999.
- [8] M. Costabel, M. Dauge, and S. Nicaise. Singularities of Maxwell interface problems. *M2AN Math. Model. Numer. Anal.*, 33(3):627–649, 1999.



- [9] M. Costabel, M. Dauge, and S. Nicaise. *Corner Singularities and Analytic Regularity for Linear Elliptic Systems. Part I: Smooth domains*. <http://hal.archives-ouvertes.fr/hal-00453934/en/> Online version of Chapters 1 to 5. (2010).
- [10] E. Creusé, Y. Le Menach, S. Nicaise, F. Piriou, and R. Tittarelli. Two guaranteed equilibrated error estimators for harmonic formulations in eddy current problems. *Comput. Math. Appl.*, 77(6):1549–1562, 2019.
- [11] E. Creusé, S. Nicaise, and Z. Tang. Goal-oriented error estimation based on equilibrated flux and potential reconstruction for the approximation of elliptic and parabolic problems. Preprint hal-03719266, Céramaths, UPHF, 2022.
- [12] E. Creusé, S. Nicaise, Z. Tang, Y. Le Menach, N. Nemitz, and F. Piriou. Residual-based *a posteriori* estimators for the  $\mathbf{A} - \varphi$  magnetodynamic harmonic formulation of the Maxwell system. *Math. Models Methods Appl. Sci.*, 22(5):1150028, 30, 2012.
- [13] E. Creusé, S. Nicaise, and R. Tittarelli. A guaranteed equilibrated error estimator for the  $\mathbf{A} - \varphi$  and  $\mathbf{T} - \Omega$  magnetodynamic harmonic formulations of the Maxwell system. *IMA J. Numer. Anal.*, 37(2):750–773, 2017.
- [14] M. Dauge. *Elliptic boundary value problems on corner domains – smoothness and asymptotics of solutions*, volume 1341 of *Lecture Notes in Mathematics*. Springer, Berlin, 1988.
- [15] A. Ern and M. Vohralík. A unified framework for a posteriori error estimation in elliptic and parabolic problems with application to finite volumes. In *Finite volumes for complex applications VI. Problems & perspectives. Volume 1, 2*, volume 4 of *Springer Proc. Math.*, pages 821–837. Springer, Heidelberg, 2011.
- [16] R. Eymard, T. Gallouët, and R. Herbin. Finite volume methods. In *Handbook of numerical analysis, Vol. VII*, *Handb. Numer. Anal.*, VII, pages 713–1020. North-Holland, Amsterdam, 2000.
- [17] M. Fratila, A. Benabou, A. Tounzi, and M. Dessoude. Iron loss calculation in a synchronous generator using finite element analysis. *IEEE Transactions on Energy Conversion*, 32(2):640–648, 2017.
- [18] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 24 of *Monographs and Studies in Mathematics*. Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [19] F. Hecht. New development in FreeFem++. *J. Numer. Math.*, 20(3-4):251–265, 2012.
- [20] T. Henneron, Y. Le Menach, F. Piriou, O. Moreau, S. Clénet, J.-P. Ducreux, and J.-C. Verite. Source field computation in NDT applications. *IEEE Transactions on Magnetics*, 43(4):1785–1788, 2007.
- [21] L. Jiang, H. Yan, J. Meng, Z. Yin, W. Wei, and Y. Wang. Analysis of eddy current effect and loss calculation of transformer winding based on finite element algorithm. In *2017 International Conference on Computer Systems, Electronics and Control (ICCSEC)*, pages 405–410, 2017.
- [22] G. Mallik, M. Vohralík, and S. Yousef. Goal-oriented *a posteriori* error estimation for conforming and nonconforming approximations with inexact solvers. *J. Comput. Appl. Math.*, 366:112367, 20, 2020.
- [23] P. Monk. *Finite element methods for Maxwell’s equations*. Numerical Mathematics and Scientific Computation. Oxford University Press, New York, 2003.

- [24] I. Mozolevski and S. Prudhomme. Goal-oriented error estimation based on equilibrated-flux reconstruction for finite element approximations of elliptic problems. *Comput. Methods Appl. Mech. Engrg.*, 288:127–145, 2015.
- [25] L. E. Payne and H. F. Weinberger. An optimal Poincaré inequality for convex domains. *Arch. Rational Mech. Anal.*, 5:286–292 (1960), 1960.
- [26] Z. Tang, Y. Le Menach, E. Creusé, S. Nicaise, F. Piriou, and N. Nemitz. A posteriori error estimator for harmonic  $\mathbf{A}$ - $\varphi$  formulation. *COMPEL*, 32(4):1219–1229, 2013.
- [27] Z. Tang, Y. Le Menach, E. Creusé, S. Nicaise, F. Piriou, and N. Nemitz. Residual and equilibrated error estimators for magnetostatic problems solved by finite element method. *IEEE Transactions on Magnetics*, 49 (5), may 2013.
- [28] Z. Tang, Y. Le Menach, E. Creusé, S. Nicaise, F. Piriou, and N. Nemitz. Residual based a posteriori error estimators for harmonic  $\mathbf{A}/\varphi$  and  $\mathbf{T}/\Omega$  formulations in eddy current problems. *IEEE Transactions on Magnetics*, 49 (5):1721–1724, may 2013.
- [29] Z. Tang, S. Lou, A. Benabou, E. Creusé, S. Nicaise, J. Korecki, and J.-C. Mipo. Guaranteed quantity of interest error estimate based on equilibrated flux reconstruction. *IEEE Transactions on Magnetics*, 57(6):1–4, 2021.
- [30] R. Tittarelli, Y. Le Menach, F. Piriou, E. Creusé, S. Nicaise, and J.-P. Ducreux. Comparison of numerical error estimators for eddy-current problems solved by FEM. *IEEE Transactions on Magnetics*, 54(3):1–4, 2018.