



**HAL**  
open science

## The effect of the 7R allele at the DRD4 locus on risk tolerance is independent of background risk in Senegalese fishermen

Gwen-Jirō Clochard, Aby Mbengue, Clément Mettling, Birane Diouf, Charlotte Faurie, Omar Sene, Emilie Chancerel, Erwan Guichoux, Guillaume Hollard, Michel Raymond, et al.

### ► To cite this version:

Gwen-Jirō Clochard, Aby Mbengue, Clément Mettling, Birane Diouf, Charlotte Faurie, et al.. The effect of the 7R allele at the DRD4 locus on risk tolerance is independent of background risk in Senegalese fishermen. *Scientific Reports*, 2023, 13 (1), pp.622. 10.1038/s41598-022-27002-3 . hal-03954770

**HAL Id: hal-03954770**

**<https://hal.science/hal-03954770>**

Submitted on 24 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN

## The effect of the 7R allele at the DRD4 locus on risk tolerance is independent of background risk in Senegalese fishermen

Gwen-Jirō Clochard<sup>1,2</sup>✉, Aby Mbengue<sup>3</sup>, Clément Mettling<sup>4,5</sup>, Birane Diouf<sup>3</sup>, Charlotte Faurie<sup>4</sup>, Omar Sene<sup>6</sup>, Emilie Chanceler<sup>7</sup>, Erwan Guichoux<sup>7</sup>, Guillaume Hollard<sup>1,9</sup>, Michel Raymond<sup>4,9</sup> & Marc Willinger<sup>8,9</sup>

It has been shown that living in risky environments, as well as having a risky occupation, can moderate risk-tolerance. Despite the involvement of dopamine in the expectation of reward described by neurobiologists, a GWAS study was not able to demonstrate a genetic contribution of genes involved in the dopaminergic pathway in risk attitudes and gene candidate studies gave contrasting results. We test the possibility that a genetic effect of the DRD4-7R allele in risk-taking behavior could be modulated by environmental factors. We show that the increase in risk-tolerance due to the 7R allele is independent of the environmental risk in two populations in Northern Senegal, one of which is exposed to a very high risk due to dangerous fishing.

Humans need to adapt their behavior as a result of risk. Previous research has shown that risk coping attitudes are partly heritable<sup>1</sup>. Genes involved in the regulation of the dopaminergic system are good candidates to explain the heritability of risk behavior. However, many reports on gene and behavior association, based on small-sample candidate gene have found contrasting results, leading to debates in the scientific community<sup>2,3</sup>.

To overcome this limitation, a genome-wide association study (GWAS), based on over 1 million individuals, identified 99 loci associated with general risk tolerance<sup>4</sup>. Surprisingly, none of identified loci were close to genes involved in the dopamine pathway. Their bioinformatic analysis pointed to the role of genes expressed in brain regions involved in decision-making, although genes involved in dopamine biosynthesis (TH) or receptors (DRD1,2,3 and 4) did not reach statistical significance.

Yet, the evidence that not only these brain regions but the dopamine neurotransmitter itself plays a role in the expectation of reward is compelling: dopaminergic neurons can code the probability of the reward in a primate model<sup>5</sup>. Moreover, a known side-effect of the treatment of Parkinson disease (known to impair dopamine production) is to dramatically increase impulsivity<sup>6</sup>. The dopamine receptor gene DRD4 fulfills many criteria as a good candidate gene: it is highly polymorphic<sup>7,8</sup>, expressed in the prefrontal cortex, it shows an unusually large variable repeat region (VNTR: variable number tandem repeat) coding for 16 amino acids in the third cytoplasmic loop, a region interacting with SH3 domain-binding proteins.

While the 4 repeat (4R) variant is the ancestral, and predominant allele in most human populations<sup>9</sup>, there exist variations between 2 and 11 repeats (2R to 11R). The different alleles have functional differences<sup>10–13</sup>. The DRD4-7R allele is under strong positive selection in human population<sup>14–16</sup>, and has been shown to be linked to more risk-tolerant attitudes<sup>17–19</sup>. However, some findings revealed a lack of differences in the domain of financial risk-taking<sup>20–23</sup>.

The discrepancy between these studies may come from the fact that GWAS studies tend to underestimate the genetic variance due to gene-gene or gene-environment interactions, or an inability to capture rare genetic variants. Furthermore, candidate-gene studies conducted in specific environments may sometimes benefit from

<sup>1</sup>Ecole polytechnique - CREST (CNRS-UMR 9194), Palaiseau, France. <sup>2</sup>University of Chicago and Universidad del CEMA's Joint Initiative for Latin American Experimental Economics, Chicago, USA. <sup>3</sup>Université Gaston Berger, Saint-Louis, Senegal. <sup>4</sup>CNRS, IRD, ISEM Univ Montpellier, Montpellier, France. <sup>5</sup>CNRS, IGH Univ Montpellier, Montpellier, France. <sup>6</sup>Université Alioune Diop, Bambey, Senegal. <sup>7</sup>INRAE, BIOGECO, Plateforme Genome Transcriptome de Bordeaux, Univ Bordeaux, 33610 Cestas, France. <sup>8</sup>CNRS, INRAE, CEE-M Univ Montpellier, Montpellier, France. <sup>9</sup>These authors contributed equally: Guillaume Hollard, Michel Raymond and Marc Willinger. ✉email: gwen-jiro.clochard@polytechnique.org

circumstances revealing a genetic variance. For instance, administration of L-DOPA to volunteers did not lead to an increase in gambling propensity unless the subjects carried at least one copy of the 7-repeat allele<sup>24</sup>. The negative association between DRD4 variation and risk-taking previously reported<sup>23</sup> might have been concealed by the association with MAOA variation, an enzyme catalysing dopamine. It is therefore likely that the effect of genetic variants of DRD4 on risk-taking behavior appears only in specific circumstances. GWAS studies, by leveling all environmental conditions or gene interactions, may mask some dopaminergic genetic contributions. Two studies have also shown the effect of DRD4 to be modulated by an interaction with maternal effects<sup>25,26</sup>.

Humans also adapt their risk attitudes as a response to the level of risk in their environment<sup>27</sup>. In particular, people have been found to be more risk-averse in the presence of unfair background risk<sup>28–31</sup>, in accordance with the “risk-vulnerability” hypothesis<sup>32,33</sup>.


The aim of the present paper is to test the interaction between the influence of the 7R allele on risk-tolerance and the level of risk to which people are exposed.

## Results

**Risk-tolerance by zone.** The village of Guet Ndar (Saint-Louis region in Northern Senegal) is famous for its fisheries. Fishing in the area is very dangerous, with authorities reporting 20 deaths due to fishing on average per year over the past 20 years<sup>34</sup>. Given the demography of the village, with 20 000 inhabitants, among which fishing represents the main occupation of approximately 80% of the adult male workforce, this corresponds to approximately 4% of the male population who died due to fishing in the last 20 years. The prevalence of deaths is strongly linked to the intersection of strong currents coming from the Senegal river and an upwelling current from the ocean<sup>35</sup>. However, these currents attract a lot of fish, making fishing more profitable than other activities in the region (fishermen in our sample declare income significantly higher than non-fishermen,  $p < 0.01$ , Table S.1).

In this paper, we compared populations from the fishing village of Guet Ndar ( $N = 601$ ), which is labelled as the *risky area*, and that of a farming village called Mouit, 23 kilometers away ( $N = 263$ ), labelled the *non-risky area*. Importantly, the two populations are mostly composed of the same ethnic group (the Wolofs, representing approximately 80% of the sample in both areas). Because fishing is an activity predominately performed by men, our sample only consists of men. Only participants over 18 years old were allowed to participate. No further restrictions on participants were placed, e.g. based on residency, activity. Descriptive statistics are provided in Table S.2.

Our experimental measure of risk-tolerance was based on a lottery task<sup>36</sup>. A description of the task is provided in the Supplementary Materials. Results indicate that risk-tolerance varied between the risky and non-risky areas. Participants from the risky area tended to exhibit less risk-tolerance than participants from the non-risky area (Figure 1, Student’s  $t$ -test  $p < 0.01$ ). The difference remains significant after controlling for age and education



**Figure 1.** Average level of risk-tolerance by zone. Note: This figure shows the average level of risk-tolerance between the risky and non-risky areas. A higher level of risk-tolerance indicates the choice of a riskier lottery by participants in the lottery choice task. Segments represent 95% confidence intervals. Student’s  $t$ -test \*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$ .

(Table S.3). Our data is consistent with field data and laboratory experiments showing that people exposed to high background risk tend to exhibit less risk-tolerance, in accordance with the “risk-vulnerability hypothesis”<sup>32,33</sup>.

**Genotypes by zone.** Genotypes at the *DRD4* locus displayed two common alleles (4R and 7R, with 4 and 7 repeats, respectively), which was expected for populations in Sub-Saharan Africa<sup>9</sup>, and 5 minor alleles with negligible frequencies (2R, 3R, 5R, 6R and 8R) leading to 21 different genotypes (Table 1). The 7R allele was more prevalent than previously found in other African populations<sup>9</sup>, possibly because the ethnic groups from participants were not sampled in previous work. In Chang et al., 1996, the groups analyzed are Bantu and San Bushmen, from South-Africa and Namibia, Biaka from the Central African Republic, Mbuti from the Democratic Republic of Congo and Falasha in Ethiopia. The closest group from our sample geographically would be the Biaka people from the Central African Republic, 5,000 km away.

Within each area, populations were not at Hardy-Weinberg equilibrium ( $p < 0.01$  in the non-risky area,  $p = 0.02$  in the risky area), and displayed an heterozygote deficiency ( $F_{IS} = 0.134$  in the non-risky area, and  $F_{IS} = 0.052$  in the risky area).

Because we were primarily interested in the effect of the 7R allele on risk-taking, we combined all other alleles into a single category, identified as allele “X”. This combination yielded three genotypes: XR/XR, XR/7R and 7R/7R. Hardy-Weinberg equilibrium was rejected ( $p = 0.01$ ) for the non-risky area, but not for the risky area ( $p = 0.40$ ), see Table 1. Deviations from HW equilibrium were  $F_{IS} = 0.181$  in the non-risky area, and  $F_{IS} = 0.037$  in the risky area.

We found evidence of limited migration. In the risky area, 81% of our sample were born in the same village, as were 72% of their parents, and 68% of grand parents. Numbers were slightly lower for the non-risky area, with 67% of participants, 58% of their parents and 50% of their grand parents (Table S.6).

Genotype	Non-risky area		Risky area	
	N	%	N	%
<b>Panel A. Without combination of genotypes</b>				
22	3	1.4	3	0.6
24	7	3.3	14	2.8
25	1	0.5	–	–
27	–	–	2	0.4
34	3	1.4	–	–
36	1	0.5	–	–
37	1	0.5	–	–
44	84	39.1	202	40
45	19	8.8	34	6.7
46	14	6.5	18	3.6
47	48	22.3	149	29.5
48	5	2.3	12	2.4
55	4	1.9	3	0.6
56	–	–	1	0.2
57	4	1.9	14	2.6
58	–	–	2	0.4
66	2	0.9	1	0.2
67	1	0.5	12	2.4
77	15	7	34	6.7
78	3	1.4	4	0.8
88	–	–	1	0.2
<b>HW equilibrium</b>				
<i>p</i>	<0.01		0.023	
<b>Panel B. Allele 7R versus other alleles</b>				
XX	143	66.5	291	57.5
X7	57	26.5	181	35.8
77	15	7.0	34	6.7
<b>HW equilibrium</b>				
<i>p</i>	0.011		0.40	

**Table 1.** Genotypic composition at the *DRD4* locus of populations from the Saint-Louis region in the non-risky and risky areas. The *p*-value (*p*) corresponds to the HW probability exact test. Genotype *ij* refers to the *DRD4* genotype *iR/jR*. For *Panel B*, all alleles not 7R are combined in the X allele.

The genotypic differentiation between the two areas was measured as  $F_{ST} = 0.0036$ , and was marginally non-significant (exact G test,  $p = 0.094$ ). This level of genotypic differentiation was compared with those displayed by 30 micro-satellite loci. One locus (032) was not polymorphic and was discarded. The other 29 loci displayed between 2 and 15 alleles. Their level of genotypic differentiation ranged between  $F_{ST} = -0.0094$  and  $F_{ST} = 0.0226$ , with an overall average value of  $F_{ST} = 0.0035$  (Figure S.2 and Table S.8).


**Risk-tolerance by genotype.** Risk-tolerance was not independent of genotype at the DRD4 locus (Figure 2 and Table 2, Column 1). The 7R allele demonstrated a significant additive effect ( $p = 0.01$ ), and no dominance effect was found ( $p = 0.31$ ). The 7R allele increased risk-tolerance. Importantly, the result held after controlling for age, education and the living area (Table 2, Column 2). Our results indicated that the 7R allele was associated with more risk-tolerance than other alleles, in line with previous literature<sup>17,18</sup>. Moreover, we did not find that the 7R allele was associated with a measure of novelty-seeking (Table S.5).

Environmental risk did not appear to significantly moderate the effect of the 7R allele. First, the additive effect held when analyzing both areas separately (Table 2, Columns 3 and 4, Figure S.1), although the significance levels dropped slightly due to sample limitations ( $p = 0.05$  and  $p = 0.08$  in the non-risky and risky area, respectively). Second, the interaction between the additive effect and the area (Table 2, Column 5) was not significant ( $p = 0.25$ ).

## Discussion

In this paper, we found that the 7R allele of DRD4 affects risk-attitudes by an additivity effect, not a dominance effect. This is in contrast with previous research<sup>37</sup> who found that heterozygotes 2R/4R had lower risk tolerance. It is unclear if this difference comes from the type of risky environment considered (background volcanic risk or risky subsidence type), or comes from the different alleles involved (2R and 4R in Indonesia, or 7R in Senegal). In addition, we did not find evidence that the 7R allele is associated with novelty seeking, as previously found<sup>38</sup>.

The main result of this paper is that there is no difference in the effect of the 7R allele of the DRD4 locus on risk-tolerance, depending on the risk to which the participants were exposed (Table 2, column 5). Two reasons could be advanced, then, to explain why the DRD4 locus did not reach statistical significance in the GWAS analysis on risk-taking behavior<sup>4</sup>. First, the measure of risk-tolerance in this study differs from the GWAS study: while ours is a behavioral measure, theirs is based on a survey question. Second, as GWAS can usually only test for the effects of Single Nucleotide Polymorphisms (SNPs), it did not directly measure the one due to the DRD4 VNTR. However, as the 7R allele is under strong positive selection, SNPs with a high linkage disequilibrium to this VNTR should have been detected<sup>14,15</sup>.



**Figure 2.** Average level of risk-tolerance by genotype. Note: This figure shows the average level of risk-tolerance between the different genotypes. A higher level of risk-tolerance indicates the choice of a riskier lottery by participants in the lottery choice task. X/X, X/7 and 7/7 represent genotypes, with all alleles not 7R combined into the X allele. Samples from both risky and non-risky areas are pooled. Segments represent 95% confidence intervals. Student's *t*-test \*  $p < 0.1$ , \*\*  $p < 0.05$ .

	(1)	(2)	(3)	(4)	(5)
	Combined	Combined	Non-risky	Risky	Gene-
	sample	sample	area	area	environment
	without	with	only	only	interaction
	controls	controls			
7R: additive effect	0.068** (0.027)	0.064** (0.028)	0.097* (0.049)	0.056* (0.032)	0.107** (0.045)
7R: dominance effect	-0.037 (0.036)	-0.026 (0.037)	-0.003 (0.069)	-0.036 (0.042)	-0.024 (0.037)
Age		-0.002* (0.001)			-0.002** (0.001)
Education		-0.004 (0.005)			-0.004 (0.005)
Risky area		-0.120*** (0.032)			-0.094** (0.038)
Risky area × 7R: additive effect					-0.062 (0.049)
Constant	0.456*** (0.017)	0.613*** (0.053)	0.507*** (0.030)	0.431*** (0.021)	0.597*** (0.054)
Observations	721	699	215	506	699
R <sup>2</sup>	0.009	0.030	0.026	0.006	0.033

**Table 2.** Differences between genotypes in risk-tolerance. *Note:* The outcome variable is risk-tolerance. A higher level of risk-tolerance indicates the choice of a riskier lottery by participants in the lottery choice task. Standard errors in parentheses. The coefficients are the results of Ordinary Least Square (OLS) estimations. In column 1, the sample is pooled (non-risky and risky areas) and no control variable is included. In column 2, the sample is pooled and controls for age, education and a dummy for living in the risky area are included. In columns 3 and 4, the sample is restricted to individuals from the non-risky and risky areas, respectively. In column 5, we interact the additive effect of the 7R allele with the dummy variable for living in the risky area. Further details on the equation used in column 5 can be found in the Methods section. Student's *t*-test \*  $p < 0.10$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$ . To see the correlations between controls and risk-tolerance, absent any genetic factor, see Table S.3.

The sample would satisfy conditions for a genetic adaptation to habitat<sup>39,40</sup>, with (1) limited migration - 68% of grandparents of the risky area were born in the same village (Table S.6), (2) strong economic benefit to live in risky area and (3) an heritable genetic trait which would help cope with risk, i.e. the 7R allele. However, we did not find evidence of genetic differentiation at the DRD4 locus relative to 29 unlinked microsatellites loci (Table S.8 and Figure S.2). Moreover, if there was genetic differentiation, it would move in the opposite direction as the risk-vulnerability hypothesis found in previous work for DRD4<sup>37</sup>, as the 7R allele, favoring more risk-tolerant attitudes, is more prevalent in the risky area. Altogether, our results indicate that no selection at the DRD4 locus is apparent in our sample. This does not mean that such selection is absent, as many generations of selection are required for gene frequencies to change. This dangerous fishing activity started perhaps around the 16th century<sup>41</sup>, thus, with 4–5 generations per century, this gives approximately 20–25 generations for which selection could have occurred, which is small. It is thus unclear if selection is acting, but during a too short period of time, or if there is currently no selection at the DRD4 locus.

Another point worth mentioning is that the observed differences between zones could also reflect the effects of occupation on risk attitudes, because of a strong correlation between the living area and the probability of being a fisherman (85% of the sample in the risky area declared their main activity as fishing, vs. 4% in the non-risky area).

There are two potential confounding factors in this study. First, our results could be driven by differences in income between fishermen and non-fishermen. Fishermen indeed are richer than non-fishermen (Table S.1). Because the proportion of missing values represented 29% of our sample (details in Supplementary Materials), imputing missing values was recommended<sup>42</sup>. For this purpose, we used both random forest methods<sup>43</sup> and Lasso regularization for imputations. No correlation was found between income and risk-tolerance in our sample (Table S.9, column 1), although this had been previously found elsewhere<sup>44</sup>. After imputing the missing values with either of these methods and controlling for income, the effect of the 7R allele on risk-tolerance remained significant (Table S.9, columns 3 and 4).

A second potential confounding factor is the experience with commerce or financial activities. Prior work has suggested that market integration of a community can impact decision-making in behavioral economics paradigms<sup>45</sup>. It is possible that the two populations differ in this aspect, but no information is available on this point in our dataset.

Further work should focus on genetic adaptation at other loci, for instance using the loci identified in the GWAS on risk attitudes<sup>4</sup>. Moreover, identifying other solutions for people to cope with risk in risky environments could also be further investigated.

## Methods

A field study was conducted in the Saint-Louis region in Northern Senegal between March 2018 and March 2020. All experiments were conducted in accordance with relevant guidelines and regulations. The protocol (including genotyping) was approved by the Senegalese National Ethics Committee (*Comité National d’Ethique en Recherche en Santé*), and informed consent was obtained from all participants. Behavioral measures were made at the same time as samples were collected for genotyping, so genotypes were not established at the time of measure. Investigators were blind to the behavioral measures during the genotyping.

**Measure of risk-tolerance.** We relied on a standard measure of risk-elicitation task from the experimental economics literature<sup>36</sup>. Instructions were displayed in French (the official written language of the country) and enumerators were present to explain the instructions in Wolof, the vernacular language of Senegal. Participants were invited to choose a card among five. On each card, two amounts were displayed, with an associated color (red or black) and the corresponding amount in coins of XOF 100, in order to have a more visual representation. At the end of the experiment, one ball was randomly drawn by a local child and gains were calculated. The cards ranged from completely risk-free (400 XOF for both balls) to extremely unequal (0 XOF if Red, 1200 XOF if Black). At each new card, the risk is increased, but so is the average amount won. Participants performed the task once. Cards used are displayed in Figure S.3.

**Genotyping.** DRD4 genotyping was done as described in<sup>37</sup>. In short, DNA was collected on FTA paper and extracted according to the manufacturer’s instruction. 506 and 211 samples from the risky and non risky area respectively were of sufficient quality to allow amplification with the appropriate primers. Relevant allele was estimated by the size of the PCR product on a 2% agarose gel.

The variable *7R: additive effect* is equal to the number of 7R alleles for the individual, while the variable *7R: dominance effect* is a dummy variable equal to 1 when the individual possesses only one allele 7R.

Microsatellite genotyping was based on high-throughput sequencing technology (SSRseq). 190 samples of each area were picked up randomly with the sample() function in R. 30 microsatellite tests were designed according to a streamlined SSRseq development workflow described in<sup>46</sup>, of which 29 gave differentiation information (one had only one allele for all individuals). The genomic localization of the 29 microsatellites and their corresponding  $F_{ST}$  between the 2 populations are presented in Table S.8. Details on the design and analysis are in supplementary materials.

**Population genetics.** DRD4 locus was tested for conformity with Hardy-Weinberg (HW) equilibrium using the exact probability test<sup>47</sup>. Deviations from HW equilibrium were measured using the  $F_{IS}$  estimator<sup>48</sup>. DRD4 and microsatellite loci genotypic differentiation between populations was tested for by calculating an unbiased estimate of the P-value of a log-likelihood (G) based exact test<sup>49</sup>, a global test over loci was calculated using Fisher’s method. Population differentiation was measured using the  $F_{ST}$  estimator<sup>48</sup>. Calculations were performed using Genepop R package (V. 1.1.7), based on<sup>50</sup>.

**Statistical analysis.** In the tables and figures, the significance levels are calculated using Student’s *t*-test, comparing the ratio of the effect size to the standard error. For column 5 of Table 2, the estimated equation is the following, where *i* denotes the individual.

$$\begin{aligned} \text{risk tolerance}_i = & \beta_0 + \beta_1 7R : \text{additive effect}_i + \beta_2 7R : \text{dominance effect}_i \\ & + \beta_3 \text{Age}_i + \beta_4 \text{Education}_i + \beta_5 \text{Risky area}_i \\ & + \beta_6 7R : \text{additive effect}_i \times \text{Risky area}_i + \epsilon_i \end{aligned} \quad (1)$$

A significant positive (resp. negative) coefficient for the interaction term ( $\beta_6$ ) would indicate that the additive effect of the 7R allele on risk-tolerance is stronger (resp. weaker) in the risky area.

A power analysis was calculated before the genetic analysis was performed. Using Dagnelie’s formula, with a 520 samples, an allelic difference of 0.1 could be detected with  $p = 0.05$  with a power of  $1 - \beta = 0.90$ .

## Data availability

The data used for this paper are available on the repository of the American Economic Association, under the identifier “openicpsr-179321”, and can be accessed using the following link (login necessary). The sequencing data are registered on the BioProject data base, under the identifier ID PRJNA879442, and are accessible using the following link (embargo until 2023-10-05).

Received: 16 August 2022; Accepted: 22 December 2022

Published online: 12 January 2023

## References

1. Cesarini, D., Dawes, C. T., Johannesson, M., Lichtenstein, P. & Wallace, B. Genetic variation in preferences for giving and risk taking. *Q. J. Econ.* **124**, 809–842 (2009).
2. Ioannidis, J. P. Why most published research findings are false. *PLoS med.* **2**, e124 (2005).

3. Hewitt, J. K. Editorial policy on candidate gene association and candidate gene-by-environment interaction studies of complex traits. *Behav. Genet.* **42**, 1 (2012).
4. Karlsson Linnér, R. *et al.* Genome-wide association analyses of risk tolerance and risky behaviors in over 1 million individuals identify hundreds of loci and shared genetic influences. *Nat. Genet.* **51**, 245–257 (2019).
5. Fiorillo, C. D., Tobler, P. N. & Schultz, W. Discrete coding of reward probability and uncertainty by dopamine neurons. *Science* **299**, 1898–1902 (2003).
6. Bowden-Jones, H. *et al.* Gambling disorder in the UK: Key research priorities and the urgent need for independent research funding. *Lancet Psychiatry* **9**, 321–329 (2022).
7. Van Tol, H. H. *et al.* Cloning of the gene for a human dopamine d4 receptor with high affinity for the antipsychotic clozapine. *Nature* **350**, 610–614 (1991).
8. Gong, S. *et al.* A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. *Nature* **425**, 917–925 (2003).
9. Chang, F.-M., Kidd, J. R., Livak, K. J., Pakstis, A. J. & Kidd, K. K. The world-wide distribution of allele frequencies at the human dopamine d4 receptor locus. *Hum. Genet.* **98**, 91–101 (1996).
10. Asghari, V. *et al.* Modulation of intracellular cyclic amp levels by different human dopamine d4 receptor variants. *J. Neurochem.* **65**, 1157–1165 (1995).
11. Jovanovic, V., Guan, H.-C. & Van Tol, H. Comparative pharmacological and functional analysis of the human dopamine d4. 2 and d4. 10 receptor variants. *Pharmacogenetics* **9**, 561–568 (1999).
12. Van Tol, H. H. *et al.* Multiple dopamine d4 receptor variants in the human population. *Nature* **358**, 149–152 (1992).
13. Oak, J. N., Oldenhof, J. & Van Tol, H. H. The dopamine d4 receptor: One decade of research. *Eur. J. Pharmacol.* **405**, 303–327 (2000).
14. Ding, Y.-C. *et al.* Evidence of positive selection acting at the human dopamine receptor d4 gene locus. *Proc. Natl. Acad. Sci.* **99**, 309–314 (2002).
15. Wang, E. *et al.* The genetic architecture of selection at the human dopamine receptor d4 (drd4) gene locus. *Am. J. Hum. Genet.* **74**, 931–944 (2004).
16. Matthews, L. J. & Butler, P. M. Novelty-seeking drd4 polymorphisms are associated with human migration distance out-of-africa after controlling for neutral population gene structure. *Am. J. Phys. Anthropol.* **145**, 382–389 (2011).
17. Dreber, A. *et al.* The 7r polymorphism in the dopamine receptor d4 gene (drd4) is associated with financial risk taking in men. *Evol. Hum. Behav.* **30**, 85–92 (2009).
18. Kuhnen, C. M. & Chiao, J. Y. Genetic determinants of financial risk taking. *PLoS One* **4**, e4362 (2009).
19. Carpenter, J. P., Garcia, J. R. & Lum, J. K. Dopamine receptor genes predict risk preferences, time preferences, and related economic choices. *J. Risk Uncertain.* **42**, 233–261 (2011).
20. Dreber, A., Rand, D. G., Wernerfelt, N., Montgomery, C. & Malhotra, D. K. *Genetic correlates of economic and social risk taking*, Tech. Rep, SSRN (2012).
21. Anderson, A., Dreber, A. & Vestman, R. Risk taking, behavioral biases and genes: Results from 149 active investors. *J. Behav. Exp. Financ.* **6**, 93–100 (2015).
22. Muda, R. *et al.* The dopamine receptor d4 gene (drd4) and financial risk-taking: Stimulating and instrumental risk-taking propensity and motivation to engage in investment activity. *Front. Behav. Neurosci.* **12**, 34 (2018).
23. Frydman, C., Camerer, C., Bossaerts, P. & Rangel, A. Maa-I carriers are better at making optimal financial decisions under risk. *Proc. R. Soc. B Biol. Sci.* **278**, 2053–2059 (2011).
24. Eisenegger, C. *et al.* Dopamine receptor d4 polymorphism predicts the effect of l-dopa on gambling behavior. *Biol. Psychiatry* **67**, 702–706 (2010).
25. Bakermans-Kranenburg, M. J. & Van Ijzendoorn, M. H. Gene-environment interaction of the dopamine d4 receptor (drd4) and observed maternal insensitivity predicting externalizing behavior in preschoolers. *Dev. Psychobiol. J. Int. Soc. Dev. Psychobiol.* **48**, 406–409 (2006).
26. King, A. P. *et al.* Dopamine receptor gene drd4 7-repeat allele x maternal sensitivity interaction on child externalizing behavior problems: Independent replication of effects at 18 months. *PLoS One* **11**, e0160473 (2016).
27. Lee, J. The effect of the background risk in a simple chance improving decision model. *J. Risk Uncertain.* **36**, 19–41 (2008).
28. Harrison, G. W., List, J. A. & Towe, C. Naturally occurring preferences and exogenous laboratory experiments: A case study of risk aversion. *Econometrica* **75**, 433–458 (2007).
29. Malmendier, U. & Nagel, S. Depression babies: Do macroeconomic experiences affect risk taking?. *Q. J. Econ.* **126**, 373–416 (2011).
30. Beaud, M. & Willinger, M. Are people risk vulnerable?. *Manag. Sci.* **61**, 624–636 (2015).
31. Cameron, L. & Shah, M. Risk-taking behavior in the wake of natural disasters. *J. Hum. Resour.* **50**, 484–515 (2015).
32. Gollier, C. & Pratt, J. W. Risk vulnerability and the tempering effect of background risk. *Econom. J. Econom. Soc.* **64**, 1109–1123 (1996).
33. Eeckhoudt, L., Gollier, C. & Schlesinger, H. Changes in background risk and risk taking behavior. *Econom. J. Econom. Soc.* **64**, 683–689 (1996).
34. Surveillance Côtière, d. S.-L. Bilan des accidents et pertes en vies humaines au niveau de l'embouchure du fleuve senegal de 2003 à 2019. Tech. Rep., Station de surveillance côtière de Saint-Louis (2020).
35. Laloë, F. & Samba, A. *La pêche artisanale au Sénégal: Ressource et stratégie de pêche*. Ph.D. thesis, Paris 11 (1989).
36. Binswanger, H. P. Attitudes toward risk: Experimental measurement in rural india. *Am. J. Agri. Econom.* **62**, 395–407 (1980).
37. Faurie, C. *et al.* Evidence of genotypic adaptation to the exposure to volcanic risk at the dopamine receptor drd4 locus. *Sci. Rep.* **6**, 1–7 (2016).
38. Kluger, A., Siegfried, Z. & Ebstein, R. A meta-analysis of the association between drd4 polymorphism and novelty seeking. *Mol. Psychiatry* **7**, 712–717 (2002).
39. Kirkpatrick, M. *Genes and adaptation: A pocket guide* (Adaptation. Academic Press, San Deigo, CA, 1996).
40. Orr, H. A. & Coyne, J. A. The genetics of adaptation: a reassessment. *Am. Nat.* **140**, 725–742 (1992).
41. Chauveau, J.-P. La pêche piroguère sénégalaise : les leçons de l'histoire. *Revue Mer Spécial*, 10–15 (1984).
42. Jakobsen, J. C., Gluud, C., Wetterslev, J. & Winkel, P. When and how should multiple imputation be used for handling missing data in randomised clinical trials—a practical guide with flowcharts. *BMC Med. Res. Methodol.* **17**, 1–10 (2017).
43. Stekhoven, D. J. & Bühlmann, P. Missforest - non-parametric missing value imputation for mixed-type data. *Bioinformatics* **28**, 112–118 (2012).
44. Hartog, J., Ferrer-i Carbonell, A. & Jonker, N. Linking measured risk aversion to individual characteristics. *Kyklos* **55**, 3–26 (2002).
45. Henrich, J. *et al.* Markets, religion, community size, and the evolution of fairness and punishment. *science* **327**, 1480–1484 (2010).
46. Lepais, O. *et al.* Fast sequence-based microsatellite genotyping development workflow. *PeerJ* **8**, e9085 (2020).
47. Rousset, F. & Raymond, M. Testing heterozygote excess and deficiency. *Genetics* **140**, 1413–1419 (1995).
48. Weir, B. S. & Cockerham, C. C. Estimating f-statistics for the analysis of population structure. *Evolution* **38**, 1358–1370 (1984).
49. Goudet, J., Raymond, M., de Meeüs, T. & Rousset, F. Testing differentiation in diploid populations. *Genetics* **144**, 1933–1940 (1996).
50. Raymond, M. & Rousset, F. Population genetics software for exact tests and ecumenicism. *J. Hered.* **86**, 248–249 (1995).



## Acknowledgements

The authors gratefully acknowledge financial support from GENES, the Key initiatives MUSE Sea & Coast and Investissements d'Avenir (ANR-11-IDEX-0003/Labex Ecodec/ANR-11-LABX-0047). The authors also acknowledge the help in data collection from field research assistants. Microsatellite design and analysis were performed at the PGTB (doi:10.15454/1.5572396583599417E12) with the help of Zoe Delporte.

## Author contributions

G.-J.C., A.M., B.D., O.S. and M.W. collected the data. G.-J.C., C.M., C.F., G.H., M.R. and M.W. analyzed the data. E.C., E.G., C.M. and M.R. performed the micro-satellite analysis. G.-J.C., C.M., C.F., G.H., M.R. and M.W. wrote the paper. All authors reviewed the manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-27002-3>.

**Correspondence** and requests for materials should be addressed to G.-J.C.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023