



HAL
open science

Minimal area of Finsler disks with minimizing geodesics

Marcos Cossarini, Stéphane Sabourau

► **To cite this version:**

Marcos Cossarini, Stéphane Sabourau. Minimal area of Finsler disks with minimizing geodesics. Journal of the European Mathematical Society, 2024, 26, pp.985-1029. 10.4171/JEMS/1339. hal-03961993

HAL Id: hal-03961993

<https://hal.science/hal-03961993v1>

Submitted on 29 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

MINIMAL AREA OF FINSLER DISKS WITH MINIMIZING GEODESICS

MARCOS COSSARINI AND STÉPHANE SABOURAU

ABSTRACT. We show that the Holmes–Thompson area of every Finsler disk of radius r whose interior geodesics are length-minimizing is at least $\frac{6}{\pi}r^2$. Furthermore, we construct examples showing that the inequality is sharp and observe that the equality case is attained by a non-rotationally symmetric metric. This contrasts with Berger’s conjecture in the Riemannian case, which asserts that the round hemisphere is extremal. To prove our theorem we discretize the Finsler metric using random geodesics. As an auxiliary result, we show that the integral geometry formulas of Blaschke and Santaló hold on Finsler manifolds with almost no trapped geodesics.

CONTENTS

1. Introduction	2
2. Finsler metrics and Holmes–Thompson volume	4
3. Integral geometry in Finsler manifolds with almost no trapped geodesics	7
4. Discretization of Finsler surfaces	14
5. Minimal area of disks: from discrete to Finsler metrics	21
6. Quasi wall systems and interval families	23
7. Inadmissible configurations in a minimal simple disk	25
8. Pairs of adjacent arcs	28
9. Proof of the discrete area lower bound	30
10. Simple discrete metric disks of minimal area	31
11. Hilbert’s fourth problem and extremal Finsler disks	33
12. Appendix A: Differentiability of distance realizing curves on Finsler surfaces with boundary	39
13. Appendix B: A weak law of large numbers for mostly independent random variables	42
14. Appendix C: A version of Levi’s extension lemma	45
References	46

2020 *Mathematics Subject Classification*. Primary 53C23; Secondary 53C60, 53C65.

Partially supported by the Bézout Labex (ANR-10-LABX-58) and the ANR project Min-Max (ANR-19-CE40-0014).

1. INTRODUCTION

Isoembolic inequalities on Riemannian manifolds are curvature-free volume estimates in terms of the injectivity radius. The first sharp isoembolic inequality valid in all dimension is due to Berger [Ber80] who showed that the volume of every closed Riemannian n -manifold M satisfies

$$\text{vol}(M) \geq \alpha_n \left(\frac{\text{inj}(M)}{\pi} \right)^n \quad (1.1)$$

where α_n is the volume of the canonical n -sphere. Furthermore, equality holds if and only if M is isometric to a round sphere. The two-dimensional case was proved earlier in [Ber76].

A long standing conjecture in Riemannian geometry also due to Berger asserts that every ball $B(r)$ of radius $r \leq \frac{1}{2} \text{inj}(M)$ in a closed Riemannian n -manifold M satisfies

$$\text{vol } B(r) \geq \frac{\alpha_n}{2} \left(\frac{2r}{\pi} \right)^n$$

with equality if and only if $B(r)$ is isometric to a round hemisphere of (intrinsic) radius r . This can be viewed as a local version of the sharp isoembolic inequality (1.1). This conjecture is open even in the two-dimensional case where the previous inequality can be written as

$$\text{area } D(r) \geq \frac{8}{\pi} r^2. \quad (1.2)$$

An account on isoembolic inequalities and Berger's conjecture is given in [CK03, §6]. A non-sharp volume estimate $\text{vol } B(r) \geq c_n r^n$ was established by Berger [Ber76], [Ber77] for $n = 2$ or 3 , and by Croke [Cro80, Proposition 14] for every n . The conjecture (with a sharp constant) is satisfied for metrics of the form $ds^2 = dr^2 + f(r, \theta)^2 d\theta^2$ in polar coordinates when $n \geq 3$; see [Cro83]. In [Cro84], Croke also showed that the optimal inequality (1.2) holds true on average over all balls $B(r)$ of M . In the two-dimensional case, the best general estimate $\text{area } D(r) \geq \frac{8-\pi}{2} r^2$ can be found in [Cro09]. The optimal lower bound (1.2) on the area of $D(r)$ has recently been obtained by Chambers–Croke–Liokumovich–Wen in the two-dimensional case for every $r \leq \frac{1}{2} \text{conv}(M)$, where $\text{conv}(M)$ is the convexity radius of M ; see [Cha+17]. Note that in this case, the radius r of $D(r)$ does not exceed $\frac{1}{4} \text{inj}(M)$ since $\text{conv}(M) \leq \frac{1}{2} \text{inj}(M)$.

The condition that $r \leq \frac{1}{2} \text{inj}(M)$ can be relaxed by requiring instead that every interior geodesic in $B(r)$ is length-minimizing. The results of [Ber76], [Ber77] and [Cro80, Proposition 14], for instance, still hold under this more general condition.

In this article, we consider the case of self-reverse Finsler metric disks whose interior geodesics are length-minimizing. (A precise definition of Finsler metrics and area can be found in Section 2.) It is natural to expect that the inequality (1.2) holds in this setting. This is the case for

isosystolic inequalities on the projective plane, where the canonical round metric minimizes the systolic area among both Riemannian and Finsler metrics; see [Iva02], [Iva11]. However, we show that the round hemisphere is not area minimizing among Finsler metric disks of the same radius whose interior geodesics are length-minimizing. More precisely, we establish a sharp isoeombolic inequality for Finsler metrics in the two-dimensional case under the assumption that every interior geodesic is length-minimizing. We observe that the extremal metric is not Riemannian and, surprisingly, not even rotationally symmetric.

Before stating our main result, let us introduce the following definition.

Definition 1.1. A *Finsler disk* D of radius r with minimizing interior geodesics is a disk with a Finsler metric such that

- every interior point of D is at distance less than r from a specified center point O ;
- every point of ∂D is at distance exactly r from O ;
- every interior geodesic of D is length-minimizing.

For instance, a ball of radius r on a complete Finsler plane with no conjugate points is a Finsler disk of radius r with minimizing interior geodesics.

The optimal version of Berger’s conjecture for Finsler surfaces with self-reverse metric is given by the following result.

Theorem 1.2. *Let D be a self-reverse Finsler metric disk D of radius r with minimizing interior geodesics. Then the Holmes–Thompson area of D satisfies*

$$\text{area}(D) \geq \frac{6}{\pi} r^2.$$

Furthermore, the inequality is optimal.

We emphasize that the disks in this theorem are not necessarily convex.

The lower bound is attained by a non-smooth metric. This extremal metric is described in Section 11, where we also show how to smooth it out, obtaining an almost extremal projective Finsler metric. We remark that none of these metrics is rotationally symmetric.

The proof of Theorem 1.2 and the construction of extremal and almost extremal metrics occupy the whole article. The approach, based on a discretization of the metric (*cf.* [Cos18]), is fairly robust and new in this context. The article is organized as follows.

In Section 2, we recall the notions of Finsler manifolds, their Holmes–Thompson measure, and their geodesics described from the Hamiltonian point of view.

In Section 3, we establish versions of the integral geometry formulas of Blaschke and Santaló that are valid for manifolds with almost no trapped geodesics. In the case of a disk as in Theorem 1.2, the formulas say that the length of a curve in the disk is proportional to the expected number of

intersections with a random geodesic, and the area of a region is proportional to the expected length of the intersection with a random geodesic.

In Section 4, we introduce the notion of a quasi wall system on a surface, generalizing the wall systems studied in [Cos18]. A quasi wall system on a surface is a 1-dimensional submanifold satisfying certain conditions. It determines a discrete metric, according to which the length of a curve is its number of intersections with the quasi wall system, and the area of the surface is the number of self-intersections of the quasi wall system. We show how to approximate a self-reverse Finsler metric with minimizing geodesics by a quasi wall system consisting of random geodesics. To prove the approximation properties we use the integral geometry formulas to compute the expected values of discrete length and area, and then we apply the law of large numbers.

In Section 5, we use this approximation result to show that Theorem 1.2 follows from an analogous theorem on simple discrete metric disks.

Sections 6, 7, 8 and 9 are devoted to the proof of this discrete theorem. The proof is based on identifying certain configurations on a quasi wall system and operating on this configurations in order to transform a simple discrete disk into a new one of less area. When the disk has minimum area, none of this configurations is present, and this implies that the quasi wall system is of a special kind where we can compute a lower bound for the area.

In Section 10, we construct a simple discrete disk of minimal discrete area and show that it is unique up to isotopy.

Finally, in Section 11, we apply Busemann's construction of projective metrics in relation with Hilbert's fourth problem to approximate the extremal simple discrete metric with a Finsler metric with minimizing interior geodesics.

Acknowledgment. The first author thanks the Laboratoire d'Analyse et de Mathématiques Appliquées (LAMA) at the Université Gustave Eiffel and the Groupe Troyanov at the École Polytechnique Fédérale de Lausanne for hosting him as a postdoc while this work was done. The second author would like to thank the Fields Institute and the Department of Mathematics at the University of Toronto, where part of this work was accomplished, for their hospitality.

2. FINSLER METRICS AND HOLMES–THOMPSON VOLUME

In this section, we recall basic definitions of Finsler geometry.

2.1. Finsler metrics. Let us recall the definition of a Finsler metric.

Definition 2.1. A *Finsler metric* on a smooth manifold M is a continuous function $F : TM \rightarrow [0, \infty)$ on the tangent bundle TM of M satisfying the following properties (here, $F_x := F|_{T_x M}$ for short):

- (1) Positive homogeneity: $F_x(tv) = t F_x(v)$ for every $v \in T_x M$ and $t \geq 0$.
- (2) Subadditivity: $F_x(v + w) \leq F_x(v) + F_x(w)$ for every $v, w \in T_x M$.
- (3) Positive definiteness: $F_x(v) > 0$ for every nonzero $v \in T_x M$.
- (4) Smoothness: F is smooth outside the zero section.
- (5) Strong convexity: for any two linearly independent vectors $v, w \in T_x M$, the Hessian value $q_v(w) = \left. \frac{d^2}{dt^2} \right|_{t=0} F_x(v + tw)$ is strictly positive.

Additionally, a Finsler metric F may be or not be

- (6) Self-reverse: $F_x(-v) = F_x(v)$ for every $v \in T_x M$.

Equivalently, one could define a Finsler metric by replacing (3) and (5) with the condition that for every nonzero vector $v \in TM$, the Hessian of F^2 at v is positive definite; see [Cos20].

In each tangent space $T_x M$, the unit ball and unit sphere determined by the norm F_x are

$$B_x M = \{v \in T_x M \mid F_x(v) \leq 1\} \text{ and } U_x M = \{v \in T_x M \mid F_x(v) = 1\}.$$

Similarly, in the cotangent space $T_x^* M$, the norm F_x^* dual to F_x determines a unit co-ball $B_x^* M$ and a unit co-sphere $U_x^* M$.

Remark 2.2. To handle technical details in case M has nonempty boundary, we extend the metric F to a manifold $M^+ \supseteq M$, of the same dimension as M but without boundary.

2.2. Length, geodesics and distance realizing arcs.

Definition 2.3. Let M be a manifold with a Finsler metric F . The *length* of a piecewise- C^1 curve $\gamma : I \rightarrow M$ is defined as the integral of its *speed* $F(\gamma'(t))$, that is,

$$\text{length}(\gamma) = \int_I F(\gamma'(t)) dt \tag{2.1}$$

and the *distance* $d_F(x, y)$ between two points x and y in M is the infimum length of a curve γ in M joining x to y .

A *distance realizing curve* is a curve $\gamma : I \rightarrow M$ such that

$$d_F(\gamma(t), \gamma(t')) = t' - t$$

for every $t < t'$.

A *geodesic* of M is a smooth, unit-speed curve $\gamma : I \rightarrow M$ that is extremal for the length functional. In case M has boundary, the extremality is defined by considering variations in M^+ , see Remark 2.2. Thus the geodesics of M are the geodesics of M^+ that are contained in M . Equivalently, the geodesics of M are the unit-speed curve curves that satisfy the Euler–Lagrange equation for the length functional; see Definition 2.6 below for an explicit equation in terms of momentum.

In a compact connected Finsler manifold, every pair of points are joined by a distance realizing arc.¹ A distance realizing arc contained in the interior of M is necessarily a geodesic and is therefore smooth. However, a distance realizing arc of M does not necessarily lie in the interior of M , even if its endpoints do. Still, if the manifold is two-dimensional, then every distance realizing arc is C^1 and has unit speed; see Theorem 12.1. Thus, in a compact Finsler surface, every pair of points x, y are joined by a C^1 arc of length $d_F(x, y)$.

2.3. Symplectic structure on the cotangent bundle. Recall also some definitions about the geodesic flow of a Finsler manifold from the Hamiltonian viewpoint; see [Arn89, Chap. 7–9], [Álv06] and [Cos20].

Definition 2.4. Let M be a manifold. The *tautological one-form* α_M on T^*M is defined as

$$\alpha_M|_{\xi}(V) = \xi(d\pi_{\xi}(V))$$

for every $\xi \in T^*M$ and $V \in T_{\xi}T^*M$, where $\pi : T^*M \rightarrow M$ is the canonical projection. The *standard symplectic form* ω_M on T^*M is given by

$$\omega_M = d\alpha_M.$$

Using canonical coordinates (x_i, ξ_i) on T^*M , these forms can be expressed as

$$\alpha_M = \sum_i \xi_i dx_i, \quad \omega_M = \sum_i d\xi_i \wedge dx_i. \quad (2.2)$$

Definition 2.5. Let (M, F) be a Finsler manifold. The *Legendre map*

$$\mathcal{L} : UM \rightarrow U^*M,$$

is defined as follows: the image of a unit vector $v \in U_xM$ is the unique unit covector $\xi \in U_x^*M$ such that $\xi(v) = 1$. Since F is strongly convex, the Legendre map is a diffeomorphism. Its inverse is the Legendre map associated to the dual metric F^* on T^*M , which is also strongly convex. The unit covectors will also be referred to as *momentums*. The *Hamiltonian lift* of a unit-speed curve γ in M is the curve $t \mapsto \mathcal{L}(\gamma'(t))$ in U^*M .

Definition 2.6. The *cogeodesic vector field* Z on U^*M is defined by the equations

$$\begin{aligned} \iota_Z(\omega_M|_{U^*M}) &= 0 \\ \iota_Z(\alpha_M) &= 1 \end{aligned}$$

Its integral curves are the Hamiltonian lifts of the geodesics in M ; see [Cos20].

It follows from the Cartan formula that the forms α_M and ω_M restricted to U^*M are invariant under the cogeodesic flow.

¹A proof for more general, complete self-reverse metrics is given [Gro07, §1.12]; see also [Men14, Theorem 9.1] for directed metrics.

2.4. Holmes–Thompson volume. We will consider the following notion of volume.

Definition 2.7. The *Holmes–Thompson volume* of a Finsler n -manifold M is defined as the symplectic volume of its unit co-ball bundle $B^*M \subseteq T^*M$, divided by the volume ϵ_n of the Euclidean unit ball in \mathbb{R}^n . That is,

$$\text{vol}(M) = \frac{1}{\epsilon_n} \int_{B^*M} \frac{1}{n!} \omega_M^n \tag{2.3}$$

where ω_M is the standard symplectic form on T^*M and $\frac{1}{n!} \omega_M^n = \frac{1}{n!} \omega_M \wedge \dots \wedge \omega_M$ is the corresponding volume form.

The Holmes–Thompson volume of a Finsler manifold is bounded from above by its Hausdorff measure, with equality if and only if the metric is Riemannian; see [Dur98]. The factor $\frac{1}{\epsilon_n}$ ensures that for Riemannian metrics, Holmes–Thompson volume agrees with the conventional definition.

3. INTEGRAL GEOMETRY IN FINSLER MANIFOLDS WITH ALMOST NO TRAPPED GEODESICS

The goal of this section is to establish versions of two classical formulas in integral geometry, namely the formulas of Blaschke [Bla35] and Santaló [San52; San76], which are in turn generalizations for manifolds of the classical Crofton formulas on the Euclidean plane. In [ÁB06], Blaschke’s formula is proved for Finsler manifolds whose space of geodesics is a smooth manifold. For non-convex Finsler disks with minimizing geodesics, the space of geodesics may not be a smooth manifold so a different version of the formula is required.

Definition 3.1. Let M be a Finsler n -manifold with nonempty boundary. A *traversing geodesic* of M is a maximal geodesic $\gamma : [0, \ell(\gamma)] \rightarrow M$ which does not intersect ∂M , except at its endpoints where it meets the boundary transversely. The Finsler manifold M has *almost no trapped geodesics* if for almost every unit tangent vector $v \in UM$, the maximal geodesic γ_v defined by $\gamma'_v(0) = v$ reaches the boundary of M in the future and in the past, that is, $\gamma_v(t) \in \partial M$ for some $t \geq 0$ and some $t \leq 0$.

For instance, a compact Finsler manifold with minimizing interior geodesics has almost no trapped geodesics. Another example is obtained by taking a closed Finsler manifold with ergodic geodesic flow and removing a smoothly bounded nonempty open set.

As we will explain below, the space Γ of traversing geodesics of M is a $(2n - 2)$ -dimensional manifold admitting a natural symplectic structure, whose corresponding natural volume measure is denoted by μ_Γ ; see Definition 3.6.

Theorem 3.2 (Blaschke’s formula). *Let M be a Finsler n -manifold with almost no trapped geodesics. Then the Holmes–Thompson volume of an*

immersed hypersurface $N \subseteq M$ is equal to

$$\text{vol}_{n-1}(N) = \frac{1}{2 \epsilon_{n-1}} \int_{\gamma \in \Gamma} \#(\gamma \cap N) \, d\mu_{\Gamma}(\gamma) \quad (3.1)$$

where $\#(\gamma \cap N)$ is the number of times that γ intersects N .

Similarly, the Holmes–Thompson volume of a co-oriented immersed hypersurface $N \subseteq M$ is equal to

$$\text{vol}_{n-1}(N) = \frac{1}{\epsilon_{n-1}} \int_{\gamma \in \Gamma} \#(\gamma \cap^+ N) \, d\mu_{\Gamma}(\gamma) \quad (3.2)$$

where $\#(\gamma \cap^+ N)$ is the number of times that γ intersects N transversely in the positive direction.

In equation (3.1), we can restrict the integral to geodesics $\gamma \in \Gamma$ which are transverse to the hypersurface N since the geodesics $\gamma \in \Gamma$ which are tangent to N form a subset of zero measure; see Proposition 3.7.(3).

Remark 3.3. Since every geodesic of Γ intersects ∂M at its two endpoints exactly (but only once in positive direction), we derive from (3.2) that the total measure of the space Γ is

$$\mu_{\Gamma}(\Gamma) = \epsilon_{n-1} \text{vol}_{n-1}(\partial M)$$

In particular, if M is compact, then $\mu_{\Gamma}(\Gamma) < \infty$.

Theorem 3.4 (Santaló’s formula). *Let M be a Finsler n -manifold with almost no trapped geodesics. Then the Holmes–Thompson volume of a smoothly-bounded domain $D \subseteq M$ is equal to*

$$\text{vol}_n(D) = \frac{1}{n \epsilon_n} \int_{\gamma \in \Gamma} \text{length}(\gamma \cap D) \, d\mu_{\Gamma}(\gamma). \quad (3.3)$$

In the case of Finsler surfaces with self-reverse metric, the Blaschke and Santaló formulas specialize as follows.

Corollary 3.5. *Let M be a self-reverse Finsler metric surface with almost no trapped geodesics. Then the length of any immersed curve c in M is*

$$\text{length}(c) = \frac{1}{4} \int_{\gamma \in \Gamma} \#(\gamma \cap c) \, d\mu_{\Gamma}(\gamma), \quad (3.4)$$

and the Holmes–Thompson area of any smoothly-bounded domain $D \subseteq M$ is

$$\text{area}(D) = \frac{1}{2\pi} \int_{\gamma \in \Gamma} \text{length}(\gamma \cap D) \, d\mu_{\Gamma}(\gamma) \quad (3.5)$$

$$= \frac{1}{8\pi} \iint_{(\gamma_0, \gamma_1) \in \Gamma \times \Gamma} \#(\gamma_0 \cap \gamma_1 \cap D) \, d\mu_{\Gamma}(\gamma_0) \, d\mu_{\Gamma}(\gamma_1). \quad (3.6)$$

The equation (3.6), obtained from (3.5) and (3.4), will be called the Santaló+Blaschke formula. In deducing this formula, we use the hypothesis that the metric is self-reverse when we equate the length of a geodesic with

its Holmes–Thompson measure. In general, the Holmes–Thompson measure of a curve is the average of its forward and backward lengths.

The rest of this section is dedicated to describing the symplectic structure on Γ and proving Theorems 3.2 and 3.4.

3.1. Symplectic manifold of traversing geodesics. Let M be a Finsler n -manifold. Recall that Γ is the space of traversing geodesics of M . This space Γ is a $(2n - 2)$ -dimensional manifold parameterized by the initial vectors $\gamma'(0) \in UM|_{\partial M}$ of the geodesics $\gamma : [0, \ell(\gamma)] \rightarrow M$ of Γ . Note that the length $\ell(\gamma)$ depends smoothly on $\gamma \in \Gamma$.

Definition 3.6. Define the open subset of U^*M

$$U_\Gamma^*M = \{\mathcal{L}(\gamma'(t)) \in U^*M \mid \gamma \in \Gamma, t \in [0, \ell(\gamma)]\}$$

consisting of the momentums of the traversing geodesics of M . Note that U_Γ^*M is a Z -invariant open subset of U^*M .

Consider the surjective submersion

$$\pi_\Gamma : U_\Gamma^*M \rightarrow \Gamma \tag{3.7}$$

taking any momentum $\xi \in U_\Gamma^*M$ to the geodesic $\gamma \in \Gamma$ that it generates. The fibers of π_Γ are the Z -orbits corresponding to the traversing geodesics of M .

There exists a unique 2-form ω_Γ on Γ such that

$$\pi_\Gamma^* \omega_\Gamma = \omega_M|_{U_\Gamma^*M}.$$

This follows from the invariance of the 2-form $\omega_M|_{U_\Gamma^*M}$ under the cogeodesic flow, and the fact that this form vanishes in the direction of Z ; see Definition 2.6. The form ω_Γ is symplectic and determines on Γ a smooth volume measure μ_Γ given by

$$d\mu_\Gamma = \frac{1}{(n-1)!} |\omega_\Gamma^{n-1}|. \tag{3.8}$$

3.2. Non-traversing geodesics are negligible. We will need the following result in order to establish our versions of Blaschke’s and Santaló’s formulas. This feature is not required in the previous versions and necessitates the manifold to have almost no trapped geodesics.

Recall that a subset A of a manifold X is *negligible* in X if the image of A in any local chart of X has zero measure.

Proposition 3.7.

- (1) *The complement of the open subset $U_\Gamma^*M \subseteq U^*M$ is negligible in U^*M .*
- (2) *Given a hypersurface $H \subseteq U^*M$ transverse to Z , the complement of $H \cap U_\Gamma^*M$ is negligible in H .*
- (3) *The set of geodesics $\gamma \in \Gamma$ tangent to an immersed hypersurface of M or passing through an immersed submanifold of M of codimension > 1 has zero measure in Γ .*

Proof. To avoid technical problems we extend the Finsler metric to an open manifold M^+ ; see Remark 2.2. This ensures the cogeodesic flow $(\xi, t) \mapsto Z^t(\xi)$ is defined on an open domain.

By definition of Γ , the complement $U^*M \setminus U_\Gamma^*M$ is formed of momentums of two types. First, momentums of U^*M that correspond to geodesics of M with at least one end trapped in M . These momentums form a negligible set since M has almost no trapped geodesics; see Definition 3.1. Second, momentums of U^*M corresponding to geodesics tangent to the boundary ∂M . These momentums are of the form $Z^t(\xi)$, where Z^t is the cogeodesic flow and ξ is the Legendre image of a unit vector v tangent to ∂M . These unit vectors form a manifold $U\partial M$ of dimension $2n - 3$. Thus, by Sard's theorem, the map from an open subset of $U\partial M \times \mathbb{R}$ to U^*M defined by $(v, t) \mapsto Z^t(\mathcal{L}(v))$ has negligible image in U^*M . Having considered both types of momentums, we conclude that the complement $U^*M \setminus U_\Gamma^*M$ is negligible in U^*M .

For the second point, simply observe that if A is a Z -invariant negligible subset of U^*M and H is a hypersurface of U^*M transverse to Z , then $A \cap H$ is negligible in H . Apply this property to $A = U^*M \setminus U_\Gamma^*M$ to conclude.

The proof of the third point is similar to the proof of the first point and relies on Sard's theorem. \square

3.3. Manifold of positive momentums across a hypersurface. We will need the following notion in the proof of Blaschke's formula.

Definition 3.8. Let N be a co-oriented embedded hypersurface in a Finsler manifold M . (For example, we can have $N = \partial M$ co-oriented so that inwards-pointing vectors are positive.) Denote by $C^*N \subseteq U^*M|_N$ the *manifold of momentums crossing positively* the hypersurface N , that is, the momentums corresponding under the Legendre map to unit vectors transverse to N pointing in the positive direction according to the co-orientation of N . Note that C^*N is an open subset of $U^*M|_N$ and therefore a differentiable manifold, with the structure of an open ball bundle over N .

Consider the *restriction map*

$$\begin{aligned} \rho_N : \quad C^*N &\longrightarrow (B^*N)^\circ \\ \xi \in T_x^*M &\longmapsto \xi|_{T_x N} \end{aligned}$$

to the interior $(B^*N)^\circ$ of the unit co-ball bundle B^*N of N .

The following statement can be found in [ÁB06, Lemma 5.4]. We simply provide the details of the proof.

Lemma 3.9. *The space C^*N is a symplectic submanifold of T^*M and the restriction map*

$$\rho_N : (C^*N, \omega_M) \rightarrow ((B^*N)^\circ, \omega_N)$$

is a symplectomorphism. Thus,

$$\rho_N^* \omega_N = \omega_M|_{C^*N}.$$

Proof. Let $\xi \in C^*N$ with basepoint $x \in N$. By definition, the norm of ξ is 1, so the norm of its restriction ξ' to T_xN is at most 1. Furthermore, by strong convexity of F_x^* , the linear form ξ attains its maximum only at its Legendre-dual unit vector, which is positive and thus not contained in T_xN . Therefore, $\|\xi'\| < 1$ and the restriction map ρ_N takes values in $(B^*N)^\circ$.

To see that ρ_N is a diffeomorphism, we employ local coordinates $(x_i)_{1 \leq i \leq n}$ in M so that the hypersurface N is given by the equation $x_n = 0$. Let $(x_i, v_i)_i$ and $(x_i, \xi_i)_i$ be the corresponding coordinates in TM and T^*M . In terms of these coordinates, the operator ρ_N acts by suppressing the last coefficient, that is, if $\xi = (\xi_i)_{1 \leq i \leq n}$, then $\xi' = (\xi_i)_{1 \leq i \leq n-1}$. Hence ρ_N is smooth.

To prove that ρ_N is bijective, consider a covector $\xi' = (\xi_i)_{1 \leq i \leq n-1} \in (B_x^*N)^\circ$ and denote its norm $c = \|\xi'\| < 1$. The covectors $\xi \in T_x^*M$ such that $\xi|_{T_xN} = \xi'$ are of the form $\xi^t = (\xi_1, \dots, \xi_{n-1}, t)$ with $t \in \mathbb{R}$. Consider the function $t \mapsto \|\xi^t\|$, where $\|\cdot\|$ is the norm F_x^* on T_x^*M that is dual to F_x . This function is bounded below by c , and by the Hahn–Banach theorem, this lower bound is attained at some $t_0 \in \mathbb{R}$. Furthermore, since the norm F_x^* is strongly convex, the set of values of t such that $\|\xi^t\| \leq 1$ is a compact interval $[t_-, t_+]$ that contains t_0 in its interior, and $\|\xi^t\| = 1$ if and only if $t = t_\pm$. Thus we are left with two candidates ξ^{t_\pm} that are the only unit covectors ξ whose restriction to T_xN is ξ' .

We claim that ξ^{t_+} is positive and ξ^{t_-} is negative. Indeed, when $t = t_0$, the covector ξ^t , as a function $B_xM \rightarrow \mathbb{R}$, is bounded above by c . As t increases towards t_+ , the values of ξ^t on the positive side of B_xM (where $v_n > 0$) increase, and the values on the negative side decrease, because the coefficient ξ_n increases. Thus any functional ξ^t with $t > t_0$, restricted to the ball B_xM , must attain its maximum value $\|\xi^t\| > c$ on a positive vector. This implies that ξ^t is positive if $t > t_0$, and similarly, ξ^t is negative if $t < t_0$. We conclude that ξ^{t_+} is the only positive unit covector ξ whose restriction to T_xN is ξ' . This proves that ρ_N is bijective. Additionally, t_+ depends smoothly on ξ' by the implicit function theorem. This finishes the proof that the restriction map $\rho_N : C^*N \rightarrow (B^*N)^\circ$ is a diffeomorphism.

Let us show that $\rho_N^*\alpha_N = \alpha_M|_{C^*N}$. In canonical coordinates, the tautological 1-form α_M on T^*M is written as $\alpha_M = \sum_{i=1}^n \xi_i dx_i$. In restricting to C^*N , the last term vanishes because $x_n = 0$ on N , thus the restricted form can be written as $\alpha_M|_{C^*N} = \sum_{i=1}^{n-1} \xi_i dx_i$. On the other hand, the tautological 1-form of N is $\alpha_N = \sum_{i=1}^{n-1} \xi_i dx_i$, and this expression is unchanged by the pullback ρ_N^* since the map $\rho_N : C^*N \rightarrow (B^*N)^\circ$ acts simply by suppressing the coordinate ξ_n . We conclude that $\rho_N^*\alpha_N = \alpha_M|_{C^*N}$. Taking the exterior differential of this expression, we obtain $\rho_N^*\omega_N = \omega_M|_{C^*N}$. This implies that C^*N is a symplectic submanifold of T^*M . \square

3.4. Coarea formula and fiber integration. In the proofs of Blaschke’s and Santaló’s formulas, we will need the following version of the coarea formula; see [Die72, (16.24.8)] (see also [Fed69, Theorem 3.2.3] and [BBI01, Theorem 5.5.8] when $n = m$).

Lemma 3.10. *Let $\pi : X \rightarrow Y$ be a submersion between two oriented manifolds of dimension n and m with $n \geq m$. Let α and β be two differential forms on X and Y of degree $n - m$ and m . Then*

$$\int_X \alpha \wedge \pi^* \beta = \int_{y \in Y} \left(\int_{\pi^{-1}(y)} \alpha \right) \beta$$

where $\pi^{-1}(y)$ is endowed with the orientation induced by π from the orientations of X and Y .

In particular, for $n = m$ and $\alpha = 1$, we have

$$\int_X \pi^* \beta = \int_{y \in Y} \#(\pi^{-1}(y)) \beta. \quad (3.9)$$

3.5. Proof of the Blaschke formula. We can now proceed to the proof of Blaschke's formula (3.1).

Proof of Theorem 3.2. We will follow the proof given in [ÁB06, Theorem 5.2] under the extra assumption that the space of oriented geodesics on M is a manifold.

The Blaschke formula (3.1) for a non-cooriented hypersurface N can be deduced from the co-oriented version (3.2) by taking the co-oriented double cover of N . Therefore it is sufficient to prove the latter formula. Furthermore, every immersed hypersurface can be decomposed into a disjoint union of embedded hypersurfaces up to a negligible set. Therefore it is sufficient to prove (3.2) for a co-oriented embedded hypersurface N .

By definition of the Holmes–Thompson volume, see (2.3), we have

$$\begin{aligned} \text{vol}_{n-1}(N) &= \frac{1}{\epsilon_{n-1}(n-1)!} \int_{B^*N} \omega_N^{n-1} \\ &= \frac{1}{\epsilon_{n-1}(n-1)!} \int_{C^*N} \omega_M^{n-1} \end{aligned}$$

where the second equality follows from Lemma 3.9.

Now, apply Proposition 3.7.(2) with $H = C^*N \subseteq U^*M$. It follows that $C^*N \cap U_\Gamma^*M$ has full measure in C^*N . Thus,

$$\int_{C^*N} \omega_M^{n-1} = \int_{C^*N \cap U_\Gamma^*M} \omega_M^{n-1}.$$

Consider the map $\pi : C^*N \cap U_\Gamma^*M \rightarrow \Gamma$ taking a unit momentum of M based at N pointing in a positive direction (with respect to the co-orientation of N) to the traversing geodesic it generates. Apply the fiber integration formula (3.9) to this map with $\beta = \omega_\Gamma^{n-1}$. This yields the relation

$$\int_{C^*N \cap U_\Gamma^*M} \omega_M^{n-1} = \int_{\gamma \in \Gamma} \#(\gamma \cap^+ N) \omega_\Gamma^{n-1}.$$

where $\#(\gamma \cap^+ N)$ is the number of times that γ crosses N transversely in the positive sense (as determined by the co-orientation of N). Taking into

account the definition of μ_Γ by equation (3.8), Blaschke's formula follows. \square

3.6. Proof of the Santaló formula. We will need the following proposition expressing the Holmes–Thompson volume of a manifold as an integral over the bundle of dual unit spheres (instead of dual unit balls).

Proposition 3.11. *The Holmes–Thompson volume of a Finsler n -manifold M is equal to*

$$\text{vol}(M) = \frac{1}{\epsilon_n n!} \int_{U^*M} \alpha_M \wedge \omega_M^{n-1}.$$

Proof. Observe that $d(\alpha_M \wedge \omega_M^{n-1}) = \omega_M^n$ and $\partial B^*M = U^*M \cup B^*M|_{\partial M}$. By Stokes' theorem, we have

$$\begin{aligned} \text{vol}_n(M) &= \frac{1}{\epsilon_n n!} \int_{B^*M} d(\alpha_M \wedge \omega_M^{n-1}) \\ &= \frac{1}{\epsilon_n n!} \int_{U^*M} \alpha_M \wedge \omega_M^{n-1} + \frac{1}{\epsilon_n n!} \int_{B^*M|_{\partial M}} \alpha_M \wedge \omega_M^{n-1} \end{aligned} \quad (3.10)$$

Now, recall that the tangent space to $B^*M|_{\partial M}$ at (x, ξ) decomposes as

$$T_{(x,\xi)}B^*M|_{\partial M} \simeq T_x\partial M \oplus T_x^*M$$

where the horizontal space $T_x\partial M$ is of dimension $n - 1$ and the vertical space T_x^*M is of dimension n . Note that the one-form α_M vanishes on the vertical space T_x^*M and the two-form ω_M vanishes at bi-vectors formed of two horizontal or two vertical vectors. This follows from the coordinate expression (2.2). The $(2n - 1)$ -form $\alpha_M \wedge \omega_M^{n-1}$ evaluated at (u_1, \dots, u_{2n-1}) , where $n - 1$ vectors u_i are horizontal and n vectors u_i are vertical, can be written as a sum of terms of the form

$$\pm \alpha_M(u_{\sigma(1)}) \cdot \omega_M(u_{\sigma(2)}, u_{\sigma(3)}) \cdots \omega_M(u_{\sigma(2n-2)}, u_{\sigma(2n-1)}) \quad (3.11)$$

where σ is a permutation. If $u_{\sigma(1)}$ is vertical then the factor $\alpha_M(u_{\sigma(1)})$ is equal to zero. If $u_{\sigma(1)}$ is horizontal then there are only $n - 2$ horizontal vectors (and n vertical ones) among the remaining vectors, which implies that one of the factors $\omega_M(u_{\sigma(2k)}, u_{\sigma(2k+1)})$ has two horizontal vectors and therefore vanishes. In both cases, the term (3.11) vanishes. Therefore, the $(2n - 1)$ -form $\alpha_M \wedge \omega_M^{n-1}$ vanishes on $B^*M|_{\partial M}$. Thus, the second term in (3.10) vanishes as required. \square

Now, let us prove Santaló's formula (3.3).

Proof of Theorem 3.4. Recall that $\omega_M = \pi_\Gamma^* \omega_\Gamma$, see Definition 3.6, and that U_Γ^*M has full measure in U^*M , see Proposition 3.7.(1). By Proposition 3.11, we have

$$\text{vol}_n(D) = \frac{1}{\epsilon_n n!} \int_{U^*D \cap U_\Gamma^*M} \alpha_M \wedge \pi_\Gamma^* \omega_\Gamma^{n-1}.$$

By Lemma 3.10, integrating along the fibers of the submersion $\pi : U^*D \cap U_\Gamma^*M \rightarrow \Gamma$ induced by π_Γ , see (3.7), we obtain

$$\text{vol}_n(D) = \frac{1}{\epsilon_n n!} \int_{\gamma \in \Gamma} \left(\int_{\pi^{-1}(\gamma)} \alpha_M \right) \omega_\Gamma^{n-1}.$$

Since all the fibers $\pi^{-1}(\gamma) = \{\mathcal{L}(\gamma'(t)) \in U^*M \mid t \in [0, \ell(\gamma)]\} \cap U^*D$ are tangent to the cogeodesic vector field Z on U^*M and $\alpha_M(Z) = 1$, we derive

$$\int_{\pi^{-1}(\gamma)} \alpha_M = \text{length}(\gamma \cap D).$$

Hence,

$$\begin{aligned} \text{vol}_n(D) &= \frac{1}{\epsilon_n n!} \int_{\gamma \in \Gamma} \text{length}(\gamma \cap D) \omega_\Gamma^{n-1} \\ &= \frac{1}{\epsilon_n n} \int_{\gamma \in \Gamma} \text{length}(\gamma \cap D) \, d\mu_\Gamma(\gamma). \end{aligned}$$

□

4. DISCRETIZATION OF FINSLER SURFACES

The goal of this section is to describe a discretization of Finsler disks with minimizing interior geodesics into simple discrete metric disks. For this, we adapt the general approach of discretization developed in [Cos18] in relation with the filling area conjecture. The main novelty is that, in our case, the discrete geometry is described by a system of curves (wall system) made of geodesics.

First, we need to fix some notation regarding intersections of maps.

Definition 4.1. The *intersections* of a map $f : X \rightarrow Y$ with a map $f' : X' \rightarrow Y$ lying in a subset $A \subseteq Y$ are the ordered pairs in the set

$$I_A(f, f') = \{(x, x') \in X \times X' \text{ such that } f(x) = f'(x') \in A\}.$$

The *number of intersections* between f and f' is defined as

$$\#(f \cap f') = \#I_Y(f, f'),$$

where $\#S$ denotes the cardinality of a set S .

Similarly, the *self-intersections* of a map $f : X \rightarrow Y$ lying in a subset $A \subseteq Y$ are the *unordered* pairs in the set

$$I_A(f) = \{\{x, x'\} \subseteq X \text{ such that } f(x) = f(x') \in A \text{ but } x \neq x'\},$$

and the *multiplicity* of a point $y \in Y$ as a self-intersection of f is the number $\#I_{\{y\}}(f)$. A self-intersection is *simple* if it has multiplicity 1.

Let us introduce the notion of wall system on a disk; see [Cos18].

Definition 4.2. A (smooth) *wall system* on a surface M is a 1-dimensional (smooth) immersed submanifold \mathcal{W} satisfying the following conditions:

- (1) the immersion map is proper (that is, the preimage of any compact subset of M is compact);
- (2) \mathcal{W} is transverse to the boundary ∂M and satisfies $\partial\mathcal{W} = \mathcal{W} \cap \partial M$;
- (3) \mathcal{W} is self-transverse and has only simple self-intersections;
- (4) no self-intersections of \mathcal{W} lie on the boundary ∂M .

As a technical remark, we note that the symbol \mathcal{W} denotes the *immersion map*, not its image $\text{Im}(\mathcal{W}) \subseteq M$, nor its domain. The domain is a 1-manifold, *i.e.*, a disjoint union of countably many intervals and circles. Hence the expression $\partial\mathcal{W} \subseteq \partial M$ involves an abuse of notation and actually means $\text{Im}(\partial\mathcal{W}) \subseteq \partial M$, where $\partial\mathcal{W}$ is the restriction of the map \mathcal{W} to the boundary of the domain of \mathcal{W} . The image of \mathcal{W} will also be denoted \mathcal{W} . Thus, the expression $M \setminus \mathcal{W}$ denotes $M \setminus \text{Im}(\mathcal{W})$.

Eventually we will need to relax the definition by dropping the last condition (4). In this case, we say that \mathcal{W} is a *quasi wall system* on M .

The curves that form a quasi wall system are called its *walls*. Note that if the surface M is compact, then \mathcal{W} consists of finitely many compact walls; each of these walls is either a loop that avoids the boundary or an arc that meets the boundary only at its two endpoints.

A quasi wall system \mathcal{W} on a disk D is *simple* if its walls are arcs that have no self-intersections and that meet each other at most once.²

In this paper, every quasi wall system \mathcal{W} is smooth unless we make it clear that it is piecewise smooth. In that case, the non-smooth points of \mathcal{W} may not coincide with the self-intersection points of \mathcal{W} . Note that a piecewise smooth quasi wall system can be turned into a smooth quasi wall system by an isotopic deformation.

Example 4.3. Let D be the unit disk in the Euclidean plane. A wall system made of the horizontal and vertical diameters of D has area 1. A quasi wall system made of the three sides of an inscribed triangle of D has area $\frac{3}{2}$.

We will also need the following definitions regarding the geometry induced by a quasi wall system.

Definition 4.4. Every quasi wall system \mathcal{W} on a compact surface M determines a *discrete length*

$$\text{length}_{\mathcal{W}}(c) = \#(c \cap \mathcal{W}) \tag{4.1}$$

for curves c in M . That is, the length of a curve is the number of times it intersects the quasi wall system (counted with multiplicity). Every quasi wall system \mathcal{W} also induces a pseudo-distance on $M \setminus \mathcal{W}$ defined by

$$d_{\mathcal{W}}(x, y) = \inf_c \text{length}_{\mathcal{W}}(c)$$

²Simple wall systems are also called pseudoline arrangements; see [Cos18]. However, some authors (*e.g.*, [FG17]) only consider complete pseudoline arrangements, which are those where every pair of walls crosses *exactly* once.

where the infimum is taken over all paths of M joining x to y . We will refer to the pseudo-distance $d_{\mathcal{W}}$ on M as the *discrete distance* induced by \mathcal{W} on D .

The *discrete area* of (M, \mathcal{W}) is the number of self-crossings of \mathcal{W} contained in the interior M° of M plus half the number of self-crossings on the boundary. That is,

$$\text{area}(M, \mathcal{W}) = \#I_{M^\circ}(\mathcal{W}) + \frac{1}{2}\#I_{\partial M}(\mathcal{W}) = \#I_{M^\circ + \frac{1}{2}\partial M}(\mathcal{W})$$

where $\#I_{M^\circ + \frac{1}{2}\partial M}$ is just an abbreviation for $\#I_{M^\circ} + \frac{1}{2}\#I_{\partial M}$.

Note that, if \mathcal{W} consists of finitely many curves γ_i , then

$$\text{area}(M, \mathcal{W}) = \sum_{i < j} \#I_{M^\circ + \frac{1}{2}\partial M}(\gamma_i, \gamma_j) + \sum_i \#I_{M^\circ + \frac{1}{2}\partial M}(\gamma_i) \quad (4.2)$$

When the quasi wall system is simple, the curves of \mathcal{W} have no self-intersections and the second sum vanishes.

We will need the following result describing the intersection of two distance realizing arcs of M . Recall that Γ is the space of traversing geodesics of M (*i.e.*, geodesic arcs of M which do not intersect ∂M except at their endpoints where they meet the boundary transversely).

Lemma 4.5. *Let M be a self-reverse Finsler metric disk with minimizing interior geodesics. Let $\gamma \in \Gamma$ be a traversing geodesic of M and let $[x, y]$ be a distance realizing arc of M with endpoints x and y not lying in γ . Then*

$$\#(\gamma \cap [x, y]) = \begin{cases} 1 & \text{if } \gamma \text{ separates } x \text{ and } y \\ 0 & \text{otherwise} \end{cases}$$

Proof. By Theorem 12.1, the distance realizing arc $[x, y]$ is C^1 .

Suppose that the arcs γ and $[x, y]$ are tangent, either at an interior point of M or at an endpoint of γ in ∂M . In both cases, this implies that $[x, y]$ contains γ since the distance realizing arc $[x, y]$ follows the geodesic flow in the interior of M and the endpoints x, y do not lie in γ . Now, since the interior geodesic γ is transverse to ∂M at its endpoints \bar{x} and \bar{y} , the distance realizing arc $[x, y]$ is not differentiable at \bar{x} and \bar{y} . In particular, it is not C^1 , which is absurd. Therefore, the arcs γ and $[x, y]$ may only have transverse intersections.

Suppose that the arcs γ and $[x, y]$ intersect at least twice, say at a and b (with a and b different from x and y). Since both arcs are distance realizing curves, the subarcs $[a, b] \subseteq [x, y]$ and $\gamma_{ab} \subseteq \gamma$ joining a and b have the same length. Construct an arc α joining x and y by replacing the subarc $[a, b]$ of $[x, y]$ with the arc γ_{ab} of the same length. By construction, the arc α is a distance realizing curve. But since the intersection between γ and $[x, y]$ is transverse, the arc α is not differentiable at a and b . In particular, it is not C^1 , which is absurd. Therefore, the arcs γ and $[x, y]$ intersect at most once, and so exactly once if γ separates x and y .

Suppose now that γ does not separate x and y . Then the arc $[x, y]$ does not intersect γ . Otherwise, it would go from one side of γ to the other (recall that γ and $[x, y]$ have transverse intersection) and, because x and y are on the same side of γ , it would have to cross γ a second time, which is excluded. Therefore, the arcs γ and $[x, y]$ do not intersect if γ does not separate x and y . \square

Let us compare the shortest paths for Finsler metrics and discrete metrics.

Definition 4.6. A quasi wall system is *geodesic* if its walls are geodesics.

Proposition 4.7. *Let M be a self-reverse Finsler metric disk with minimizing interior geodesics, and let \mathcal{W} be a geodesic quasi wall system on M . Then, every distance realizing arc $[x, y]$ of M with endpoints x, y not lying in \mathcal{W} is also length minimizing with respect to \mathcal{W} .*

Thus, for every $x, y \in M \setminus \mathcal{W}$, we have

$$d_{\mathcal{W}}(x, y) = \text{length}_{\mathcal{W}}([x, y]). \tag{4.3}$$

Proof. The quasi wall system \mathcal{W} is made of finitely many geodesics γ_i that are transverse to ∂M . By Lemma 4.5, the arc $[x, y]$ crosses only those geodesics γ_i that separate x from y , exactly once. Therefore, no curve from x to y can be shorter than $[x, y]$ with respect to \mathcal{W} . \square

Before proceeding we derive a useful consequence of the last lemma.

Lemma 4.8. *Let M be a self-reverse Finsler metric disk with minimizing interior geodesics. Then*

$$d(x, y) \leq \frac{1}{2} \text{length}(\partial M)$$

for any pair of points $x, y \in M$. The same inequality holds if the distance and length are taken with respect to a geodesic quasi wall system \mathcal{W} , that is,

$$d_{\mathcal{W}}(x, y) \leq \frac{1}{2} \text{length}_{\mathcal{W}}(\partial M)$$

for points $x, y \in M \setminus \mathcal{W}$.

Proof. Join the points $x, y \in M$ by a distance realizing path $[x, y]$. By Lemma 4.5, each traversing geodesic γ of M intersects $[x, y]$ at most once and meets ∂M exactly twice. Then the inequality $d(x, y) \leq \frac{1}{2} \text{length}(\partial M)$ follows from Blaschke's formula (3.4) applied to $[x, y]$.

The claim regarding the geodesic quasi wall system \mathcal{W} is proved in a similar way. By Proposition 4.7, the distance realizing arc $[x, y]$ is also length-minimizing with respect to \mathcal{W} . Since each wall of \mathcal{W} crosses $[x, y]$ at most once and meets ∂M exactly twice, we derive the desired second inequality from the definition of $\text{length}_{\mathcal{W}}$; see (4.1). \square

Simple wall systems can be used to discretize Finsler disks M with minimizing interior geodesics.

For every $a, b \in \mathbb{R}$ and every $\varepsilon > 0$, we write $a \simeq b \pm \varepsilon$ if $|a - b| < \varepsilon$.

Theorem 4.9. *Let (M, F) be a self-reverse Finsler metric disk with minimizing interior geodesics. Then, for every $\varepsilon > 0$ and every integer n large enough, there exists a wall system \mathcal{W} , made of n geodesics of M , such that for every $x, y \in M \setminus \mathcal{W}$, we have*

$$\frac{1}{n}d_{\mathcal{W}}(x, y) \simeq \frac{2}{L}d_F(x, y) \pm \varepsilon \quad (4.4)$$

$$\frac{2}{n^2 - n} \text{area}(M, \mathcal{W}) \simeq \frac{2\pi}{L^2} \text{area}(M, F) \pm \varepsilon \quad (4.5)$$

where $L = \text{length}_F(\partial M)$. Furthermore, the wall system \mathcal{W} is necessarily simple.

Note that [Cos18, Theorem 7.1] states the existence of a theorem with similar approximation properties but not necessarily made of geodesics.

Proof. The wall system \mathcal{W} will be made of random geodesics. Recall that Γ is the space of traversing geodesics of M (i.e., geodesic arcs of M which do not intersect ∂M except at their endpoints where they meet the boundary transversely) and has a natural measure μ_{Γ} ; see (3.8). Furthermore, this space has finite total measure $\mu_{\Gamma}(\Gamma) = 2L$; see 3.3. Thus we may define on Γ the probability measure $\mathbb{P} = \frac{\mu_{\Gamma}}{2L}$.

Take n independent identically distributed (i.i.d.) random geodesics $\gamma_1, \dots, \gamma_n$ of Γ with probability distribution \mathbb{P} . Almost surely, these geodesics form a wall system \mathcal{W} of M ; see Definition 4.2; because they are pairwise different and form only simple crossings located in the interior of M . Moreover, this wall system is simple, since the geodesics are minimizing and therefore they cannot cross each other more than once by Lemma 4.5. At this point, Theorem 4.9 follows from the next two lemmas.

The proof of the first lemma requires to establish a uniform version of the weak law of large numbers for the Blaschke formula (3.4).

Lemma 4.10. *With arbitrarily large probability for n large enough, the relation*

$$\frac{1}{n}d_{\mathcal{W}}(x, y) \simeq \frac{2}{L}d_F(x, y) \pm \varepsilon$$

holds for every $x, y \in M \setminus \mathcal{W}$

Proof. Let \mathcal{D} be a finite covering of M by smoothly bounded disks D with perimeter $\text{length}_F(\partial D) < \varepsilon$. Fix a basepoint p in each disk $D \in \mathcal{D}$ and denote by P the collection of all basepoints. Almost surely, the geodesics of \mathcal{W} avoid the points of P and are transverse to the boundaries of the disks $D \in \mathcal{D}$.

The following claim shows that the conclusion of the lemma holds in some finite cases.

Claim 4.11. *The following assertions hold with arbitrarily large probability for n large enough.*

(1) For every pair of points $p, q \in P$, we have

$$\frac{1}{n}d_{\mathcal{W}}(p, q) \simeq \frac{2}{L}d_F(p, q) \pm \varepsilon. \quad (4.6)$$

(2) For every disk $D \in \mathcal{D}$ and every pair of points $x, y \in D \setminus \mathcal{W}$, we have

$$\frac{1}{n}d_{\mathcal{W}}(x, y) \leq \left(\frac{1}{L} + \frac{1}{2} \right) \varepsilon. \quad (4.7)$$

Proof. (1) Recall that the distance realizing arc $[p, q]$ is C^1 embedded in M ; see Theorem 12.1.

The intersection function $f = f_{p,q} : \Gamma \rightarrow \mathbb{N}$ defined by

$$f(\gamma) = \#(\gamma \cap [p, q])$$

is a nonnegative measurable function. By Blaschke's formula (3.4), the random variables $X_i = f(\gamma_i)$ with $1 \leq i \leq n$ are i.i.d. with finite expected value

$$E(X_i) = \int_{\Gamma} \#(\gamma \cap [p, q]) \, d\mathbb{P} = \frac{2}{L}d_F(p, q).$$

Note that $E|X_i| = E(X_i) < \infty$. By the weak law of large numbers applied to $\{X_i\}$, see [Dur19, Theorem 2.2.14] or [Tao09], we derive that

$$\left| \frac{1}{n} \sum_{i=1}^n \#(\gamma_i \cap [p, q]) - \frac{2}{L}d_F(p, q) \right| < \varepsilon$$

with arbitrarily large probability for n large enough. By Proposition 4.7, we have

$$d_{\mathcal{W}}(p, q) = \text{length}_{\mathcal{W}}([p, q]) = \sum_{i=1}^n \#(\gamma_i \cap [p, q]),$$

hence (1) follows.

(2) The proof of the second assertion is similar. For a disk $D \in \mathcal{D}$, the intersection function $f(\gamma) = \#(\gamma \cap \partial D)$ has expected value $\frac{2}{L} \text{length}_F(\partial D)$ by Blaschke's formula (3.4). Applying the weak law of large numbers to the random variables $X_i = f(\gamma_i)$ as previously, we derive

$$\left| \frac{1}{n} \sum_{i=1}^n \#(\gamma_i \cap \partial D) - \frac{2}{L} \text{length}_F(\partial D) \right| < \varepsilon$$

with arbitrarily large probability for n large enough. Thus,

$$\begin{aligned} \frac{1}{n} \text{length}_{\mathcal{W}}(D) &\simeq \frac{2}{L} \text{length}_F(D) \pm \varepsilon \\ &\leq \left(\frac{2}{L} + 1 \right) \varepsilon. \end{aligned}$$

Since D is a disk with minimizing interior geodesics, the discrete part of Lemma 4.8 yields (2). \square

Without loss of generality, we can assume that the conclusion of the previous claim is satisfied. Let $x, y \in M \setminus \mathcal{W}$. The points x and y lie in some disks D_x and D_y of \mathcal{D} . Denote by p_x and p_y the basepoints of D_x and D_y . Since D_x is a disk with minimizing interior geodesics, by Lemma 4.8 we have

$$d_F(x, p_x) \leq \frac{1}{2} \text{length}_F(\partial D_x) < \frac{1}{2} \varepsilon,$$

thus by the triangle inequality, we obtain

$$|d_F(x, y) - d_F(p_x, p_y)| \leq d_F(x, p_x) + d_F(y, p_y) < \varepsilon. \quad (4.8)$$

Combining the triangle inequality with (4.7), we obtain

$$\begin{aligned} \left| \frac{1}{n} d_{\mathcal{W}}(x, y) - \frac{1}{n} d_{\mathcal{W}}(p_x, p_y) \right| &\leq \frac{1}{n} d_{\mathcal{W}}(x, p_x) + \frac{1}{n} d_{\mathcal{W}}(y, p_y) \\ &\leq \left(\frac{2}{L} + 1 \right) \varepsilon. \end{aligned} \quad (4.9)$$

Thus, the following equalities

$$\frac{1}{n} d_{\mathcal{W}}(x, y) \underset{(4.9)}{\simeq} \frac{1}{n} d_{\mathcal{W}}(p_x, p_y) \underset{(4.6)}{\simeq} \frac{2}{L} d_F(p_x, p_y) \underset{(4.8)}{\simeq} \frac{2}{L} d_F(x, y)$$

hold up to additive constants which are universal multiples of ε (namely, $(\frac{2}{L} + 1) \varepsilon$ for the first one, ε for the second and $\frac{2}{L} \varepsilon$ for the third one). Therefore,

$$\left| \frac{1}{n} d_{\mathcal{W}}(x, y) - \frac{2}{L} d_F(x, y) \right| < C_0 \varepsilon$$

where $C_0 = \frac{4}{L} + 2$. Hence the first lemma. \square

The proof of the second lemma is simpler and rests on the weak law of large numbers for the Santaló+Blaschke formula (3.6).

Lemma 4.12. *With arbitrarily large probability for n large enough, we have*

$$\frac{2}{n^2 - n} \text{area}(M, \mathcal{W}) \simeq \frac{2\pi}{L^2} \text{area}(M) \pm \varepsilon.$$

Proof. The intersection counting function $f : \Gamma \times \Gamma \rightarrow \mathbb{N}$ defined by

$$f(\gamma, \gamma') = \#(\gamma \cap \gamma')$$

is a measurable function that takes value 0 or 1 almost surely. The $\frac{n(n-1)}{2}$ random variables $X_{i,j} = f(\gamma_i, \gamma_j)$ with $i < j$ are identically distributed but not completely independent. In fact $X_{i,j}$ is independent of $X_{k,l}$ if and only if $\{i, j\} \cap \{k, l\} = \emptyset$. To apply the generalized weak law of large numbers, Theorem 13.1, we must check that the variables $X_{i,j}$ are sufficiently independent. There are $\sim n^2$ variables $X_{i,j}$, which yield $\sim n^4$ pairs $(X_{i,j}, X_{k,l})$, of which only $\sim n^3$ are not independent. Therefore the proportion of nonindependent pairs $p \sim \frac{n^3}{n^4} \sim \frac{1}{n}$ goes to zero as $n \rightarrow \infty$. Thus, by Theorem 13.1, the average value of the variables $X_{i,j}$,

$$\frac{\sum_{i < j} X_{i,j}}{\frac{n(n-1)}{2}} = \frac{\sum_{i < j} \#(\gamma_i \cap \gamma_j)}{\frac{n(n-1)}{2}} = \frac{\text{area}(M, \mathcal{W})}{\frac{n(n-1)}{2}}$$

converges in probability to the expected value, which, by the Santaló+Blaschke formula (3.6), is equal to

$$\mathbb{E}(X_{i,j}) = \iint_{\Gamma \times \Gamma} \#(\gamma \cap \gamma') \, d\mathbb{P}(\gamma) \, d\mathbb{P}(\gamma') = \frac{2}{L^2} \pi \operatorname{area}(M). \quad \square$$

This concludes the proof of Theorem 4.9. □

5. MINIMAL AREA OF DISKS: FROM DISCRETE TO FINSLER METRICS

The goal of this section is to state a discrete version of the area lower bound on Finsler disks with minimizing interior geodesics and to show how to derive the area lower bound for Finsler metrics from its discrete version.

Let us recall the area lower bound for Finsler metrics we want to prove.

Theorem 5.1. *Let M be a self-reverse Finsler metric disk of radius r with minimizing interior geodesics. Then the Holmes–Thompson area of M satisfies*

$$\operatorname{area}(M) \geq \frac{6}{\pi} r^2.$$

In order to state the discrete version of this result, we need to introduce the notion of simple discrete metric disks.

Definition 5.2. A topological disk D with a quasi wall system \mathcal{W} is a *simple discrete metric disk* of radius r centered at an interior point $O \in D \setminus \mathcal{W}$ if all the points of $D \setminus \mathcal{W}$ are at $d_{\mathcal{W}}$ -distance at most r from O , all the points of $\partial D \setminus \mathcal{W}$ are at distance exactly r from O and the quasi wall system \mathcal{W} is simple; see Definition 4.2.

The following result, which will be proved in the subsequent sections, can be seen as a discrete version of Theorem 5.1.

Theorem 5.3. *The discrete area of every simple discrete metric disk (D, \mathcal{W}) of radius r satisfies*

$$\operatorname{area}(D, \mathcal{W}) \geq \frac{3}{2} r^2.$$

Furthermore, the equality is attained.

Assuming this discrete area lower bound, we can derive Theorem 5.1 as follows.

Proof of Theorem 5.1 (assuming Theorem 5.3). Let M be a Finsler disk of radius r centered at O with minimizing interior geodesics. By Theorem 4.9, for every $\varepsilon > 0$, there exists a simple wall system \mathcal{W}_M , made of n interior geodesics of M , satisfying the estimates (4.4) and (4.5). The simple wall system \mathcal{W}_M decomposes M into convex polygonal cells. By definition, all the points in a cell are at the same distance from the center of M with respect to the discrete distance $d_{\mathcal{W}_M}$. Since M has minimizing interior geodesics, the

geodesic rays of length r arising from its center O form a geodesic foliation \mathcal{F} of the punctured disk $M \setminus \{O\}$. The sides of the cells of M , which lie in the geodesics of \mathcal{W}_M , are transverse to the foliation \mathcal{F} , otherwise the origin O would lie in \mathcal{W} .

Consider a convex polygonal cell Δ of M not containing O . Denote by d the $d_{\mathcal{W}_M}$ -distance from O to the interior of the cell Δ . The geodesic rays of the foliation intersecting Δ form a spray \mathcal{F}_Δ , where each ray of \mathcal{F}_Δ intersects Δ along an interval with nonempty interior, except for the two extremal rays of the spray which intersect the convex polygonal cell Δ at two vertices; see Figure 1. Denote by β_Δ the broken line made of two segments joining the center of Δ to these two extremal vertices. Note that every geodesic ray of the spray \mathcal{F}_Δ intersects the broken line β_Δ at a single point; see Figure 1. Since the rays of the spray are length-minimizing with respect to $d_{\mathcal{W}_M}$, see Proposition 4.7, all the cells intersecting the spray between O and β_Δ are at $d_{\mathcal{W}_M}$ -distance at most d from O , and all the cells intersecting the spray after β_Δ are at $d_{\mathcal{W}_M}$ -distance at least d from O .




FIGURE 1. Spray \mathcal{F}_Δ intersecting the convex polygonal cell Δ

Denote by r_0 the integral part of $n \left(\frac{2r}{L} - \varepsilon \right)$. By (4.4), every boundary point $p \in \partial M \setminus \mathcal{W}_M$ is at $d_{\mathcal{W}_M}$ -distance greater than r_0 from O , that is, $d_{\mathcal{W}_M}(O, p) > r_0$. A cell of M whose interior points are at $d_{\mathcal{W}_M}$ -distance r_0 from O will be referred to as an outermost cell. The broken lines β_Δ , where Δ runs over all outermost cells of M , form a piecewise geodesic closed curve delimiting a topological disk $D \subseteq M$ containing O . This curve can be smoothed to ensure that D is a smoothly bounded manifold. The restriction $\mathcal{W} = \mathcal{W}_M \cap D$ of \mathcal{W}_M to D defines a simple quasi wall system on D . By construction, all the points of $D \setminus \mathcal{W}$ are at $d_{\mathcal{W}}$ -distance at most r_0 from O and all the points of $\partial D \setminus \mathcal{W}$ are at distance exactly r_0 from O . Hence, (D, \mathcal{W}) is a simple discrete metric disk of radius r_0 . By Theorem 5.3 and by definition of the discrete area (4.2), we have

$$\text{area}(M, \mathcal{W}_M) \geq \text{area}(D, \mathcal{W}) \geq \frac{3}{2} r_0^2.$$

Dividing by n^2 , using (4.4) and (4.5), and letting ε go to zero, we obtain

$$\frac{\pi}{L^2} \text{area}(M) \geq \frac{3}{2} \left(\frac{2}{L} r \right)^2.$$

Hence, $\text{area}(M) \geq \frac{6}{\pi} r^2$. □

Sections 6–9 are devoted to the proof of Theorem 5.3.

6. QUASI WALL SYSTEMS AND INTERVAL FAMILIES

In this section, we show how to encode a simple discrete disk as a 1-dimensional object.

Let (D, \mathcal{W}) be a simple discrete disk of radius r and center O . Identify the boundary ∂D with the circle S^1 , and identify the punctured disk $D \setminus \{O\}$ with the flat cylinder $\mathcal{C} = S^1 \times [0, \infty)$. Under this identification, the point O of D corresponds to the point at infinity in the one-point compactification of the cylinder \mathcal{C} . Note that the universal cover of \mathcal{C} is the half plane $\mathcal{H} = \mathbb{R} \times [0, +\infty)$.

Definition 6.1. Given a simple arc α in the cylinder $M = D \setminus \{O\}$ (or in the half plane $M = \mathbb{R} \times [0, +\infty)$) with endpoints on the boundary ∂M , denote by $\bar{\alpha}$ the segment of ∂D with the same endpoints, homotopic to α in M . The arc α *covers a point* p of ∂M if p lies in $\bar{\alpha}$. Similarly, the arc α *covers another arc* β if $\bar{\beta}$ lies in $\bar{\alpha}$. Two arcs α and α' are *adjacent* if the intervals $\bar{\alpha}$ and $\bar{\alpha}'$ are adjacent, meaning that they have exactly one point in common.

Definition 6.2. An arc in the flat cylinder $M = S^1 \times [0, \infty)$ (or in the half plane $M = \mathbb{R} \times [0, +\infty)$) with endpoints on the boundary ∂M is *standard* if it consists of a segment of slope 1 followed by a segment of slope -1 ; see Figure 2. A quasi wall system \mathcal{W} is standard if its walls are standard arcs. For two boundary points $a, b \in S^1 = \partial M$ (or $a < b \in \mathbb{R}$ if M is the half plane), we denote by $[a, b]$ the arc of S^1 that goes from a to b in the positive (*i.e.*, counterclockwise) sense, and we denote by \widehat{ab} the standard arc in M that is homotopic to $[a, b]$.

Let (D, \mathcal{W}) be a simple discrete disk of radius r centered at O . Denote by $\mathcal{I} = \mathcal{I}_{\mathcal{W}}$ the set of boundary intervals $\bar{\alpha}$ homotopic to the walls α of \mathcal{W} . The family \mathcal{I} of intervals of S^1 contains all the information about \mathcal{W} that is relevant to our problem of finding simple discrete disks of minimum area. For instance, two walls α, β of \mathcal{W} meet on ∂D if and only if the intervals $\bar{\alpha}, \bar{\beta}$ have a common endpoint. That is,

$$\#I_{\partial D}(\alpha, \beta) = 1 \iff \#(\partial\bar{\alpha} \cap \partial\bar{\beta}) = 1. \tag{6.1}$$

Furthermore, assuming $\bar{\alpha}$ and $\bar{\beta}$ have no common endpoints, the arcs α and β cross in the interior of D if and only if the interval $\bar{\alpha}$ contains *exactly*

one endpoint of $\bar{\beta}$. That is,

$$\#I_{D^\circ}(\alpha, \beta) = 1 \iff \#(\bar{\alpha} \cap \partial\bar{\beta}) = 1. \quad (6.2)$$

One consequence of these formulas is that the discrete area of (D, \mathcal{W}) given by (4.2) may be computed from \mathcal{I} .

The following result characterizes the relation between the quasi wall system \mathcal{W} and the interval family \mathcal{I} . Before stating this result, we need to introduce a definition. A point p of S^1 is *generic* with respect to a finite interval family \mathcal{I} of S^1 if p is not an endpoint of any interval of \mathcal{I} . Alternatively, the endpoints of the intervals of \mathcal{I} are the non-generic points of S^1 .

Proposition 6.3. *Let (D, \mathcal{W}) be a simple discrete disk of radius r centered at O . The family $\mathcal{I} = \mathcal{I}_{\mathcal{W}}$ of intervals of S^1 has the following properties:*

- (1) *no pair of intervals of \mathcal{I} cover S^1 ;*
- (2) *every generic point of S^1 is contained in exactly r intervals of \mathcal{I} ;*
- (3) *every non-generic point of S^1 is an endpoint of exactly two, adjacent intervals of \mathcal{I} .*

Moreover, if a finite family \mathcal{I} of intervals of S^1 satisfies the conditions (1)–(3), then $\mathcal{I} = \mathcal{I}_{\mathcal{W}}$ for some quasi wall system \mathcal{W} that makes D a simple discrete metric disk of radius r and center O . For instance, one may let \mathcal{W} be the unique standard quasi wall system homotopic to \mathcal{I} on $D \setminus \{O\}$.

Proof.

(1) If two intervals $\bar{\alpha}, \bar{\beta} \in \mathcal{I}$ cover S^1 , then the corresponding walls α, β of \mathcal{W} would form a bigon containing the point O , which implies they cross twice, contradicting the hypothesis that \mathcal{W} is simple.

(2) Consider a generic point $p \in S^1$. Since \mathcal{W} is a simple quasi wall system on D , the distance between any pair of points of D is the number of walls that separate them; see Proposition 14.1 of Appendix 14. On the other hand, the walls that separate O from p are the walls that cover p . Hence the result.

(3) This follows from the previous property: if $p \in S^1$ is the endpoint of some interval $\bar{\alpha} \in \mathcal{I}$, it must also be the startpoint of some other interval so that every generic point near p is contained in the same number r of intervals of \mathcal{I} . This means that p is the endpoint of two walls, and it cannot be the endpoint of more walls because \mathcal{W} can only have simple self-intersections on ∂D since it is a quasi wall system; see Definition 4.2.

Now, let \mathcal{I} be a finite family of intervals of S^1 satisfying conditions (1)–(3), and let \mathcal{W} be the unique standard quasi wall system homotopic to \mathcal{I} on $D \setminus \{O\}$. Clearly, \mathcal{W} is a quasi wall system, and it is simple because it is made of arcs that intersect each other at most once. Also, every point $p \in D \setminus \mathcal{W}$ is at distance at most r from O , and exactly r if $p \in \partial D$. (A shortest path is the vertical ray from p to O .) This shows that (D, \mathcal{W}) is a simple discrete disk of radius r centered at O . \square

7. INADMISSIBLE CONFIGURATIONS IN A MINIMAL SIMPLE DISK

In this section, we rule out some intersection patterns for extremal quasi wall system on a disk.

Consider a quasi wall system \mathcal{W} on D defining a simple discrete metric disk of radius r with minimal discrete area. By Proposition (6.3), we can assume that \mathcal{W} is formed of standard arcs; see Definition 6.2.

Lemma 7.1. *No arc of \mathcal{W} covers two (possibly adjacent) intersecting arcs of \mathcal{W} .*

Proof. By contradiction, suppose that an arc γ of \mathcal{W} covers two intersecting arcs $\alpha = \widehat{ac}$ and $\beta = \widehat{bd}$ of \mathcal{W} . Switching the roles of the two arcs if necessary, we may assume that the points a, b, c, d appear in that order in the interval $\overline{\gamma}$ (with possibly $b = c$). See Figure 2. Let \mathcal{W}' be the collection of curves obtained from \mathcal{W} by replacing α and β with the standard arcs $\alpha' = \widehat{ad}$ and $\beta' = \widehat{bc}$ (with no β' if $b = c$). See Figure 2. Note that, like \mathcal{W} , the immersed 1-submanifold \mathcal{W}' is a quasi wall system on D . Moreover, we claim that \mathcal{W}' also makes D a simple discrete metric disk of radius r centered at O . This is because none of the properties (1)–(3) of Proposition 6.3 is affected by the replacement. For instance, there is no arc δ of \mathcal{W}' such that the intervals $\overline{\delta}$ and $\overline{\alpha'}$ cover the boundary ∂D , because in that case $\overline{\delta}$ and $\overline{\gamma}$ would also cover ∂D , however the arcs δ and $\overline{\gamma}$ are already present in \mathcal{W} , contradicting by Proposition 6.3 the fact that \mathcal{W} is simple. Also, the fact that every generic point of ∂D is covered by exactly r arcs of the quasi wall system is clearly maintained, as well as the fact that each non-generic boundary point is the common endpoint of two adjacent walls.

Let us show that the area of (D, \mathcal{W}') is less than the area of (D, \mathcal{W}) by comparing the number of self intersections of the quasi wall systems \mathcal{W} and \mathcal{W}' according to the discrete area formula (4.2). First, observe that every pair of arcs of \mathcal{W} different from α and β belongs to \mathcal{W}' . Therefore, these pairs of arcs give the same contribution to the discrete areas of \mathcal{W} and \mathcal{W}' . Let $\delta = \widehat{pq}$ be an arc of \mathcal{W} different from α and β . By considering cases regarding the location of the endpoints p and q with respect to the points a, b, c and d , we see that

$$\#I_{D^\circ + \frac{1}{2}\partial D}(\delta, \alpha' \cup \beta') \leq \#I_{D^\circ + \frac{1}{2}\partial D}(\delta, \alpha \cup \beta).$$

In fact, equality holds unless p and q lie in the interiors of $[a, b]$ and $[c, d]$, in which case the inequality is strict. Finally, note that

$$\#I_{D^\circ}(\alpha', \beta') = 0 \quad \text{and} \quad \#I_{D^\circ}(\alpha, \beta) = 1.$$

We conclude that

$$\text{area}(D, \mathcal{W}') \leq \text{area}(D, \mathcal{W}) - 1,$$

which contradicts the minimality of the discrete area of (D, \mathcal{W}) . □

Lemma 7.2. *No arc of \mathcal{W} intersects two adjacent arcs of \mathcal{W} .*




FIGURE 2. Replacing two intersecting arcs covered by a third arc

Proof. By contradiction, suppose that an arc γ of \mathcal{W} intersects two adjacent arcs α and β of \mathcal{W} . We choose γ so that it is minimal with respect to the covering relation, among arcs that intersects α and β (i.e., no arc of \mathcal{W} covered by γ intersects α and β). Denote by a, b, c, d, e the endpoints of the three arcs, in the order in which they are found on the interval $\bar{\alpha} \cup \bar{\beta}$. Thus, $\alpha = \widehat{ac}$, $\beta = \widehat{ce}$ and $\gamma = \widehat{bd}$, and no arc of \mathcal{W} that covers c is covered by γ (other than γ itself). See Figure 3. Let c^- and c^+ be two points of ∂D close to c such that $[c^-, c^+] \cap \partial \mathcal{W} = \{c\}$. Let \mathcal{W}' be the collection of curves obtained from \mathcal{W} by replacing the three arcs α, β and γ with the four arcs $\alpha' = \widehat{ac^+}$, $\beta' = \widehat{c^-e}$, $\gamma^- = \widehat{bc^-}$ and $\gamma^+ = \widehat{c^+d}$. See Figure 3.

Note that \mathcal{W}' is a quasi wall system on the disk D . In fact, \mathcal{W}' makes D a simple discrete disk of radius r centered at O . To see this we argue as in the proof of Lemma 7.1. By Proposition 6.3, it is enough to check that the family $\mathcal{I} = \mathcal{I}_{\mathcal{W}'}$ of boundary segments $\bar{\delta}$ corresponding to the walls δ of \mathcal{W}' satisfies the properties (1)–(3) of Proposition 6.3. To check Property (2) (that each generic point of ∂D is covered r times by the walls of \mathcal{W}') note that both $\alpha \cup \beta \cup \gamma$ and $\alpha' \cup \beta' \cup \gamma^- \cup \gamma^+$ cover twice the generic points of $[b, d]$ and once the remaining generic points of $[a, e]$. Property (3) regarding non-generic boundary points is also maintained, with the wall endpoint c replaced by the two points c^- and c^+ . Finally, to check the property (1), suppose δ and ε are two arcs of \mathcal{W}' that cover the whole boundary ∂D . It is impossible that both δ and ε are among the new arcs α', β' and γ^\pm because that would mean that α and β already cover ∂D , contradicting the fact that \mathcal{W} is

simple. Similarly, the arcs δ and ε cannot be both among the unchanged arcs (those in $\mathcal{W} \cap \mathcal{W}'$) either, otherwise \mathcal{W} would not be simple. Therefore, δ is one of the unchanged arcs and ε is one of the new arcs α' , β' , γ^\pm . In the case $\varepsilon = \alpha'$, we see that δ and α' cannot cover ∂D since this would imply that δ and α already cover ∂D . This is because $\overline{\alpha'} \setminus \overline{\alpha}$ is contained in the interval $[c^-, c^+]$ which contains no endpoints of δ since $[c^-, c^+] \cap \mathcal{W} = \{c\}$. The case $\varepsilon = \beta'$ is analogous and the cases $\varepsilon = \gamma^\pm$ are easier to rule out since the arcs γ^\pm are covered by γ . We conclude that the property (1) is satisfied, thus (D, \mathcal{W}') is a simple discrete metric disk of radius r .




FIGURE 3. Replacing a configuration of one arc intersecting two adjacent arcs.

Let us show that the area of (D, \mathcal{W}') is less than the area of (D, \mathcal{W}) . Again, we use the discrete area formula (4.2), which says

$$\text{area}(D, \mathcal{W}) = \sum_{\{\delta, \varepsilon\}} \#I_{D^\circ + \frac{1}{2}\partial D}(\delta, \varepsilon)$$

where the sum runs over pairs $\{\delta, \varepsilon\}$ of different walls of \mathcal{W} . The pairs $\{\delta, \varepsilon\}$ of walls that are contained in $\mathcal{W} \cap \mathcal{W}'$ make the same contribution to $\text{area}(D, \mathcal{W})$ and to $\text{area}(D, \mathcal{W}')$. To evaluate the contribution of pairs $\{\delta, \varepsilon\}$ with $\delta \in \mathcal{W} \cap \mathcal{W}'$ and $\varepsilon \notin \mathcal{W} \cap \mathcal{W}'$, we note that any arc $\delta = \widehat{pq}$ with no endpoints in $[c^-, c^+]$ satisfies

$$\#I_{D^\circ + \frac{1}{2}\partial D}(\delta, \alpha' \cup \beta' \cup \gamma^+ \cup \gamma^-) = \#I_{D^\circ + \frac{1}{2}\partial D}(\delta, \alpha \cup \beta \cup \gamma)$$

unless $p \in [b, c]$ and $q \in [c, d]$. This is seen by considering case by case the possible locations of p and q with respect to a, b, c, d, e . The equality holds for all arcs $\delta = \widehat{pq} \in \mathcal{W} \cap \mathcal{W}'$, because the exceptional case $p \in [b, c]$ and $q \in [c, d]$ is excluded by how γ was chosen: the arc $\gamma = \widehat{cd}$ covers no other arc $\delta = \widehat{pq}$ of \mathcal{W} that in turn covers c . Finally, to compute the contribution

of the pairs $\{\delta, \varepsilon\}$ where none of the two arcs δ and ε is in $\mathcal{W} \cap \mathcal{W}'$, we note that

$$\#I_{D^\circ + \frac{1}{2}\partial D}(\alpha' \cup \beta' \cup \gamma^- \cup \gamma^+) = 2$$

while

$$\#I_{D^\circ + \frac{1}{2}\partial D}(\alpha \cup \beta \cup \gamma) = \frac{5}{2}.$$

We conclude that $\text{area}(D, \mathcal{W}') = \text{area}(D, \mathcal{W}) - \frac{1}{2}$, contradicting the minimality of \mathcal{W} . \square

8. PAIRS OF ADJACENT ARCS

In this section we show that the sequences of adjacent arcs in an extremal quasi wall system on a disk have a periodic structure.

Consider a quasi wall system \mathcal{W} on the disk D , made of standard arcs, defining a simple metric disk of radius r centered at O with minimal discrete area as in Section 7. Recall that the upper half plane $\mathcal{H} = \mathbb{R} \times [0, \infty)$ is the universal cover of the cylinder $\mathcal{C} = S^1 \times [0, \infty) = D \setminus \{O\}$. We identify its boundary $\partial\mathcal{H}$ with the real line \mathbb{R} . Let $\mathcal{W}_{\mathcal{H}}$ be the quasi wall system on \mathcal{H} formed of all the lifts of the arcs of \mathcal{W} .

Definition 8.1. Since every generic point of ∂D is covered by exactly r arcs of \mathcal{W} , the quasi wall system $\mathcal{W}_{\mathcal{H}}$ is composed of exactly r bi-infinite sequences $(\alpha_i)_{i \in \mathbb{Z}}$ of adjacent arcs

$$\alpha_i = \widehat{a_i a_{i+1}},$$

called the *strands* of $\mathcal{W}_{\mathcal{H}}$. The *width* of α_i is the number $a_{i+1} - a_i$. The points a_i where the strand $(\alpha_i)_i$ meets the boundary $\partial\mathcal{H}$ are called the *stops* of the strand. Note that every arc α of $\mathcal{W}_{\mathcal{H}}$ is contained in exactly one strand.

The following result describes how each strand intersects a pair of adjacent arcs of $\mathcal{W}_{\mathcal{H}}$.

Lemma 8.2. *Let $\alpha_0 = \widehat{a_0 a_1}$ and $\alpha_1 = \widehat{a_1 a_2}$ be two adjacent arcs of $\mathcal{W}_{\mathcal{H}}$. Then every strand of $\mathcal{W}_{\mathcal{H}}$ has exactly one arc with endpoints on the boundary interval $I = [a_0, a_2)$. This arc is covered by α_0 or by α_1 .*

Proof. The strand that contains the arcs α_0 and α_1 clearly satisfies the proposition. Thus let $(\beta_i)_{i \in \mathbb{Z}}$ be any other strand of $\mathcal{W}_{\mathcal{H}}$, numbered so that the arc β_0 covers the point a_1 . This strand has a stop in I , otherwise β_0 would cover the two adjacent arcs α_0 and α_1 , in contradiction with Lemma 7.1. Also, the strand $(\beta_i)_i$ cannot have stops in both intervals $[a_0, a_1)$ and $[a_1, a_2)$, otherwise the arc β_0 would intersect the two adjacent arcs α_0 and α_1 , in contradiction with Lemma 7.2. Thus the strand $(\beta_i)_i$ has stops in exactly one of the intervals $[a_0, a_1)$ and $[a_1, a_2)$, say, the second one; see Figure 4. Furthermore, it cannot have just one stop in this interval, otherwise the two adjacent arcs β_0, β_1 that share this stop would intersect α_1 , in contradiction with Lemma 7.2. Also, it cannot have three stops in the

interval, otherwise the adjacent arcs β_1 and β_2 would be covered by α_1 , in contradiction with Lemma 7.1. We conclude that the strand $(\beta_i)_i$ has exactly two stops (and therefore one arc) in the interval $[a_0, a_2)$, and both of these stops are covered by one of the arcs α_0 or α_1 . See Figure 4.




FIGURE 4. Leaping over every other arc.

□

Let n be the number of walls of the quasi wall system \mathcal{W} on the disk D . From now on, changing the parameterization of the boundary circle $S^1 = \partial D$, we assume that S^1 is a circle of length n , thus $S^1 = \mathbb{R}/n\mathbb{Z}$, and that the endpoints of the walls of \mathcal{W} are located at the semi integer points. Therefore, on the universal cover of the cylinder $\mathcal{C} = D \setminus \{O\}$, which is the upper half plane \mathcal{H} , we have $\partial\mathcal{W}_{\mathcal{H}} = \mathbb{Z} + \frac{1}{2} \subseteq \mathbb{R} = \partial\mathcal{H}$.

Note that the quasi wall system $\mathcal{W}_{\mathcal{H}}$ is periodic of period n (where n is the number of walls of \mathcal{W}) in the sense that it is invariant by horizontal translations of displacement length n . However, the following result implies that $\mathcal{W}_{\mathcal{H}}$ is also periodic with period $2r$, where r is the number of strands of $\mathcal{W}_{\mathcal{H}}$; see Definition 8.1.

Lemma 8.3. *The sum of the widths of two adjacent arcs α_0, α_1 of $\mathcal{W}_{\mathcal{H}}$ is equal to $2r$.*

Proof. Consider two adjacent arcs $\alpha_0 = \widehat{a_0 a_1}$ and $\alpha_1 = \widehat{a_1 a_2}$ as in Lemma 8.2. According to that lemma, each of the r strands of $\mathcal{W}_{\mathcal{H}}$ has exactly two stops in the interval $[a_0, a_2)$. Therefore there are $2r$ semi-integers in that interval. It follows that $a_2 - a_0 = 2r$. □

Denote by $\mathcal{S}_{[t, t+2r)} = [t, t+2r) \times [0, +\infty)$ a strip of width $2r$ of the half-plane \mathcal{H} . The following result describes the arcs of the wall system $\mathcal{W}_{\mathcal{H}}$ that are contained in such a strip.

Proposition 8.4.

- (1) *Each strip $\mathcal{S}_{[t, t+2r)}$ contains exactly one arc of each strand (and each of these arcs determines its strand completely).*
- (2) *The r arcs contained in a strip $\mathcal{S}_{[t, t+2r)}$ do not intersect each other.*
- (3) *Any pair of strands intersects each other exactly twice in the strip $\mathcal{S}_{[t, t+2r)}$.*

Proof.

(1) Consider a strand $(\alpha_i)_{i \in \mathbb{Z}}$, with $\alpha_i = \widehat{a_i a_{i+1}}$. According to Lemma 8.3, we have the relation $a_{i+2} = a_i + 2r$ for all i . This implies that the strip $\mathcal{S}_{[t, t+2r]}$ contains exactly two stops and thus exactly one arc of the strand $(\alpha_i)_i$. The same relation implies that two consecutive stops determine the strand.

(2) Consider a second strand $(\beta_j)_{j \in \mathbb{Z}}$, with $\beta_j = \widehat{b_j b_{j+1}}$. Assuming that two arcs α_0 and β_0 of $\mathcal{W}_{\mathcal{H}}$ intersect, we want to show that their are not contained in a strip $\mathcal{S}_{[t, t+2r]}$. We may assume without loss of generality that $a_0 < b_0$, therefore $b_0 \in (a_0, a_1)$. Since the strand $(\beta_j)_j$ has a stop in the interval $[a_0, a_1)$ by Lemma 8.2 it cannot have a stop in $[a_1, a_2)$. It follows that $b_1 > a_2 = a_0 + 2r$, hence the arcs $\alpha_0 = \widehat{a_0, a_1}$ and $\beta_0 = \widehat{b_0, b_1}$ are not contained in a strip of width $2r$.

(3) Consider two strands $(\alpha_i)_{i \in \mathbb{Z}}$ and $(\beta_j)_{j \in \mathbb{Z}}$ as above. We want to show that they cross exactly twice in a strip $\mathcal{S}_{[t, t+2r]}$. Since both strands are invariant by horizontal translation of displacement length $2r$, we may choose t arbitrarily. We set $t = a_0$. By Lemma 8.2, the strand $(\beta_i)_i$ has stops in exactly one of the intervals (a_0, a_1) and (a_1, a_2) . Thus, it intersects (twice) exactly one of the arcs $\alpha_0 = \widehat{a_0 a_1}$, $\alpha_1 = \widehat{a_1 a_2}$. \square

We also note the following.

Lemma 8.5. *In the wall system $\mathcal{W}_{\mathcal{H}}$, there is an arc of width 1.*

Proof. Let $\alpha_0 = \widehat{a_0 a_1}$ be an arc that is minimal with respect to covering (i.e., α_0 does not cover any arc of $\mathcal{W}_{\mathcal{H}}$). We want to show that $a_1 - a_0 = 1$. By Lemma 8.2, each strand other than the one generated by α_0 has two stops in the interval (a_0, a_2) , both contained either in (a_0, a_1) or in (a_1, a_2) . Thus, if the interval (a_0, a_1) has any stop, it has in fact two stops of a strand, and therefore there is an arc of $\mathcal{W}_{\mathcal{H}}$ covered by α_0 . However, this possibility is excluded by the minimality of α_0 . Therefore, the interval (a_0, a_1) has no stops and hence its endpoints a_0 and a_1 are consecutive semi-integers. \square

9. PROOF OF THE DISCRETE AREA LOWER BOUND

We can now proceed to the proof of the discrete area lower bound for simple discrete metric disks, see Theorem 5.3, making use of the previous notations and constructions. Namely, let us prove the following.

Theorem 9.1. *The discrete area of every simple discrete metric disk of radius r is at least $\frac{3}{2}r^2$.*

Proof. Let (D, \mathcal{W}') be a simple discrete metric disk of radius r and center O that has minimal area. Recall that the punctured disk $D \setminus \{O\}$ is identified with the flat cylinder $\mathcal{C} = S^1 \times [0, +\infty)$. As shown in Section 6, \mathcal{W}' is homotopic in \mathcal{C} to a quasi wall system \mathcal{W} made of standard arcs, such that (D, \mathcal{W}) is also a discrete disk of radius r centered at O and has the same area as (D, \mathcal{W}') . Thus we must show that $\text{area}(D, \mathcal{W}) \geq \frac{3}{2}r^2$. Also,

we may assume that the lift of \mathcal{W} to the universal cover $\mathcal{H} = \mathbb{R} \times [0, +\infty)$ is a quasi wall system $\mathcal{W}_{\mathcal{H}}$ such that $\partial\mathcal{W}_{\mathcal{H}} = \mathbb{Z} + \frac{1}{2} \subseteq \mathbb{R} = \partial\mathcal{H}$ as in Section 8.

Let $t \in \mathbb{R}$ be a generic number. By Proposition 8.4, the weighted number of self-intersections of the quasi wall system $\mathcal{W}_{\mathcal{H}}$ that lie in the strip $\mathcal{S}_{[t, t+2r]}$ is

$$\#I_{\mathcal{H}^\circ + \frac{1}{2}\partial\mathcal{H}}(W_{\mathcal{H}}|_{\mathcal{S}_{[t, t+2r]}}) = 2\frac{r(r-1)}{2} + \frac{1}{2}2r = r^2. \quad (9.1)$$

The first term counts the crossings between the different strands: each pair of strands crosses twice, and the crossings are located in the interior of the half-plane \mathcal{H} . The second term counts, with weight $\frac{1}{2}$, the intersections that lie in the boundary $\partial\mathcal{H}$; these are the intersections between adjacent arcs, that belong to the same strand. Thus, the discrete area of the disk (D, \mathcal{W}) is

$$\text{area}(D, \mathcal{W}) = \#I_{\mathcal{H}^\circ + \frac{1}{2}\partial\mathcal{H}}(W_{\mathcal{H}}|_{\mathcal{S}_{[t, t+n]}}) = \frac{n}{2r} \#I_{\mathcal{H}^\circ + \frac{1}{2}\partial\mathcal{H}}(W_{\mathcal{H}}|_{\mathcal{S}_{[t, t+2r]}}) = \frac{n}{2r} r^2,$$

where n is the number of walls of \mathcal{W} .

To finish we will show that $n \geq 3r$. Let $(\alpha_i = \widehat{a_i a_{i+1}})_{i \in \mathbb{Z}}$ be a strand of $\mathcal{W}_{\mathcal{H}}$ such that $a_0 - a_{-1} = 1$. Such a strand exists by Lemma 8.5. Moreover, we may assume that $a_0 = \frac{1}{2}$ and $a_{-1} = -\frac{1}{2}$. The interval (a_0, a_1) has width $2r - 1$ (by Lemma 8.3) and contains $2r - 2$ semi-integers.

Each of these semi-integers is either the startpoint or the endpoint of one of the $r - 1$ arcs that are covered by α_0 ; see Proposition 8.4. Let b_0 be the rightmost of the $r - 1$ startpoints. Note that

$$b_0 \geq a_0 + (r - 1). \quad (9.2)$$

This point b_0 is a stop of a strand $(\beta_j = \widehat{b_j b_{j+1}})_{j \in \mathbb{Z}}$. The arc β_0 is covered by α_0 and the arc $\beta_1 = \widehat{b_1 b_2}$ intersects the arc α_0 . The arcs α_0 and β_1 cannot extend over a whole fundamental domain $\mathcal{S}_{[t, t+n]}$ of the universal cover, by the property (1) of Proposition 6.3. Therefore, $n > b_2 - a_0$. On the other hand, by Lemma 8.3 and the inequality (9.2), we have

$$b_2 = b_0 + 2r \geq a_0 + 3r - 1.$$

We conclude that $n > 3r - 1$, or, equivalently, $n \geq 3r$, as we had to prove. \square

10. SIMPLE DISCRETE METRIC DISKS OF MINIMAL AREA

In this section, we analyze the equality case of Theorem 9.1.

Proposition 10.1. *For every positive integer r , there is a simple discrete metric disk of radius r and area $\frac{3}{2}r^2$. It is unique up to isotopy of the disk with the center fixed.*

Proof. Recall the proof of Theorem 9.1. Let \mathcal{W}' be a simple quasi wall system such that (D, \mathcal{W}') is a simple discrete metric disk of radius r with minimal discrete area. Consider the simple quasi wall system \mathcal{W} homotopic to \mathcal{W}' made of standard arcs. To attain the lower bound on $\text{area}(D, \mathcal{W})$ and so on $\text{area}(D, \mathcal{W}')$, we must have $n = 3r$, therefore the inequality (9.2)

must be an equality. This implies that, for the $r - 1$ arcs covered by α_0 , the $r - 1$ startpoints must precede the $r - 1$ endpoints in the interval (a_0, a_1) . In consequence, these $r - 1$ arcs together with the arc α_0 form a chain with respect to the covering relation; see Figure 5. This implies that the r arcs are completely determined, and by Proposition 8.4, so are the quasi wall systems $\mathcal{W}_{\mathcal{H}}$ and \mathcal{W} , which are made of standard arcs. Thus, the wall system $\mathcal{W}_{\mathcal{H}}$ contains all arcs of the form $kr - \widehat{s}, kr + s$ with k integer and $s \in (0, r)$ semi integer; see Figure 5. Similarly, the quasi wall system \mathcal{W} is obtained from $\mathcal{W}_{\mathcal{H}}$ by taking the quotient of \mathcal{H} under the horizontal translation of displacement length $3r$; see Figure 6. This proves the uniqueness of the simple discrete metric disk of minimal area, but only up to homotopy of the quasi wall system. The uniqueness up to isotopy of the disk follows from the next result.

Lemma 10.2. *Let \mathcal{W} and \mathcal{W}' be two simple quasi wall systems on the disk D , homotopic in $D \setminus \{0\}$ and forming no triangle in $D \setminus \{O\}$. Then there is an isotopy of D which fixes O and carries \mathcal{W} to \mathcal{W}' .*

Proof. We proceed by induction in the number n of walls. The case $n = 1$ is trivial. In general, we argue as follows.

Let γ be a wall of \mathcal{W} that covers no other wall of \mathcal{W} ; see Definition 6.1. The curve γ divides the disk D into two topological closed disks A and B which intersect along γ , with $O \in A$. The part of \mathcal{W} that lies in B consists of $k \geq 0$ arcs going from γ to $\partial B \setminus \gamma$. These arcs are pairwise disjoint, otherwise they would form a triangle in $B \subseteq D \setminus \{O\}$. The part of \mathcal{W} that lies in A , excluding γ , is a quasi wall system on A with $n - 1$ walls.

Let γ' be the wall of \mathcal{W}' homotopic to γ in $D \setminus \{O\}$. We apply to \mathcal{W}' a first isotopy of $D \setminus \{O\}$ to ensure that $\gamma' = \gamma$. The wall γ' does not cover any other wall β' of \mathcal{W}' , otherwise the wall β of \mathcal{W} homotopic to β' would cross γ twice. Similarly as in \mathcal{W} , the part of \mathcal{W}' lying in B consists of k pairwise disjoint arcs going from γ to $\partial B \setminus \gamma$. Thus, by applying a second isotopy, we may ensure that $\mathcal{W}' \cap B = \mathcal{W} \cap B$. Finally, we get $(\mathcal{W}' \setminus \gamma') \cap A = (\mathcal{W} \setminus \gamma) \cap A$ by applying an isotopy of the disk A fixing O , whose existence is guaranteed by the inductive hypothesis. \square

Now, the walls of \mathcal{W} do not delimit any triangle in $D \setminus \{O\}$ (where each side lies in a wall); see Figures 6 and 5. Since two arcs of \mathcal{W} intersect each other if and only if the same holds with the corresponding homotopic arcs of \mathcal{W}' , we deduce that the walls of \mathcal{W}' do not form any triangle in $D \setminus \{O\}$ either. The uniqueness of the simple discrete metric disk of minimal area up to isotopy of the disk fixing O follows from Lemma 10.2. \square

Remark 10.3. The isotopy between \mathcal{W} and \mathcal{W}' can also be derived from [GS97], where it is proved that two wall systems on a closed surface which are homotopic to each other and are both in minimally crossing position (*i.e.*, they attain the minimum number of self-intersections possible in their




FIGURE 5. The lift $W_{\mathcal{H}}$ corresponding to an area minimizing simple discrete disk (D, \mathcal{W}) of radius $r = 5$, where the quasi wall system \mathcal{W} consists of standard arcs.




FIGURE 6. An area minimizing simple discrete disk (D, \mathcal{W}) of radius $r = 5$ where the topological disk D is a hexagon and the quasi wall system \mathcal{W} consists of straight lines.

homotopy class) can be obtained one from the other by isotopies and triangle flip moves (called “type III moves” in [GS97]). Strictly speaking, we first need to adapt this result to quasi wall systems on surfaces with boundary. Since \mathcal{W} and \mathcal{W}' do not form any triangle in $D \setminus \{0\}$, we conclude that they are isotopic in D .

11. HILBERT’S FOURTH PROBLEM AND EXTREMAL FINSLER DISKS

In this section, we construct a Finsler disk of radius r with minimizing interior geodesics whose area is arbitrarily close to the lower bound $\frac{6}{\pi}r^2$ given by Theorem 1.2.

Let us first go over Busemann’s construction of projective metrics in relation with Hilbert’s fourth problem. We refer to [Pog79], [Ale78], [Sza86], [Pap14] and references therein for an account on the subject.

The space Γ of oriented lines in \mathbb{R}^2 can be identified with $S^1 \times \mathbb{R}$. Under this identification, an oriented line γ is represented by a pair $(e^{i\theta}, p)$ where $e^{i\theta}$ is the direction of the oriented line γ and $p = \langle \overrightarrow{OH} \times e^{i\theta}, \vec{e}_z \rangle$ is the signed distance from the origin O to γ . Here, H is a point of γ , the vector \vec{e}_z is the third vector in the canonical basis of \mathbb{R}^3 , thus it is a unit vector orthogonal to \mathbb{R}^2 , and “ \times ” is the vector product in \mathbb{R}^3 .

Definition 11.1. Let μ be a (nonnegative) Borel measure on Γ . Consider the following conditions:

- (1) the measure is invariant under the involution of Γ reversing the orientation of lines;
- (2) the measure of every compact subset of Γ is finite;
- (3) the set of all oriented lines passing through any given point of \mathbb{R}^2 has measure zero;
- (4) the set of all oriented lines passing through any given line segment in \mathbb{R}^2 has positive measure.

A Borel measure μ satisfying (1)-(3) induces a length function

$$\text{length}_\mu(\alpha) = \frac{1}{4} \int_{\gamma \in \Gamma} \#(\gamma \cap \alpha) \, d\mu(\gamma)$$

defined for any curve α in the plane \mathbb{R}^2 . For this kind of length function, straight segments are shortest paths, therefore the pseudo-distance associated to this length function is

$$d_\mu(x, y) = \frac{1}{4} \int_{\gamma \in \Gamma} \#(\gamma \cap [x, y]) \, d\mu(\gamma) = \frac{1}{4} \mu(\Gamma_{[x, y]}),$$

where Γ_A denotes the set of lines $\gamma \in \Gamma$ that intersect a subset or point A contained in the plane \mathbb{R}^2 . The pseudo-distance d_μ is *projective*, which means that $d(x, z) = d(x, y) + d(y, z)$ for every $x, y, z \in \mathbb{R}^2$ with $y \in [x, z]$, and in fact every continuous projective distance is obtained from a unique measure μ ; see [Ale78]. If μ also satisfies (4) then d_μ is a projective distance (and vice-versa).

For example, the product measure λ , given by $d\lambda = d\theta \, dp$, yields the Euclidean distance $d_\lambda(x, y) = |y - x|$.

The projective distance induced by a Borel measure satisfying the conditions (1)–(4) is not Finsler in general. Borel measures inducing a Finsler metric can be characterized as follows; see [Álv05] for a presentation of this result due to Pogorelov [Pog79] and [ÁB10] for a generalization.

Theorem 11.2. *Let μ be a Borel measure on Γ satisfying (1)–(4). The distance d_μ is Finsler if and only if μ is a positive smooth measure. In this case, the smooth measure on Γ induced by the symplectic form associated to the Finsler metric, see (3.8), coincides with μ .*

Here, a measure μ on Γ is (*positive*) *smooth* if it admits a (positive) smooth function h as density, that is, $d\mu = h \, d\lambda$.

Remark 11.3. The geodesics of a plane with a projective Finsler metric d_μ are the straight lines parametrized by μ -length. Therefore, a plane with a projective Finsler metric has minimizing geodesics.

We may define the area of a Borel set D in the plane with a measure μ on Γ satisfying (1)–(3) by the Santaló+Blaschke formula (3.6)

$$\text{area}_\mu(D) = \frac{1}{8\pi} \int_{\gamma_0 \in \Gamma} \int_{\gamma_1 \in \Gamma} \#(\gamma_0 \cap \gamma_1 \cap D) \, d\mu(\gamma_1) \, d\mu(\gamma_0). \quad (11.1)$$

In other terms, the area measure is the normalized pushforward measure

$$\text{area}_\mu = \frac{1}{8\pi} i_* (\mu \times \mu) \quad (11.2)$$

where $i : \Gamma \times \Gamma \setminus \Delta_\Gamma \rightarrow \mathbb{R}P^2$ maps each ordered pair of different lines to its intersection point in the projective plane $\mathbb{R}P^2 \supseteq \mathbb{R}^2$. (Note that the diagonal Δ_Γ has measure zero because μ has no atoms.) This area function coincides with Holmes–Thompson area if the metric is Finsler; see (3.6).

11.1. Construction of a non-Finsler extremal disk. Let us construct a non-Finsler projective pseudo-metric disk satisfying the equality case in Theorem 1.2. Consider the three pairs of one-parameter families L_k^\pm of oriented lines in \mathbb{R}^2 defined as

$$\begin{aligned} L_k^+ : \mathbb{R}_+ &\rightarrow \Gamma = S^1 \times \mathbb{R} & \text{and} & & L_k^- : \mathbb{R}_+ &\rightarrow \Gamma = S^1 \times \mathbb{R} \\ t &\mapsto (e^{i\frac{2k\pi}{3}}, t) & & & t &\mapsto (e^{i(\frac{2k\pi}{3} + \pi)}, -t) \end{aligned}$$

where $k \in \{0, 1, 2\}$; see Figure 7. Note that the lines $L_k^+(t)$ and $L_k^-(t)$ only differ by their orientation. We will sometimes denote these families of lines by L_k when the orientation does not matter. Consider the (nonsmooth) Borel measure on Γ

$$\mu_{\text{ext}} = \nu_0 + \nu_1 + \nu_2$$

where

$$\nu_k = \frac{1}{2} [(L_k^+)_*(\mathcal{L}) + (L_k^-)_*(\mathcal{L})]$$

is the average of the push-forwards to Γ of the Lebesgue measure \mathcal{L} on \mathbb{R}_+ . Let D_k be the line passing through O orthogonal to L_k . Let $\overline{D}_k \subseteq D_k$ be the ray from O that intersects orthogonally every line $L_k(t)$. Denote by π_k the orthogonal projection of \mathbb{R}^2 to D_k . By construction, the d_{ν_k} -pseudo-distance between two points $x, y \in \mathbb{R}^2$ is equal to one quarter times the Euclidean length of the projection of $[x, y]$ to D_k lying in \overline{D}_k . That is,

$$d_{\nu_k}(x, y) = \frac{1}{4} \text{length}(\pi_k([x, y]) \cap \overline{D}_k) \leq \frac{1}{4} |x - y|$$

for every $x, y \in \mathbb{R}^2$. Furthermore,

$$d_{\mu_{\text{ext}}}(x, y) = \sum_{k=0,1,2} d_{\nu_k}(x, y) = \frac{1}{4} \sum_{k=0,1,2} \text{length}(\pi_k([x, y]) \cap \overline{D}_k).$$

Observe also that the measure μ_{ext} satisfies (1)–(3), but not (4). Thus, $d_{\mu_{\text{ext}}}$ is a projective pseudo-distance on \mathbb{R}^2 .




FIGURE 7. Extremal pseudo-metric disk

The disk $D_{\mu_{\text{ext}}}(r)$ of radius r for the pseudo-distance $d_{\mu_{\text{ext}}}$ with center the origin O of \mathbb{R}^2 is the minimal regular hexagon containing the Euclidean disk of radius $4r$, whose vertices are $\frac{2\sqrt{3}}{3}4re^{ik\frac{\pi}{3}}$ for $k \in \{0, \dots, 5\}$; see Figure 7. A direct computation using (11.1) shows that its area is $\frac{6}{\pi}r^2$. Thus, the disk $D_{\mu_{\text{ext}}}(r)$ is a non-Finsler projective pseudo-metric disk satisfying the equality case in Theorem 1.2. One can think of it as an extremal (degenerate) metric for the problem considered. Observe also that $D_{\mu_{\text{ext}}}(r)$ is not rotationally symmetric.

Remark 11.4. By identifying all pairs of points at zero pseudo-distance, the pseudo-metric disk $D_{\mu_{\text{ext}}}(r)$ identifies with the closed ball $D(r)$ of radius r centered at the tip of a cone composed of three copies of a quadrant of the ℓ^1 -plane glued together. It follows from a direct computation that the Holmes–Thompson area of the disk $D(r)$ is equal to $\frac{6}{\pi}r^2$. Defined in this way, the metric on $D(r)$ is non-Finsler (*e.g.*, it has a singularity at the origin and the tangent norms are neither smooth nor strongly convex) but can still be thought of as an extremal (degenerate) metric. As previously, we observe that $D(r)$ is not rotationally symmetric. Note that the (pseudo)-metrics on $D_{\mu_{\text{ext}}}(r)$ and $D(r)$ can be viewed as continuous versions of the extremal simple discrete disk; see Section 10.

11.2. Construction of a Finsler nearly extremal disk. In the rest of this section, we explain how to modify the pseudo-metric $d_{\mu_{\text{ext}}}$ so as to obtain a projective Finsler disk of radius r whose area is arbitrarily close to $\frac{6}{\pi}r^2$. First, the projective pseudo-metric $d_{\mu_{\text{ext}}}$ can be approximated by a projective metric by simply adding to μ a multiple $\varepsilon\lambda$ of the uniform measure λ (given by $d\lambda = d\theta dp$) so that the point (4) is also satisfied; this changes $d_{\mu_{\text{ext}}}$ by adding ε times the Euclidean distance. This projective metric is not Finsler, but in turn it can be approximated by a Finsler metric; see [Pog79]. More generally, every projective distance d_μ , where μ is a Borel measure satisfying (1)-(4), can be approximated by a projective Finsler distance on every compact set of \mathbb{R}^2 . Thus, by Theorem 11.2, there exists a sequence μ_n

of positive smooth measures on Γ such that the corresponding sequence of Finsler distances d_{μ_n} uniformly converges to $d_{\mu_{\text{ext}}}$ on every compact set of \mathbb{R}^2 . This approximation result is obtained by a convolution argument on the distance function d_μ . Although it is possible that the measures μ_n weakly converge to μ_{ext} , this issue is not addressed in [Pog79]. This leads us to consider a slightly different approach. Instead of regularizing the distance function, we smooth out the measure μ_{ext} and show that the corresponding projective Finsler distance converges to $d_{\mu_{\text{ext}}}$. This alternative approach to the regularization of a projective distance provides a weak convergence of measure by construction, which allows us to estimate areas as well as distances.

We proceed as follows. First, we truncate the measure μ_{ext} by setting a bound for the absolute value of the p coordinate of the lines $\gamma \in \Gamma$. In this way, we obtain a probability measure μ_0 on Γ , without changing the corresponding distance function in a neighborhood of the origin. Similarly, we truncate the uniform measure λ to a probability measure λ_0 . This enables us to use standard theorems on weak convergence of probability measures.

Let us now describe the convolution process. For $\varepsilon > 0$, let h_ε be a smooth nonnegative function on $\Gamma = \mathbb{R}/2\pi\mathbb{Z} \times \mathbb{R}$, with support in $(-\varepsilon, \varepsilon) \times (-\varepsilon, \varepsilon)$, such that $\int_\Gamma h_\varepsilon(\theta, p) \, d\theta \, dp = 1$. For each $\varepsilon > 0$, consider the positive smooth measure μ_ε on Γ with density $h_\varepsilon * \mu_0$, that is,

$$d\mu_\varepsilon = (h_\varepsilon * \mu_0) \, d\lambda$$

where $h_\varepsilon * \mu_0$ is the smooth function on Γ defined by the convolution

$$h_\varepsilon * \mu_0(\gamma) = \int_\Gamma h_\varepsilon(\gamma - \gamma') \, d\mu_0(\gamma')$$

and λ is the standard product measure on $\Gamma = \mathbb{R}/2\pi\mathbb{Z} \times \mathbb{R}$, given by $d\lambda = d\theta \, dp$. By [Bog18, §1.4.3], the smooth measure μ_ε weakly converges to μ_0 as ε goes to zero. Define also the measure

$$\mu_\varepsilon^+ = (1 - \varepsilon)\mu_\varepsilon + \varepsilon \lambda_0,$$

which also converges to μ_0 as $\varepsilon \rightarrow 0$. By Theorem 11.2, the distance $d_{\mu_\varepsilon^+}$ induced by μ_ε^+ is a projective Finsler distance on a neighborhood of the origin in \mathbb{R}^2 .

To approximate distances and areas, we have the following tools.

Lemma 11.5. *Let μ and μ_n be probability measures on Γ satisfying the conditions (1)–(3) of Definition 11.1. If μ_n weakly converges to μ , then the distance d_{μ_n} converges uniformly to d_μ on every compact subset of \mathbb{R}^2 .*

Proof. Note first that the distance between two points is

$$d_\mu(x, y) = \mu(\Gamma_{[x,y]})$$

where $\Gamma_{[x,y]}$ denotes the set of lines that intersect the segment $[x, y]$. Let \mathcal{A} be the family of sets $\Gamma_{[x,y]} \subseteq \Gamma$, where x and y lie in a fixed compact set $K \subseteq \mathbb{R}^2$.

According to Theorems 2 and 3 from [BT67], to show uniform convergence $\mu_n(A) \rightarrow \mu(A)$ for all sets $A \in \mathcal{A}$, it is sufficient to show that $\mu(B_\delta(\partial A)) \rightarrow 0$ uniformly as $\delta \rightarrow 0$, where $B_\delta(S)$ denotes the δ -neighborhood of a set $S \subseteq \Gamma$ (say, with respect to the supremum distance in terms of the coordinates θ, p). Note also that

$$\partial\Gamma_{[x,y]} = \Gamma_x \cup \Gamma_y,$$

where Γ_z is the set of lines that contain a point z . (In the coordinates θ, p , the set Γ_z is the graph of a sinusoidal function with amplitude proportional to the Euclidean distance of z to the origin.) Therefore it suffices to show that $\mu(B_\delta(\Gamma_x)) \rightarrow 0$ uniformly with respect to $x \in K$, as $\delta \rightarrow 0$. Suppose that this is not the case. Then there are sequences $\delta_m \rightarrow 0$ and $x_m \rightarrow x \in K$ such that $\mu(B_{\delta_m}(\Gamma_{x_m}))$ does not tend to zero. However, we also have

$$B_{\delta_m}(\Gamma_{x_m}) \subseteq B_{\delta'_m}(\Gamma_x)$$

for some sequence $\delta'_m \rightarrow 0$, which yields $B_{\delta'_m}(\Gamma_x) \rightarrow \Gamma_x$ and therefore

$$\mu(B_{\delta_m}(\Gamma_{x_m})) \leq \mu(B_{\delta'_m}(\Gamma_x)) \rightarrow 0.$$

since $\mu(\Gamma_x) = 0$ by the condition (3) on μ . This contradiction finishes the proof. \square

Lemma 11.6. *Let μ and μ_n be probability measures on Γ satisfying the conditions (1)–(3) of Definition 11.1. If μ_n weakly converges to μ , then $\text{area}_{\mu_n}(K)$ converges to $\text{area}_\mu(K)$ for every compact set $K \subseteq \mathbb{R}^2$ such that $\mu(\partial K) = 0$.*

Proof. We will use some properties of weak convergence of probability measures; see [Bil99]. Since μ_n weakly converges to μ , it follows from [Bil99, Theorem 3.2] that the product measure $\mu_n \times \mu_n$ converges weakly to $\mu \times \mu$ on $\Gamma \times \Gamma$. The fact that μ has no atoms implies that the diagonal $\Delta_\Gamma \subseteq \Gamma \times \Gamma$ has zero measure, that is, $\mu \times \mu(\Delta_\Gamma) = 0$, and the same holds for the measures μ_n . Thus, we may restrict the measures $\mu \times \mu$ and $\mu_n \times \mu_n$ to the set $\Gamma \times \Gamma \setminus \Delta_\Gamma$ and they remain probability measures. Moreover, since the diagonal Δ_Γ is a closed set, the product measure $\mu_n \times \mu_n$ weakly converges to $\mu \times \mu$ on $\Gamma \times \Gamma \setminus \Delta_\Gamma$; see condition (iv) of the portmanteau theorem [Bil99, Theorem 2.1]. Furthermore, since the function $i : \Gamma \times \Gamma \setminus \Delta_\Gamma \rightarrow \mathbb{R}P^2$ is continuous, the pushforward measure $i_*(\mu_n \times \mu_n)$ weakly converges to $\mu \times \mu$. This follows from the definition of weak convergence; see [Bil99, p. 29]. Therefore, the area measure $\text{area}_{\mu_n} = \frac{1}{8\pi} i_*(\mu_n \times \mu_n)$ weakly converges to area_μ , with both area measures considered as probability measures on the projective plane $\mathbb{R}P^2$; see (11.2). Finally, to show that $\text{area}_{\mu_n}(K) \rightarrow \text{area}_\mu(K)$, we must check, according to part (v) of the portmanteau theorem [Bil99, Theorem 2.1], that K is a continuity set of area_μ , which by definition means that $\mu(\partial_{\mathbb{R}P^2} K) = 0$. This follows from the facts that K is compact and $\mu(\partial_{\mathbb{R}^2} K) = 0$. \square

Consider the disk $D_{\mu_\varepsilon^+}(r)$ centered at O of radius r for the distance $d_{\mu_\varepsilon^+}$. The numbers $r > 0$ and $\varepsilon > 0$ are small enough so that the truncations

of μ_{ext} and λ have no effect on the disk $D_{\mu_\varepsilon^+}(r)$. The number r is fixed while ε goes to 0.

Proposition 11.7. *The disk $D_{\mu_\varepsilon^+}(r)$ is a projective Finsler disk with minimizing interior geodesics, whose area converges to $\frac{6}{\pi}r^2$, as ε goes to zero. Therefore, the area lower bound in Theorem 1.2 is sharp.*

Proof. The fact that $d_{\mu_\varepsilon^+}$ is a projective Finsler metric follows from Theorem 11.2 and the fact that its geodesics are minimizing was stated in Remark 11.3.

To compute its area we proceed as follows. By uniform convergence of the metrics, see Lemma 11.5, for every $\delta > 0$ and every $\varepsilon > 0$ small enough, we have

$$D_{\mu_0}(r - \delta) \subseteq D_{\mu_\varepsilon^+}(r) \subseteq D_{\mu_0}(r + \delta).$$

Therefore,

$$\text{area}_{\mu_\varepsilon^+}(D_{\mu_0}(r - \delta)) \leq \text{area}_{\mu_\varepsilon^+}(D_{\mu_\varepsilon^+}(r)) \leq \text{area}_{\mu_\varepsilon^+}(D_{\mu_0}(r + \delta)).$$

Since the sets $D_{\mu_0}(r \pm \delta)$ are compact and have boundary of μ_0 -measure zero, Lemma 11.6 shows that

$$\text{area}_{\mu_\varepsilon^+}(D_{\mu_0}(r \pm \delta)) \rightarrow \text{area}_{\mu_0}(D_{\mu_0}(r \pm \delta))$$

as $\varepsilon \rightarrow 0$. We conclude that

$$\text{area}_{\mu_\varepsilon^+}(D_{\mu_\varepsilon^+}(r)) \rightarrow \text{area}_{\mu_0}(D_{\mu_0}(r))$$

as $\varepsilon \rightarrow 0$. □

12. APPENDIX A: DIFFERENTIABILITY OF DISTANCE REALIZING CURVES ON FINSLER SURFACES WITH BOUNDARY

Consider a manifold M with nonempty boundary endowed with a Finsler metric F . Recall that a distance realizing curve is a curve $\alpha : I \rightarrow M$ defined on an interval $I \subseteq \mathbb{R}$ such that

$$d_F(\alpha(t), \alpha(t')) = t' - t$$

for every $t < t'$.

Regularity and convexity assumptions on the metric ensuring that distance realizing arcs are C^1 have been studied by Busemann and Mayer for manifolds without boundary; see [BM41]. Here, we are not primarily interested in metrics of low regularity. Instead, our concern lies in the possible lack of convexity of the manifold at the boundary. Without convexity, the distance realizing arcs do not necessarily satisfy a differential equation, therefore the standard techniques to prove that geodesics are smooth do not apply. This is exemplified by a Euclidean plane with an open disk removed, where distance realizing arcs do not have more than one derivative in general and are not determined by their initial velocity vector.

In the case of Riemannian manifolds, it was shown in [Wol79] and [AA81] that distance realizing paths are C^1 . Here, by adapting the argument of [AA81], we prove that the same result holds for Finsler surfaces.

Theorem 12.1. *On a Finsler surface M , every distance realizing arc $\alpha : I \rightarrow M$ is C^1 . Furthermore, the velocity vectors $\alpha'(t)$ have unit norm.*

Let us introduce some technical definitions. Without loss of generality, using a coordinate system, we will assume that the manifold M is the closed upper half of \mathbb{R}^n .

Definition 12.2. Let $\alpha : I \rightarrow M$ be a continuous curve, where $I \subseteq \mathbb{R}$ is an interval. Fix $t_0 \in I$ and denote $x_0 = \alpha(t_0)$. An *arrival velocity* of α at t_0 is a vector $v \in T_{x_0}M$ that is an accumulation point of the set of vectors

$$V^- = \left\{ \frac{\alpha(t) - \alpha(t_0)}{t - t_0} \mid t < t_0 \right\}$$

as t goes to t_0 . Similarly, a *departure velocity* of α at t_0 is a vector $v \in T_{x_0}M$ that is an accumulation point of the set of vectors

$$V^+ = \left\{ \frac{\alpha(t) - \alpha(t_0)}{t - t_0} \mid t > t_0 \right\}$$

as t goes to t_0 . Note that if α is differentiable on the left (resp. right) at t_0 , then α has exactly one arrival (resp. departure) velocity at t_0 .

We begin by proving a weak differentiability result.

Lemma 12.3. *Let (M, F) be a Finsler manifold and let $\alpha : I \rightarrow M$ be a distance realizing arc. Fix $t_0 \in I$ and denote $x_0 = \alpha(t_0)$. Then*

- (1) *The curve α has at least one arrival velocity and one departure velocity at t_0 (unless $t_0 = \min I$ or $t_0 = \max I$, respectively).*
- (2) *Every arrival or departure velocity v has norm $F_{x_0}(v) = 1$.*
- (3) *If the curve α is differentiable on one side at an interior point t_0 of I , then α is differentiable at t_0 .*

Proof. By continuity of the Finsler metric at x_0 , we can bound F_x below and above by two multiples of the norm $F_{x_0} = |\cdot|$ for every x close enough to x_0 . That is,

$$\lambda^- |v| \leq F_x(v) \leq \lambda^+ |v|$$

for every $v \in \mathbb{R}^n$, which in turn implies that

$$\lambda^- |x - x_0| \leq d_F(x_0, x) \leq \lambda^+ |x - x_0|.$$

This implies that the sets of vectors V^\pm are bounded when t goes to t_0 , which implies the first claim. In fact, as x goes to x_0 , the optimal coefficients λ^\pm converge to 1, which implies the second claim.

To prove the last claim, we assume that the curve α is differentiable on the left at an interior point t_0 of I . (The argument is similar if α is differentiable on the right at t_0 .) Let v^- be the arrival tangent vector. Let us prove that α is differentiable on the right at t_0 and has departure tangent vector $v^+ = v^-$.

By contradiction, assume that the set of vectors V^+ has an accumulation point $v^+ \neq v^-$ as t goes to t_0 . As already noticed in the second claim, we have $|v^-| = |v^+| = 1$. Since the norm $F_{x_0} = |\cdot|$ is strictly convex, we also have $|v^- + v^+| < 2$. Let $\tau_m \rightarrow 0$ be a decreasing sequence of positive numbers such that

$$y_m^+ = \alpha(t_0 + \tau_m) = \alpha(t_0) + \tau_m v^+ + o(\tau_m).$$

Since α is differentiable on the left at t_0 , we also have

$$y_m^- = \alpha(t_0 - \tau_m) = \alpha(t_0) - \tau_m v^- + o(\tau_m).$$

Thus,

$$d_F(y_m^+, y_m^-) \leq \lambda^+ |y_m^+ - y_m^-| = \lambda^+ \tau_m |v^+ + v^- + o(1)|.$$

For m large enough, we can take λ^+ arbitrarily close to 1. It follows from the inequality $|v^+ + v^-| < 2$ that

$$d_F(y_m^-, y_m^+) < 2\tau_m$$

contradicting that α is a distance realizing arc. \square

Before proceeding to the proof of Theorem 12.1, we extend the Finsler metric F to a surface $M^+ \supseteq M$ with empty boundary; see Remark 2.2. As for any Finsler surface with empty boundary, every point of M^+ has a normal neighborhood, that is, an open neighborhood U such that for any two points $x, y \in U$, there is a unique geodesic from x to y contained in U and this geodesic is the unique distance-realizing path from x to y in M^+ ; see [BCS00, p. 160]. Note that if this geodesic is contained in M , then it is also the unique distance-realizing path from x to y in M .

Proof of Theorem 12.1. We assume first that the metric is self-reverse.

Let $\alpha : I \rightarrow M$ be a distance realizing arc. Let $t_0 \in I$ and let $x_0 = \alpha(t_0)$. If $x_0 = \alpha(t_0)$ lies in the interior of M then the arc α coincides with a geodesic in a neighborhood of t_0 , where it is C^1 (and we are done). Thus, we can assume that x_0 lies in ∂M .

Again, we assume without loss of generality by working in a small enough neighborhood of x_0 that M is a closed half-space of $M^+ = \mathbb{R}^2$ and that every geodesic is a unique distance-realizing path.

Suppose that the arc α is not differentiable on the right at some $t_0 \in I$. (The argument is similar if α is not differentiable on the left at t_0 .) The arc α has two departure velocities v and w . Let K_v and K_w be two convex cones based at x_0 that contain the points $x_0 + v$ and $x_0 + w$ in their interior and only meet at x_0 . Take a unit vector $u \in T_{x_0}M$ not tangent to ∂M that points in the interior of M and separates K_v from K_w , and denote by γ_u the geodesic with initial velocity $\gamma'_u(t_0) = u$. This geodesic does not visit K_v nor K_w in some interval (t_0, t_2) . On the other hand, the arc $\alpha(t)$ visits the cones K_v and K_w infinitely many times in any interval (t_0, τ) , with $\tau > t_0$. Therefore, it must cross the geodesic γ_u at some time $t_1 \in (t_0, t_2)$. Since γ_u is the unique distance-realizing path between any of its points, the arc α

coincides with γ_u in $[t_0, t_1]$. Thus α does not visit K_v and K_w in (t_0, t_1) . This contradiction proves that α is differentiable on the right at t_0 . It follows from Lemma 12.3 that α is differentiable at every interior point $t_0 \in I$.

Suppose α is not C^1 on the right at t_0 . (The argument is similar in case it is not C^1 on the left.) The vector $v = \alpha'(t_0)$ points inside M or is tangent to the boundary of M . Since the velocities $\alpha'(t)$ are unit vectors and the curve α is not C^1 on the right at t_0 , its derivative α' has an accumulation point $w \neq v$ when t goes to t_0 from the right. Let u be a unit vector spanning a line that separates v from w . Consider three disjoint neighbourhoods U, V, W of u, v, w such that for every u', v', w' in U, V, W respectively, the line spanned by u' separates v' from w' . Let K_V be the union of the rays contained in M starting at x_0 with direction $v' \in V$, and let R be any of these rays. Note that u is transverse to all these rays. Working in a small enough neighbourhood of x_0 , we can assume that the family Γ of geodesics that visit R with velocity u foliates the cone K_V , and that their tangent vectors do not deviate too much from u and thus lie in U . Since the velocity of α at t_0 lies in the open set V , the arc α restricted to some nontrivial interval $[t_0, t_3)$ lies in K_V . Now, since w is an accumulation point for α' when t goes to t_0 from the right, there exists $t_2 \in (t_0, t_3)$ such that $w' = \alpha'(t_2)$ lies in W . Let $x_2 = \alpha(t_2)$, and let v' be the direction from x_0 to x_2 . Let γ be the geodesic of Γ passing through x_2 , and let $u' \in U$ be its velocity at x_2 . The vector $w' = \alpha'(t_2)$ points strictly inside the region of M delimited by γ containing x_0 , since the vector v' points outside, and the line generated by the vector u' separates v' from w' .

Therefore, the arc α starting at x_0 must cross γ a first time at $t_1 \in (t_0, t_2)$ before crossing it again at t_2 . Since γ is the unique distance-realizing path between $\alpha(t_1)$ and $\alpha(t_2)$, the arc α coincides with γ in $[t_1, t_2]$, which contradicts the fact that α' is transverse to γ at t_2 (or t_1). This finishes the proof of 12.1 for self-reverse metrics.

In the case of directed metrics we adapt the argument as follows. Apart from the foliation Γ , we need a second foliation Γ^- of K_V by geodesics transverse to the ray R with initial velocity $-u$. Then we proceed as in the proof and after choosing the point x_1 in K_V , we let γ and γ^- be the two geodesics of Γ and Γ^- passing through x_1 . We keep only the part of each geodesic before it reaches x_1 and discard the rest. These two half geodesics delimit a region of K_V containing x_0 . The curve α points strictly inside this region at x_1 . Therefore, it must cross either γ or γ^- a first time before reaching x_1 . We derive a contradiction as in the previous proof. \square

13. APPENDIX B: A WEAK LAW OF LARGE NUMBERS FOR MOSTLY INDEPENDENT RANDOM VARIABLES

Let us prove the following generalization of the weak law of large numbers.

Theorem 13.1 (Weak law of large numbers for identically distributed, mostly independent random variables). *Fix a real valued random variable X*

with finite expected absolute value $\mathbb{E}(|X|) < \infty$. Then the average $\bar{X} = \frac{1}{n} \sum_i X_i$ of n random variables X_i , each with the same distribution as X , is near the expected value $\mathbb{E}(X)$ with high probability if the proportion of nonindependent pairs

$$p = \frac{\#\{(i, j) \mid X_i \text{ and } X_j \text{ are not independent}\}}{n^2}$$

is low. More precisely, for every $\varepsilon, \delta > 0$, we have $\mathbb{P}(|\bar{X} - \mathbb{E}(X)| \geq \varepsilon) \leq \delta$ if $p \leq p_0(X, \delta, \varepsilon)$.

Note that we do not explicitly require n to be large, but this is generally necessary for p to be low, because each variable X_i is in general correlated with itself,³ which implies that $p \geq \frac{n}{n^2} = \frac{1}{n}$. If these are the only correlations and n goes to infinity, then $p = \frac{1}{n} \rightarrow 0$ and therefore \bar{X} converges to $\mathbb{E}(X)$ in probability. In this way, we recover the usual weak law of large numbers.

Proof. The proof is not harder than the standard proof of the weak law of large numbers; see [Tao09, Theorem 1.5.1] for instance. It combines the moment method with a truncation argument. We proceed by cases; only the first one requires attention to the non-independent pairs.

Case $\mathbb{E}(X^2) < \infty$ and $\mathbb{E}(X) = 0$. Fix $\varepsilon > 0$. We have to prove that the probability of deviation $\mathbb{P}(|\bar{X}| > \varepsilon)$ gets arbitrarily low if p is sufficiently small. To apply Chebyshev's inequality, we compute

$$\mathbb{E}(\bar{X}^2) = \frac{1}{n^2} \sum_i \sum_j \mathbb{E}(X_i X_j) \leq p \mathbb{E}(X^2).$$

Here we used the Cauchy–Schwartz inequality $\mathbb{E}(X_i X_j) \leq \mathbb{E}(X^2)$ and the fact that $\mathbb{E}(X_i X_j) = \mathbb{E}(X_i) \mathbb{E}(X_j) = \mathbb{E}(X)^2 = 0$ if X_i and X_j are independent. Applying Chebyshev's inequality, we obtain

$$\mathbb{P}(|\bar{X}| \geq \varepsilon) \leq \frac{\mathbb{E}(\bar{X}^2)}{\varepsilon^2} \leq \frac{p \mathbb{E}(X^2)}{\varepsilon^2} \xrightarrow{p \rightarrow 0} 0$$

as we had to prove.

The remaining cases are reduced to the first one in the same way as in the usual proof of the weak law of large numbers.

Case $\mathbb{E}(X^2) < \infty$. This case follows from the previous one applied to the random variable $Y = X - \mathbb{E}(X)$, which satisfies $\mathbb{E}(Y^2) < \infty$ and $\mathbb{E}(Y) = 0$.

General case $\mathbb{E}(|X|) < \infty$. Here we apply a standard truncation method. It is sufficient to show that

$$\mathbb{P}(|\bar{X} - \mathbb{E}(X)| \geq 3\varepsilon) \leq 2\delta \tag{13.1}$$

if p is small enough with respect to ε, δ and X . We may assume $\delta \leq 1$.

³A random variable is independent of itself if and only if its probability distribution is concentrated in one value.

We proceed as follows. For any cutoff value $M \geq 0$, we decompose the random variable X as a sum of a bounded part and a tail

$$X = X^{<M} + X^{\geq M} \quad (13.2)$$

where the bounded part is

$$X^{<M} = \mathbb{1}_{|X| < M} X = \begin{cases} X & \text{if } |X| < M \\ 0 & \text{otherwise,} \end{cases}$$

and the tail is

$$X^{\geq M} = \mathbb{1}_{|X| \geq M} X = \begin{cases} X & \text{if } |X| \geq M \\ 0 & \text{otherwise.} \end{cases}$$

In the same way we decompose the variables $X_i = X_i^{<M} + X_i^{\geq M}$ and define two separate average values: one for the bounded parts, $\overline{X^{<M}} = \frac{1}{n} \sum_i X_i^{<M}$, and one for the tails, $\overline{X^{\geq M}} = \frac{1}{n} \sum_i X_i^{\geq M}$. These averages satisfy $\overline{X} = \overline{X^{<M}} + \overline{X^{\geq M}}$.

A key fact about the decomposition (13.2) is that the expected absolute value $\mathbb{E}(|X^{\geq M}|)$ of the tail part gets arbitrarily small if M is sufficiently large. This follows from the pointwise convergence $|X^{\geq M}| \rightarrow 0$ as $M \rightarrow +\infty$, which is dominated by $|X|$, or from the formula

$$\mathbb{E}(|X^{\geq M}|) = \mathbb{E}(|X|^{\geq M}) = \int_M^{+\infty} x \, d\mathbb{P}_{|X|}(x),$$

where $\mathbb{P}_{|X|}$ is the probability distribution of $|X|$ on \mathbb{R} . We choose M so that

$$\mathbb{E}(|X^{\geq M}|) \leq \delta \varepsilon. \quad (13.3)$$

This implies that the average $\overline{X^{\geq M}}$ of the tail parts also has small expected absolute value

$$\mathbb{E}(|\overline{X^{\geq M}}|) \leq \mathbb{E}(|X^{\geq M}|) \leq \delta \varepsilon.$$

By Markov's inequality, this implies that $\overline{X^{\geq M}}$ is small in absolute value with high probability

$$\mathbb{P}(|\overline{X^{\geq M}}| \geq \varepsilon) \leq \frac{\mathbb{E}(|\overline{X^{\geq M}}|)}{\varepsilon} \leq \frac{\delta \varepsilon}{\varepsilon} = \delta. \quad (13.4)$$

Now, the bounded part $X^{<M}$ has finite second moment $\mathbb{E}((X^{<M})^2) < \infty$. Therefore, we may apply the previous case of the theorem, which yields

$$\mathbb{P}\left(|\overline{X^{<M}} - \mathbb{E}(X^{<M})| \geq \varepsilon\right) \leq \delta \quad (13.5)$$

if p is small enough. Conveniently, $\mathbb{E}(X^{<M})$ is near $\mathbb{E}(X)$ because

$$|\mathbb{E}(X) - \mathbb{E}(X^{<M})| = |\mathbb{E}(X^{\geq M})| \leq \mathbb{E}(|X^{\geq M}|) \leq \delta \varepsilon \leq \varepsilon.$$

Here we used (13.3) and the assumption $\delta \leq 1$. Combining this with (13.5) and (13.4) by the triangle inequality, the result (13.1) follows. \square

14. APPENDIX C: A VERSION OF LEVI’S EXTENSION LEMMA

In this section, we prove the following version of Levi’s extension (or enlargement) lemma for pseudoline arrangements. It concerns arrangements on a disk, rather than on the projective plane as in the more standard version of the lemma; see [FG17].

Proposition 14.1. *Let M be a disk with a simple quasi wall system \mathcal{W} . Then*

$$d_{\mathcal{W}}(x, y) = \text{number of walls of } \mathcal{W} \text{ that separate } x \text{ from } y. \quad (14.1)$$

for any two points $x, y \in M \setminus \mathcal{W}$.

Note that if M is a Finsler disk with minimizing interior geodesics and \mathcal{W} is geodesic, then this proposition follows from Proposition 4.7.

Proof. It is clear that

$$d_{\mathcal{W}}(x, y) \geq \text{number of walls of } \mathcal{W} \text{ that separate } x \text{ from } y.$$

To prove the reverse inequality we will show the following.

Claim 14.2. *There exists a smooth path α from x to y that is in general position with respect to $\mathcal{W} \cup \partial M$ and crosses each wall of \mathcal{W} at most once.*

Here, we say that a smooth curve α is in general position with respect to an immersed 1-submanifold N if it is regular, transverse to N and avoids the self-intersections of N . If α is piecewise smooth, we require in addition that none of its non-smooth points lie in N .

We prove the claim by induction on the number of walls. Suppose the claim is valid for any quasi wall system \mathcal{W} made of n walls. Consider a simple quasi wall system \mathcal{W}' obtained by adding an extra w' to \mathcal{W} . By inductive hypothesis, there is a smooth path α that satisfies all the conditions of the claim with respect to \mathcal{W} . By perturbing α , we ensure that it is transverse to w' as well. If α crosses w' at most once, then we are done. Otherwise, let x' and y' be the first and last points of α where α crosses w' . Note that they are generic points of w' : they are neither on \mathcal{W} , nor on ∂M . Replace the segment of α from x' to y' by the segment $[x', y']$ of w' , and let α' be the resulting curve. We claim that α' is a piecewise smooth curve, in general position with respect to \mathcal{W} , that crosses each wall of \mathcal{W} at most once. This is because the segment $[x', y']$ that we inserted only crosses the walls of \mathcal{W} that separate x' from y' (since it is part of a wall of the simple quasi wall system \mathcal{W}'), and these walls are necessarily crossed as well by the piece of α between x' and y' that we replaced.

The next step is to perturb the curve α' so that the segment $[x', y']$ is displaced sideways and away from w' and the resulting curve α'' is in general position with respect to $\mathcal{W} \cup \partial M$ and crosses \mathcal{W} the same number of times as α' does, and, in addition, is transverse to w' and crosses w' at most once. Thus, α'' is in general position with respect to $\mathcal{W}' \cup \partial M$ and crosses each wall

of \mathcal{W}' at most once, but is non-smooth at two points. To make it smooth, we modify it near these two points. \square

REFERENCES

- [Ale78] R. Alexander. “Planes for which the lines are the shortest paths between points”. *Illinois J. Math.* 22.2 (1978), 177–190. DOI: 10.1215/ijm/1256048729 (cit. on pp. 33, 34).
- [AA81] R. Alexander and S. Alexander. “Geodesics in Riemannian manifolds-with-boundary”. *Indiana Univ. Math. J.* 30.4 (1981), 481–488. DOI: 10.1512/iumj.1981.30.30039 (cit. on p. 40).
- [Álv05] J.-C. Álvarez-Paiva. “Symplectic geometry and Hilbert’s fourth problem”. *J. Differential Geom.* 69.2 (2005), 353–378. DOI: 10.4310/jdg/1121449109 (cit. on p. 34).
- [Álv06] J. Álvarez-Paiva. “Some problems on Finsler geometry”. *Handbook of differential geometry, Vol. 2.* 2006, 1–33 (cit. on p. 6).
- [ÁB06] J.-C. Álvarez-Paiva and G. Berck. “What is wrong with the Hausdorff measure in Finsler spaces”. *Adv. Math.* 204.2 (2006), 647–663. DOI: 10.1016/j.aim.2005.06.007. arXiv: math/0408413 (cit. on pp. 7, 10, 12).
- [ÁB10] J.-C. Álvarez-Paiva and G. Berck. “Finsler surfaces with prescribed geodesics” (2010). arXiv: 1002.0243 (cit. on p. 34).
- [Arn89] V. I. Arnol’d. *Mathematical methods of classical mechanics.* Vol. 60. Graduate Texts in Mathematics. Corrected reprint of the second edition. Springer-Verlag, 1989 (cit. on p. 6).
- [BCS00] D. Bao, S.-S. Chern, and Z. Shen. *An introduction to Riemann–Finsler geometry.* Vol. 200. Graduate Texts in Mathematics. Springer-Verlag, New York, 2000, xx+431. DOI: 10.1007/978-1-4612-1268-3 (cit. on p. 41).
- [Ber76] M. Berger. “Some relations between volume, injectivity radius, and convexity radius in Riemannian manifolds” (1976), 33–42 (cit. on p. 2).
- [Ber77] M. Berger. “Volume et rayon d’injectivité dans les variétés riemanniennes de dimension 3”. *Osaka Math. J.* 14.1 (1977), 191–200. URL: <http://projecteuclid.org/euclid.ojm/1200770220> (cit. on p. 2).
- [Ber80] M. Berger. “Une borne inférieure pour le volume d’une variété riemannienne en fonction du rayon d’injectivité”. *Ann. Inst. Fourier (Grenoble)* 30.3 (1980), 259–265. DOI: 10.5802/aif.802 (cit. on p. 2).
- [Bil99] P. Billingsley. *Convergence of probability measures.* Second Edition. Wiley Series in Probability and Statistics: Probability and Statistics. A Wiley-Interscience Publication. John Wiley & Sons, 1999. DOI: 10.1002/9780470316962 (cit. on p. 38).
- [BT67] P. Billingsley and F. Topsøe. “Uniformity in weak convergence”. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 7 (1967), 1–16. DOI: 10.1007/BF00532093 (cit. on p. 38).
- [Bla35] W. Blaschke. “Zur Variationsrechnung”. *Abh. Math. Sem. Univ. Hamburg* 11.1 (1935), 359–366. DOI: 10.1007/BF02940732 (cit. on p. 7).
- [Bog18] V. I. Bogachev. *Weak convergence of measures.* Vol. 234. Mathematical Surveys and Monographs. Amer. Math. Soc., 2018 (cit. on p. 37).

- [BBI01] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*. Vol. 33. Graduate Studies in Mathematics. Amer. Math. Soc., 2001. DOI: 10.1090/gsm/033 (cit. on p. 11).
- [BM41] H. Busemann and W. Mayer. “On the foundations of calculus of variations”. *Trans. Amer. Math. Soc.* 49 (1941), 173–198. DOI: 10.2307/1990020 (cit. on p. 39).
- [Cha+17] G. R. Chambers, C. Croke, Y. Liokumovich, and H. Wen. “Area of convex disks”. *Proc. Amer. Math. Soc. (to appear)* (2017). arXiv: 1701.06594 (cit. on p. 2).
- [Cos18] M. Cossarini. “Discrete Surfaces with Length and Area and Minimal Fillings of the Circle”. PhD thesis. IMPA - Instituto Nacional de Matemática Pura e Aplicada, 2018. arXiv: 2009.02415 (cit. on pp. 3, 4, 14, 15, 18).
- [Cos20] M. Cossarini. “Finsler geodesics: a square-free approach”. *In preparation* (2020) (cit. on pp. 5, 6).
- [Cro80] C. B. Croke. “Some isoperimetric inequalities and eigenvalue estimates”. *Ann. Sci. École Norm. Sup. (4)* 13.4 (1980), 419–435. URL: http://www.numdam.org/item?id=ASENS_1980_4_13_4_419_0 (cit. on p. 2).
- [Cro83] C. B. Croke. “On the volume of metric balls”. *Proc. Amer. Math. Soc.* 88.4 (1983), 660–664. DOI: 10.2307/2045458 (cit. on p. 2).
- [Cro84] C. B. Croke. “Curvature free volume estimates”. *Invent. Math.* 76.3 (1984), 515–521. DOI: 10.1007/BF01388471 (cit. on p. 2).
- [Cro09] C. B. Croke. “Area of small disks”. *Bull. Lond. Math. Soc.* 41.4 (2009), 701–708. DOI: 10.1112/blms/bdp044 (cit. on p. 2).
- [CK03] C. B. Croke and M. Katz. “Universal volume bounds in Riemannian manifolds”. *Surveys in differential geometry*. Vol. 8. Surv. Differ. Geom. Int. Press, 2003, 109–137. DOI: 10.4310/SDG.2003.v8.n1.a4 (cit. on p. 2).
- [Die72] J. Dieudonné. *Treatise on analysis. Vol. III*. Translated from the French by I. G. MacDonald, Pure and Applied Mathematics, Vol. 10-III. Academic Press, 1972 (cit. on p. 11).
- [Dur98] C. E. Durán. “A volume comparison theorem for Finsler manifolds”. *Proc. Amer. Math. Soc.* 126.10 (1998), 3079–3082. DOI: 10.1090/S0002-9939-98-04629-2 (cit. on p. 7).
- [Dur19] R. Durrett. *Probability—theory and examples*. Vol. 49. Cambridge Series in Statistical and Probabilistic Mathematics. Fifth edition. Cambridge University Press, 2019. DOI: 10.1017/9781108591034 (cit. on p. 19).
- [Fed69] H. Federer. *Geometric measure theory*. Vol. 153. Die Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 1969 (cit. on p. 11).
- [FG17] S. Felsner and J. E. Goodman. “Pseudoline Arrangements”. *Handbook of Discrete and Computational Geometry*. Third edition. 2017 (cit. on pp. 15, 45).
- [GS97] M. de Graaf and A. Schrijver. “Making curves minimally crossing by Reidemeister moves”. *J. Combin. Theory Ser. B* 70.1 (1997), 134–156. DOI: 10.1006/jctb.1997.1754 (cit. on pp. 32, 33).
- [Gro07] M. Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Modern Birkhäuser Classics. With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates. Birkhäuser, 2007 (cit. on p. 6).

- [Iva02] S. V. Ivanov. “On two-dimensional minimal fillings”. *St. Petersburg Math. J.* 13.1 (2002), 1–25 (cit. on p. 3).
- [Iva11] S. V. Ivanov. “Filling minimality of Finslerian 2-discs”. *Proc. Steklov Inst. Math.* 273.1 (2011), 176–190. DOI: 10.1134/S0081543811040079. arXiv: 0910.2257 (cit. on p. 3).
- [Men14] A. C. G. Mennucci. “Geodesics in asymmetric metric spaces”. *Anal. Geom. Metr. Spaces* 2.1 (2014), 115–153. DOI: 10.2478/agms-2014-0004 (cit. on p. 6).
- [Pap14] A. Papadopoulos. “Hilbert’s fourth problem”. *Handbook of Hilbert geometry*. Vol. 22. IRMA Lect. Math. Theor. Phys. Eur. Math. Soc., 2014, 391–431. arXiv: 1312.3172 (cit. on p. 33).
- [Pog79] A. V. Pogorelov. *Hilbert’s fourth problem*. Translated by Richard A. Silverman, Scripta Series in Mathematics. Winston & Sons, 1979 (cit. on pp. 33, 34, 36, 37).
- [San52] L. A. Santaló. “Measure of sets of geodesics in a Riemannian space and applications to integral formulas in elliptic and hyperbolic spaces”. *Summa Brasil. Math.* 3 (1952), 1–11 (cit. on p. 7).
- [San76] L. A. Santaló. *Integral geometry and geometric probability*. With a foreword by Mark Kac, Encyclopedia of Mathematics and its Applications, Vol. 1. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976, xvii+404 (cit. on p. 7).
- [Sza86] Z. I. Szabó. “Hilbert’s fourth problem. I”. *Adv. in Math.* 59.3 (1986), 185–301. DOI: 10.1016/0001-8708(86)90056-3 (cit. on p. 33).
- [Tao09] T. Tao. *Poincaré’s legacies, pages from year two of a mathematical blog. Part I*. Amer. Math. Soc., 2009 (cit. on pp. 19, 43).
- [Wol79] F. Wolter. *Interior metric, shortest paths and loops in Riemannian manifolds with not necessarily smooth boundary*. Diplomarbeit, Technische Universität Berlin. 1979 (cit. on p. 40).

SECTION DE MATHÉMATIQUES, ECOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE,
STATION 8, 1015 LAUSANNE, SWITZERLAND

E-mail address: marcos.cossarini@epfl.ch

UNIV PARIS EST CRETEIL, CNRS, LAMA, F-94010 CRETEIL, FRANCE

UNIV GUSTAVE EIFFEL, LAMA, F-77447 MARNE-LA-VALLÉE, FRANCE

E-mail address: stephane.sabourau@u-pec.fr