



HAL
open science

Editorial: Extracellular matrix in homeostasis and cancer

Ana Rita Carlos, Valérie Allamand

► **To cite this version:**

Ana Rita Carlos, Valérie Allamand. Editorial: Extracellular matrix in homeostasis and cancer. *Frontiers in Genetics*, 13, 2022, Extracellular matrix in homeostasis and cancer, 10.3389/fgene.2022.1107969 . hal-03968194

HAL Id: hal-03968194

<https://hal.science/hal-03968194>

Submitted on 1 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



OPEN ACCESS

EDITED AND REVIEWED BY
Michael Baudis,
University of Zurich, Switzerland

*CORRESPONDENCE
Ana Rita Carlos,
✉ arcarlos@fc.ul.pt

SPECIALTY SECTION
This article was submitted to Cancer
Genetics and Oncogenomics,
a section of the journal
Frontiers in Genetics

RECEIVED 25 November 2022
ACCEPTED 01 December 2022
PUBLISHED 09 December 2022

CITATION
Carlos AR and Allamand V (2022),
Editorial: Extracellular matrix in
homeostasis and cancer.
Front. Genet. 13:1107969.
doi: 10.3389/fgene.2022.1107969

COPYRIGHT
© 2022 Carlos and Allamand. This is an
open-access article distributed under
the terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Editorial: Extracellular matrix in homeostasis and cancer

Ana Rita Carlos^{1*} and Valérie Allamand^{2,3}

¹EC3c & CHANGE, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal, ²Sorbonne Université-Inserm, Centre de Recherche en Myologie, Institut de Myologie, Paris, France, ³Unit of Muscle Biology, Department of Experimental Medical Science, Lund University, Lund, Sweden

KEYWORDS

extracellular matrix, cancer, tissue homeostasis, tissue regeneration, cancer biomarkers, stress and damage

Editorial on the Research Topic Extracellular matrix in homeostasis and cancer

Organs are composed not only of a variety of cells, but also of extracellular matrices (ECMs), which provide a physical scaffold and specialized microenvironment for cells and tissues, and allow biochemical signaling. Particularly since the 1970s, the ECM started to be recognized as a highly dynamic component of tissues and the importance of ECM remodeling was found to play an important role in embryonic development, tissue repair and regeneration. In addition to the key ECM components, namely collagens, laminins, fibronectin, elastin, hyaluronan and proteoglycans, our current understanding of ECM dynamics and its tissue specificity allowed us to expand the list of ECM components, collectively termed the matrisome (Naba et al., 2012). In addition to these “core” components (around 300 in mammals), there is a variety of ECM-related molecules, which include proteins required for ECM remodeling, cytokines, chemokines and growth factors, together revealing the dynamic nature of the ECM.

For this Research Topic, we thought of bringing together contributions on how the ECM plays a critical role in tissue homeostasis and disease, with special emphasis on how ECM remodeling and mutations contribute to cancer.

In recognition of the key importance of ECM in tissue development and homeostasis, there are emerging studies using *in vitro* three-dimensional models, where the ECM composition and structure directly impacts cell proliferation, migration, differentiation or survival, mimicking *in vivo* conditions. In keeping with this notion, Carvalho and others have shown that ECM is able to modulate proliferation of mesenchymal stromal cells (MSC), a multipotent stem/progenitor cell population capable of proliferating and differentiating into a variety of lineages, such as adipogenic, chondrogenic, and osteogenic (Carvalho et al.). Moreover, the authors show that ECM derived from young donors was able to rescue the impaired proliferative and osteogenic potential of aged MSC, therefore highlighting the importance of ECM composition and its potential for therapeutic intervention.

Considering the key role of ECM in maintaining tissue integrity, and promoting tissue repair and regeneration, it is not surprising that when the composition of ECM is altered, for

example due to mutations or changes in the levels of ECM components this can also lead to diseases. These changes in the levels of ECM components may be triggered by several endogenous and exogenous insults, that can promote increased oxidative stress and DNA damage, as reviewed by Martins *et al.* In this review, the authors go over how DNA damage and oxidative stress affect the expression and deposition of ECM molecules, and conversely how mutations in the genes that code for ECM components cause an increase in DNA damage and oxidative (Martins *et al.*). Both circumstances make it challenging for cells and tissues to return to a state of homeostasis, which has an adverse effect on organ and tissue function and may be a trigger for several disorders. An example of these disorders includes cancer, in which cancer cells manipulate a number of elements involved in ECM remodeling to serve their own purposes, promoting cancer cell migration, proliferation, and epithelial-to-mesenchymal transition. In accordance, several studies have revealed the potential of using ECM and associated components as prognostic signatures for different types of cancer. This is the case of *CTHRC1* (collagen triple helix repeat containing 1) and *LRFN4* (leucine rich repeat and fibronectin type III domain containing 4), as prognostic signature in stomach adenocarcinoma (Han *et al.*), *COL4A6* (collagen IV alpha six chain), *FGA* (fibrinogen alpha chain) and *FSCN1* (fascin actin-bundling protein 1), as potential biomarkers for lung adenocarcinoma (Zeng *et al.*) or *FNI* (fibronectin), *LAMB4* (laminin beta four chain), *LAMB3* (laminin beta three chain), *DMP1* (dentin matrix acidic phosphoprotein 1), *CHAD* (chondroadherin), and *MMRN1* (multimerin 1, also called elastin microfibril interfacer 4), as a matrisome-related gene signature of head and neck squamous cell carcinoma (Huang *et al.*).

Altogether these findings stress how crucial ECM is for preserving tissue homeostasis. Improving our understanding regarding the matrisome and other ECM-related molecules will be critical to improve therapeutic strategies in cancer. In addition, and considering that mutations in genes that encode ECM components have been linked to a variety of genetic disorders, such as different neuromuscular diseases, neurodegenerative diseases, Ehlers Danlos Syndromes, and osteogenesis imperfecta, this knowledge will have a wide clinical application.

Reference

Naba, A., Clauser, K. R., Hoersch, S., Liu, H., Carr, S. A., and Hynes, R. O. (2012). The matrisome: *In silico* definition and *in vivo* characterization by proteomics of

Author contributions

ARC and VA wrote and discussed the editorial.

Funding

ARC was supported by Fundação para a Ciência e Tecnologia (FCT, Portugal; CEECIND/01589/2017 and PTDC/BTM-ORG/1383/2020); L'Oréal Portugal Medals of Honor for Women in Science 2019; Unit Funding (Ref. UIDB/00329/2020); the generosity of the donor Henrique Meirelles, who chose to support the MATRIHEALTH Project. VA was supported by INSERM (Institut National de la Santé et de la Recherche Médicale) and by Stiftelsen Crafoord and Wenner-Gren.

Acknowledgments

The guest editors would like to express their gratitude to all contributors and reviewers for their important work on this Research Topic. We also hope that the medical and genetics communities will find this Research Topic to be of interest.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

normal and tumor extracellular matrices. *Mol. Cell. Proteomics* 11, M111.014647. doi:10.1074/MCP.M111.014647