

Amoebophrya ceratii

Laure Guillou, Jeremy Szymczak, Catharina Alves-De-Souza

▶ To cite this version:

Laure Guillou, Jeremy Szymczak, Catharina Alves-De-Souza. Amoebophrya ceratii. Trends in Parasitology, 2023, 39 (2), pp.152-153. 10.1016/j.pt.2022.11.009. hal-03971751

HAL Id: hal-03971751 https://hal.science/hal-03971751v1

Submitted on 20 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Amoebophrya ceratii

- 2
- 3 Laure Guillou¹*, Jeremy Szymczak¹, Catharina Alves-de-Souza²
- 4 ¹Sorbonne Université, CNRS, UMR7144 Adaptation et Diversité en Milieu Marin, Ecology of Marine
- 5 Plankton (ECOMAP), Station Biologique de Roscoff, 29680 Roscoff, France
- 6 ²Algal Resources Collection, Center for Marine Science, University of North Carolina Wilmington,
- 7 Wilmington, NC 28409, USA
- 8 *Correspondance: <u>lguillou@sb-roscoff.fr</u> (L. Guillou)
- 9 ORCID and Twitter:
- 10 L. Guillou: 0000-0003-1032-7958; @guillou_laure
- 11 J. Szymczak: 0000-0002-7112-7207
- 12 C. Alves-de-Souza: 0000-0001-9577-8090; @C_AlvesDeSouza
- 13
- 14 SUMMARY (1004/1000 characters)

15 Dinoflagellates are one of the most abundant and diverse components of marine plankton. Many of 16 them, if not all, are infected by the endoparasitoid Amoebophrya ceratii (Syndiniales, Marine 17 Alveolate Group II), an early emerging dinoflagellate group. As these parasitoids frequently display 18 narrow host ranges, they are believed as diversified as their hosts and composed of cryptic species. 19 Infections last 2-4 days and end with the host death and the release of a transient fast-swimming 20 colony (the vermiform) that fragments within a few hours into hundreds of infective cells 21 (dinospores). Due to this exponential growth, A. ceratii exerts striking potential on top-down control 22 of planktonic dinoflagellate blooms in natural waters. Both A. ceratii and their dinoflagellate hosts 23 share the same common ancestor, a red myzozoan microalga. A. ceratii is rather atypical by having 24 lost its plastid and an extremely reduced mitochondrial DNA. It can be co-cultivated with its host but 25 not cryopreserved.

26

- 27 (1439/1450 characters)
- 28 KEY FACTS
- 29 A. ceratii is haploid, with V-shaped permanently condensed chromosomes during most of its life
- 30 cycle. Genomes are rather small (hundreds of Mb) compared to other dinoflagellates (~3-245 Gb).
- 31 Genes are encoded into unidirectional clusters and mRNA are trans-spliced. In two strains, introns
- 32 were predominantly non-canonicals (differing from the GT-AG motif) and partly composed of
- 33 introners acting as transposable elements.
- 34 The OXPHOS pathway, as complex III is missing, breaks the mitochondrial electron transport chain
- 35 into two independent operating sub-chains.

36 DISEASE FACTS

- 37 Infection occurs either in the nucleus or cytoplasm. Once inside the host, A. ceratii nutrition mode
- 38 starts with osmotrophy and then shifts to phagotrophy. Pre-digested host chromosomes are sucked
- into feeding tubes and consumed in food vacuoles.
- 40 During the infection, the host is "zombified": it still swims, with functional mitochondria and
- 41 chloroplasts despite the complete digestion of its nucleus. The energy produced by the host
- 42 organelles likely benefits the parasitoid.
- 43 The lifespan of spores is of only a few days. A. ceratii is able to enter dormancy simultaneously with
- 44 its host resting cyst, emerging months later to propagate novel infections.
- 45 TAXONOMY AND CLASSIFICATION
- 46 **PHYLUM:** Myzozoa
- 47 CLASS: Dinoflagellata
- 48 **ORDER:** Syndiniales
- 49 **FAMILY:** Amoebophryidae
- 50 **GENUS:** Amoebophrya
- 51 SPECIES: A. ceratii (species complex composed of multiple cryptic species)

52 Acknowledgments

- 53 This research is founded by the ANR (ANR-21-CE02-0030-01 EPHEMER). Electron microscopy images
- 54 were provided by Sophie Lepanse (MERIMAGE platform, Station Biologique of Roscoff) and Gérard
- 55 Prensier (in memory). Estelle Bigeard (Station Biologique of Roscoff) is in charge of parasitic strains
- 56 within the Roscoff Culture Collection (RCC).
- 57 Declaration of interests
- 58 The authors declare no competing interests
- 59 Resources
- 60 <u>http://www.aquasymbio.fr/en/amoebophrya-ceratii-species-complex</u>
- 61 https://www.roscoff-culture-collection.org/ (Roscoff Culture Collection)
- 62 Literature
- 1. Cachon J. 1964 Contribution à l'étude des péridiniens parasites. Cytologie, cycles évolutifs. Ann.
- 64 Sci. Nat. Zool. 6, 1–158
- 65 2. Cai R. et al. 2020 Cryptic species in the parasitic Amoebophrya species complex revealed by a
- 66 polyphasic approach. Sci. Rep. 10, 2531 https://doi.org/10.1038/s41598-020-59524-z
- 67 3. Chambouvet A. et al. 2008 Control of toxic marine dinoflagellate blooms by serial parasitic killers
- 68 Science 322, 1254–1257
- 69 4. Chambouvet A. et al. 2011 Interplay between the parasite *Amoebophrya* sp. (Alveolata) and the
- 70 cyst formation of the red tide dinoflagellate Scrippsiella trochoidea. Protist 162, 637–649
- 5. Coats D.W. and Park M.G. 2002 Parasitism of photosynthetic dinoflagellates by three strains of
- 72 Amoebophyra (Dinophyta): parasite survival, infectivity, generation time and host specificity. J.
- 73 Phycol. 38, 520–528.
- 6. Decelle J. et al. 2022 Intracellular development and impact of a marine eukaryotic parasite on its
- 75 zombified microalgal host. The ISME J. https://doi.org/10.1038/s41396-022-01274-z
- 76 7. Farhat S. *et al.* 2021 Rapid protein evolution, organellar reductions, and invasive intronic elements
- in the marine aerobic parasite dinoflagellate Amoebophrya spp.. BMC Biol. 19
- 78 https://doi.org/10.1186/s12915-020-00927-9
- 79 8. John U. et al. 2019 An aerobic eukaryotic parasite with functional mitochondria that likely lacks a
- 80 mitochondrial genome. Science Advances April 24, 10.1126/sciadv.aav1110
- 9. Kayal E. and David R.S. 2021 Is the Dinoflagellate *Amoebophrya* Really Missing an mtDNA? Mol.
- 82 Biol. Evol. doi:10.1093/molbev/msab041
- 83 10. Kayal E. et al. 2020 Dinoflagellate host chloroplasts and mitochondria remain functional during
- 84 *Amoebophrya* infection. Frontiers in Microbiol. 11:600823 doi: 10.3389/fmicb.2020.600823.

85