



HAL
open science

Trivialisable control-affine systems revisited

Timothée Schmoderer, Witold Respondek

► **To cite this version:**

Timothée Schmoderer, Witold Respondek. Trivialisable control-affine systems revisited. 2023. hal-03971788

HAL Id: hal-03971788

<https://hal.science/hal-03971788>

Preprint submitted on 3 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

Trivialisable control-affine systems revisited

Timothée Schmoderer*and Witold Respondek†

February 3, 2023

Abstract

The purpose of this paper is to explore the concept of trivial control systems, namely systems whose dynamics depends on the controls only. Trivial systems have been introduced and studied by Serres in the the context of control-nonlinear systems on the plane with a scalar control. In our work, we begin by proposing an extension of the notion of triviality to control-affine systems with arbitrary number of states and controls. Next, our first result concerns two novel characterisations of trivial control-affine systems, one of them is based on the study of infinitesimal symmetries and is thus geometric. Second, we derive a normal form of trivial control-affine systems whose Lie algebra of infinitesimal symmetries possesses a transitive almost abelian Lie subalgebra. Third, we study and propose a characterisation of trivial control-affine systems on 3-dimensional manifolds with scalar control. In particular, we give a novel proof of the previous characterisation obtained by Serres. Our characterisation is based on the properties of two functional feedback invariants: the curvature (introduced by Agrachev) and the centro-affine curvature (used by Wilkens). Finally, we give several normal forms of control-affine systems, for which the curvature and the centro-affine curvature have special properties.

Résumé

L'objectif de cet article est d'explorer le concept de système de contrôles trivial, c'est-à-dire un système dont la dynamique dépend uniquement des commandes. Les systèmes triviaux ont été introduits et étudiés par Serres dans le contexte des systèmes bidimensionnels et non-linéaires par rapport à un contrôle scalaire. Dans notre travail, nous commençons par proposer une extension de la notion de trivialité aux systèmes affines par rapport aux contrôles et avec un nombre arbitraire d'états et de contrôles. Ensuite, notre premier résultat concerne deux nouvelles caractérisations des systèmes affines triviaux, l'une d'entre elles est basée sur l'étude des symétries infinitésimales et est donc géométrique. Deuxièmement, nous donnons une forme normale pour les systèmes affines triviaux dont l'algèbre de Lie des symétries infinitésimales possède une sous-algèbre de Lie presque abélienne. Troisièmement, nous étudions et proposons une caractérisation des systèmes affines triviaux sur des variétés à 3 dimensions et avec un contrôle scalaire. En particulier, nous donnons une nouvelle preuve de la caractérisation précédemment obtenue par Serres. Notre caractérisation est basée sur les propriétés de deux invariants fonctionnels du bouclage : la courbure (introduite par Agrachev) et la courbure centro-affine (utilisée par Wilkens). Enfin, nous donnons plusieurs formes normales de systèmes affines, pour lesquelles la courbure et la courbure centro-affine ont des propriétés particulières.

Keywords: Control-affine system - Feedback equivalence - Trivial control systems - Control curvature - Normal forms - Infinitesimal symmetries

2020 Mathematics Subject Classification: 93A10 - 93B52 - 93B10 - 93B27 - 37N35 - 37C79

*Laboratoire de Mathématiques de l'INSA UR 3226 - FR CNRS 3335, INSA Rouen Normandie, Avenue de l'Université, 76800 St Etienne du Rouvray, France; timothee.schmoderer@insa-rouen.fr & witold.respondek@insa-rouen.fr

†Institute of Automatic Control, Łódź University of Technology, Poland

1 Introduction

In this paper, we consider control-affine systems Σ of the form

$$\Sigma : \dot{\xi} = f(\xi) + \sum_{i=1}^m g_i(\xi)u_i, \quad u_i \in \mathbb{R}, \quad (1)$$

where the state ξ belongs to a smooth n -dimensional manifold \mathcal{M} (or an open subset of \mathbb{R}^n , since most of our results are local), and f and g are smooth vector fields on \mathcal{M} , i.e. smooth sections of the tangent bundle $T\mathcal{M}$. Throughout the paper, the word "smooth" will always mean C^∞ -smooth, and all objects (manifolds, vector fields, differential forms, functions) are assumed to be smooth. We denote a control-affine system by the pair $\Sigma = (f, g)$, where $g = (g_1, \dots, g_m)$. To any control-affine system $\Sigma = (f, g)$ we attach two distributions:

$$\mathcal{G} = \text{span} \{g_1, \dots, g_m\} \quad \text{and} \quad \mathcal{G}^1 = \mathcal{G} + [f, \mathcal{G}] = \text{span} \{g_1, \dots, g_m, [f, g_1], \dots, [f, g_m]\}. \quad (2)$$

We call two control-affine systems $\Sigma = (f, g)$ and $\tilde{\Sigma} = (\tilde{f}, \tilde{g})$ *feedback equivalent*, if there exists a diffeomorphism $\phi : \mathcal{M} \rightarrow \tilde{\mathcal{M}}$ and smooth functions $\alpha : \mathcal{M} \rightarrow \mathbb{R}^m$ and $\beta : \mathcal{M} \rightarrow GL_m(\mathbb{R})$ such that

$$\tilde{f} = \phi_* \left(f + \sum_{i=1}^m g_i \alpha_i \right) \quad \text{and} \quad \tilde{g}_i = \phi_* \left(\sum_{j=1}^m g_j \beta_i^j \right),$$

where ϕ_* denotes the tangent map of ϕ . If ϕ is defined locally around ξ_0 and $\tilde{\xi}_0 = \phi(\xi_0)$, then we say that Σ and $\tilde{\Sigma}$ are locally feedback equivalent at ξ_0 and $\tilde{\xi}_0$, respectively. Feedback equivalence of control-affine systems means equivalence of the affine distributions $\mathcal{A} = f + \mathcal{G}$ and $\tilde{\mathcal{A}} = \tilde{f} + \tilde{\mathcal{G}}$ attached to Σ and $\tilde{\Sigma}$, respectively.

In the thesis [17], Serres proposed the notion of a trivial system of the form

$$(\mathcal{T}) : \dot{x} = F(w), \quad x \in \mathcal{X}, \quad w \in \mathbb{R}^m,$$

where w is the control that enters nonlinearly. The dynamics $F(w)$ of a trivial system does not depend on the state variables x and thus depends on control variables w only. Actually, (\mathcal{T}) is called flat in [17] but that name can be misleading because, first, there is a well established notion of flat control systems [7] and, second, the class of trivial control systems does not coincide with control systems of zero-curvature [2], which thus can be considered as geometrically flat, as we will discuss in Section 3. For those reasons, following [18], we call (\mathcal{T}) a trivial system and we say that a general control-nonlinear system $\dot{x} = F(x, w)$ is trivialisable if it is equivalent, via a feedback of the form $\tilde{x} = \phi(x)$, $\tilde{w} = \psi(x, w)$, to a trivial system (\mathcal{T}) , where $(\phi, \psi) : \mathcal{X} \times \mathbb{R}^m \rightarrow \tilde{\mathcal{X}} \times \mathbb{R}^m$ is a diffeomorphism. Inspired by the above considerations, we adapt the concept of triviality to control-affine systems as follows.

Definition 1 (Trivial control-affine systems). *We say that a control-affine system $\Sigma = (f, g)$ is trivialisable if it is feedback equivalent to a trivial system of the form:*

$$(\mathcal{T}) : \begin{cases} \dot{x} &= F(w) \\ \dot{w} &= u \end{cases}, \quad (x, w) \in \mathcal{M} = \mathcal{X} \times \mathbb{R}^m, \quad u \in \mathbb{R}^m,$$

whose \dot{x} -dynamics depend on the controlled w -variables only.

The notions of trivial and trivialisable general control-nonlinear versus control-affine systems are two sides of the same coin. Indeed, two control-nonlinear systems $\dot{x} = F(x, w)$ and $\dot{\tilde{x}} = \tilde{F}(\tilde{x}, \tilde{w})$ are feedback equivalent if and only if their control-affine extensions $\dot{x} = F(x, w)$, $\dot{w} = u$ and $\dot{\tilde{x}} = \tilde{F}(\tilde{x}, \tilde{w})$, $\dot{\tilde{w}} = \tilde{u}$ are equivalent via control-affine feedback transformations, see [10, equation 3.6]. Therefore a control-nonlinear system $\dot{x} = F(x, w)$ is trivialisable if and only if $\dot{x} = F(x, w)$, $\dot{w} = u$ is trivialisable in the sense of Definition 1 and the latter class is the object of our studies in this paper.

Trivial control systems are interesting to study because they model trajectories of dynamical systems under a nonholonomic constraint that does not depend on the point. Indeed, under the additional regularity assumption that $\text{rk} \frac{\partial F}{\partial w}(w) = m$, equivalently, the distribution \mathcal{G}^1 of (T) satisfies $\text{rk} \mathcal{G}^1 = 2m$, there exist local coordinates $x = (z, y)$, with $\dim z = n - 2m$ and $y = (y_1, \dots, y_m)$ such that the equations of (T) can be rewritten

$$\begin{cases} \dot{z} &= f(w) \\ \dot{y} &= w \\ \dot{w} &= u \end{cases}$$

and we conclude that the trajectories of (T) satisfy the nonholonomic constraints $\dot{z} = f(y)$, whose shape is independent of the point $x = (z, y)$. Denoting by \mathcal{X} the (locally defined) quotient manifold \mathcal{M}/\mathcal{G} , we see that a trajectory $x(t) \in \mathcal{X}$ satisfies the nonholonomic constraint $\dot{z} = f(y)$ if and only if there exists a smooth control $u(t)$ such that $(x(t), w(t))$ is a trajectory of (T) . Connections between equations on the tangent bundle and control systems are explored in [13, 14]. Examples of trivial systems can be found in the literature; e.g. in [14] we characterise trivial elliptic, hyperbolic, and parabolic control systems, Dubin's car [6] is a very simple model of system that is trivial, and, finally, trivial control-nonlinear system on surfaces (i.e. $n = 2$) and with scalar control have been studied, characterised (and normal forms in particular cases have been given) in [17, 18, 16].

1.1 Outline of the paper

In the next subsection, we develop the main notions of differential geometry and of control theory that we will need in the rest of the paper. Next, in Section 2, we study trivial control-affine systems on manifolds of arbitrary dimension and with an arbitrary number of controls. We propose two novel characterisations of trivial systems, one of them is based on the Lie algebra of infinitesimal symmetries. Moreover, using our characterisation of trivial systems via symmetries, we will give a normal form of trivial systems whose Lie algebra of infinitesimal symmetries possesses a transitive almost abelian Lie subalgebra. Afterwards, in Section 3, we will be interested in revisiting the characterisation of trivial systems discovered by Serres [17] in the context of control-nonlinear systems on surfaces. We propose a characterisation of trivial control-affine systems on 3-dimensional manifolds with scalar control. Our characterisation exhibits a discrete invariant, and two fundamental functional invariants: the control curvature introduced by Agrachev [2, 1], and the centro-affine curvature used by Wilkens [19]. Both functional invariants can be computed for any control-affine system. We will provide another proof and interpretation of Serres results. Finally, in Section 4 we discuss several normal forms (some new and some existing in the literature) of control-affine systems, for which the control curvature and the centro-affine curvature have special properties.

1.2 Preliminaries

In this subsection, we recall the main definitions and notions of differential geometry and of control theory that we need in the paper. The main notations that we use are summarised in Table 1.

Differential Geometry. For a manifold \mathcal{M} we will denote by $T\mathcal{M}$ and $T^*\mathcal{M}$ the tangent and cotangent bundle, respectively. The space of all smooth vector fields (smooth sections of $T\mathcal{M}$) will be denoted $V^\infty(\mathcal{M})$ and the space of all smooth differential p -forms by $\Lambda^p(\mathcal{M})$, except for smooth functions (0-forms) whose space is denoted $C^\infty(\mathcal{M})$. For a diffeomorphism $\phi : \mathcal{M} \rightarrow \tilde{\mathcal{M}}$, a vector field $f \in V^\infty(\mathcal{M})$, and a differential p -form $\omega \in \Lambda^p(\tilde{\mathcal{M}})$, we denote by $\phi_*f \in V^\infty(\tilde{\mathcal{M}})$ the push-forward of f , and by $\phi^*\omega \in \Lambda^p(\mathcal{M})$ the pull-back of ω . The (local) flow of a vector field $f \in V^\infty(\mathcal{M})$ is denoted by γ_t^f (for any t for which it is defined). The Lie derivative of a differential p -form ω along a vector field f will be denoted by $L_f(\omega)$. In particular, for a function $\lambda \in C^\infty(\mathcal{M})$ and its differential $d\lambda$ (an exact 1-form) we have

$$L_f(\lambda) = \langle d\lambda, f \rangle \quad \text{and} \quad L_f(d\lambda) = dL_f(\lambda).$$

For any smooth functions α , λ , and μ , the Lie derivative possesses the following properties: $L_{\alpha f}(\lambda) = \alpha L_f(\lambda)$, and $L_f(\lambda\mu) = L_f(\lambda)\mu + \lambda L_f(\mu)$. Iterative Lie derivatives are defined by $L_f^k(\lambda) = L_f(L_f^{k-1}(\lambda))$, for any $k \geq 2$. For any two vector fields $f, g \in V^\infty(\mathcal{M})$, we define their Lie bracket as a new vector field, denoted $[f, g] \in V^\infty(\mathcal{M})$, such that for any smooth function λ we have

$$L_{[f, g]}(\lambda) = L_f(L_g(\lambda)) - L_g(L_f(\lambda)).$$

The Lie bracket possesses the following properties: it is bilinear over \mathbb{R} , it is skew-commutative, i.e. $[f, g] = -[g, f]$, and it satisfies the Jacobi identity:

$$[f, [g, h]] + [g, [h, f]] + [h, [f, g]] = 0, \quad \forall f, g, h \in V^\infty(\mathcal{M}).$$

Moreover, for any smooth function α , and any vector fields f , g , and h , we have

$$[f, \alpha g + h] = \alpha [f, g] + L_f(\alpha)g + [f, h].$$

Two vector fields f and g satisfying $[f, g] = 0$ are said to be commuting; since under diffeomorphisms $\phi : \mathcal{M} \rightarrow \tilde{\mathcal{M}}$ the Lie bracket is transformed by $[\phi_*f, \phi_*g] = \phi_*[f, g]$, the commutativity property does not depend on coordinates. The celebrated *Flow-box* theorem (also called the "Straightening-out theorem" or the "Local linearisation lemma") asserts that on a given n -dimensional manifold \mathcal{M} there exists a local coordinate system (x_1, \dots, x_n) such that $f = \frac{\partial}{\partial x_1}$ in a neighbourhood of any point p where $f(p) \neq 0$. This can simultaneously be done for a family of (locally) independent vector fields (f_1, \dots, f_m) if and only if they are mutually commuting. We set $\text{ad}_f^0 g = g$, $\text{ad}_f g = [f, g]$, and the iterated Lie bracket is denoted by $\text{ad}_f^k g = [f, \text{ad}_f^{k-1} g]$ for $k \geq 1$; see [9, chapter 1] for a detailed introduction and proofs of the above properties.

Infinitesimal symmetries. We briefly introduce the notion of symmetries of control-affine systems (see [12, 8] for a detailed introduction). For a control-affine system $\Sigma = (f, g)$, see (1), with state \mathcal{M} a smooth n -dimensional manifold, we define the field of admissible velocities \mathcal{A} as

$$\mathcal{A}(\xi) = \left\{ f(\xi) + \sum_{i=1}^m g_i(\xi)u_i : u_i \in \mathbb{R} \right\} \subset T_\xi \mathcal{M}.$$

We say that a diffeomorphism $\phi : \mathcal{M} \rightarrow \mathcal{M}$ is a *symmetry* of Σ if it preserves the field of affine m -planes \mathcal{A} (equivalently, the affine distribution $\mathcal{A} = f + \mathcal{G}$), that is, $\phi_*\mathcal{A} = \mathcal{A}$. We say that a vector field v on \mathcal{M} is an infinitesimal symmetry of $\Sigma = (f, g)$ if the (local) flow γ_t^v of v is a local symmetry, for any t for which it exists, that is, $(\gamma_t^v)_*\mathcal{A} = \mathcal{A}$. Consider the system $\Sigma = (f, g)$ and recall that \mathcal{G} is the distribution spanned by the vector fields g_1, \dots, g_m . We have the following characterisation of infinitesimal symmetries.

Proposition 1. *A vector field v is an infinitesimal symmetry of the control-affine system $\Sigma = (f, g)$ if and only if*

$$[v, g] = 0 \pmod{\mathcal{G}} \quad \text{and} \quad [v, f] = 0 \pmod{\mathcal{G}}.$$

By the Jacobi identity, it is easy to see that if v_1 and v_2 are infinitesimal symmetries, then so is $[v_1, v_2]$, hence the set of all infinitesimal symmetries forms a real Lie algebra. Notice that the Lie algebra of infinitesimal symmetries is attached to the affine distribution $\mathcal{A} = f + \mathcal{G}$ and not to a particular pair $(f, g) = (f, g_1, \dots, g_m)$. Different pairs (f, g) related via feedback transformations (α, β) define the same \mathcal{A} and thus have the same Lie algebra of infinitesimal symmetries which, therefore, is a feedback invariant object attached to Σ .

$\mathcal{M}, T\mathcal{M}, \xi = (x, w)$	Smooth n -dimensional manifold, its tangent bundle, and its local coordinates with $\dim w = m$.
ϕ, ϕ_*, ϕ^*	A diffeomorphism, its tangent map, its cotangent map.
$\Sigma = (f, g)$	A control-affine system; see (1).
\mathcal{G} and \mathcal{G}^1	Distribution spanned by the vector fields g_1, \dots, g_m and the distribution spanned by the vector fields g_1, \dots, g_m and $[f, g_1], \dots, [f, g_m]$; see (2).
(T)	Trivial control-affine system; see Definition 1.
$\mathfrak{L}, \mathfrak{A}, \mathfrak{I}$	A real Lie (sub)algebra, a subalgebra, an ideal.
$\Sigma_\lambda, \Sigma_\lambda^{0,k}$	Normal forms of trivial systems having an almost abelian subalgebra of infinitesimal symmetries; see Theorem 2 and Proposition 2.
$\Sigma_s = (f_s, g)$	Control affine system given by a semi-canonical pair; see Definition 3.
$\Sigma_c = (f_c, g_c)$	Control affine system given by the canonical pair; see Definition 3.
(k_1, k_2, k_3) and $(\lambda_1, \lambda_2, \lambda_3)$	Structure functions attached to any control-affine system on a 3-dimensional manifold with scalar control; see (6).
$(\varepsilon, \kappa, \nu)$	Feedback invariants of control-affine systems; defined for the canonical pair by (6') and expressed for any control-affine system by (11).

Table 1: Main notations for the paper

2 Trivial control-affine systems

In this section, we first propose two new characterisations of trivialisable control-affine systems (with the state-space of arbitrary dimension and with an arbitrary number of controls); see Theorem 1 below. Second, we give a normal form of trivial systems whose Lie algebra of infinitesimal symmetries possesses a transitive almost abelian Lie subalgebra; see Theorem 2 and Proposition 2 of this section.

2.1 Characterisations of trivial systems

The following theorem gives two characterisations of trivialisable systems. The first one is technical and shows that triviality is a property that depends on the coordinates (like being a linear control system depends on the choice of coordinates), and the second one is based on infinitesimal symmetries and is thus geometric. Recall that to a control-affine system $\Sigma = (f, g)$ we attach two distributions $\mathcal{G} = \text{span}\{g_1, \dots, g_m\}$ and $\mathcal{G}^1 = \mathcal{G} + [f, \mathcal{G}]$, see (2).

Theorem 1 (Two characterisations of trivialisable systems). *Consider a control-affine system $\Sigma = (f, g)$ with state on a n -dimensional manifold \mathcal{M} and with $m \geq 1$ controls. The following assertions hold locally around ξ_0 :*

- (i) *Suppose that $\text{rk } \mathcal{G}^1 = m + k$ is constant. The system Σ is locally trivialisable if and only if Σ is locally feedback equivalent to*

$$\Sigma_T : \begin{cases} \dot{x}_i &= h_i(x, w) \\ \dot{w} &= u \end{cases}, \quad \text{for } 1 \leq i \leq n - m,$$

where the smooth scalar functions h_1, \dots, h_{n-m} satisfy

$$\text{rk span}\{dh_1, \dots, dh_{n-m}\} = k. \quad (3)$$

- (ii) Σ is, locally around ξ_0 , trivialisable if and only if the distribution \mathcal{G} is involutive and of constant rank m and, additionally, the Lie algebra of infinitesimal symmetries of Σ possesses an abelian subalgebra \mathfrak{A} such that $\mathfrak{A}(\xi_0) \oplus \mathcal{G}(\xi_0) = T_{\xi_0}\mathcal{M}$.

Observe that the assumption on the rank of the distribution \mathcal{G}^1 in statement (i) implies that the dimension n of the manifold \mathcal{M} is greater than or equal to $m+k$. If $n = m+k$, then the trivialisation (T) of Σ_T (and thus of Σ) can be taken (for suitable w and u) as $\dot{x}_i = w_i$, $1 \leq i \leq k$, $\dot{w}_j = u_j$, $1 \leq j \leq m$. On the other hand, if $n > m+k$, then $\dot{x}_i = w_i$, $1 \leq i \leq k$, $\dot{x}_i = F_i(w_1, \dots, w_k)$, $k+1 \leq i \leq m+k$, and $\dot{w}_j = u_j$, $1 \leq j \leq m$. Notice that $k \leq m$, so if $n > 2m$, then there are always nonlinear equations $\dot{x}_i = F_i(w_1, \dots, w_k)$. In item (ii), there are no particular relations between the dimension of the state space and on the number of controls (other than the obvious $n \geq m$).

Remark 1. For the system Σ_T , define $H = (h_1, \dots, h_{n-m})^t$. Then, under the assumption that $\text{rk } \mathcal{G}^1$ is constant, condition (3) can be equivalently reformulated as

$$\text{rk } \frac{\partial H}{\partial w}(x, w) = \text{rk } \frac{\partial H}{\partial(x, w)}(x, w),$$

in a neighbourhood of (x_0, w_0) .

Proof.

(i) Suppose that Σ is locally trivialisable, i.e. by Definition 1, Σ is locally feedback equivalent to (T), which is of the form of Σ_T with $h_i(x, w) = F_i(w)$, for $1 \leq i \leq n-m$, and we now show that those functions satisfy (3). On one hand, the condition $\text{rk } \mathcal{G}^1 = m+k$ implies that the Jacobian matrix $\frac{\partial F}{\partial w}$ is of constant rank k , where $F = (F_1, \dots, F_{n-m})^T$. On the other hand, we obtain that $dh_i = dF_i = \sum_{j=1}^m \frac{\partial F_i}{\partial w_j} dw_j$. Hence the rank of $\text{span}\{dh_1, \dots, dh_{n-m}\}$ is the same as that of $\frac{\partial F}{\partial w}$ and the conclusion follows. Conversely, assume that Σ is feedback equivalent to Σ_T . Using the assumption $\text{rk } \mathcal{G}^1 = m+k$ we can reorder the x -coordinates such that $h = (\hat{h}_1, \dots, \hat{h}_k, \tilde{h}_{k+1}, \dots, \tilde{h}_{n-m})$, where $\text{rk } \frac{\partial \hat{h}}{\partial w} = k$. We set $\hat{w}_i = \hat{h}_i(x, w)$, for $1 \leq i \leq k$, completed by $\hat{w}_{k+1}, \dots, \hat{w}_m$ (chosen among the w_i 's) in such a way that $\hat{w}_1, \dots, \hat{w}_m$ form a local coordinate system. We conclude, by condition (3), that the functions $\tilde{h}_{k+1}, \dots, \tilde{h}_{n-m}$ depend on the variables $\hat{w}_1, \dots, \hat{w}_k$ only. Using a feedback transformation that yields $\hat{w}_i = \hat{u}_i$, $1 \leq i \leq k$, we conclude that Σ_T is, indeed, a trivial system in coordinates (x, \hat{w}) .

(ii) Suppose that $\Sigma = (f, g)$ is locally trivialisable, then for (T) we have $\mathcal{G} = \text{span}\left\{\frac{\partial}{\partial w_1}, \dots, \frac{\partial}{\partial w_m}\right\}$, which clearly is involutive and of constant rank m . Moreover, the vector fields $v_i = \frac{\partial}{\partial x_i}$, for $1 \leq i \leq n-m$, are commuting symmetries of (T) that span the abelian Lie algebra \mathfrak{A} satisfying $\mathfrak{A}(\xi_0) \oplus \mathcal{G}(\xi_0) = T_{\xi_0}\mathcal{M}$. Conversely, suppose that the Lie algebra of infinitesimal symmetries of $\Sigma = (f, g)$ possesses an abelian subalgebra $\mathfrak{A} = \text{vect}_{\mathbb{R}}\{v_1, \dots, v_{n-m}\}$. By $\mathfrak{A}(\xi_0) \oplus \mathcal{G}(\xi_0) = T_{\xi_0}\mathcal{M}$ the vector fields v_1, \dots, v_{n-m} are linearly independent, so we choose local coordinates $\tilde{\xi} = (\tilde{x}, \tilde{w})$ such that $v_i = \frac{\partial}{\partial \tilde{x}_i}$, for $1 \leq i \leq n-m$. In those coordinates, we have $g_j = A_j(\tilde{x}, \tilde{w})\frac{\partial}{\partial \tilde{x}} + B_j(\tilde{x}, \tilde{w})\frac{\partial}{\partial \tilde{w}}$. Since \mathcal{G} is of constant rank m and satisfies $\mathfrak{A}(\tilde{\xi}_0) \oplus \mathcal{G}(\tilde{\xi}_0) = T_{\tilde{\xi}_0}\mathcal{M}$, via a suitable feedback transformation we choose generators of \mathcal{G} as $\tilde{g}_j = A_j(\tilde{x}, \tilde{w})\frac{\partial}{\partial \tilde{x}} + \frac{\partial}{\partial \tilde{w}_j}$ (to simplify notations, we skip the "tildes" and denote \tilde{g}_j by g_j). Using that v_i are symmetries of Σ , that is $[v_i, g_j] \in \mathcal{G}$, we deduce that $A_j = A_j(\tilde{w})$, therefore we actually have $[v_i, g_j] = 0$. Moreover, \mathcal{G} is involutive so we deduce that $[g_j, g_k] = 0$. Therefore, all vector fields v_i and g_j commute and thus there exist coordinates $\xi = (x, w)$ such that $v_i = \frac{\partial}{\partial x_i}$, for $1 \leq i \leq n-m$, and $g_j = \frac{\partial}{\partial w_j}$, for $1 \leq j \leq m$. The fields v_i are symmetries of Σ so $[v_i, f] \in \mathcal{G}$ implying that $f = F\frac{\partial}{\partial x} + F\frac{\partial}{\partial w}$, where $F = F(w)$ and we achieve $F = 0$ by a suitable feedback transformation. □

2.2 Normal form of trivial systems possessing an almost abelian Lie subalgebra of infinitesimal symmetries

Theorem 1 of the previous subsection asserts that the Lie algebra of infinitesimal symmetries of trivialisable systems possesses an abelian subalgebra complementary to the distribution \mathcal{G} . In this subsection, we study a particular case when the abelian subalgebra \mathfrak{A} is actually an abelian ideal of codimension

one of a subalgebra \mathfrak{L} of symmetries, the latter acting transitively on \mathcal{M} . We give a normal form of control-affine systems (with a, necessarily, scalar control) possessing such Lie algebra of symmetries.

Definition 2 (Almost abelian Lie algebra). *Let \mathfrak{L} be a real Lie algebra; following the definition of [5], we call \mathfrak{L} almost abelian if it has an abelian ideal \mathfrak{I} of codimension one.*

It is a simple application of Lie algebra homology to deduce that an almost abelian Lie algebra (possibly of infinite dimension) is isomorphic to the semi-direct product $\mathfrak{L} \cong \mathfrak{I} \rtimes \text{vect}_{\mathbb{R}}\{v_0\}$ and that its structure is determined by the action of v_0 on \mathfrak{I} , namely by

$$\begin{aligned} \text{ad}_{v_0} : \mathfrak{I} &\longrightarrow \mathfrak{I} \\ v &\longmapsto [v_0, v]. \end{aligned}$$

Moreover, two almost abelian Lie algebras $\mathfrak{L} = \mathfrak{I} \rtimes \text{vect}_{\mathbb{R}}\{v_0\}$ and $\tilde{\mathfrak{L}} = \tilde{\mathfrak{I}} \rtimes \text{vect}_{\mathbb{R}}\{\tilde{v}_0\}$ are isomorphic if and only if there exists a real invertible transformation $P : \mathfrak{I} \rightarrow \tilde{\mathfrak{I}}$ such that $P \text{ad}_{v_0} = \mu \text{ad}_{\tilde{v}_0} P$ for some $\mu \in \mathbb{R}^*$; see [4, Proposition 11]. Therefore, isomorphism classes of almost abelian Lie algebras correspond to similarity classes of the linear operator ad_{v_0} (up to multiplication by a scalar). In particular, if \mathfrak{L} is finite dimensional, the similarity classes of ad_{v_0} are given by the Jordan normal forms. In the following theorem, we consider the simplest case where ad_{v_0} is diagonalisable over \mathbb{R} and give a normal form of control-affine systems that have \mathfrak{L} as Lie subalgebra of infinitesimal symmetries.

Theorem 2 (Almost abelian infinitesimal symmetries). *Consider a control-affine system $\Sigma = (f, g)$ on an n -dimensional state manifold \mathcal{M} and with scalar control. Assume that the Lie algebra of infinitesimal symmetries possesses a Lie subalgebra \mathfrak{L} for which the following conditions hold at ξ_0 : $f(\xi_0) \notin \mathcal{G}(\xi_0)$, \mathfrak{L} acts transitively on \mathcal{M} , $\mathfrak{L} = \mathfrak{I} \rtimes \text{vect}_{\mathbb{R}}\{v_0\}$ is almost abelian, its abelian ideal satisfies $\mathfrak{I}(\xi_0) \oplus \mathcal{G}(\xi_0) = T_{\xi_0}\mathcal{M}$, and the operator ad_{v_0} is non-singular and diagonalisable over \mathbb{R} . Then, Σ is, locally around ξ_0 , feedback equivalent to a trivial system of the form*

$$\Sigma_{\lambda} : \begin{cases} \dot{x}_i &= \eta_i(w+1)^{\lambda_i} & 1 \leq i \leq n-1 \\ \dot{w} &= u \end{cases}, \quad u \in \mathbb{R}$$

around $(x_0, 0) \in \mathbb{R}^{n-1} \times \mathbb{R}$, where the λ_i 's are the eigenvalues of ad_{v_0} and η_i are constants equal 0 or 1, with $\eta_1 = 1$.

Observe that the assumptions: \mathfrak{L} is almost abelian, acts transitively on \mathcal{M} , and satisfies $\mathfrak{I}(\xi_0) \oplus \mathcal{G}(\xi_0) = T_{\xi_0}\mathcal{M}$, are quite restrictive on the number of controls of Σ . Namely, they imply that $\text{rk } \mathcal{G} = 1$, i.e. Σ has a scalar control.

Remark 2 (Converse implication). *By a straightforward computation, we see that the vector fields*

$$v_i = \frac{\partial}{\partial x_i}, \quad \forall 1 \leq i \leq n-1, \quad \text{and} \quad v_0 = \sum_{i=1}^{n-1} \lambda_i x_i \frac{\partial}{\partial x_i} + (w+1) \frac{\partial}{\partial w}$$

are infinitesimal symmetries of Σ_{λ} . Thus the Lie algebra of infinitesimal symmetries of Σ_{λ} possesses an almost abelian subalgebra $\mathfrak{L} = \text{vect}_{\mathbb{R}}\{v_1, \dots, v_{n-1}, v_0\}$. Therefore, the theorem is actually an "if and only if" statement since all other assumptions are feedback invariant.

Proof. Consider the control-affine system $\Sigma = (f, g)$ given by vector fields f and g , and let n vector fields v_1, \dots, v_{n-1}, v_0 generate the n -dimensional Lie subalgebra $\mathfrak{L} = \text{vect}_{\mathbb{R}}\{v_1, \dots, v_{n-1}, v_0\}$ of the algebra of infinitesimal symmetries, which by assumption is almost abelian, whose abelian ideal is $\mathfrak{I} = \text{vect}_{\mathbb{R}}\{v_1, \dots, v_{n-1}\}$ and since ad_{v_0} is diagonalisable over \mathbb{R} we conclude that

$$[v_0, v_i] = \lambda_i v_i, \quad \forall 1 \leq i \leq n-1. \quad (4)$$

By statement (ii) of Theorem 1, Σ is locally trivialisable and following the proof of that theorem we deduce that there exists local coordinates (x, \hat{w}) around $(x_0, 0)$ such that $v_i = \frac{\partial}{\partial x_i}$, for $1 \leq i \leq n-1$, $g = \frac{\partial}{\partial \hat{w}}$, and $f = \sum_{i=1}^{n-1} F_i \frac{\partial}{\partial x_i}$, where $F_i = F_i(\hat{w})$. Express the infinitesimal symmetry $v_0 =$

$\sum_{i=1}^{n-1} \gamma_i \frac{\partial}{\partial x_i} + \bar{\delta} \frac{\partial}{\partial \hat{w}}$, where $\gamma_i = \gamma_i(x)$ since v_0 is a symmetry of $\mathcal{G} = \text{span} \left\{ \frac{\partial}{\partial \hat{w}} \right\}$, and $\bar{\delta} = \bar{\delta}(x, \hat{w})$. Using the commutativity relations (4) that have not been changed by applying diffeomorphisms, we obtain

$$v_0 = - \sum_{i=1}^{n-1} (\lambda_i x_i + a_i) \frac{\partial}{\partial x_i} + \bar{\delta}(\hat{w}) \frac{\partial}{\partial \hat{w}}, \quad a_i \in \mathbb{R}.$$

To avoid unnecessary computations, we replace v_0 by $-v_0 - \sum_{i=1}^{n-1} a_i v_i \in \mathfrak{L}$, having the same properties, so we can assume $v_0 = \sum_{i=1}^{n-1} \lambda_i x_i \frac{\partial}{\partial x_i} + \delta(\hat{w}) \frac{\partial}{\partial \hat{w}}$, where $\delta(\hat{w}) = -\bar{\delta}(\hat{w})$. Using the fact that v_0 is a symmetry of f , i.e. $[v_0, f] \in \mathcal{G}$, we deduce the following equations:

$$\delta(\hat{w}) \frac{dF_i}{d\hat{w}}(\hat{w}) - \lambda_i F_i(\hat{w}) = 0, \quad \forall 1 \leq i \leq n-1. \quad (5)$$

The assumption $f(\xi_0) \notin \mathcal{G}(\xi_0)$ implies that $(F_1, \dots, F_{n-1})(0) \neq 0 \in \mathbb{R}^{n-1}$, so suppose that $F_1(0) \neq 0$ (renumber the x_i 's if necessary) and thus $F_1(\hat{w}) = c + h(\hat{w})$ with $c = F_1(0)$ for a function h satisfying $h(0) = 0$. Replacing x_1 by $\frac{x_1}{c}$ we may assume that $F_1(\hat{w}) = 1 + h(\hat{w})$. Moreover, by (5) we conclude that $h'(0) \neq 0$; recall that by assumption ad_{v_0} is non-singular and hence $\lambda_i \neq 0$ for all $1 \leq i \leq n-1$. Thus, $w = (1 + h(\hat{w}))^{1/\lambda_1} - 1$ is a diffeomorphism around $\hat{w}_0 = 0$ and, to simplify notations, we keep the symbols f and v_0 for those vector fields represented using the coordinate w . We have $F_1(w) = (1+w)^{\lambda_1}$, thus relation (5) gives, with \hat{w} renamed w , $\delta(w) = w + 1$ and implies that the functions F_2, \dots, F_{n-1} satisfy

$$(1+w) \frac{dF_i}{dw}(w) = \lambda_i F_i(w), \quad \text{for } 2 \leq i \leq n-1.$$

Solving those equations around $w_0 = 0$ gives $F_i = c_i (w+1)^{\lambda_i}$, with $c_i \in \mathbb{R}$. Thus, normalising x_i with the non-zero c_i we get that Σ takes the form of Σ_λ around $(x_0, 0) \in \mathbb{R}^n$. \square

The above theorem generalises our previous results on the Lie algebra of infinitesimal symmetries of hyperbolic and parabolic systems for which we have $n = 3$ and, respectively, $(\lambda_1, \lambda_2) = (1, -1)$ and $(\lambda_1, \lambda_2) = (2, 1)$, see [15].

The situation is much more involved when $f(\xi_0) \in \mathcal{G}(\xi_0)$ as the following proposition illustrates.

Proposition 2. *Consider a control-affine system $\Sigma = (f, g)$ with a n -dimensional state manifold and a scalar control, and let \mathfrak{L} be a Lie subalgebra of infinitesimal symmetries. Suppose that the following conditions hold at ξ_0 : $f(\xi_0) \in \mathcal{G}(\xi_0)$, there exists $k \geq 1$ the smallest integer such that $g \wedge \text{ad}_g^k f(\xi_0) \neq 0$, \mathfrak{L} acts transitively on \mathcal{M} , $\mathfrak{L} \cong \mathfrak{I} \rtimes \text{vect}_{\mathbb{R}} \{v_0\}$ is almost abelian, $\mathfrak{I}(\xi_0) \oplus \mathcal{G}(\xi_0) = T_{\xi_0} \mathcal{M}$, and the action of ad_{v_0} is non-singular and diagonalisable over \mathbb{R} . Then, the eigenvalues λ_i of ad_{v_0} are such that $\frac{\lambda_i}{\lambda_1} k$ are positive integers greater or equal than k , where λ_1 is the smallest, in absolute value, eigenvalue of ad_{v_0} . Moreover, the system Σ is locally feedback equivalent to*

$$\Sigma_\lambda^{0,k} : \begin{cases} \dot{x}_1 &= w^k \\ \dot{x}_i &= \eta_i (w^k)^{\lambda_i/\lambda_1} \quad 2 \leq i \leq n-1 \\ \dot{w} &= u \end{cases}, \quad u \in \mathbb{R},$$

around $(x_0, 0) \in \mathbb{R}^n$, where η_i are constants equal to 0 or 1.

Observe that the normal form $\Sigma_\lambda^{0,k}$ defines a polynomial system since $k \frac{\lambda_i}{\lambda_1}$ are positive integers. Moreover, although the eigenvalues λ_i can be any, the conditions of the above proposition imply severe restrictions on them. Namely, all λ_i 's have the same sign and $\lambda_i = \frac{l_i}{k} \lambda_1$, where the l_i 's are integers satisfying $l_i \geq k$. Furthermore, it is a classical fact that (under the above assumptions) the integer k is an invariant of feedback transformations. Hence if $k \neq k'$, then $\Sigma_\lambda^{0,k}$ and $\Sigma_\lambda^{0,k'}$ are not locally feedback invariant around $(x_0, 0)$.

Proof. The beginning of the proof is the same as that of the previous theorem up to equation (5), so we start from there. By $f(\xi_0) \in \mathcal{G}(\xi_0)$ we have $(F_1, \dots, F_{n-1})(0) = 0 \in \mathbb{R}^{n-1}$ and due to $g \wedge \text{ad}_g^k f(0) \neq 0$ we assume that $\frac{d^k F_1}{d\hat{w}^k}(0) \neq 0$, if necessary relabel the x_i -coordinates. Using the Taylor expansion we

can write $F_1 = \hat{w}^k H(\hat{w})$, where $H(0) \neq 0$ and we can suppose $H(0) > 0$, if not, replace x_1 by $-x_1$. We apply the local diffeomorphism $w = \hat{w} (H(\hat{w}))^{1/k}$, that maps f and v_0 into vector fields that, for simplicity, we denote again by f and v_0 , respectively. We have $F_1(w) = w^k$, so equation (5), expressed in the w -coordinate, implies, for $i = 1$, that $\delta(w) = \frac{\lambda_1}{k} w$ and

$$\lambda_1 w \frac{dF_i}{dw}(w) = \lambda_i k F_i(w), \quad \text{for } 2 \leq i \leq n-1.$$

Solving those equations around $w_0 = 0$ implies $|F_i| = c_i |w_i|^{k\lambda_i/\lambda_1}$, with $c_i \in \mathbb{R}$. The only C^∞ -solutions are those given by either $c_i = 0$ or by $c_i \neq 0$ with $\frac{\lambda_i}{\lambda_1} k$ being a positive integer. The corresponding smooth solution is $F_i = c_i w^{k\lambda_i/\lambda_1}$ and by definition of k we have $k \frac{\lambda_i}{\lambda_1} \geq k$ (otherwise, $g \wedge \text{ad}_f^{k'} g(\xi_0) \neq 0$, where $k' = \frac{\lambda_i}{\lambda_1} k < k$, contradicting the definition of k). Thus $|\lambda_i| \geq |\lambda_1|$ and λ_1 is, indeed, the eigenvalue of ad_{v_0} of minimal absolute value.. Finally, normalising the coordinates x_i , with $c_i \neq 0$, we obtain the desired form $\Sigma_\lambda^{0,k}$. \square

The previous proposition describes all smooth systems having an almost abelian Lie subalgebra of symmetries for which k exists, in particular all analytic systems. Notice that for a single-input analytic system either k exists or, if not, then it is locally feedback equivalent to a trivial system $\dot{x} = c$, $\dot{w} = u$, where $c \in \mathbb{R}^{n-1}$. In the C^∞ category there are, however, systems for which k does not exist but the symmetry algebra possesses an almost abelian subalgebra. For example, consider around $(x_0, 0)$ the system

$$\begin{cases} \dot{x}_1 = \mathfrak{f}(w) \\ \dot{x}_i = \mathfrak{f}(w)^{\lambda_i/\lambda_1} & 2 \leq i \leq n-1 \\ \dot{w} = u \end{cases}, \quad \text{with } \mathfrak{f}(w) = \exp\left(-\frac{1}{w^2}\right), \mathfrak{f}(0) = 0,$$

and $\lambda_i/\lambda_1 \in \mathbb{N}^*$. By a straightforward calculation, one may check that the system possesses an almost abelian Lie subalgebra of infinitesimal symmetries but, obviously, k does not exist at $(x_0, 0)$ and thus it is not feedback equivalent to $\Sigma_\lambda^{0,k}$.

3 Trivial systems on 3D-manifolds

In this section, we study trivial system on 3-dimensional manifolds with scalar control. Our aim is to give a new version of the results of [16] and to extend them by giving several normal forms.

Throughout this section, we consider a control-affine system $\Sigma = (f, g)$ of the form

$$\Sigma : \dot{\xi} = f(\xi) + g(\xi)u, \quad u \in \mathbb{R},$$

where the state ξ belongs to a 3-dimensional manifold \mathcal{M} and the vector fields f and g satisfy, at any $\xi \in \mathcal{M}$, the following regularity assumptions

$$(A1) \quad f \wedge g \wedge [g, f] \neq 0,$$

$$(A2) \quad g \wedge [g, f] \wedge [g, [g, f]] \neq 0.$$

To any control-affine system $\Sigma = (f, g)$ we attach 6 structure functions uniquely defined by the following decompositions:

$$\begin{aligned} [f, [f, g]] &= k_1 g + k_2 [g, f] + k_3 [g, [g, f]], \\ [g, [g, f]] &= \lambda_1 f + \lambda_2 g + \lambda_3 [g, f]. \end{aligned} \tag{6}$$

Observe that assumption (A2) implies that $\lambda_1 \neq 0$. We now define two different classes of pairs (f, g) .

Definition 3 ((Semi)-canonical pairs). *We call the pair (f, g) semi-canonical if $k_3 \equiv 0$, and we will denote it by (f_s, g) . If, additionally $\lambda_1 \equiv \pm 1$, then we call (f, g) a canonical pair and we denote it by (f_c, g_c) .*

Observe that a semi-canonical pair is characterised by the inclusion $[f, [f, g]] \in \text{span}\{g, [g, f]\}$, which is the property of the singular vector field of Σ (justifying the notation (f_s, g) for a semi-canonical pair). Under assumption (A2), that singular vector field is unique and can be computed using the singular control, thus we are not surprised that the following proposition shows that a semi-canonical pair exists (but observe that our proof does not require using the machinery of singular controls).

The following proposition shows that a control-affine system is always feedback equivalent to a system given by a semi-canonical and even by a canonical pair. Moreover, those pairs can be *explicitly* constructed, meaning that the feedback transformations bringing (f, g) into (f_s, g) or into (f_c, g_c) are constructed with differentiation and algebraic operations only (no differential equations to be solved). Furthermore, a canonical pair is *unique* (up to multiplying g_c by -1) hence its structure functions are feedback equivariants (up to a discrete involution, see Remark 3 below).

Proposition 3 (Existence of semi-canonical and canonical pairs). *Consider a control-affine system $\Sigma = (f, g)$ satisfying assumptions (A1) and (A2). Then, the following statements hold globally:*

- (i) Σ is globally feedback equivalent to $\Sigma_s = (f_s, g)$, where (f_s, g) is a semi-canonical pair;
- (ii) Σ is globally feedback equivalent to $\Sigma_c = (f_c, g_c)$, where (f_c, g_c) is a canonical pair.

Moreover (f_s, g) and (f_c, g_c) can be explicitly constructed via the following feedback transformations

$$f_s = f + gk_3, \quad f_c = f_s, \quad \text{and} \quad g_c = |\lambda_1|^{-1/2} g.$$

Furthermore, the canonical pair (f_c, g_c) is unique up to $g_c \mapsto -g_c$.

The proof of the above proposition is based on the following lemma, which gives some relations between the structure functions $k_1, k_2, k_3, \lambda_1, \lambda_2$, and λ_3 , and shows how they change under feedback transformations. Its proof is a straightforward computation that we detail in Appendix A.

Lemma 1. *Consider a control-affine system $\Sigma = (f, g)$ with structure functions (k_1, k_2, k_3) and $(\lambda_1, \lambda_2, \lambda_3)$. Then, the following relations hold:*

$$L_f(\lambda_1) = -k_2\lambda_1 - L_g(\lambda_1 k_3), \tag{7a}$$

$$L_f(\lambda_2) - \lambda_3 k_1 + L_g(k_1) = -k_2\lambda_2 - L_g(\lambda_2 k_3), \tag{7b}$$

$$L_f(\lambda_3) - \lambda_2 = -k_3\lambda_1 - L_g(k_2) - L_g(\lambda_3 k_3). \tag{7c}$$

Under a feedback transformation $f \mapsto \tilde{f} = f + g\alpha$ and $g \mapsto \tilde{g} = g\beta$, where α and β are smooth scalar functions satisfying $\beta \neq 0$, the structure functions are transformed by

$$\begin{aligned} \tilde{k}_1 &= k_1 + L_{[g,f]}(\alpha) + \frac{1}{\beta} (L_f(\gamma) + \alpha L_g(\gamma) - \gamma L_g(\alpha)) + \tilde{k}_2 \frac{\gamma}{\beta} \\ &\quad + \tilde{k}_3 (L_{[g,f]}(\beta) + L_g(\gamma) - \gamma L_g(\ln(\beta))), \\ \tilde{k}_2 &= k_2 - L_f(\ln(\beta)) - \frac{\gamma}{\beta} - \alpha L_g(\ln(\beta)) + k_3 L_g(\beta), \\ \tilde{k}_3 &= \frac{1}{\beta} (k_3 - \alpha), \end{aligned} \tag{8}$$

$$\tilde{\lambda}_1 = \beta^2 \lambda_1, \quad \tilde{\lambda}_2 = \beta \lambda_2 - \beta \lambda_1 \alpha + \gamma \lambda_3 - L_{[g,f]}(\beta) - L_g(\gamma) + 2\gamma L_g(\ln(\beta)), \quad \tilde{\lambda}_3 = \beta \lambda_3 + L_g(\beta), \tag{9}$$

where $\gamma = L_f(\beta) + \alpha L_g(\beta) - \beta L_g(\alpha)$.

Proof of Proposition 3. Consider $\Sigma = (f, g)$ whose structure functions are (k_1, k_2, k_3) and $(\lambda_1, \lambda_2, \lambda_3)$. By equations (8) and (9) we have that under feedback transformations $\beta \tilde{k}_3 = k_3 - \alpha$ and $\tilde{\lambda}_1 = \beta^2 \lambda_1$. Hence, choosing $\alpha = k_3$ we obtain that the transformed pair (\tilde{f}, \tilde{g}) , where $\tilde{f} = f + gk_3$, is semi-canonical. Moreover, additionally choosing $\beta = |\lambda_1|^{-1/2}$, recall that $\lambda_1 \neq 0$ from assumption (A2), yields a canonical pair $(f_c, g_c) = (\tilde{f}, \tilde{g})$, where $\tilde{g} = \beta g$. Clearly, the singular vector field $f_s = f_c$ is uniquely defined, but the canonical vector field g_c is unique up to $g_c \mapsto \pm g_c$. \square

Observe that for the canonical pair (f_c, g_c) we additionally have $k_2 \equiv 0$, due to (7a). Thus, the canonical pair (f_c, g_c) satisfies the following decomposition (renaming k_1 to κ , λ_1 to ε , λ_2 to μ , λ_3 to ν)

$$\begin{aligned} [f_c, [f_c, g_c]] &= \kappa g_c, \\ [g_c, [g_c, f_c]] &= \varepsilon f_c + \mu g_c + \nu [g_c, f_c], \end{aligned} \quad (6')$$

where $\varepsilon = \pm 1$. Moreover, using equations (7b) and (7c) we deduce that κ , μ , and ν are related by

$$L_{f_c}(\mu) - \nu\kappa + L_{g_c}(\kappa) = 0 \quad (7b')$$

$$L_{f_c}(\nu) - \mu = 0, \quad (7c')$$

from which we deduce that the feedback invariants κ and ν are associated via

$$L_{f_c}^2(\nu) - \nu\kappa + L_{g_c}(\kappa) = 0. \quad (10)$$

Therefore, a canonical pair identifies explicitly a discrete invariant $\varepsilon = \pm 1$ and two constructible feedback invariant functions κ and ν called, respectively, the *curvature* and the *centro-affine curvature* by analogy with Serres' work [18]; see also [19]. Observe that due to (7c') above, μ is determined by ν . Moreover, the curvature κ determines the centro-affine curvature ν up to an affine part; i.e. if f_c is rectified on $\frac{\partial}{\partial x}$, then ν is determined by κ via (10) up to two functions ν_1 and ν_0 satisfying $L_{f_c}(\nu_i) = 0$.

Remark 3. A canonical pair is unique up to $g_c \mapsto -g_c$. Hence the centro-affine curvature ν is a feedback equivariant up to the involution $\nu \mapsto -\nu$ (which does not influence our conditions below). We will get back to that subtlety in Proposition 6, where we will construct several normal forms. On the other hand, the curvature κ is a true feedback invariant (actually, a feedback equivariant that changes as $\phi^*\kappa$ under a diffeomorphism ϕ).

For a given control-affine system $\Sigma_c = (f_c, g_c)$, given by a canonical pair, we will denote by $(\varepsilon, \kappa, \nu)$ the triple of invariants. Although the canonical pair can be constructed without much work, for the sake of completeness, we give the expression of $(\varepsilon, \kappa, \nu)$ for an arbitrary control-affine system. In particular, observe that our formula for the curvature κ generalises the one given in [3, p. 376], where k_3 is already normalised to 0.

Proposition 4 (Invariants of control-affine systems). *Consider a control-affine system $\Sigma = (f, g)$ on a 3-dimensional state-space manifold, and with scalar control, and let (k_1, k_2, k_3) and $(\lambda_1, \lambda_2, \lambda_3)$ be structure functions defined by (6). Then, the invariants $(\varepsilon, \kappa, \nu)$ are given by:*

$$\begin{aligned} \varepsilon &= \operatorname{sgn}(\lambda_1), \quad \kappa = k_1 + \frac{1}{2}L_f(k_2 - L_g(k_3)) + \frac{1}{4}(k_2 - L_g(k_3))^2 + L_{[g,f]}(k_3) + \frac{1}{2}k_3L_g(k_2 - L_g(k_3)), \\ \text{and } \nu &= |\lambda_1|^{-1/2} \left(\lambda_3 - \frac{1}{2}L_g(\ln|\lambda_1|) \right). \end{aligned} \quad (11)$$

Our formula for the curvature κ is, indeed, a generalisation of that in [3] because if $k_3 = 0$ (i.e. we suppose that f is the singular vector field f_s), then (11) reads

$$\kappa = k_1 + \frac{1}{2}L_f(k_2) + \frac{1}{4}(k_2)^2,$$

that is to say, exactly as the formula given by Agrachev [1, 3].

Proof. Consider a pair (f, g) with structure functions (k_1, k_2, k_3) and $(\lambda_1, \lambda_2, \lambda_3)$. To deduce the expression of the invariants $(\varepsilon, \kappa, \nu)$, we apply equations (8) and (9) with $\alpha = k_3$ and $\beta = |\lambda_1|^{-1/2}$, namely the feedback transformation that constructs the canonical pair. We detail the computation for κ and left the computation for ν to the reader. Recall, that applying a feedback to construct the canonical pair, we obtain as a by-product $\tilde{k}_2 = 0$ (see the proof of Proposition 3). First we have,

$$\gamma = -\frac{1}{2}|\lambda_1|^{-1/2}L_f(\Lambda) - \frac{1}{2}k_3|\lambda_1|^{-1/2}L_g(\Lambda) - |\lambda_1|^{-1/2}L_g(k_3) = |\lambda_1|^{-1/2} \left(-\frac{1}{2}L_f(\Lambda) - \frac{1}{2}k_3L_g(\Lambda) - L_g(k_3) \right),$$

where $\Lambda = \ln |\lambda_1|$. Second, using $\tilde{k}_2 = 0$ we deduce

$$k_2 = -\frac{1}{2}L_f(\Lambda) - \frac{1}{2}k_3L_g(\Lambda) + |\lambda|^{1/2}\gamma = -L_f(\Lambda) - k_3L_g(\Lambda) - L_g(k_3).$$

Therefore, inserting the last expression of k_2 into γ , we get $\gamma = \frac{1}{2}|\lambda_1|^{-1/2}(k_2 - L_g(k_3))$. Now the curvature reads $\kappa = \tilde{k}_1$, i.e

$$\begin{aligned} \kappa &= k_1 + L_{[g,f]}(k_3) + |\lambda_1|^{1/2} \left(\frac{1}{2}L_f \left(|\lambda_1|^{-1/2} (k_2 - L_g(k_3)) \right) + \frac{1}{2}k_3L_g \left(|\lambda_1|^{-1/2} (k_2 - L_g(k_3)) \right) \right. \\ &\quad \left. - \frac{1}{2}|\lambda_1|^{-1/2} (k_2 - L_g(k_3)) L_g(k_3) \right), \\ &= k_1 + L_{[g,f]}(k_3) - \frac{1}{2}(k_2 - L_g(k_3))L_g(k_3) \\ &\quad + \frac{1}{2}|\lambda_1|^{1/2} \left(|\lambda_1|^{-1/2} L_f(k_2 - L_g(k_3)) - \frac{1}{2}(k_2 - L_g(k_3))|\lambda_1|^{-3/2} L_f(|\lambda_1|) \right) \\ &\quad + \frac{1}{2}|\lambda_1|^{1/2} k_3 \left(|\lambda_1|^{-1/2} L_g(k_2 - L_g(k_3)) - \frac{1}{2}(k_2 - L_g(k_3))|\lambda_1|^{-3/2} L_g(|\lambda_1|) \right) \\ &= k_1 + L_{[g,f]}(k_3) - \frac{1}{2}(k_2 - L_g(k_3))L_g(k_3) + \frac{1}{2}L_f(k_2 - L_g(k_3)) \\ &\quad - \frac{1}{4}(k_2 - L_g(k_3))L_f(\Lambda) + \frac{1}{2}k_3L_g(k_2 - L_g(k_3)) - \frac{1}{4}(k_2 - L_g(k_3))k_3L_g(\Lambda) \\ &= k_1 + L_{[g,f]}(k_3) + \frac{1}{2}L_f(k_2 - L_g(k_3)) + \frac{1}{2}k_3L_g(k_2 - L_g(k_3)) \\ &\quad - \frac{1}{4}(k_2 - L_g(k_3))(L_f(\Lambda) + L_g(\Lambda) + 2L_g(k_3)) \\ &= k_1 + \frac{1}{2}L_f(k_2 - L_g(k_3)) + \frac{1}{4}(k_2 - L_g(k_3))^2 + L_{[g,f]}(k_3) + \frac{1}{2}k_3L_g(k_2 - L_g(k_3)). \end{aligned}$$

□

Now consider a trivial system, whose state (x, y, w) belongs to a 3-dimensional manifold \mathcal{M} ,

$$(T) : \begin{cases} \dot{x} &= F_1(w) \\ \dot{y} &= F_2(w) \\ \dot{w} &= u \end{cases}, \quad (x, y, w) \in \mathcal{M}, \quad u \in \mathbb{R}.$$

Notice that (T) is, in general, not given by a canonical pair but is given by a semi-canonical pair since $[f, [f, g]] = 0$. Clearly, for trivial systems we have $\kappa = 0$, but the converse is not true as discovered in [16] and as we will show in the following theorem.

Theorem 3 (Characterisation of trivial systems). *Consider a control-affine system $\Sigma = (f, g)$ together with its structure functions κ and ν . Then, Σ is locally trivialisable if and only if its canonical form $\Sigma_c = (f_c, g_c)$ satisfies*

$$\kappa = 0, \quad L_{f_c}(\nu) = 0, \quad \text{and} \quad L_{[f_c, g_c]}(\nu) = 0. \quad (12)$$

Observe that the conditions of (12) can explicitly be tested on the control-affine system $\Sigma = (f, g)$. Indeed, with the help of Proposition 3, we explicitly construct the canonical pair (f_c, g_c) of Σ for which the invariants κ and ν can be computed by algebraic operations only. Another way to test condition (12) on an arbitrary control-affine system $\Sigma = (f, g)$ is to compute the invariants κ and ν using (11) and then to evaluate (12) with $f_c = f + gk_3$ and $g_c = |\lambda_1|^{-1/2}g$.

Proof. We begin with necessity and suppose that Σ is trivialisable. Then, (T) is given by $f = F_1(w)\frac{\partial}{\partial x} + F_2(w)\frac{\partial}{\partial y}$ and $g = \frac{\partial}{\partial w}$ (which, a priori, is not a canonical pair), whose structure functions are $k_1 = k_2 = k_3 = 0$, $\lambda_1 = \lambda_1(w)$, $\lambda_2 = 0$, and $\lambda_3 = \lambda_3(w)$. In particular, observe that λ_1 and λ_3 satisfy $L_f(\lambda_1) = L_{[g,f]}(\lambda_1) = 0$ and $L_f(\lambda_3) = L_{[g,f]}(\lambda_3) = 0$. As in the proof of Proposition 3, to transform

the pair (f, g) of (T) into the canonical pair (f_c, g_c) we use $\beta = |\lambda_1|^{-1/2}$, which therefore satisfies $L_f(\beta) = 0$ and $L_{[g, f]}(\beta) = 0$. Now, using equations (8) and (9) of Lemma 1, we calculate the structure functions of $(f_c, g_c) = (f, g\beta)$ which are $\tilde{\kappa} = 0$, $\tilde{\varepsilon} = \pm 1$, $\tilde{\mu} = 0$, and $\tilde{\nu} = \tilde{\lambda}_3 = \beta\lambda_3 + L_g(\beta)$. Hence, for the canonical pair (f_c, g_c) of (T) we have

$$\begin{aligned} L_{f_c}(\tilde{\nu}) &= \beta L_f(\lambda_3) + L_f(L_g(\beta)) = \beta L_f(\lambda_3) + L_g(L_f(\beta)) - L_{[g, f]}(\beta) = 0, \\ L_{[g_c, f_c]}(\tilde{\nu}) &= \beta L_{[g, f]}(\tilde{\nu}) = \beta (L_{[g, f]}(\lambda_3) + L_{[g, f]}(L_g(\beta))) = \beta (L_g(L_{[g, f]}(\beta)) - L_{[g, [g, f]]}(\beta)) \\ &= \beta (-\lambda_1 L_f(\beta) - \lambda_3 L_{[g, f]}(\beta)) = 0, \end{aligned}$$

and the necessity of (12) is proved.

Now, conversely, suppose that Σ_c , given by its canonical pair (f_c, g_c) , satisfies (12). First, due to Lemma 2 of Appendix B, we apply a diffeomorphism $(x, y, w) = \phi(\xi)$ that simultaneously rectifies the distribution $\mathcal{F} = \text{span}\{f_c, [g_c, f_c]\}$ and the vector field g_c , that is $\phi_*\mathcal{F} = \text{span}\left\{\frac{\partial}{\partial x}, \frac{\partial}{\partial y}\right\}$ and $\phi_*g_c = \frac{\partial}{\partial w}$. In those coordinates, we have $f_c = f_1\frac{\partial}{\partial x} + f_2\frac{\partial}{\partial y}$, with $f_i = f_i(x, y, w)$, and we have $\nu = \nu(w)$ since $L_{f_c}(\nu) = L_{[g_c, f_c]}(\nu) = 0$ and $f_c \wedge g_c \wedge [g_c, f_c] \neq 0$. Therefore, using relation (6') we deduce that f_c satisfies the following two equations (notice that equation (7c') together with $L_{f_c}(\nu) = 0$ imply that $\mu = 0$)

$$[f_c, [f_c, g_c]] = 0 \quad \text{and} \quad [g_c, [g_c, f_c]] = \varepsilon f_c + \nu(w) [g_c, f_c].$$

The second equation reads

$$\frac{\partial^2 f_c}{\partial w^2} = \varepsilon f_c + \nu(w) \frac{\partial f_c}{\partial w}, \quad (13)$$

and, interpreted as a second order linear ODE with respect to w and with parameters (x, y) , admits local solutions of the form

$$f_c(x, y, w) = F_1(w) \begin{pmatrix} a_1 \\ a_2 \\ 0 \end{pmatrix} + F_2(w) \begin{pmatrix} b_1 \\ b_2 \\ 0 \end{pmatrix} = F_1(w)A + F_2(w)B. \quad (14)$$

In (14), $F_1(w)$ and $F_2(w)$ are smooth fundamental solutions functions of (13) (i.e. $F_1(w_0) = 1$, $F_1'(w_0) = 0$, $F_2(w_0) = 0$, and $F_2'(w_0) = 1$) and $a_i = a_i(x, y)$ and $b_i = b_i(x, y)$, for $i = 1, 2$, so $A = a_1\frac{\partial}{\partial x} + a_2\frac{\partial}{\partial y}$ and $B = b_1\frac{\partial}{\partial x} + b_2\frac{\partial}{\partial y}$ are smooth vector fields on \mathbb{R}^2 equipped with coordinates (x, y) .

Using the commutativity of f_c and $[g_c, f_c]$ we deduce that

$$[F_1A + F_2B, F_1'A + F_2'B] = (F_1F_2' - F_1'F_2) [A, B] = 0.$$

By $F_1F_2' - F_1'F_2 \neq 0$ (since $f_c \wedge [g_c, f_c] \neq 0$), we conclude that $[A, B] = 0$ and, therefore, there exists a local diffeomorphism $\psi(x, y)$ that simultaneously rectify A and B (seen as vector fields on \mathbb{R}^2). For simplicity, we still denote the new coordinates by (x, y) , i.e. we have $\psi_*A = \frac{\partial}{\partial x}$ and $\psi_*B = \frac{\partial}{\partial y}$. In coordinates (x, y, w) , the vector fields (f_c, g_c) take the form

$$f_c = F_1(w) \frac{\partial}{\partial x} + F_2(w) \frac{\partial}{\partial y} \quad \text{and} \quad g_c = \frac{\partial}{\partial w}$$

and therefore we conclude that the system $\Sigma_c = (f_c, g_c)$ is trivial. \square

Remark that in our proof we start with a canonical pair (f_c, g_c) and we render it trivial by constructing a suitable local coordinate system.

Remark 4. *The previous theorem was first discovered by Serres in [17]. In the proof of [17, Theorem 4.3.3] (but also in [18, Theorem 4.3] and in [16, Theorem 3.4]), he shows, using his notation, that $\alpha_2 = a_2(u, q_2) - q_1$ and $\frac{\partial a_2}{\partial q_2} = b(u)$ and then consider the case $\alpha_2 = a_2(u) - q_1$ and not the general case $\alpha_2 = b(u)q_2 + a_2(u) - q_1$. The proof of [17, Theorem 4.3.3], given for the case $b \equiv 0$ (which, using our notation, is equivalent to $\nu \equiv 0$), still provides an inspiring intuition to treat the general case, as done in our proof.*

In the following proposition, we express the structure functions of a trivial system (T) and give two canonical forms of control-affine system that are trivialisable. Both canonical forms are expressed using the canonical pair but in different coordinate systems. For two smooth scalar functions $F(w)$ and $G(w)$, we define their Wronskian as $\mathbb{W}(F, G) = F'G - FG'$. Recall that for any control-affine system $\Sigma = (f, g)$ satisfying (A1) and (A2) we defined, via (6), structure functions k_1, k_2, k_3 and $\lambda_1, \lambda_2, \lambda_3$.

Proposition 5. *Consider a control-affine system $\Sigma = (f, g)$ and suppose that it satisfies conditions (12) of Theorem 3. Then, locally, the following hold*

(i) Σ admits the normal form (T), that is,

$$\Sigma^T : \begin{cases} \dot{x} &= F_1(w) \\ \dot{y} &= F_2(w) \\ \dot{w} &= u \end{cases}$$

whose structure functions are $k_1 = k_2 = k_3 = 0$ and $\lambda_1 = -\frac{\mathbb{W}(F_1', F_2')}{\mathbb{W}(F_1, F_2)}$, $\lambda_2 = 0$, $\lambda_3 = \frac{\mathbb{W}'(F_1, F_2)}{\mathbb{W}(F_1, F_2)}$.

(ii) Σ admits the canonical forms $\Sigma_c^{T,1}$ and $\Sigma_c^{T,2}$ given, respectively, by

$$\Sigma_c^{T,1} : \begin{cases} \dot{x} &= F_{c,1}(w) \\ \dot{y} &= F_{c,2}(w) \\ \dot{w} &= u \end{cases} \quad \text{and} \quad \Sigma_c^{T,2} : \begin{cases} \dot{x} &= 1 + \varepsilon y u \\ \dot{y} &= (x - \nu(w)y) u \\ \dot{w} &= u \end{cases},$$

where $\frac{\mathbb{W}(F_{c,1}', F_{c,2}')}{\mathbb{W}(F_{c,1}, F_{c,2})} \equiv \pm 1$ and whose invariants are $(\varepsilon^1, \kappa^1, \nu^1) = \left(-\frac{\mathbb{W}(F_{c,1}', F_{c,2}')}{\mathbb{W}(F_{c,1}, F_{c,2})}, 0, \frac{\mathbb{W}'(F_{c,1}, F_{c,2})}{\mathbb{W}(F_{c,1}, F_{c,2})}\right)$ and $(\varepsilon^2, \kappa^2, \nu^2) = (\varepsilon, 0, \nu(w))$, respectively.

Remark 5. *Neither the structure functions k_i nor λ_i are feedback invariant. Item (i) asserts that for the normal form $\Sigma^T = (f, g)$, all $k_i = 0$, so the pair (f, g) is semi-canonical (thus, actually, $f = f_s$) but, in general, it is not canonical since λ_1 is a non trivial function. Item (ii) assures that, given $\Sigma^T = (f, g)$, we can always choose w_c -coordinate, as $w = \phi(w_c)$, such that $F_{c,i} = \phi^* F_i$ satisfy $\frac{\mathbb{W}(F_{c,1}', F_{c,2}')}{\mathbb{W}(F_{c,1}, F_{c,2})} = \pm 1$ and the corresponding pair (f_c, g_c) , where $f_c = F_{c,1} \frac{\partial}{\partial x} + F_{c,2} \frac{\partial}{\partial y}$ and $g_c = \beta g$, with $\beta = \phi'$, is canonical.*

Proof. The normal form presented in item (i) is a direct consequence of Theorem 1 and it is a straightforward computation to derive the expressions of the structure functions. To obtain the canonical form $\Sigma_c^{T,1}$ of item (ii), we consider Σ^T and define $g_c = \beta g$, where $\beta = |\lambda_1|^{-1/2}$, see Proposition 3. We choose $w = \phi(\hat{w})$ satisfying $(\phi^{-1})' \beta = 1$. Then in the coordinates (x, y, \hat{w}) , the system Σ^T takes the form $\Sigma_c^{T,1}$, where $F_{c,i} = \phi^* F_i$ and whose third equation reads $\hat{w} = \hat{u}$.

Finally, the canonical form $\Sigma_c^{T,2}$ is a special case of item (i) of Proposition 6 presented in the next section. \square

The two presented canonical forms are somehow *dual* to each other. Indeed, both are given in terms of the canonical pair (f_c, g_c) of Σ and for $\Sigma_c^{T,1}$ we adopt coordinates for which the vector field g_c^1 is rectified, whereas for $\Sigma_c^{T,2}$ the coordinates are chosen so that f_c^2 is rectified. The two normal forms carry complementary informations about the control-affine system Σ . The canonical form $\Sigma_c^{T,1}$ exhibits the trivial nature of Σ , but its invariants ε and ν are not immediately visible, and the canonical form $\Sigma_c^{T,2}$ explicitly identifies the invariants $\varepsilon = \pm 1$ and ν but hides the triviality of the system. The two canonical forms show that trivial systems depend on a smooth function of one variable: for $\Sigma_c^{T,2}$ it is clearly $\nu(w)$ and for $\Sigma_c^{T,1}$ it is the function $F_{c,2}(w)$ that determines $F_{c,1}(w)$ (or, equivalently, the other way around) through the ODE $\frac{\mathbb{W}(F_{c,1}', F_{c,2}')}{\mathbb{W}(F_{c,1}, F_{c,2})} = \pm 1$.

4 Normal forms of flat and centro-flat control-affine systems on 3D-manifolds

We have shown that the curvature κ and the centro-affine curvature ν are two functional feedback equivariants of control-affine systems, hence, their properties define non-equivalent classes of systems.

In this section, we propose a normal form for each class of control-systems that is presented in [Table 2](#) below. The presented classes describe all the cases for which the curvature κ and the centro-affine curvature ν satisfy $\kappa\nu \equiv 0$ together with the particular sub-cases for which, additionally, either κ or ν is constant.

Notation	Name	Properties
$\Sigma^{\varepsilon, \kappa=0, \nu}$	Flat	Curvature κ vanishes
$\Sigma^{\varepsilon, \kappa, \nu=0}$	Centro-flat	Centro-affine curvature ν vanishes
$\Sigma^{\varepsilon, \kappa'=0, \nu=0}$	Flat-constant	Curvature κ is constant and the centro-affine curvature ν vanishes
$\Sigma^{\varepsilon, \kappa=0, \nu'=0}$	Centro-flat-constant	Curvature κ vanishes, and the centro-affine curvature ν is constant
$\Sigma^{\varepsilon, \kappa=0, \nu=0}$	Completely flat	Curvatures κ and ν vanish

Table 2: Nomenclature of subclasses of flat and centro-flat control-affine systems

Each class of control system presented in the above table is denoted by an upper index $I = (a, b, c)$, which is defined as follows. The first element is always $\varepsilon = \pm 1$ and emphasises the dependence of the normal forms on the invariant ε ; the second element is either $\kappa = 0$ to say that the curvature vanishes or $\kappa' = 0$ to express that the curvature is constant (this notation is a bit abusive because κ is not a function of one variable in general); finally, the third index is either $\nu = 0$ or $\nu' = 0$ with the same interpretation as previously. The following proposition provides a normal form Σ_c^I for each class of control-affine systems Σ^I , where the upper multi-index I is one of the five given in [Table 2](#). The lower index c indicates that all normal forms Σ_c^I are expressed using their canonical pairs. Recall that the structure function ν is unique up to its sign, i.e. changing $g_c \mapsto -g_c$ yields $\nu \mapsto -\nu$, hence in normal forms below we suppose that $\nu \geq 0$.

Proposition 6 (Normal forms of flat control-affine systems). *Consider a control-affine system $\Sigma = (f, g)$ together with its invariants ε , κ , and ν . Then, the following statements hold locally (all normal forms below are represented by a canonical pair (f_c, g_c) and considered around an arbitrary point $(x_0, y_0, w_0) \in \mathbb{R}^3$).*

(i) *If $\kappa = 0$, then Σ is locally feedback equivalent to*

$$\Sigma_c^{\varepsilon, \kappa=0, \nu} : \begin{cases} \dot{x} &= 1 + a(y, w)u \\ \dot{y} &= (x + b(y, w))u \\ \dot{w} &= c(y, w)u \end{cases},$$

whose invariants are ε , $\kappa = 0$, and $\nu = \nu_1(y, w)x + \nu_0(y, w)$, and the functions a, b and c satisfy the following differential equations

$$\begin{aligned} \frac{\partial a}{\partial y} &= \varepsilon + \nu_1(y, w)a(y, w), & a(y_0, w) &= 0, \\ \frac{\partial b}{\partial y} &= \nu_1(y, w)b(y, w) - \nu_0(y, w), & b(y_0, w) &= 0, \\ \frac{\partial c}{\partial y} &= \nu_1(y, w)c(y, w), & c(y_0, w) &= 1, \end{aligned}$$

and thus are given by

$$\begin{aligned} a(y, w) &= \left[\varepsilon \int_{y_0}^y \exp \left(- \int_{y_0}^{\tau} \nu_1(t, w) dt \right) d\tau \right] \exp \left(\int_{y_0}^y \nu_1(\tau, w) d\tau \right), \\ b(y, w) &= \left[- \int_{y_0}^y \nu_0(\tau, w) \exp \left(- \int_{y_0}^{\tau} \nu_1(t, w) dt \right) d\tau \right] \exp \left(\int_{y_0}^y \nu_1(\tau, w) d\tau \right), \\ c(y, w) &= \exp \left(\int_{y_0}^y \nu_1(\tau, w) d\tau \right). \end{aligned}$$

(ii) If $\nu = 0$, then Σ is locally feedback equivalent to

$$\Sigma_c^{\varepsilon, \kappa, \nu=0} : \begin{cases} \dot{x} &= r(x, y) c_\varepsilon(w) \\ \dot{y} &= r(x, y) s_\varepsilon(w) \\ \dot{w} &= \varepsilon \frac{\partial r}{\partial y} c_\varepsilon(w) + \frac{\partial r}{\partial x} s_\varepsilon(w) + u \end{cases},$$

where

$$c_\varepsilon(w) = \frac{e^{w\sqrt{\varepsilon}} + e^{-w\sqrt{\varepsilon}}}{2} \quad \text{and} \quad s_\varepsilon(w) = \frac{e^{w\sqrt{\varepsilon}} - e^{-w\sqrt{\varepsilon}}}{2\sqrt{\varepsilon}},$$

whose invariants are ε , $\kappa = \kappa(x, y)$, $\nu = 0$, and the function $r(x, y)$ satisfies $r > 0$ and the following non-linear partial differential equation

$$-r(x, y)^2 \left(\frac{\partial^2}{\partial x^2} - \varepsilon \frac{\partial^2}{\partial y^2} \right) (\ln r(x, y)) = \kappa(x, y). \quad (15)$$

(iii) If κ and ν are constant then

$$\kappa\nu = 0, \quad (16)$$

i.e. at least one of them vanishes.

(iv) If $\kappa = 0$ and ν is constant, then Σ is locally feedback equivalent to

(a) If $\varepsilon = 1$, then

$$\Sigma_c^{+, \kappa=0, \nu'=0} : \begin{cases} \dot{x} &= e^{\nu w} e^{w\sqrt{\nu^2+4}} \\ \dot{y} &= e^{\nu w} e^{-w\sqrt{\nu^2+4}} \\ \dot{w} &= \frac{1}{2}u \end{cases}, \quad \text{where } \nu \geq 0;$$

(b) If $\varepsilon = -1$ and $\nu > 2$, then

$$\Sigma_c^{-, \kappa=0, \nu'=0, +} : \begin{cases} \dot{x} &= e^{\nu w} e^{w\sqrt{\nu^2-4}} \\ \dot{y} &= e^{\nu w} e^{-w\sqrt{\nu^2-4}} \\ \dot{w} &= \frac{1}{2}u \end{cases};$$

(c) If $\varepsilon = -1$ and $\nu = 2$, then

$$\Sigma_c^{-, \kappa=0, \nu'=0, 0} : \begin{cases} \dot{x} &= e^w \\ \dot{y} &= we^w \\ \dot{w} &= u \end{cases};$$

(d) If $\varepsilon = -1$ and $0 \leq \nu < 2$, then

$$\Sigma_c^{-, \kappa=0, \nu'=0, -} : \begin{cases} \dot{x} &= e^{\nu w} \cos\left(w\sqrt{4-\nu^2}\right) \\ \dot{y} &= e^{\nu w} \sin\left(w\sqrt{4-\nu^2}\right) \\ \dot{w} &= \frac{1}{2}u \end{cases}$$

Moreover, for the four normal forms above, ε , $\kappa = 0$, and ν are complete invariants.

(v) If $\nu = 0$ and κ is constant, then Σ is locally feedback equivalent to

$$\Sigma_c^{\varepsilon, \kappa'=0, \nu=0} : \begin{cases} \dot{x} &= \left(1 - \frac{\kappa}{4}(x^2 - \varepsilon y^2)\right) c_\varepsilon(w) \\ \dot{y} &= \left(1 - \frac{\kappa}{4}(x^2 - \varepsilon y^2)\right) s_\varepsilon(w) \\ \dot{w} &= \frac{-\kappa}{2}(y c_\varepsilon(w) - x s_\varepsilon(w)) + u \end{cases},$$

whose complete invariants are ε , κ , and $\nu = 0$.

(vi) If $\kappa = 0$ and $\nu = 0$, then Σ is locally feedback equivalent to

$$\Sigma_c^{\varepsilon, \kappa=0, \nu=0} : \begin{cases} \dot{x} &= c_\varepsilon(w) \\ \dot{y} &= s_\varepsilon(w) \\ \dot{w} &= u \end{cases} .$$

Before presenting a proof for those normal forms, we give some remarks about them. For item (i) we adopt coordinates, where the vector field f_c is rectified, whereas for the other normal forms we choose coordinates in which the vector field g_c is rectified. The first normal form $\Sigma_c^{\varepsilon, \kappa=0, \nu}$ of flat control-affine systems describes the most general form of a system for which the curvature κ vanishes. On the other hand, the normal form $\Sigma_c^{\varepsilon, \kappa, \nu=0}$ of item (ii), describes systems for which the centro-affine curvature ν vanishes. All other items are then special cases of those two general normal forms.

Recall that ν is unique up to its sign, that is why in item (iv) we have $\nu \geq 0$ for (iv)-(a) to (iv)-(d). It is remarkable that if κ and ν are constant (hence true invariants) then at least one of them is zero as asserted in item (iii). Moreover, relation (16) already appeared in [19], where the four families of normal forms given by $\kappa = 0$ and ν constant were listed (but the non-invariance of the sign of ν was not discussed there). The two normal forms of (vi) with $\varepsilon = \pm 1$ and $\kappa = \nu = 0$ are, respectively, given by

$$\Sigma_c^{+, \kappa=0, \nu=0} : \begin{cases} \dot{x} &= \cosh(w) \\ \dot{y} &= \sinh(w) \\ \dot{w} &= u \end{cases} \quad \text{and} \quad \Sigma_c^{-, \kappa=0, \nu=0} : \begin{cases} \dot{x} &= \cos(w) \\ \dot{y} &= \sin(w) \\ \dot{w} &= u \end{cases}$$

correspond to hyperbolic and elliptic systems without parameters and have been extensively analysed and differently characterised in [14, 15].

Proof. For each item, we consider a control-affine system $\Sigma_c = (f_c, g_c)$ given by the canonical pair and with $\varepsilon = \pm 1$ and structure functions κ and ν .

(i) Since $\kappa = 0$, using relation (6'), we conclude that the vector fields f_c and $[f_c, g_c]$ are commuting. Therefore, we can rectify them simultaneously to get $f_c = \frac{\partial}{\partial x}$ and $[f_c, g_c] = \frac{\partial}{\partial y}$. Afterwards, we determine the form of the vector field g_c . First, it satisfies $[\frac{\partial}{\partial x}, g_c] = \frac{\partial}{\partial y}$ and thus we immediately conclude

$$g_c = a(y, w) \frac{\partial}{\partial x} + (x + b(y, w)) \frac{\partial}{\partial y} + c(y, w) \frac{\partial}{\partial w} .$$

Moreover, assumptions (A1) and (A2) imply that $c \neq 0$ and $\frac{\partial}{\partial y} (\frac{a}{c}) \neq 0$. Second, for g_c we have $[g_c, -\frac{\partial}{\partial y}] = \varepsilon \frac{\partial}{\partial x} + \mu g_c - \nu \frac{\partial}{\partial y}$, where the functions μ and ν satisfy (7b') and (7c') and therefore $\nu = \nu_1(y, w)x + \nu_0(y, w)$ and $\mu = \nu_1(y, w)$. Hence, the functions a , b , and c of g_c satisfy

$$\begin{aligned} \frac{\partial a}{\partial y}(y, w) &= \varepsilon + \nu_1(y, w)a(y, w), \\ \frac{\partial b}{\partial y}(y, w) &= \nu_1(y, w)b(y, w) - \nu_0(y, w), \\ \frac{\partial c}{\partial y}(y, w) &= \nu_1(y, w)c(y, w). \end{aligned}$$

Solutions of those equations are, respectively,

$$a(y, w) = \left[\varepsilon \int_{y_0}^y \exp \left(- \int_{y_0}^\tau \nu_1(t, w) dt \right) d\tau + A(w) \right] \exp \left(\int_{y_0}^y \nu_1(\tau, w) d\tau \right), \quad (17a)$$

$$b(y, w) = \left[- \int_{y_0}^y \nu_0(\tau, w) \exp \left(- \int_{y_0}^\tau \nu_1(t, w) dt \right) d\tau + B(w) \right] \exp \left(\int_{y_0}^y \nu_1(\tau, w) d\tau \right), \quad (17b)$$

$$c(y, w) = C(w) \exp \left(\int_{y_0}^y \nu_1(\tau, w) d\tau \right). \quad (17c)$$

Updating the coordinates, we can set $C(w) = 1$, $A(w) = B(w) = 0$; and in those coordinates we obtain the normal form $\Sigma_c^{\varepsilon, \kappa=0, \nu}$.

(ii) Suppose that $\nu = 0$ and choose coordinates $(\bar{x}, \bar{y}, \bar{w})$ such that $g_c = \frac{\partial}{\partial \bar{w}}$. Then, by relation (6') we conclude that

$$f_c = \bar{A}(\bar{x}, \bar{y})c_\varepsilon(\bar{w}) + \bar{B}(\bar{x}, \bar{y})s_\varepsilon(\bar{w}), \quad c_\varepsilon(\bar{w}) = \frac{e^{\bar{w}\sqrt{\varepsilon}} + e^{-\bar{w}\sqrt{\varepsilon}}}{2}, \quad s_\varepsilon(\bar{w}) = \frac{e^{\bar{w}\sqrt{\varepsilon}} - e^{-\bar{w}\sqrt{\varepsilon}}}{2\sqrt{\varepsilon}},$$

where $\bar{A} = a_1 \frac{\partial}{\partial \bar{x}} + a_2 \frac{\partial}{\partial \bar{y}} + a_3 \frac{\partial}{\partial \bar{w}}$, with $a_i = a_i(\bar{x}, \bar{y})$, and $\bar{B} = b_1 \frac{\partial}{\partial \bar{x}} + b_2 \frac{\partial}{\partial \bar{y}} + b_3 \frac{\partial}{\partial \bar{w}}$, with $b_i = b_i(\bar{x}, \bar{y})$ are smooth vector fields. By assumption (A1), we conclude that $a_1 b_2 - a_2 b_1 \neq 0$, hence $\bar{\bar{A}} = a_1 \frac{\partial}{\partial \bar{x}} + a_2 \frac{\partial}{\partial \bar{y}}$ and $\bar{\bar{B}} = b_1 \frac{\partial}{\partial \bar{x}} + b_2 \frac{\partial}{\partial \bar{y}}$ form a moving frame of the tangent bundle of $\mathcal{X} = \mathcal{O}/\mathcal{G}$, where \mathcal{O} is an open subset of \mathbb{R}^3 , in which the rectifying coordinates $(\bar{x}, \bar{y}, \bar{w})$ are defined, and $\mathcal{G} = \text{span} \left\{ \frac{\partial}{\partial \bar{w}} \right\}$. We define a metric $\bar{\bar{g}}$ on \mathcal{X} by declaring $(\bar{\bar{A}}, \bar{\bar{B}})$ orthonormal, i.e.

$$\bar{\bar{g}}(\bar{\bar{A}}, \bar{\bar{A}}) = 1, \quad \bar{\bar{g}}(\bar{\bar{A}}, \bar{\bar{B}}) = 0, \quad \text{and} \quad \bar{\bar{g}}(\bar{\bar{B}}, \bar{\bar{B}}) = -\varepsilon.$$

Notice that the signature of $\bar{\bar{g}}$ is $(+, -\text{sgn}(\varepsilon))$, hence $\bar{\bar{g}}$ is definite for $\varepsilon = -1$ and indefinite for $\varepsilon = 1$. Since all metrics on 2-dimensional manifolds are locally conformally flat, we conclude that there exists an isometry $(x, y) = \psi(\bar{x}, \bar{y})$ such that $\bar{\bar{g}} = \psi^* \mathbf{g}$, where $\mathbf{g} = \rho(x, y)(dx^2 - \varepsilon dy^2)$ with $\rho > 0$. In the coordinates (x, y) , both the pair (\tilde{A}, \tilde{B}) , with $\tilde{A} = \psi_* \bar{\bar{A}}$ and $\tilde{B} = \psi_* \bar{\bar{B}}$, and $\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right)$ form an orthonormal frame for \mathbf{g} so we have

$$(\tilde{A}, \tilde{B}) = r(x, y)I(x, y) \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right),$$

where $r = \frac{1}{\sqrt{\rho}}$ and $I(x, y)$ is a linear isometry, i.e. it belongs to the (pseudo)-orthonormal group $O(1, -\varepsilon)$. Using a suitable change of the variable $w = \bar{w} + h(x, y)$ we can get rid of $I(x, y)$. Finally, in coordinates (x, y, w) , the vector field f_c of the control system takes the form

$$f_c = r(x, y)c_\varepsilon(w) \frac{\partial}{\partial x} + r(x, y)s_\varepsilon(w) \frac{\partial}{\partial y} + (a(x, y)c_\varepsilon(w) + b(x, y)s_\varepsilon(w)) \frac{\partial}{\partial w}.$$

We now use the structure equations (6') and deduce that necessarily

$$a(x, y) = \varepsilon \frac{\partial r}{\partial y} \quad \text{and} \quad b(x, y) = \frac{\partial r}{\partial x}$$

and that r satisfies

$$\varepsilon r \frac{\partial^2 r}{\partial y^2} - \varepsilon \left(\frac{\partial r}{\partial y} \right)^2 + \left(\frac{\partial r}{\partial x} \right)^2 - r \frac{\partial^2 r}{\partial x^2} = \kappa(x, y).$$

which can be expressed in the form of (15).

(iii) If κ and ν are constants, then due to relation (10), we immediately conclude (16).

(iv) Suppose that $\kappa = 0$ and ν is constant, then Σ satisfies condition (12) of Theorem 3 and thus Σ is locally trivialisable. Using the results of item (ii) of Proposition 5, we take Σ in the form of $\Sigma_c^{T,1}$ for which $f_c = F_{c,1}(w) \frac{\partial}{\partial x} + F_{c,2}(w) \frac{\partial}{\partial y}$ and $g_c = \frac{\partial}{\partial w}$ form a canonical pair. Using (6'), we conclude that the functions $F_{c,i}$, for $i = 1, 2$, satisfy the following second order linear ordinary differential equation

$$F_{c,i}''(w) = \varepsilon F_{c,i}(w) + \nu F_{c,i}'(w). \quad (18)$$

Solutions are dictated by the sign of the discriminant $\Delta = \nu^2 + 4\varepsilon$ of the characteristic polynomial of the ODE. Moreover, the roots of the characteristic polynomial are $r_{1/2} = \frac{\nu \pm \sqrt{\Delta}}{2}$. Recall that the sign of ν is not invariant and thus by choosing w suitably we can always get $\nu \geq 0$. Moreover, it is a trivial calculation to check that the solutions given below are fundamental solutions of (18), i.e. we just need to compute the Wronskian at w_0 .

- (a) If $\varepsilon = +1$, then $\Delta > 0$ for all $\nu \geq 0$. Solutions of (18) are given by (after normalising w with $\frac{1}{2}$)

$$F_{c,1}(w) = e^{\nu w} e^{w\sqrt{\nu^2+4}} \quad \text{and} \quad F_{c,2}(w) = e^{\nu w} e^{-w\sqrt{\nu^2+4}},$$

and we obtain the normal form $\Sigma_c^{+, \kappa=0, \nu'=0}$.

- (b) If $\varepsilon = -1$ and $\nu > 2$, then $\Delta > 0$, and solving (18) gives

$$F_{c,1}(w) = e^{\nu w} e^{w\sqrt{\nu^2-4}}, \quad \text{and} \quad F_{c,2}(w) = e^{\nu w} e^{-w\sqrt{\nu^2-4}},$$

and we obtain the normal form $\Sigma_c^{-, \kappa=0, \nu'=0, +}$.

- (c) If $\varepsilon = -1$ and $\nu = 2$, then $\Delta = 0$, and the solutions of (18) are

$$F_{c,1}(w) = we^w, \quad \text{and} \quad F_{c,2}(w) = e^w,$$

which gives $\Sigma_c^{-, \kappa=0, \nu'=0, 0}$.

- (d) If $\varepsilon = -1$ and $0 \leq \nu < 2$, then $\Delta < 0$, and the solutions of (18) are

$$F_{c,1}(w) = e^{\nu w} \cos\left(w\sqrt{4-\nu^2}\right), \quad \text{and} \quad F_{c,2}(w) = e^{\nu w} \sin\left(w\sqrt{4-\nu^2}\right),$$

which gives $\Sigma_c^{-, \kappa=0, \nu'=0, -}$.

- (v) Assume that κ is constant and $\nu = 0$, then we refine the normal form $\Sigma_c^{\varepsilon, \kappa, \nu=0}$ of item (ii). We recognize that equation (15) satisfied by $r(x, y)$ describes the curvature (in the usual differential geometry sense) of the metric $\mathbf{g} = \frac{1}{r^2}(dx^2 - \varepsilon dy^2)$. By assumption, the curvature of \mathbf{g} is constant (equal to $-\kappa$) and by Minding's theorem, surfaces with the same constant curvature are locally isometric. Therefore, there exists an isometry $(\tilde{x}, \tilde{y}) = \psi(x, y)$ such that $\mathbf{g} = \psi^* \tilde{\mathbf{g}}$ with

$$\tilde{\mathbf{g}} = \left(\frac{1}{1 - \frac{\kappa}{4}(\tilde{x}^2 - \varepsilon \tilde{y}^2)} \right)^2 (d\tilde{x}^2 - \varepsilon d\tilde{y}^2),$$

which is also of curvature $-\kappa$. The action of the isometry on (\tilde{x}, \tilde{y}) can be compensated by applying $w \mapsto w + h(x, y)$, for a suitable function h , thus we obtain that the system takes the form of $\Sigma_c^{\varepsilon, \kappa, \nu=0}$ with $r(x, y) = 1 - \frac{\kappa}{4}(x^2 - \varepsilon y^2)$, i.e. we get $\Sigma_c^{\varepsilon, \kappa'=0, \nu=0}$.

- (vi) The normal forms $\Sigma_c^{\varepsilon, \kappa=0, \nu=0}$ is a special case of item (v) with $\kappa = 0$.

□

5 Conclusions and Perspectives

In this paper, we have analysed in details the notion of triviality adapted to the context of control-affine systems. We proposed two new characterisations of trivial control-affine system, one of them is based on the existence of an abelian subalgebra of the Lie algebra of infinitesimal symmetries. In particular, we gave a normal form of trivial control-affine systems for which the Lie algebra of infinitesimal symmetries has a transitive almost abelian Lie subalgebra. In the future, we will be interested in extending our result to the case of multi-input systems and we will try to propose other characterisation of triviality that are purely geometric. In the second part of the paper, we have revisited results due to Serres [17] and we give novel proof of his characterisation of trivial systems on 3-dimensional manifolds with scalar inputs. In particular, our characterisation uses a discrete invariant $\varepsilon = \pm 1$ and two well-defined functional invariants of feedback transformations: the curvature κ (introduced by Agrachev [2]) and the centro-affine curvature ν (studied by Wilkens [19]). We show that those invariants can explicitly be computed for any control-affine system and that a canonical pair of vector fields (f_c, g_c) , on which κ and ν appear explicitly, can also be constructed with a purely algebraically defined feedback transformation.

Then, we extended the results of Serres and Wilkens by giving several normal forms of control-affine systems. In the future, our goal is two-folds: first we will be interested in the question of how to enlarge the triple $(\varepsilon, \kappa, \nu)$ to a set of *complete invariants* of control-affine systems (on 3D manifolds with scalar control). Identifying a set of complete invariants would be helpful in obtaining normal forms of control-affine system in dimension three. Second, we will be interested in generalising our characterisation of trivial control-affine systems to the multi-input case, in particular the notion of curvature of dynamics pairs, as proposed in [11], seems promising.

References

- [1] Andrei A. Agrachev. “Feedback-Invariant Optimal Control Theory and Differential Geometry, II. Jacobi Curves for Singular Extremals”. *Journal of Dynamical and Control Systems* 4.4 (1998). DOI: [10.1023/A:1021871218615](https://doi.org/10.1023/A:1021871218615).
- [2] Andrei A. Agrachev and Revaz V. Gamkrelidze. “Feedback-Invariant Optimal Control Theory and Differential Geometry—I. Regular Extremals”. *Journal of Dynamical and Control Systems* 3.3 (1997). DOI: [10.1007/BF02463256](https://doi.org/10.1007/BF02463256).
- [3] Andrei A. Agrachev and Yuri Sachkov. *Control Theory from the Geometric Viewpoint*. Vol. 87. Springer Science & Business Media, 2013. DOI: [10.1007/978-3-662-06404-7](https://doi.org/10.1007/978-3-662-06404-7).
- [4] Zhirayr Avetisyan. “Structure of Almost Abelian Lie Algebras”. *arXiv:1610.05365 [math]* (2016). URL: <http://arxiv.org/abs/1610.05365>.
- [5] Dietrich Burde and Manuel Ceballos. “Abelian Ideals of Maximal Dimension for Solvable Lie Algebras”. *arXiv:0911.2995 [math]* (2011). URL: <http://arxiv.org/abs/0911.2995>.
- [6] Lester E. Dubins. “On Curves of Minimal Length with a Constraint on Average Curvature, and with Prescribed Initial and Terminal Positions and Tangents”. *American Journal of Mathematics* 79.3 (1957). DOI: [10.2307/2372560](https://doi.org/10.2307/2372560).
- [7] Michel Fliess et al. “On Differentially Flat Nonlinear Systems”. French. *IFAC Proceedings Volumes* 25.13 (1992). DOI: [10.1016/S1474-6670\(17\)52275-2](https://doi.org/10.1016/S1474-6670(17)52275-2).
- [8] Jessy Grizzle and Steven Marcus. “The Structure of Nonlinear Control Systems Possessing Symmetries”. *IEEE Transactions on Automatic Control* 30.3 (1985).
- [9] Alberto Isidori. *Nonlinear Control Systems*. Communications and Control Engineering, Nonlinear Control Systems. Springer-Verlag, 1995. DOI: [10.1007/978-1-84628-615-5](https://doi.org/10.1007/978-1-84628-615-5).
- [10] Bronislaw Jakubczyk. “Equivalence and Invariants of Nonlinear Control Systems”. *NonLinear Controllability and Optimal Control*. Vol. 133. Routledge, 1990.
- [11] Bronislaw Jakubczyk and Wojciech Kryński. “Vector Fields with Distributions and Invariants of ODEs”. *Journal of Geometric Mechanics* 5.1 (2013). DOI: [10.3934/jgm.2013.5.85](https://doi.org/10.3934/jgm.2013.5.85).
- [12] Witold Respondek and Issa A. Tall. “Nonlinearizable Single-Input Control Systems Do Not Admit Stationary Symmetries”. *Systems & Control Letters* 46.1 (2002). DOI: [10.1016/S0167-6911\(01\)00197-9](https://doi.org/10.1016/S0167-6911(01)00197-9).
- [13] Timothée Schmoderer. “Study of Control Systems under Quadratic Nonholonomic Constraints. Motion Planning, Introduction to the Regularised Continuation Method.” PhD thesis. INSA Rouen Normandie, 2022. URL: <http://theses.fr/s221665>.
- [14] Timothée Schmoderer and Witold Respondek. *Conic Nonholonomic Constraints on Surfaces and Control Systems*. 2021. URL: <http://arxiv.org/abs/2106.08635>.
- [15] Timothée Schmoderer and Witold Respondek. “Null-forms of conic systems in \mathbb{R}^3 are determined by their symmetries”. *Systems & Control Letters* 170 (2022). DOI: <https://doi.org/10.1016/j.sysconle.2022.105397>.
- [16] Ulysse Serres. “Control Systems of Zero Curvature Are Not Necessarily Trivializable”. *arXiv:0902.2332 [math]* (2009). URL: <http://arxiv.org/abs/0902.2332>.

- [17] Ulysse Serres. “Geometry and Feedback Classification of Low-Dimensional Non-Linear Control Systems”. PhD thesis. Université de Bourgogne, 2006. URL: <https://tel.archives-ouvertes.fr/tel-00172902>.
- [18] Ulysse Serres. “On Curvature and Feedback Classification of Two-Dimensional Optimal Control Systems”. *Journal of Mathematical Sciences* 144.1 (2007). DOI: [10.1007/s10958-007-0237-8](https://doi.org/10.1007/s10958-007-0237-8).
- [19] George R. Wilkens. “Centro-Affine Geometry in the Plane and Feedback Invariants of Two-State Scalar Control Systems”. *Proceedings of Symposia in Pure Mathematics*. Ed. by G. Ferreyra et al. Vol. 64. American Mathematical Society, 1998. DOI: [10.1090/pspum/064/1654544](https://doi.org/10.1090/pspum/064/1654544).

A Detailed computations for Lemma 1

In this appendix, we detail the computation to obtain relations (7a)-(7c) between structure functions and we prove transformation rules (8) and (9) that show how the structure functions are changed under a feedback transformation. Consider a control-affine system $\Sigma = (f, g)$ with structure functions (k_1, k_2, k_3) and $(\lambda_1, \lambda_2, \lambda_3)$.

First, by applying the Jacobi identity to $[f, [g, [g, f]]]$ we deduce that $[f, [g, [g, f]]] = -[g, [f, [f, g]]]$. We compute the left-hand-side and the right-hand-side separately:

$$\begin{aligned}
[f, [g, [g, f]]] &= L_f(\lambda_1) f + L_f(\lambda_2) g - \lambda_2 [g, f] + L_f(\lambda_3) - \lambda_3 (k_1 g + k_2 [g, f] + k_3 [g, [g, f]]), \\
&= L_f(\lambda_1) f + (L_f(\lambda_2) - \lambda_3 k_1) g + (L_f(\lambda_3) - \lambda_2 - \lambda_3 k_2) [g, f] \\
&\quad - \lambda_3 k_3 (\lambda_1 f + \lambda_2 g + \lambda_3 [g, f]), \\
&= (L_f(\lambda_1) - \lambda_3 k_3 \lambda_1) f + (L_f(\lambda_2) - \lambda_3 k_1 - \lambda_3 k_3 \lambda_2) g \\
&\quad + (L_f(\lambda_3) - \lambda_2 - \lambda_3 k_2 - \lambda_3^2 k_3) [g, f].
\end{aligned}$$

And on the other hand we have

$$\begin{aligned}
[g, [f, [f, g]]] &= L_g(k_1) g + L_g(k_2) [g, f] + k_2 (\lambda_1 f + \lambda_2 g + \lambda_3 [g, f]) \\
&\quad + L_g(k_3) (\lambda_1 f + \lambda_2 g + \lambda_3 [g, f]) + k_3 [g, \lambda_1 f + \lambda_2 g + \lambda_3 [g, f]], \\
&= (k_2 \lambda_1 + \lambda_1 L_g(k_3)) f + (L_g(k_1) + k_2 \lambda_2 + L_g(k_3) \lambda_2) g \\
&\quad + (L_g(k_2) + k_2 \lambda_3 + L_g(k_3) \lambda_3) [g, f] \\
&\quad + k_3 (L_g(\lambda_1) f + \lambda_1 [g, f] + L_g(\lambda_2) g + L_g(\lambda_3) [g, f] + \lambda_3 (\lambda_1 f + \lambda_3 g + \lambda_3 [g, f])), \\
&= (k_2 \lambda_1 + \lambda_1 L_g(k_3) + k_3 L_g(\lambda_1) + k_3 \lambda_3 \lambda_1) f \\
&\quad + (L_g(k_1) + k_2 \lambda_2 + L_g(k_3) \lambda_2 + k_3 L_g(\lambda_2) + k_3 \lambda_3 \lambda_2) g \\
&\quad + (L_g(k_2) + k_2 \lambda_3 + L_g(k_3) \lambda_3 + k_3 \lambda_1 + k_3 L_g(\lambda_3) + k_3 \lambda_3^2) [g, f]
\end{aligned}$$

Identifying the terms in front of f , g , and $[g, f]$ we obtain equations (7a) to (7c).

Now, we apply a feedback transformation of the form $\tilde{f} = f + g\alpha$ and $\tilde{g} = g\beta$ and we get first $[\tilde{f}, \tilde{g}] = \beta [f, g] + \gamma g$, where $\gamma = L_f(\beta) + \alpha L_g(\beta) - \beta L_g(\alpha)$. Second,

$$\begin{aligned}
[\tilde{g}, [\tilde{g}, \tilde{f}]] &= \beta^2 [g, [g, f]] + \beta L_g(\beta) [g, f] + (-\beta L_{[g, f]}(\beta) - \beta L_g(\gamma) + \gamma L_g(\beta)) g \\
&= \beta^2 \lambda_1 f + (\beta^2 \lambda_2 - \beta L_{[g, f]}(\beta) - \beta L_g(\gamma) + \gamma L_g(\beta)) g + (\beta^2 \lambda_3 + \beta L_g(\beta)) [g, f] \\
&= \beta^2 \lambda_1 \tilde{f} + (-\beta^2 \lambda_1 \alpha + \beta^2 \lambda_2 - \beta L_{[g, f]}(\beta) - \beta L_g(\gamma) + \gamma L_g(\beta) + \gamma (\beta \lambda_3 + L_g(\beta))) g \\
&\quad + (\beta \lambda_3 + L_g(\beta)) [\tilde{g}, \tilde{f}],
\end{aligned}$$

implying that $\tilde{\lambda}_1 = \beta^2 \lambda_1$, $\tilde{\lambda}_2 = \beta \lambda_2 - \beta \lambda_1 \alpha + \gamma \lambda_3 - L_{[g, f]}(\beta) - L_g(\gamma) + 2\gamma L_g(\ln|\beta|)$, and $\tilde{\lambda}_3 =$

$\beta\lambda_3 + \mathbf{L}_g(\beta)$. Third, we have

$$\begin{aligned}
[\tilde{f}, [\tilde{f}, \tilde{g}]] &= [f + g\alpha, \beta[f, g] + \gamma g] \\
&= \beta[f, [f, g]] + \mathbf{L}_f(\beta)[f, g] + \mathbf{L}_f(\gamma)g + \gamma[f, g] \\
&\quad + \alpha\beta[g, [f, g]] + \alpha\mathbf{L}_g(\beta)[f, g] - \beta\mathbf{L}_{[f, g]}(\alpha)g + \alpha\mathbf{L}_g(\gamma)g - \gamma\mathbf{L}_g(\alpha)g, \\
&= (\beta k_1 + \mathbf{L}_f(\gamma) + \beta\mathbf{L}_{[g, f]}(\alpha) + \alpha\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\alpha))g + (\beta k_2 - \mathbf{L}_f(\beta) - \gamma - \alpha\mathbf{L}_g(\beta))[g, f] \\
&\quad + (\beta k_3 - \alpha\beta)[g, [f, g]], \\
&= (\beta k_1 + \mathbf{L}_f(\gamma) + \beta\mathbf{L}_{[g, f]}(\alpha) + \alpha\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\alpha))g + (\beta k_2 - \mathbf{L}_f(\beta) - \gamma - \alpha\mathbf{L}_g(\beta))[g, f] \\
&\quad + \frac{1}{\beta^2}(\beta k_3 - \alpha\beta) \left\{ [\tilde{g}, [\tilde{g}, \tilde{f}]] - \beta\mathbf{L}_g(\beta)[g, f] + \beta\mathbf{L}_{[g, f]}(\beta)g + \beta\mathbf{L}_g(\gamma)g - \gamma\mathbf{L}_g(\beta)g \right\},
\end{aligned}$$

implying $\tilde{k}_3 = \frac{1}{\beta}(k_3 - \alpha)$. Next, continuing the computation (denoting $\tilde{h} = \tilde{k}_3[\tilde{g}, [\tilde{g}, \tilde{f}]]$):

$$\begin{aligned}
[\tilde{f}, [\tilde{f}, \tilde{g}]] &= \left(\beta k_1 + \mathbf{L}_f(\gamma) + \beta\mathbf{L}_{[g, f]}(\alpha) + \alpha\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\alpha) + \tilde{k}_3(\beta\mathbf{L}_{[g, f]}(\beta) + \beta\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\beta)) \right) g + \tilde{h} \\
&\quad + \left(\beta k_2 - \mathbf{L}_f(\beta) - \gamma - \alpha\mathbf{L}_g(\beta) - \tilde{k}_3\beta\mathbf{L}_g(\beta) \right) [g, f] + \tilde{h} \\
&= \left(\beta k_1 + \mathbf{L}_f(\gamma) + \beta\mathbf{L}_{[g, f]}(\alpha) + \alpha\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\alpha) + \tilde{k}_3(\beta\mathbf{L}_{[g, f]}(\beta) + \beta\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\beta)) \right) g \\
&\quad + \frac{1}{\beta} \left(\beta k_2 - \mathbf{L}_f(\beta) - \gamma - \alpha\mathbf{L}_g(\beta) - \tilde{k}_3\beta\mathbf{L}_g(\beta) \right) \left\{ [\tilde{g}, \tilde{f}] + \gamma g \right\} + \tilde{h},
\end{aligned}$$

implying $\tilde{k}_2 = k_2 - \mathbf{L}_f(\ln|\beta|) - \frac{\gamma}{\beta} - \alpha\mathbf{L}_g(\ln|\beta|) - \tilde{k}_3\mathbf{L}_g(\beta)$ and finally

$$\begin{aligned}
\tilde{k}_1 &= \frac{1}{\beta} \left(\beta k_1 + \mathbf{L}_f(\gamma) + \beta\mathbf{L}_{[g, f]}(\alpha) + \alpha\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\alpha) + \tilde{k}_3(\beta\mathbf{L}_{[g, f]}(\beta) + \beta\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\beta)) + \tilde{k}_2\gamma \right) \\
&= k_1 + \mathbf{L}_{[g, f]}(\alpha) + \frac{1}{\beta} \left(\mathbf{L}_f(\gamma) + \alpha\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\alpha) + \tilde{k}_3(\beta\mathbf{L}_{[g, f]}(\beta) + \beta\mathbf{L}_g(\gamma) - \gamma\mathbf{L}_g(\beta)) + \tilde{k}_2\gamma \right).
\end{aligned}$$

B Technical lemma for the proof of Theorem 3

The sufficiency part of the proof of Theorem 3 relies on the existence of a diffeomorphism that simultaneously rectifies the distribution $\text{span}\{f_c, [g_c, f_c]\}$ and the vector field g_c as proven by the following lemma.

Lemma 2. *Consider a control-affine system $\Sigma_c = (f_c, g_c)$ given by its canonical pair and assume set $\mathcal{F} = \text{span}\{f_c, [g_c, f_c]\}$. If, the structure functions of Σ_c satisfy the condition (12) of Theorem 3, then, there exists a diffeomorphism $(x, y, w) = \phi(\xi)$ such that $\phi_*\mathcal{F} = \text{span}\left\{\frac{\partial}{\partial x}, \frac{\partial}{\partial y}\right\}$ and $\phi_*g_c = \frac{\partial}{\partial w}$.*

Proof. First, we prove that there exists smooth solutions h for the system

$$\mathbf{L}_{f_c}(h) = 0, \quad \mathbf{L}_{[g_c, f_c]}(h) = 0, \quad \text{and} \quad \mathbf{L}_{g_c}(h) = 1.$$

We need to check three integrability conditions:

1. $\mathbf{L}_{[f_c, [g_c, f_c]]}(h) = \mathbf{L}_{f_c}(\mathbf{L}_{[g_c, f_c]}(h)) - \mathbf{L}_{[g_c, f_c]}(\mathbf{L}_{f_c}(h)) = 0$ and $[f_c, [g_c, f_c]] = 0$, so $0 = 0$ and the first condition holds.
2. $\mathbf{L}_{[f_c, g_c]}(h) = \mathbf{L}_{f_c}(\mathbf{L}_{g_c}(h)) - \mathbf{L}_{g_c}(\mathbf{L}_{f_c}(h)) = 0$ and $\mathbf{L}_{[f_c, g_c]}(h) = 0$ so $0 = 0$ and the second integrability condition holds.
3. $\mathbf{L}_{[[g_c, f_c], g_c]}(h) = \mathbf{L}_{[g_c, f_c]}(\mathbf{L}_{g_c}(h)) - \mathbf{L}_{g_c}(\mathbf{L}_{[g_c, f_c]}(h)) = 0$ and $[[g_c, f_c], g_c] = -\varepsilon f_c - \nu[g_c, f_c]$. Therefore $\mathbf{L}_{[[g_c, f_c], g_c]}(h) = -\varepsilon\mathbf{L}_{f_c}(h) - \nu\mathbf{L}_{[g_c, f_c]}(h) = 0$ and $0 = 0$ the third condition holds.

Take a smooth solution h of the above system, rename it $\phi_3 = h$, and choose ϕ_1, ϕ_2 such that $d\phi_1$ and $d\phi_2$ annihilate g_c and are independent (they exist since $g_c \neq 0$). The diffeomorphism $\phi = (\phi_1, \phi_2, \phi_3) = (x, y, w)$ is such that $\phi_*\mathcal{F} = \text{span}\left\{\frac{\partial}{\partial x}, \frac{\partial}{\partial y}\right\}$ and $\phi_*g_c = \frac{\partial}{\partial w}$. \square