



HAL
open science

Behavioral effect of selective deletion of beta2* nicotinic acetylcholine receptors in neuropeptide Y expressing interneurons

Alice Abbondanza, Alexandra Tyshkevich, Jigar Dhabalia Trisha, Jan Elias, Radislav Sedláček, Stéphanie Daumas, Véronique Bernard, Helena Janickova

► To cite this version:

Alice Abbondanza, Alexandra Tyshkevich, Jigar Dhabalia Trisha, Jan Elias, Radislav Sedláček, et al.. Behavioral effect of selective deletion of beta2* nicotinic acetylcholine receptors in neuropeptide Y expressing interneurons. NeuroFrance 2021, May 2021, Online, France. hal-03972951

HAL Id: hal-03972951

<https://hal.science/hal-03972951>

Submitted on 3 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Behavioral effect of selective deletion of beta2 nicotinic acetylcholine receptors in neuropeptide Y expressing interneurons*

Abbondanza A.¹, Tyshkevich A.¹, Dhabalia T. J.¹, Elias J.², Sedlacek R.², Daumas S.³, Bernard V.³, Janickova H.¹

¹ Department of Neurochemistry, Institute of Physiology of the Czech Academy of Sciences - Prague, Czech Republic

² Czech Centre for Phenogenomics, Institute of Molecular Genetics ASCR - Vestec, Czech Republic

³ Neuroscience Paris Seine, Sorbonne Université - CNRS UMR 8246 Paris, France

Neuropeptide Y (NPY) is an abundant neuropeptide in the neocortex involved in numerous processes. NPY-expressing neurons differ in electrophysiological, molecular and morphological properties but overall, they are GABAergic interneurons (GINs). All cortical layers receive cholinergic inputs from the basal forebrain, which are crucial for cognition. Cholinergic receptors are differentially distributed across the cortex and the activation of nicotinic acetylcholine receptors (nAChRs) located on principal neurons and GINs modulates synaptic plasticity and behavior.

Functional consequences of the cholinergic activation of NPY-expressing cortical interneurons (NPY+GINs) have not been specifically clarified yet. By using the CRISPR/Cas9 approach, we deleted the most abundant beta2* subunit of nAChRs expressed by NPY+GINs in the prefrontal cortex (PFC) and investigated the effect on behavior.

By in situ hybridization, we characterized the expression of beta2* nAChRs on NPY+GINs. To induce a cell specific deletion of beta2* nAChRs, we crossed a mouse line expressing Cas9-GFP in a Cre-dependent manner with a NPY-Cre line. The resulting mice were injected with AAV vector carrying sgRNA targeting CHRN2, gene coding for beta2 nicotinic subunit, into PFC. Mutant mice and littermate controls, injected with scrambled gRNA, were tested in a battery of behavioral tasks including social preference task and novel object recognition. Immunohistochemical evaluation of neuronal markers and activity indicators showed changes in activation/inhibition balance in cortical circuits.

The successful induction of indels in CHRN2 gene in AAV-transduced cells was confirmed by FACS sorting followed by DNA sequencing. Mice with presumed deletion of nAChRs showed increased social interaction compared to controls.

The achieved efficiency of CRISPR/Cas9-induced indels in CHRN2 gene in NPY+GINs of the PFC was sufficient to induce behavioral changes in mice. Specifically, the increased sociability agrees with *Avale et al., 2011*.

This work was supported by the Grant Agency of the Czech Republic Grant #19-07983Y.