



HAL
open science

IMPACT OF THE ANGULAR VARIATION OF THE TRUNK DURING A PROPULSION CYCLE USING INERTIAL MEASUREMENT UNITS (IMU)

Florian Brassart, Didier Pradon, Eric Watelain, Sadate Bakatchina, Ilona Alberca, Thierry Weissland, Arnaud Faupin

► **To cite this version:**

Florian Brassart, Didier Pradon, Eric Watelain, Sadate Bakatchina, Ilona Alberca, et al.. IMPACT OF THE ANGULAR VARIATION OF THE TRUNK DURING A PROPULSION CYCLE USING INERTIAL MEASUREMENT UNITS (IMU). 19ème congrès de l'association des chercheurs en activités physiques et sportives, Oct 2021, Montpellier, France. 10.1051/sm/2020016. . hal-03978236

HAL Id: hal-03978236

<https://hal.science/hal-03978236>

Submitted on 8 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IMPACT OF THE ANGULAR VARIATION OF THE TRUNK DURING A PROPULSION CYCLE USING INERTIAL MEASUREMENT UNITS (IMU)

FLORIAN BRASSART^{1,3} ; DIDIER PRADON² ; ERIC WATELAIN¹ ; SADATE BAKATCHINA¹ ; ILONA ALBERCA¹ ; THIERRY WEISSLAND³ ; ARNAUD FAUPIN¹

¹Université de Toulon, IAPS - N°201723207F, 83957 La Garde, France ; ²Pole Parasport - ISPC Synergies, CHU Raymond Poincaré, APHP, Garches, France ; ³Laboratoire IMS, UMR 5218, PMH_DySCo, Pessac, France.

florian.brassart@univ-tln.fr

Introduction : Different propulsion strategies exist in wheelchair sports depending on the sport, equipment, and classification of the athlete. In athletes with high classifications, and therefore greater trunk muscles strength, have greater involvement of the trunk during a sprint, whose varies with pushing velocity [1]. One study showed that isometric trunk strength in wheelchair athletes was not directly correlated with the acceleration as the latter depended on rolling resistance [2]. But in this study, the dynamic action of the trunk was not measured during propulsion. For this reason, it would be interesting to study the influence of trunk movements during propulsion in sports requiring more mobility such as wheelchair basketball.

Hypothese: The flexion/extension range of motion of the trunk (FEROMT) have an impact on the variations in velocity during a sprint.

Protocol : 13 elite female wheelchair basketball players 5 with a classification at 3.5 and above (high points) and 8 with a classification lower than 3.5 (small points) with their own sports wheelchair.

The players ran 6 x 20 m sprints successively in a straight line with 20 seconds of recovery between each.

Results :

- **Correlation** between the **mean acceleration** per pushing phase and **FEROMT** ($p=0.015$).
- **No correlation** between the **mean deceleration** per recovery phase and **FEROMT** ($p=0.103$).
- **No correlation** between the **classification** of athletes and **FEROMT** ($p=0.068$).

However, on average high points had a greater FEROMT than small points ($21.52 \pm 6^\circ$ and $14.7 \pm 2.6^\circ$, $p=0.00017$) as seen in the literature.




Figure 2 : Relationship between the mean acceleration (blue) / deceleration (purple) and the mean FEROMT for each cycle of propulsion during the sprint for each players.

IMUs were placed :

- on the spokes of each wheel [3]
- in the center of the back at the spine level, **between Th2 and Th8**

IMU measured the rotation velocity of the wheels and the trunk with integrated gyrometers

→ Compute of wheels linear velocity and FEROMT



Figure 1 : Female wheelchair basketball player during a sprint with 3 IMUs.




Figure 3 : Wheelchairs velocity and trunk variation during 3 cycles of propulsion of a 20 m sprint.

Discussion : FEROMT seems to differ according to classification however differences are not systematic. The movement of the trunk during wheelchair propulsion appears to be beneficial in order to accelerate without increasing the deceleration during the recovery phase. This could indicate that trunk motion is beneficial to performance, bringing further insight to the study of Vanlandewijck et al. (2011) [2].

Bibliography

- [1] Veeger, H. E., van der Woude, L. H., & Rozendal, R. H. (1989). Wheelchair propulsion technique at different speeds. *Scandinavian Journal of Rehabilitation Medicine*, 21(4), 197-203.
- [2] Vanlandewijck, Yves C., Verellen, J., Beckman, E., Connick, M., & Tweedy, S. M. (2011). Trunk strength effect on track wheelchair start : Implications for classification. *Medicine and Science in Sports and Exercise*, 43(12), 2344-2351. <https://doi.org/10.1249/MSS.0b013e318223af14>
- [3] Bakatchina, S., Weissland, T., & Faupin, A. (2021). Les dispositifs d'évaluation des parasportifs pratiquant des sports de petits terrains en fauteuil roulant manuel. *Movement & Sport Sciences - Science & Motricité*. <https://doi.org/10.1051/sm/2020016>.