



HAL
open science

The multi oxygen isotope analyses on black crust from Sicily highlight the volcanic emission influence from Mount Etna on urban areas

Adeline Aroskay, Erwan Martin, Slimane Bekki, Giuseppe Montana, Luciana Randazzo, Pierre Cartigny

► To cite this version:

Adeline Aroskay, Erwan Martin, Slimane Bekki, Giuseppe Montana, Luciana Randazzo, et al.. The multi oxygen isotope analyses on black crust from Sicily highlight the volcanic emission influence from Mount Etna on urban areas. EGU General Assembly 2020, May 2020, Vienne, Austria. 2020, 10.5194/egusphere-egu2020-20136 . hal-03979023

HAL Id: hal-03979023

<https://hal.science/hal-03979023v1>

Submitted on 8 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU2020-20136

<https://doi.org/10.5194/egusphere-egu2020-20136>

EGU General Assembly 2020

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



The multi oxygen isotope analyses on black crust from Sicily highlight the volcanic emission influence from Mount Etna on urban areas

Adeline Aroskay¹, Erwan Martin¹, Slimane Bekki², Giuseppe Montana³, Luciana Randazzo⁴, and Pierre Cartigny⁵

¹Institut des Sciences de la Terre de Paris (ISTeP), Sorbonne Université, Paris, France

²Laboratoire Atmosphères, Milieux, Observations Spatiales (LATMOS), Sorbonne Université, Paris, France

³Dipartimento di Scienze della Terra e del Mare (DiSTeM), Università degli Studi di Palermo, Palermo, Italy

⁴Department of Biology, Ecology and Earth Sciences (DiBEST), Università della Calabria, Italy

⁵Institut de Physique du Globe de Paris (IPGP), Université de Paris, Paris, France

This study reports on measurements of $\Delta^{17}\text{O}$ (derived from the triple oxygen isotopes) in sulphate from black crust sampled in Sicily. Atmospheric oxidants, such as O_3 , H_2O_2 , OH and O_2 carry specific ^{17}O -anomalies, which are partly transferred to the sulphate during sulphur gas (e.g. SO_2) oxidation. Hence, the $\Delta^{17}\text{O}$ in sulphate can be used as a tracer of sulphur oxidation pathways. So far, this method has been mostly applied on sulphate from aerosols, rainwaters, volcanic deposits and ice cores. Here we propose a new approach, that aims to investigate the dominant oxidants of gaseous sulphur precursors into sulphate extracted from black crust material. Black crusts are mostly found on building/monument/sculpture and are the result of the reaction between sulphur compounds (SO_2 , H_2SO_4) and carbonate (CaCO_3) from the substrate, which leads to the formation of gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$). Sicilian black crust from sites under different emission influences (anthropogenic, marine and volcanic) were collected. Multi oxygen and sulphur isotope analyses were performed to better assess the origins of black crust sulphate in these different environments. This is crucial for both a better understanding of the sulphur cycle and the preservation of historical monument.

Multi sulphur isotopes show mostly negative values ranging from -0.4‰ to $0.02\text{‰} \pm 0.01$ and from -0.59‰ to $0.41\text{‰} \pm 0.3$ for $\Delta^{33}\text{S}$ and $\Delta^{36}\text{S}$ respectively. This is unique for natural samples and different from sulphate aerosols measured around the world ($\Delta^{33}\text{S} > 0\text{‰}$). This tends to indicate that sulphate from black crust is not generated by the same processes as sulphate aerosols in the atmosphere. Instead of SO_2 oxidation in the atmosphere, dry deposition of SO_2 and its oxidation on the substratum is preferred. The multi oxygen isotopes show a clear dependence with the geographical repartition of the samples. Indeed, black crusts from Palermo (the biggest Sicilian city) show small ^{17}O -anomalies ranging between -0.16‰ to 1.02‰ with an average value of $0.45\text{‰} \pm 0.26$ ($n=12$; 2σ). This is consistent with $\Delta^{17}\text{O}$ values measured in black crust from the Parisian Basin (Genot et al., 2020), which are also formed in an environment influenced by anthropogenic and marine emissions. On the other hand, samples from the eastern part of the Mount Etna

region, which are downwind of the volcanic emissions, show the highest ^{17}O -anomalies ranging from 0.48 ‰ to 3.87 ‰ with an average value of $2.7 \text{ ‰} \pm 0.6$ ($n=11$; 2σ).

These results indicate that volcanic emissions influence the oxygen isotopic signature of black crust sulphate. In standard urban areas, SO_2 deposited on the substratum is mostly oxidised by O_2 -TMI and H_2O_2 to generate the black crust. Yet, under the influence of volcanic emissions, O_3 may play the main role in the SO_2 oxidation.