



HAL
open science

Some remarks on the ergodic theorem for U -statistics

Herold G. Dehling, Davide Giraud, Dalibor Volny

► **To cite this version:**

Herold G. Dehling, Davide Giraud, Dalibor Volny. Some remarks on the ergodic theorem for U -statistics. 2023. hal-03979769v1

HAL Id: hal-03979769

<https://hal.science/hal-03979769v1>

Preprint submitted on 8 Feb 2023 (v1), last revised 4 Apr 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SOME REMARKS ON THE ERGODIC THEOREM FOR U -STATISTICS

HEROLD DEHLING, DAVIDE GIRAUDDO AND DALIBOR VOLNÝ

ABSTRACT. In this note, we investigate the convergence of a U -statistic of order two having stationary ergodic data. We will find sufficient conditions for the almost sure and L^1 convergence and present some counter-examples showing that the U -statistic itself might fail to converge: centering is needed as well as boundedness of $\sup_{j \geq 2} \mathbb{E}[|h(X_1, X_j)|]$.

1. INTRODUCTION

In this note, we investigate the validity of the U -statistics ergodic theorem, i.e. the almost sure convergence

$$(1) \quad \frac{1}{\binom{n}{2}} \sum_{1 \leq j < j' \leq n} h(X_i, X_{j'}) \longrightarrow \iint h(x, y) dF(x) dF(y),$$

where $(X_i)_{i \geq 1}$ is a stationary ergodic process with marginal distribution F , and $h(x, y)$ is a symmetric kernel that is $F \times F$ integrable. Birkhoff's ergodic theorem establishes the analogous result for the time averages $\frac{1}{n} \sum_{i=1}^n f(X_i)$, while Hoeffding [6] established (1) for i.i.d. processes $(X_i)_{i \geq 1}$. These two classical results naturally lead to the conjecture that (1) should hold without further assumptions, i.e. for all stationary ergodic processes $(X_i)_{i \geq 1}$ and all $L_1(F \times F)$ functions $h(x, y)$. Aaronson et al. [1] proved a partial result in this direction, namely showing that (1) holds for all $F \times F$ almost everywhere continuous and bounded kernels $h(x, y)$. At the same time, they presented counterexamples showing that (1) does not hold in full generality. One of their counterexamples is a bounded kernel where the set of discontinuities has positive $F \times F$ measure, while the other counterexample is an $F \times F$ almost everywhere continuous, but unbounded kernel.

The U -statistic ergodic theorem has subsequently been addressed by various authors, e.g. Arcones [2], Borovkova, Burton and Dehling [4]; see also the review paper by Borovkova, Burton and Dehling [5]. These papers provide both sufficient conditions for (1) to hold, as well as further counterexamples, both for stationary ergodic processes as well as under stronger mixing assumptions. Most of the positive results also address other forms of convergence in (1) such as convergence in probability and L^1 -convergence. Arcones [2] proved the ergodic theorem for absolutely regular processes under some moment assumptions. Borovkova, Burton and Dehling

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, 44780 BOCHUM, GERMANY

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE UMR 7501, UNIVERSITÉ DE STRASBOURG AND CNRS
7 RUE RENÉ DESCARTES 67000 STRASBOURG, FRANCE

UNIVERSITY OF ROUEN, LMRS AND CNRS UMR 6085.

E-mail address: herold.dehling@ruhr-uni-bochum.de, dgirauddo@unistra.fr,

dalibor.volny@univ-rouen.fr.

Date: February 8, 2023.

Key words and phrases. U -statistics, ergodic theorem, stationary sequences.

[5] investigated convergence in probability in (1), with a special focus on the kernel $h(x, y) = \log(|x - y|)$, which arises in connection with the Takens estimator for the correlation dimension.

A common feature of all these examples is that they satisfy a modified version of the U -statistics ergodic theorem, namely

$$(2) \quad \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} (h(X_i, X_j) - \mathbb{E}[h(X_i, X_j)]) \longrightarrow 0,$$

assuming that $\mathbb{E}[|h(X_i, X_j)|] < \infty$ for all i, j .

It might thus seem natural to conjecture that (2) holds without further assumptions. In this note, we present a counterexample that disproves this conjecture. In addition, we will give a short proof of the U -statistics ergodic theorem for bounded $F \times F$ -almost everywhere continuous kernels, and give a new condition for L^1 -convergence.

2. A SHORT PROOF OF THE ERGODIC THEOREM FOR U -STATISTICS

In this note, we present a short proof of the U -statistics ergodic theorem that was first established in Aaronson et al [1]. For the special case, when the process has values in \mathbb{R}^k , this proof is contained in Borovkova, Burton and Dehling [5]. Here, we give the proof for processes with values in an arbitrary separable metric space.

Theorem 2.1. *Let $(X_k)_{k \geq 0}$ be a stationary ergodic process with values in the separable metric space S and marginal distribution F , and let $h : S \times S \rightarrow \mathbb{R}$ be a symmetric kernel that is bounded and $F \times F$ -almost everywhere continuous. Then, as $n \rightarrow \infty$*

$$\frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j) \longrightarrow \iint h(x, y) dF(x) dF(y),$$

almost surely.

Proof. We define the empirical distribution of the first n random variables

$$F_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i},$$

where δ_x denotes the Dirac delta measure in x . For any $L_1(F)$ -function $f : S \rightarrow \mathbb{R}$, we obtain by Birkhoff's ergodic theorem

$$\int_S f(x) dF_n(x) = \frac{1}{n} \sum_{i=1}^n f(X_i) \rightarrow \int_S f(x) dF(x),$$

almost surely. This convergence holds in particular for any bounded measurable function $f \in C_b(S)$. Since S is separable, there exists a countably family of functions $f_i \in C_b(S)$, $i \geq 1$, that is convergence determining, i.e. that convergence of the integrals $\int f_i(x) d\mu_n(x) \rightarrow \int f_i(x) d\mu(x)$, for all $i \geq 1$, implies weak convergence of the probability measures μ_n to μ . Now, up to a set of measure 0, we get

$$\int_S f_i(x) dF_n(x) = \frac{1}{n} \sum_{j=1}^n f_i(X_j) \rightarrow \int_S f_i(x) dF(x),$$

for all $i \geq 1$, and thus $F_n \Rightarrow F$ weakly. This is in fact Varadarajan's argument [8] for the fact that the empirical distribution of i.i.d. data X_1, \dots, X_n converges weakly almost surely to the true distribution F .

By Theorem 3.2 (page 21) of Billingsley [3], we obtain convergence of the empirical product measure

$$F_n \times F_n \Rightarrow F \times F,$$

except on a set of measure 0. Thus, for any bounded $F \times F$ -a.e. continuous function $h: S \times S \rightarrow \mathbb{R}$, we obtain by the portmanteau theorem

$$\frac{1}{n^2} \sum_{1 \leq i, j \leq n} h(X_i, X_j) = \iint h(x, y) dF_n(x) dF_n(y) \rightarrow \iint h(x, y) dF(x) dF(y),$$

almost surely. Since h is bounded, we obtain $\frac{1}{n^2} \sum_{i=1}^n h(X_i, X_i) \rightarrow 0$, and thus

$$\frac{1}{n^2} \sum_{1 \leq i \neq j \leq n} h(X_i, X_j) \rightarrow \iint h(x, y) dF(x) dF(y),$$

almost surely. □

3. CONVERGENCE IN L^1 IN THE ERGODIC THEOREM FOR U -STATISTICS

In this section, we present two sufficient conditions for the convergence in L^1 of a U -statistic to $\iint h(x, y) dF(x) dF(y)$, where F denotes the distribution of X_0 . The first sufficient condition imposes a restriction on the continuity points of the kernel combined with a uniform integrability assumption. The second sufficient condition imposes a restriction on the joint distribution of vectors (X_0, X_k) , $k \geq 1$, but no other assumption is required for the kernel h .

Theorem 3.1. *Let $(X_i)_{i \geq 1}$ be a stationary ergodic sequence taking values in \mathbb{R}^d and let $h: \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a measurable function such that the family $\{h(X_1, X_j), j \geq 1\}$ is uniformly integrable. Let F be the distribution of X_1 . Assume that one of the following assumptions is satisfied:*

(A.1) *the function h is $F \times F$ almost everywhere continuous.*

(A.2) *$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |h(x, y)| dF(x) dF(y)$ is finite, the random variable X_0 has a bounded density with respect to the Lebesgue measure on \mathbb{R}^d and for each $k \geq 1$, the vector (X_0, X_k) has a density f_k with respect to the Lebesgue measure of $\mathbb{R}^d \times \mathbb{R}^d$ and $\sup_{k \geq 1} \sup_{s, t \in \mathbb{R}^d} f_k(s, t)$ is finite.*

Then

$$(3) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j) - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h(x, y) dF(x) dF(y) \right| \right] = 0.$$

Proof. Let us prove Theorem 3.1 under assumption (A.1). By Theorem 1 in [4], we know that $\frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j) \rightarrow \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h(x, y) dF(x) dF(y)$ in probability. Then it suffices to notice that uniform integrability of $\{h(X_1, X_j), j \geq 1\}$ implies that of $\left\{ \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j), n \geq 2 \right\}$.

We will prove Theorem 3.1 under assumption (A.2) in three steps: first we will show that (3) holds when h is a product of indicator functions of Borel subsets of \mathbb{R}^d . Then we will show the result by approximating the map $(x, y) \in \mathbb{R}^d \times \mathbb{R}^d \mapsto h(x, y) \mathbf{1}_{[-R, R]^d}(x) \mathbf{1}_{[-R, R]^d}(y) \mathbf{1}_{|h(x, y)| \leq R}$

in $L^1(\mathbb{P}_{(X_0, X_k)})$ uniformly with respect to k by a linear combination of products of indicator functions. Then we will conclude by uniform integrability.

First step: assume that $h(x, y) = \mathbf{1}_A(x) \mathbf{1}_B(y)$, where A and B are Borel subsets of \mathbb{R}^d . Observe that

$$(4) \quad \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j) = \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} \mathbf{1}_A(X_i) \mathbf{1}_B(X_j)$$

$$(5) \quad = \frac{1}{\binom{n}{2}} \sum_{j=2}^n \mathbf{1}_B(X_j) \sum_{i=1}^{j-1} \mathbf{1}_A(X_i)$$

$$(6) \quad = \frac{1}{\binom{n}{2}} \sum_{j=2}^n (j-1) \mathbf{1}_B(X_j) Y_j,$$

where

$$(7) \quad Y_j = \frac{1}{j-1} \sum_{i=1}^{j-1} \mathbf{1}_A(X_i).$$

Therefore, the following decomposition takes place:

$$(8) \quad \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j) \\ = \frac{1}{\binom{n}{2}} \sum_{j=2}^n (j-1) \mathbf{1}_B(X_j) (Y_j - \mathbb{P}(X_0 \in A)) + \mathbb{P}(X_0 \in A) \frac{1}{\binom{n}{2}} \sum_{j=2}^n (j-1) \mathbf{1}_B(X_j)$$

Observe that by the ergodic theorem and the Lebesgue dominated convergence theorem, the first term of the right hand side of (8) converges to 0 in L^1 . Moreover, by the ergodic theorem and a summation by parts,

$$(9) \quad \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{j=2}^n (j-1) \mathbf{1}_B(X_j) - \mathbb{P}(X_0 \in B) \right| \right] \rightarrow 0,$$

hence we derive that

$$(10) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} \mathbf{1}_A(X_i) \mathbf{1}_B(X_j) - \mathbb{P}(X_0 \in A) \mathbb{P}(X_0 \in B) \right| \right] = 0$$

and $\mathbb{P}(X_0 \in A) \mathbb{P}(X_0 \in B) = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h(x, y) dF(x) dF(y)$.

Second step. Let $R > 0$ be fixed and define

$$(11) \quad h^{(R)}(x, y) = h(x, y) \mathbf{1}_{[-R, R]^d}(x) \mathbf{1}_{[-R, R]^d}(y) \mathbf{1}_{|h(x, y)| \leq R},$$

which is integrable. By a standard result in measure theory, we know that for each positive ε , there exists an integer N , constants c_1, \dots, c_N and sets $A_{\varepsilon, \ell}, B_{\varepsilon, \ell}$, $1 \leq \ell \leq N$, such that

$$(12) \quad \int_{\mathbb{R}^d \times \mathbb{R}^d} |h^{(R)}(x, y) - h_\varepsilon(x, y)| d\lambda_d(x) d\lambda_d(y) \leq \varepsilon,$$

where

$$(13) \quad h_\varepsilon(x, y) = \sum_{\ell=1}^N c_\ell \mathbf{1}_{A_{\varepsilon, \ell}}(x) \mathbf{1}_{B_{\varepsilon, \ell}}(y).$$

Therefore, using stationarity and the fact that (X_i, X_j) has a density f_{j-i} which is bounded by a constant M independent of (i, j) ,

$$\begin{aligned} \mathbb{E} [|h^{(R)}(X_i, X_j) - h_\varepsilon(X_i, X_j)|] &= \mathbb{E} [|h^{(R)}(X_0, X_{j-i}) - h_\varepsilon(X_0, X_{j-i})|] \\ &= \int_{\mathbb{R}^d \times \mathbb{R}^d} |h^{(R)}(x, y) - h_\varepsilon(x, y)| f_{j-i}(x, y) d\lambda_d(x) d\lambda_d(y) \leq M\varepsilon \end{aligned}$$

and

$$(14) \quad \mathbb{E} \left[\left| \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h^{(R)}(x, y) dF(x) dF(y) - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h_\varepsilon(x, y) dF(x) dF(y) \right| \right] \\ \leq \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |h^{(R)}(x, y) - h_\varepsilon(x, y)| f_{X_0}(x) f_{X_0}(y) \leq \sup_{t \in \mathbb{R}^d} f_{X_0}(t) \varepsilon.$$

Consequently,

$$(15) \quad \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h^{(R)}(X_i, X_j) - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h^{(R)}(x, y) dF(x) dF(y) \right| \right] \\ \leq \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h_\varepsilon(X_i, X_j) - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h_\varepsilon(x, y) dF(x) dF(y) \right| \right] + \left(M + \sup_{t \in \mathbb{R}^d} f_{X_0}(t) \right) \varepsilon.$$

By the first step and the triangle inequality, we deduce that for each positive ε ,

$$(16) \quad \limsup_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h^{(R)}(X_i, X_j) - \int_{\mathbb{R}^d \times \mathbb{R}^d} h^{(R)}(x, y) dF(x) dF(y) \right| \right] \\ \leq \left(M + \sup_{t \in \mathbb{R}^d} f_{X_0}(t) \right) \varepsilon.$$

hence (3) holds with h replaced by h_R . Third step: by uniform integrability, for each positive ε , there exists δ such that for each A satisfying $\mathbb{P}(A) < \delta$, $\sup_{1 \leq i < j} \mathbb{E} [|h(X_i, X_j)| \mathbf{1}_A] < \varepsilon$. Let R be such that $\mathbb{P}(X_1 \notin [-R, R]^d) < \delta$, $\sup_{j \geq 2} \mathbb{E} [|h(X_1, X_j)| \mathbf{1}_{\{|h(X_1, X_j)| > R\}}] < \varepsilon$ and $\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |h(x, y) - h^{(R)}(x, y)| dF(x) dF(y) < \varepsilon$. Then for $h^{(R)}$ defined as in (11),

$$(17) \quad \mathbb{E} [|h(X_i, X_j) - h^{(R)}(X_i, X_j)|] \\ \leq \mathbb{E} \left[|h(X_i, X_j)| \left(\mathbf{1}_{\{X_i \notin [-R, R]^d\}} + \mathbf{1}_{\{X_j \notin [-R, R]^d\}} + \mathbf{1}_{\{|h(X_i, X_j)| > R\}} \right) \right] \leq 3\varepsilon$$

and it follows that

$$(18) \quad \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h(X_i, X_j) - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h(x, y) dF(x) dF(y) \right| \right] \\ \leq \mathbb{E} \left[\left| \frac{1}{\binom{n}{2}} \sum_{1 \leq i < j \leq n} h^{(R)}(X_i, X_j) - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} h^{(R)}(x, y) dF(x) dF(y) \right| \right] + 4\varepsilon,$$

and we conclude by the second step. This ends the proof of Theorem 3.1. \square

4. EXAMPLES OF FAILURE OF THE CONVERGENCE OF U -STATISTICS

Example 4.1 given in [1] shows that there exists a stationary ergodic sequence $(X_i)_{i \geq 1}$ and a bounded measurable function for which $\left(\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} h(X_i, X_j)\right)_{n \geq 2}$ converges, but not to the integral of $h(x, y)$ with respect to the product of the law of X_1 .

In a similar setting, we are able to formulate two examples, the first showing that the sequence $\left(\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} h(X_i, X_j)\right)_{n \geq 2}$ may fail to converge in probability even if $|h(X_i, X_j)|$ is bounded by 1 and the second one showing that a centered U -statistic $\left(\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} (h(X_i, X_j) - \mathbb{E}[h(X_i, X_j)])\right)_{n \geq 2}$ may also fail to converge in probability.

Suppose that $Tx = 2x \bmod 1$ is a transformation of the unit interval $[0, 1)$ equipped with the Borel sigma field \mathcal{B} and Lebesgue measure λ . We define $X_0(x) = x$, $X_k(x) = T^k x$ and $U: L^1 \rightarrow L^1$ by $UY = Y \circ T$, $Y \in L^1$.

4.1. Example 1: non-convergence of the U -statistics.

Proposition 4.1. *There exists a strictly stationary ergodic sequence $(X_i)_{i \geq 1}$ and a measurable function $h: \mathbb{R}^2 \rightarrow \mathbb{R}$ such that for each $i < j$, $\mathbb{E}[|h(X_i, X_j)|] \leq 1$ but the sequence $\left(\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} h(X_i, X_j)\right)_{n \geq 2}$ does not converge in probability.*

Proof. Let $(N_\ell)_{\ell \geq 1}$ and $(N'_\ell)_{\ell \geq 0}$ be sequences of positives integers such that $N'_0 = 1$ and for $\ell \geq 1$, $N_\ell < N'_\ell < N_{\ell+1}$ and

$$(19) \quad N'_\ell / N_\ell \geq \ell, \quad N_{\ell+1} / N'_\ell \rightarrow \infty,$$

$$(20) \quad I = \bigcup_{\ell \geq 0} I_\ell, \quad I_\ell := \{k \in \mathbb{N} : N'_\ell < k \leq N_{\ell+1}\}$$

$$(21) \quad G = \bigcup_{k \in I} \{(x, T^k x) : x \in [0, 1)\}$$

and for $x, y \in [0, 1)$,

$$h(x, y) = \mathbf{1}_G(x, y).$$

For $1 \leq i < j$ we then have

$$h(X_i, X_j) = \begin{cases} 1 & \text{if } j - i \in I, \\ 0 & \text{if } j - i \in \mathbb{N} \setminus I. \end{cases}$$

In particular, $\mathbb{E}[|h(X_i, X_j)|] \leq 1$. By (19) we have

$$(22) \quad \frac{1}{N_\ell(N_\ell - 1)} \sum_{1 \leq i < j \leq N_\ell} h(X_i, X_j) \rightarrow \frac{1}{2}, \quad \frac{1}{N'_\ell(N'_\ell - 1)} \sum_{1 \leq i < j \leq N'_\ell} h(X_i, X_j) \rightarrow 0.$$

Indeed, first observe that for each integer n ,

$$(23) \quad \sum_{1 \leq i < j \leq n} h(X_i, X_j) = \sum_{j=2}^n \sum_{i=1}^{j-1} \mathbf{1}_{j-i \in I}$$

$$(24) \quad = \sum_{j=2}^n \sum_{k=1}^{j-1} \mathbf{1}_{k \in I}$$

$$(25) \quad = \sum_{k=1}^{n-1} \sum_{j=k+1}^n \mathbf{1}_{k \in I}$$

$$(26) \quad \sum_{1 \leq i < j \leq n} h(X_i, X_j) = \sum_{k=1}^{n-1} (n-k) \mathbf{1}_{k \in I}$$

hence denoting $I_\ell = \{k \in \mathbb{N}, N'_\ell + 1 \leq k \leq N_{\ell+1}\}$, we get that for $\ell \geq 3$,

$$(27) \quad \frac{1}{N_\ell(N_\ell - 1)} \sum_{1 \leq i < j \leq N_\ell} h(X_i, X_j) = \frac{1}{N_\ell(N_\ell - 1)} \sum_{u=1}^{\ell-2} \sum_{k \in I_u} (N_\ell - k) + \frac{1}{N_\ell(N_\ell - 1)} \sum_{k \in I_{\ell-1}} (N_\ell - k) \\ =: A_\ell + B_\ell.$$

Note that bounding for $1 \leq u \leq \ell - 2$ the term $\sum_{k \in I_u} (N_\ell - k)$ by $N_\ell \text{Card}(I_u)$, and $\text{Card}(I_u)$ by $N_{u+1} - N_u$, we get

$$(28) \quad A_\ell \leq \frac{1}{N_\ell - 1} \sum_{u=1}^{\ell-2} (N_{u+1} - N_u) \leq \frac{N_{\ell-1} - N_1}{N_\ell - 1}$$

and using (19), we get $A_\ell \rightarrow 0$. Moreover,

$$(29) \quad B_\ell = \frac{1}{N_\ell(N_\ell - 1)} \sum_{k=N'_{\ell-1}+1}^{N_\ell} (N_\ell - k) = \frac{1}{N_\ell(N_\ell - 1)} \sum_{j=0}^{N_\ell - N'_{\ell-1} - 1} j \sim \frac{1}{2} \frac{(N_\ell - N'_{\ell-1} - 1)^2}{N_\ell^2}$$

hence $B_\ell \rightarrow 1/2$, which proves the first part of (22). The second one follows from the observation that $\{1, \dots, N'_\ell - 1\} \cap I_u$ is empty if $u \geq \ell - 1$, which gives in view of (26),

$$\begin{aligned} \frac{1}{N'_\ell(N'_\ell - 1)} \sum_{1 \leq i < j \leq N'_\ell} h(X_i, X_j) &= \frac{1}{N'_\ell(N'_\ell - 1)} \sum_{k=1}^{N'_\ell-1} (N'_\ell - k) \mathbf{1}_{k \in I} \\ &= \frac{1}{N'_\ell(N'_\ell - 1)} \sum_{u=1}^{\ell-2} \sum_{k \in I_u} (N'_\ell - k) \\ &\leq \frac{1}{N'_\ell} \sum_{u=1}^{\ell-2} (N_{u+1} - N'_u) \\ &\leq \frac{1}{N'_\ell} \sum_{u=1}^{\ell-2} (N_{u+1} - N_u) \leq \frac{N'_{\ell-1}}{N'_\ell}, \end{aligned}$$

where the second inequality follows from $N_u < N'_u$, and $N'_{\ell-1}/N'_\ell$ goes to 0 by (19). \square

4.2. Example 2: non-convergence of a centered U -statistic.

Proposition 4.2. *There exists a strictly stationary ergodic sequence $(X_i)_{i \geq 1}$ and a measurable function $h: \mathbb{R}^2 \rightarrow \mathbb{R}$ such that for each $i < j$, $\mathbb{E}[|h(X_i, X_j)|]$ is finite but the sequence $\left(\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} (h(X_i, X_j) - \mathbb{E}[h(X_i, X_j)])\right)_{n \geq 2}$ does not converge in probability.*

Note that in this example, $\sup_{j \geq 2} \mathbb{E}[|h(X_1, X_j)|]$ is infinite. Moreover, the sequence $\left(\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} (h(X_i, X_j) - \mathbb{E}[h(X_i, X_j)])\right)_{n \geq 2}$ converges in distribution to a centered non-degenerated Gaussian random variable.

Proof. We take the same probability space and transformation as above. For $k = 1, 2, \dots$ define

$$\bar{G}_k = T^{-k}([1/2, 1]), \quad G_k = \{(x, T^k x) : x \in \bar{G}_k\}, \quad h = \sum_{k=1}^{\infty} a_k \mathbf{1}_{G_k},$$

where

$$(30) \quad a_k = k^{3/2} - (k-1)^{3/2} \text{ for } k \geq 2 \text{ and } a_1 = 1.$$

For $1 \leq i < j$ we thus have

$$h(X_i, X_j) = a_{j-i} T^{-j}([1/2, 1]) = a_{j-i} U^j f, \quad f = \mathbf{1}_{[1/2, 1]},$$

hence $\mathbb{E}[h(X_i, X_j)] = a_{j-i}/2$ and

$$(31) \quad \sum_{1 \leq i < j \leq n} h(X_i, X_j) = \sum_{i=1}^{n-1} \sum_{j=i+1}^n a_{j-i} U^j f = \sum_{j=2}^n \sum_{i=1}^{j-1} a_{j-i} U^j f = \sum_{j=2}^n (j-1)^{3/2} U^j f$$

where $f = \mathbf{1}_{[1/2, 1]}$. In order to have a better understanding of $U^j f$, we introduce the intervals

$$(32) \quad I_{j, \ell} = \left[\frac{\ell-1}{2^j}, \frac{\ell}{2^j} \right), \quad j \geq 1, 1 \leq \ell \leq 2^j.$$

Lemma 4.3. *The sequence $(U^j(f - 1/2))_{j \geq 1}$ is a martingale difference sequence with respect to the filtration $(\mathcal{F}_j)_{j \geq 0}$, where $\mathcal{F}_j = \sigma(I_{j, \ell}, 1 \leq \ell \leq 2^j)$ and $\mathcal{F}_0 = \{\emptyset, \Omega\}$.*

Proof. We show by induction on $j \geq 1$ that

$$(33) \quad f \circ T^j(x) = \mathbf{1}_{\bigcup_{\ell=1}^{2^{j-1}} I_{j, 2\ell}}(x).$$

For $j = 1$, notice that if $x \in [0, 1/2)$, then $f \circ T(x) = f(2x) = \mathbf{1}_{[1/2, 1]}(2x) = \mathbf{1}_{[1/4, 1/2)}(x)$ and if $x \in [1/2, 1)$, then $f \circ T(x) = f(2x-1) = \mathbf{1}_{[1/2, 1]}(2x-1) = \mathbf{1}_{[3/2, 2)}(2x) = \mathbf{1}_{[3/4, 1)}(x)$ hence for each $x \in [0, 1)$, $f \circ T(x) = \mathbf{1}_{[1/4, 1/2)}(x) + \mathbf{1}_{[3/4, 1)}(x)$.

Assume now that (33) holds true for some $j \geq 1$ and let us show that

$$(34) \quad f \circ T^{j+1}(x) = \mathbf{1}_{\bigcup_{\ell=1}^{2^j} I_{j+1, 2\ell}}(x).$$

By (33) with x replaced by Tx , we derive that

$$(35) \quad f \circ T^{j+1}(x) = \mathbf{1}_{\bigcup_{\ell=1}^{2^{j-1}} I_{j, 2\ell}}(Tx).$$

If $x \in [0, 1/2)$, then

$$\mathbf{1}_{\bigcup_{\ell=1}^{2^{j-1}} I_{j, 2\ell}}(Tx) = \mathbf{1}_{\bigcup_{\ell=1}^{2^{j-1}} I_{j, 2\ell}}(2x) = \mathbf{1}_{\bigcup_{\ell=1}^{2^{j-1}} I_{j+1, 2\ell}}(x)$$

and (34) holds, and if $x \in [1/2, 1)$, then

$$\mathbf{1}_{\bigcup_{j=1}^{2^j-1} I_{j,2\ell}}(Tx) = \mathbf{1}_{\bigcup_{\ell=1}^{2^{j-1}} I_{j,2\ell}}(2x-1) = \sum_{\ell=1}^{2^{j-1}} \mathbf{1}_{I_{j,2\ell}}(2x-1) = \sum_{\ell=1}^{2^{j-1}} \mathbf{1}_{I_{j,2\ell+2^j}}(x) = \mathbf{1}_{\bigcup_{\ell=2^{j-1}+1}^{2^j} I_{j+1,2\ell}}(x)$$

hence (34) also holds.

By (33), it is clear that $U^j f$ is \mathcal{F}_j -measurable. Moreover,

$$(36) \quad \mathbb{E}[U^{j+1}(f - 1/2) \mid \mathcal{F}_j] = \sum_{\ell=1}^{2^j} \mathbb{E}[\mathbf{1}_{I_{j+1,2\ell}} - 1/2 \mid \mathcal{F}_j] = 0.$$

□

Notice that $\binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} (h(X_i, X_j) - \mathbb{E}[h(X_i, X_j)]) = \sum_{j=1}^n d_{n,j}$, where

$$(37) \quad d_{n,j} = \binom{n}{2}^{-1} (j-1)^{3/2} \left(U^j f - \frac{1}{2} \right), j \geq 2, d_{n,1} = 0.$$

Then $(d_{n,j})_{j \geq 1}$ is a martingale difference sequence with respect to the filtration $(\mathcal{F}_j)_{j \geq 0}$ given as in Lemma 4.3.

Recall that by [7], if $(d_{n,j})_{n \geq 1, 1 \leq j \leq n}$ is an array of martingale differences, such that

$$(38) \quad \max_{1 \leq j \leq n} |d_{n,j}| \rightarrow 0 \text{ in probability,}$$

$$(39) \quad \text{there exists } M > 0 \text{ such that } \sup_{n \geq 1} \max_{1 \leq j \leq n} \mathbb{E}[d_{n,j}^2] \leq M \text{ and}$$

$$(40) \quad \sum_{j=1}^n d_{n,j}^2 \rightarrow \sigma^2 \text{ in probability,}$$

then $\sum_{j=1}^n d_{n,j}$ converges in distribution to a centered normal distribution with variance σ^2 .

Noticing that $|U^j f(x) - 1/2| = 1/2$, we can see that (38) and (39) are satisfied as well as (40) with $\sigma^2 = 1/4$.

Letting $Y_n = \binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} (h(X_i, X_j) - \mathbb{E}[h(X_i, X_j)])$, we thus get that $Y_n \rightarrow N(0, 1/4)$. Expressing $Y_{2n} - Y_n$ as a sum of a martingale difference array, the same argument as above gives that $Y_{2n} - Y_n$ converges in distribution to a non-degenerated normal random variable hence $(Y_n)_{n \geq 1}$ cannot converge in probability. □

REFERENCES

- [1] J. Aaronson, R. Burton, H. Dehling, D. Gilat, T. Hill, and B. Weiss, *Strong laws for L - and U -statistics*, Trans. Amer. Math. Soc. **348** (1996), no. 7, 2845–2866. MR 1363941
- [2] M. A. Arcones, *The law of large numbers for U -statistics under absolute regularity*, Electron. Comm. Probab. **3** (1998), 13–19. MR 1624866
- [3] P. Billingsley, *Convergence of probability measures*, John Wiley & Sons, Inc., New York-London-Sydney, 1968. MR 0233396
- [4] S. Borovkova, R. Burton, and H. Dehling, *Consistency of the Takens estimator for the correlation dimension*, Ann. Appl. Probab. **9** (1999), no. 2, 376–390. MR 1687339
- [5] ———, *From dimension estimation to asymptotics of dependent U -statistics*, Limit theorems in probability and statistics, Vol. I (Balatonlelle, 1999), János Bolyai Math. Soc., Budapest, 2002, pp. 201–234. MR 1979966

- [6] W. Hoeffding, *The strong law of large numbers for U -statistics.*, University of North Carolina Institute of Statistics Mimeograph Series **302** (1961).
- [7] D. L. McLeish, *Dependent central limit theorems and invariance principles*, Ann. Probability **2** (1974), 620–628. MR 358933
- [8] V. S. Varadarajan, *Weak convergence of measures on separable metric spaces*, Sankhyā **19** (1958), 15–22. MR 94838