



**HAL**  
open science

# Gradient estimates for a nonlinear elliptic equation on a smooth metric measure space

Xiaoshan Wang, Linfen Cao

► **To cite this version:**

Xiaoshan Wang, Linfen Cao. Gradient estimates for a nonlinear elliptic equation on a smooth metric measure space. 2023. hal-03984127v1

**HAL Id: hal-03984127**

**<https://hal.science/hal-03984127v1>**

Preprint submitted on 12 Feb 2023 (v1), last revised 14 Feb 2023 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# GRADIENT ESTIMATES FOR A NONLINEAR ELLIPTIC EQUATION ON A SMOOTH METRIC MEASURE SPACE

XIAOSHAN WANG<sup>A</sup>, LINFEN CAO<sup>B</sup>

ABSTRACT. Let  $(M, g, e^{-f} dv)$  be a smooth metric measure space. We consider local gradient estimates for positive solutions to the following elliptic equation

$$\Delta_f u + au \log u + bu = 0$$

where  $a, b$  are two real constants and  $f$  be a smooth function defined on  $M$ . As an application, we obtain a Liouville type result for such equation in the case  $a < 0$  under the  $m$ -dimensions Bakry-Émery Ricci curvature.

## 1. INTRODUCTION

In this paper, we study the local gradient estimate for the positive solution to the following weighted nonlinear elliptic equation

$$\Delta_f u + au \log u + bu = 0 \tag{1.1}$$

on a smooth metric measure space  $(M, g, e^{-f} dv)$ , where  $a, b$  are two real constants and  $f$  be a smooth function defined on  $M$ . The motivation to study (1.1) comes from understanding the Ricci flow. Moreover, the (1.1) is closely related to the famous Gross Logarithmic Sobolev inequality, see [2]. It is well known that Yau has proved in [14] that every positive or bounded harmonic function is constant if  $M$  has nonnegative Ricci curvature by establishing gradient estimates for the solutions to Laplacian equation, see also [6, 8, 11].

It is natural to consider similar Liouville type results for positive solutions to the nonlinear elliptic equation (1.1). In [10], Qian considered positive solutions to

$$\Delta u + au \log u = 0 \tag{1.2}$$

and proved the following

**Theorem 1.1.** *(B. Qian) Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold with the Ricci curvature  $\text{Ric}(B(x, R)) \geq -K$ , where*

---

2010 *Mathematics Subject Classification.* Primary 58J35, Secondary 35B45.

*Key words and phrases.* gradient estimates, nonlinear equation,  $m$ -dimensions Bakry-Émery Ricci curvature, Liouville type property.

The second author is supported by NSFC (No.11671121, 11971153).

Corresponding author: linfencao6@163.com (Linfen Cao).

$K \geq 0$  is a constant. Let  $u$  be a positive solution to (1.2) on  $B(x, R)$ , then for any  $\alpha > 0$ ,

$$\sup_{y \in B(x, \frac{R}{2})} \frac{|\nabla u|}{u} \leq C(n) \sqrt{(1 + \alpha) \left( (a + K) + \frac{(1 + \alpha)}{R^2} \right) + \sqrt{\frac{1 + \alpha}{\alpha}} |a|} L(x, R), \quad (1.3)$$

where  $L(x, R) = \sup_{y \in B(x, R)} |\log u| < \infty$  and  $C(n)$  is a constant depending only on the dimension  $n$ .

In particular, by letting  $R \rightarrow \infty$  in (1.3), we obtain the following gradient estimates on complete noncompact Riemannian manifolds:

$$\frac{|\nabla u|}{u} \leq C(n) \sqrt{(1 + \alpha)(a + K) + \sqrt{\frac{1 + \alpha}{\alpha}} |a|} L, \quad (1.4)$$

where  $L = \sup_M |\log u|$ .

*Remark 1.1.* Clearly, from 1.4, it is easy to see that if  $a + K < 0$  and  $a < 0$ , then any bounded positive solution to (1.2) must be a constant  $u \equiv 1$ . On the other hand, in [5], Huang and Ma also obtained the similar Liouville type result by a different method.

Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold and  $f$  be a smooth function defined on  $M$ . In general, the triple  $(M, g, e^{-f} dv)$  is called a smooth metric measure space. The  $f$ -Laplacian operator is defined by

$$\Delta_f = e^f \operatorname{div}(e^{-f} \nabla) = \Delta - \nabla f \nabla,$$

which is symmetric in  $L^2(M, g, e^{-f} dv)$ .

It is well-known that the  $m$ -dimensions Bakry-Émery Ricci curvature associated with the  $f$ -Laplacian is defined by (see [3, 4, 13])

$$\operatorname{Ric}_f^m = \operatorname{Ric} + \nabla^2 f - \frac{df \otimes df}{m - n},$$

where  $m \geq n$  is a constant and  $m = n$  if and only if  $f$  is a constant. Define

$$\operatorname{Ric}_f = \operatorname{Ric} + \nabla^2 f.$$

Then  $\operatorname{Ric}_f$  can be seen as  $\infty$ -dimensions Bakry-Émery Ricci curvature. Recently, the  $\infty$ -dimensions Bakry-Émery Ricci curvature has become an important object of study in Riemannian geometry. The equation  $\operatorname{Ric}_f = \rho \langle \cdot, \cdot \rangle$  for some constant  $\rho$  is just the gradient Ricci soliton equation, which plays an important role in the study of Ricci flow (see [1]).

The aim of this paper is to generalize the results of Qian in [10] to the weighted nonlinear elliptic equation (1.1) under the assumption that the  $m$ -dimensions Bakry-Émery Ricci curvature is bounded from blow. Our main results are as follows:

**Theorem 1.2.** *Let  $(M, g, e^{-f} dv)$  be an  $n$ -dimensional complete smooth metric measure space with  $m$ -dimensions Bakry-Émery Ricci curvature*

$Ric_f^m(B(x, R)) \geq -K$ , where  $K \geq 0$  is a constant. Let  $u$  be a positive solution to the nonlinear equation (1.1) on  $B(x, R)$ , then there exists a constant  $C = C(m)$ , such that

$$\sup_{y \in B(x, \frac{R}{2})} \frac{|\nabla u|}{u} \leq C \sqrt{a + K + \frac{1}{R^2} + |b| + |a|L}. \quad (1.5)$$

where  $L(x, R) := \sup_{y \in B(x, R)} |\log u| < \infty$ .

In particular, by letting  $R \rightarrow \infty$  in (1.5), we obtain the following gradient estimates on complete noncompact Riemannian manifolds:

$$\frac{|\nabla u|}{u} \leq C \sqrt{a + K + |b| + |a|L}, \quad (1.6)$$

where  $L = \sup_M |\log u|$ .

From (1.6), it is easy to obtain the following results:

**Corollary 1.3.** *Let  $(M, g, e^{-f} dv)$  be an  $n$ -dimensional complete smooth metric measure space with  $Ric_f^m \geq -K$ . If  $u$  is a bounded positive solution to (1.1) with  $a + K < 0$  and  $a < 0$ , then  $u \equiv 1$ . Furthermore, if  $Ric_f^m \geq 0$  and  $a \leq 0$ , then any bounded positive solution to (1.1) must be  $u \equiv 1$ .*

*Remark 1.2.* When  $m = n$  in Theorem 1.2, we have  $Ric_f^m = Ric$  and  $\Delta_f = \Delta$ . Hence, in this case, our Theorem 1.2 becomes Theorem 1.1 of Qian. That is, our results of this paper generalize those of Qian in [10].

## 2. PROOF OF THEOREM 1.2

**Lemma 2.1.** *Let  $u$  be a bounded positive solution to the nonlinear equation (1.1), then we have,*

$$\begin{aligned} |\nabla u| \Delta_f |\nabla u| &\geq \frac{|\nabla(|\nabla u|)|^2}{m} - (au \log u + bu)^2 + Ric_f^m(\nabla u, \nabla u) \\ &\quad - (a + a \log u + b)|\nabla u|^2. \end{aligned} \quad (2.1)$$

*Proof.* Since we have the Bochner-weitzenböck formula with respect to  $f$ -Laplacian, for any  $u \in C^2(M)$ , we have

$$\frac{1}{2} \Delta_f |\nabla u|^2 = |\nabla^2 u|^2 + Ric_f(\nabla u, \nabla u) + (\nabla u, \nabla \Delta_f u).$$

On the other hand,

$$\Delta_f |\nabla u|^2 = 2|\nabla u| \Delta_f |\nabla u| + 2|\nabla(|\nabla u|)|^2,$$

hence

$$|\nabla u| \Delta_f |\nabla u| = |\nabla^2 u|^2 + Ric_f(\nabla u, \nabla u) + (\nabla u, \nabla \Delta_f u) - |\nabla(|\nabla u|)|^2.$$

Since  $u$  is a solution to (1.1), we obtain

$$|\nabla u| \Delta_f |\nabla u| = |\nabla^2 u|^2 + Ric_f(\nabla u, \nabla u) - (a + a \log u + b)|\nabla u|^2 - |\nabla(|\nabla u|)|^2.$$

If we consider a local normal chart at  $x$  in which  $u_1(x) = |\nabla u|(x)$  and  $u_j(x) = 0$  for  $j \geq 2$ , then  $\nabla_i(|\nabla u|) = u_{1i}$ , hence  $|\nabla(|\nabla u|)|^2 = \sum_{i \geq 2} u_{1i}^2$ . Since  $u$  is a solution to (1.1), in the above local chart we have at  $x$

$$\sum_{i \geq 2} u_{ii} = -u_{11} - au \log u - bu + \nabla f \cdot \nabla u.$$

Therefore,

$$\begin{aligned} |\nabla^2 u|^2 - |\nabla(|\nabla u|)|^2 &= \sum_{i \geq 1, j \geq 1} u_{ij}^2 - \sum_{j \geq 1} u_{1j}^2 = \sum_{i \geq 2, j \geq 1} u_{ij}^2 \geq \sum_{i \geq 2} u_{i1}^2 + \sum_{i \geq 2} u_{ii}^2 \\ &\geq \sum_{i \geq 2} u_{i1}^2 + \frac{1}{n-1} \left( \sum_{i \geq 2} u_{ii} \right)^2 \\ &= \sum_{i \geq 2} u_{i1}^2 + \frac{1}{n-1} (-u_{11} - au \log u - bu + \nabla f \cdot \nabla u)^2 \\ &\geq \frac{1}{(n-1)(1+\alpha)} \sum_{i \geq 1} u_{i1}^2 - \frac{1}{(n-1)\alpha} (au \log u + bu - \nabla f \cdot \nabla u)^2 \\ &\geq -\frac{1}{(n-1)\alpha} \left[ \left(1 + \frac{1}{\epsilon}\right) (au \log u + bu)^2 + (1+\epsilon) (\nabla f \cdot \nabla u)^2 \right] \\ &\quad + \frac{1}{(n-1)(1+\alpha)} \sum_{i \geq 1} u_{i1}^2 \\ &= \frac{1}{m} \sum_{i \geq 1} u_{i1}^2 - (au \log u + bu)^2 - \frac{(\nabla f \cdot \nabla u)^2}{m-n}, \end{aligned}$$

where we use the elementary inequality (see [12])  $(a+b)^2 \geq \frac{1}{1+\alpha} a^2 - \frac{1}{\alpha} b^2$  and  $(a+b)^2 \leq (1+\epsilon)a^2 + (1+\frac{1}{\epsilon})b^2$  which holds for any  $\alpha > 0$ ,  $\epsilon > 0$ . The last equality we choose  $\alpha = \frac{m-n+1}{n-1}$  and  $\epsilon = \frac{1}{m-n}$ .

Hence

$$\begin{aligned} |\nabla u| \Delta_f |\nabla u| &\geq \frac{|\nabla(|\nabla u|)|^2}{m} - (au \log u + bu)^2 - \frac{(\nabla f \cdot \nabla u)^2}{m-n} \\ &\quad + Ric_f(\nabla u, \nabla u) - (a + a \log u + b) |\nabla u|^2 \\ &= \frac{|\nabla(|\nabla u|)|^2}{m} - (au \log u + bu)^2 + \left( Ric_f - \frac{df \otimes df}{m-n} \right) u_i u_j \quad (2.2) \\ &\quad - (a + a \log u + b) |\nabla u|^2 \\ &= \frac{|\nabla(|\nabla u|)|^2}{m} - (au \log u + bu)^2 + Ric_f^m(\nabla u, \nabla u) \\ &\quad - (a + a \log u + b) |\nabla u|^2. \end{aligned}$$

This completes the prove of Lemma 2.1. □

### Proof of Theorem 1.2.

*Proof.* Denote  $\psi = |\nabla \log u| = \frac{|\nabla u|}{u}$ , i.e.,

$$|\nabla u| = \psi u.$$

Direct computation gives

$$\nabla \psi = \frac{\nabla(|\nabla u|)}{u} - \frac{|\nabla u| \nabla u}{u^2}. \quad (2.3)$$

At the point where  $\nabla u \neq 0$ , we have

$$\begin{aligned} \Delta_f |\nabla u| &= \Delta_f (\psi u) = u \Delta_f \psi + \psi \Delta_f u + 2 \nabla \psi \cdot \nabla u \\ &= u \Delta_f \psi - \psi (a u \log u + b u) + 2 \nabla \psi \cdot \nabla u. \end{aligned}$$

This yields

$$\begin{aligned} \Delta_f \psi &= \frac{\Delta_f |\nabla u|}{u} + \psi (a \log u + b) - 2 \frac{\nabla \psi \cdot \nabla u}{u} \\ &= \frac{|\nabla u| \Delta_f |\nabla u|}{|\nabla u| u} + \psi (a \log u + b) - 2 \frac{\nabla \psi \cdot \nabla u}{u}. \end{aligned}$$

By Lemma 2.1, we can derive,

$$\begin{aligned} \Delta_f \psi &\geq \frac{\frac{|\nabla(|\nabla u|)|^2}{m} - (a u \log u + b u)^2 + Ric_f^m(\nabla u, \nabla u) - (a + a \log u + b) |\nabla u|^2}{|\nabla u| u} \\ &\quad + \psi (a \log u + b) - 2 \frac{\nabla \psi \cdot \nabla u}{u} \\ &\geq \frac{|\nabla(|\nabla u|)|^2}{m |\nabla u| u} - \frac{(a \log u + b)^2}{\psi} - (a + K) \psi - 2 \frac{\nabla \psi \cdot \nabla u}{u}. \end{aligned} \quad (2.4)$$

For any  $\delta > 0$ , by (2.3)

$$\begin{aligned} 2 \frac{\nabla \psi \cdot \nabla u}{u} &= (2 - \delta) \frac{\nabla \psi \cdot \nabla u}{u} + \delta \frac{\nabla \psi \cdot \nabla u}{u} \\ &= (2 - \delta) \frac{\nabla \psi \cdot \nabla u}{u} + \delta \frac{\nabla u}{u} \left( \frac{\nabla(|\nabla u|)}{u} - \frac{|\nabla u| \nabla u}{u^2} \right) \\ &= (2 - \delta) \frac{\nabla \psi \cdot \nabla u}{u} + \delta \frac{\nabla(|\nabla u|) \cdot \nabla u}{u^2} - \delta \psi^3 \\ &\leq (2 - \delta) \frac{\nabla \psi \cdot \nabla u}{u} + \delta \frac{|\nabla(|\nabla u|)| \cdot |\nabla u|}{u^2} - \delta \psi^3 \\ &\leq (2 - \delta) \frac{\nabla \psi \cdot \nabla u}{u} + \frac{\delta}{2} \left( \frac{|\nabla(|\nabla u|)|^2}{|\nabla u| u} + \frac{|\nabla u|^3}{u^3} \right) - \delta \psi^3 \\ &= (2 - \delta) \frac{\nabla \psi \cdot \nabla u}{u} + \frac{\delta}{2} \frac{|\nabla(|\nabla u|)|^2}{|\nabla u| u} - \frac{\delta}{2} \psi^3. \end{aligned}$$

Choosing  $\delta = \frac{2}{m}$  and substituting into (2.4), we obtain

$$\begin{aligned} \Delta_f \psi \geq & -(a + K)\psi - \left(2 - \frac{2}{m}\right) \frac{\nabla \psi \cdot \nabla u}{u} + \frac{\psi^3}{m} \\ & - \frac{(a \log u + b)^2}{\psi}. \end{aligned} \quad (2.5)$$

Now we define

$$F(y) := (R^2 - d^2(x, y)) \frac{|\nabla u|(y)}{u(y)} = (R^2 - d^2)\psi(y),$$

and

$$\psi(y) = \frac{|\nabla u|(y)}{u(y)}, \quad y \in B(x, R).$$

Since  $F|_{\partial B(x, R)} = 0$ , if  $\nabla u = 0$ , then  $F$  can only achieve its maximum at some point  $x_0 \in B(x, R)$ , if  $|\nabla u|(x_0) = 0$ , the desired result holds. Then, without loss of generality, we can suppose  $|\nabla u|(x_0) \neq 0$ . Assume  $x_0 \notin \text{cut}(x)$ , by the maximum principle we have  $\Delta_f F(x_0) \leq 0$  and  $\nabla F(x_0) = 0$ . It yields, at  $x_0$ ,

$$\nabla F = -\psi \nabla d^2 + (R^2 - d^2) \nabla \psi = 0.$$

It holds,

$$\frac{\nabla \psi}{\psi} = \frac{\nabla d^2}{R^2 - d^2} = \frac{2d \nabla d}{R^2 - d^2} \quad (2.6)$$

and

$$\Delta_f F = \Delta_f((R^2 - d^2)\psi) = (R^2 - d^2)\Delta_f \psi - \psi \Delta_f d^2 - 2\nabla d^2 \nabla \psi \leq 0. \quad (2.7)$$

Hence, dividing by  $(R^2 - d^2)\psi$  to both sides of (2.7) and combining (2.6), we have at  $x_0$ ,

$$0 \geq \frac{\Delta_f \psi}{\psi} - \frac{\Delta_f d^2}{R^2 - d^2} - \frac{8d^2}{(R^2 - d^2)^2}.$$

By the  $f$ -Laplacian comparison theorem in [12] (see also [7] or [9]), we have

$$\Delta_f d^2 \leq C\sqrt{K}d \coth(\sqrt{K}d) \leq C\sqrt{K}d,$$

where  $C$  only depends on  $m$ . Together with (2.5), we have at  $x_0$ ,

$$\begin{aligned} 0 \geq & -(a + K) - \left(2 - \frac{2}{m}\right) \frac{\nabla \psi \cdot \nabla u}{\psi u} + \frac{\psi^2}{m} \\ & - \frac{(a \log u + b)^2}{\psi^2} - \frac{C\sqrt{K}d}{R^2 - d^2} - \frac{8d^2}{(R^2 - d^2)^2}. \end{aligned} \quad (2.8)$$

By (2.6),

$$\frac{\nabla \psi \cdot \nabla u}{\psi u} = \frac{2d}{R^2 - d^2} \frac{\nabla d \cdot \nabla u}{u} \leq \frac{2d\psi}{R^2 - d^2},$$

then multiplying both sides of (2.8) by  $\psi^2(R^2 - d^2)^4$ , combining with  $\psi = \frac{F}{R^2 - d^2}$ , we can derive

$$\begin{aligned} 0 &\geq \frac{1}{m}F^4 - \frac{4(m-1)}{m}dF^3 - (R^2 - d^2)^4(a \log u + b)^2 \\ &\quad - ((a+K)(R^2 - d^2)^2 + 8d^2 + C\sqrt{K}d(R^2 - d^2))F^2 \\ &\geq \frac{1}{m}F^4 - 4RF^3 - ((a+K)R^2 + C\sqrt{K}R + 8)R^2F^2 \\ &\quad - R^8(a \log u + b)^2. \end{aligned}$$

Since  $\frac{1}{2m}F^4 - 4RF^3 \geq -8mR^2F^2$ , it holds

$$\begin{aligned} \frac{1}{2m}F^4 - ((a+K)R^2 + C\sqrt{K}R + C)R^2F^2 \\ - R^8(a \log u + b)^2 \leq 0. \end{aligned}$$

It follows

$$\begin{aligned} \sup_{y \in B(x, R)} (R^2 - d^2(x, y)) |\nabla \log u| &\leq F(x_0) \\ &\leq \sqrt{2m((a+K)R^2 + C\sqrt{K}R + C)R^2 + \sqrt{2m}(|b| + |a|L)R^4}. \end{aligned}$$

Restricting on the ball  $B(x, \frac{R}{2})$ , we have

$$\begin{aligned} \sup_{y \in B(x, \frac{R}{2})} \frac{3R^2}{4} |\nabla \log u| &\leq \sup_{y \in B(x, \frac{R}{2})} (R^2 - d^2(x, y)) |\nabla \log u| \leq F(x_0) \\ &\leq \sqrt{2m((a+K)R^2 + C\sqrt{K}R + C)R^2 + \sqrt{2m}(|b| + |a|L)R^4}. \end{aligned}$$

Therefore, we can derive

$$\sup_{y \in B(x, \frac{R}{2})} \frac{|\nabla u|}{u} \leq \sqrt{4m(a+K + C\frac{\sqrt{K}}{R} + \frac{C}{R^2}) + \sqrt{8m}(|b| + |a|L)}.$$

Then use the Cauchy-Schwarz inequality, we obtain

$$\sup_{y \in B(x, \frac{R}{2})} \frac{|\nabla u|}{u} \leq C\sqrt{a+K + \frac{1}{R^2} + |b| + |a|L}.$$

Now let  $R \rightarrow \infty$ , this yields, for any  $x \in M$ ,

$$\psi(x) \leq \psi(x_0) \leq C\sqrt{a+K + |b| + |a|L}.$$

This completes the prove of theorem 1.2.  $\square$



## REFERENCES

- [1] H.-D. Cao, Recent progress on Ricci solitons, Recent advances in geometric analysis, Adv. Lect. Math. 11, Int. Press, Somerville, MA, (2010), 1-38.
- [2] L. Gross, Logarithmic Sobolev inequality and contractivity properties of semigroups, in: Dirichlet Forms, Varenna, 1992, in: Lecture Notes in Math. vol. 1563, Springer, Berlin, 1993, pp. 54-88.
- [3] G. Y. Huang, B. Q. Ma, Gradient estimates for a nonlinear parabolic equation on Riemannian manifolds, Arch. Math. 94 (2010), 265-275.
- [4] G. Y. Huang, H. Z. Li, Gradient estimates and entropy formulae of porous medium and fast diffusion equations for the Witten laplacian, Pacific J. Math. 268 (2014), 47-78.
- [5] G. Y. Huang, B. Q. Ma, Gradient estimates and Liouville type theorems for a nonlinear elliptic equation, Arch. Math. 105 (2015), 491-499.
- [6] P. Li, S. T. Yau, On the parabolic kernel of the Schrödinger operator, Acta Math. 156 (1986), 153-201.
- [7] J. Li, Gradient estimates and Harnack inequalities for nonlinear parabolic and nonlinear elliptic equations on Riemannian manifold, J. Funct. Anal. 100 (1991), 233-256.
- [8] X. D. Li, Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds, J. Math. Pures Appl. 84 (2005), 1295-1361.
- [9] Y. Li, Li-Yau-Hamilton estimates and Bakry-Emery Ricci curvature, Nonlinear Anal. 113 (2015), 1-32.
- [10] B. Qian, Yau's gradient estimates for a nonlinear elliptic equation, Arch. Math. 108 (2017), 427-435.
- [11] R. Schoen, S. T. Yau, Lectures on Differential Geometry, International Press, Cambridge, MA, 1994.
- [12] J. Y. Wu, Upper bounds on the first eigenvalue for a diffusion operator via Bakry-Émery Ricci curvature, J. Math. Anal. Appl. 361 (2010), 10-18.
- [13] J. Y. Wu, Elliptic gradient estimates for a nonlinear heat equation and applications, Nonlinear Anal. 151 (2017), 1-17.
- [14] S. T. Yau, Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry, Indiana Math. J. 25 (1976), 659-670.

HENAN ENGINEERING LABORATORY FOR BIG DATA STATISTICAL ANALYSIS AND OPTIMAL CONTROL, COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, HENAN NORMAL UNIVERSITY, XINXIANG, 453007 CHINA

*E-mail address:* xswang2017@126.com (Xiaoshan Wang<sup>A</sup>), linfencao6@163.com (Linfen Cao<sup>B</sup>)

A. DEPARTMENT OF MATHEMATICS, LUOYANG NORMAL UNIVERSITY, LUOYANG 471934, CHINA;

B. COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, HENAN NORMAL UNIVERSITY, XINXIANG, 453007 HENAN, CHINA