



HAL
open science

N-Boc and N-CBz Ethyl Oxamates: New Gabriel Reagents

Fabienne Berrée, Jean-Pierre Bazureau, Gwendal Michelot, Maurice Le Cone

► **To cite this version:**

Fabienne Berrée, Jean-Pierre Bazureau, Gwendal Michelot, Maurice Le Cone. N-Boc and N-CBz Ethyl Oxamates: New Gabriel Reagents. *Synthetic Communications*, 1999, 29 (15), pp.2685-2693. 10.1080/00397919908086430 . hal-03985910

HAL Id: hal-03985910

<https://hal.science/hal-03985910>

Submitted on 13 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


N-BOC AND N-CBZ ETHYL OXAMATES : NEW GABRIEL REAGENTS.

Fabienne Berrée,* Jean-Pierre Bazureau,
Gwendal Michelot and Maurice Le Corre

Laboratoire de Synthèse Organique, Associé au CNRS.
Avenue du Général Leclerc, 35042 Rennes, France.


Abstract : N-Boc and N-CBz ethyl oxamates can be directly coupled with various halides under Gabriel conditions which afford the corresponding N-protected amines after mild deprotection with LiOH.

We recently reported the use of the N-Boc ethyl oxamate **1a** in Mitsunobu reactions.¹ The condensation products **2** could be easily deprotected in N-Boc amines **3** by using LiOH (Scheme 1).



Scheme 1


Despite of the high sensitivity of ethoxalyl group to weak nucleophiles, we observed that it was also possible to generate the potassium salt by treatment of **1a** with t-BuOK. Then this salt was used in Gabriel reactions^{2,3} (Scheme 2).



Scheme 2

The reactions were carried out in DMF at 60°C during 1 hour and led to the N-alkylated ethyl oxamates **2** in good yields (Table 1). For the reaction with 1,4-dibromobutane, we have observed that two equivalents of the salt have reacted with one equivalent of 1,4-dibromobutane to give the bis-condensed product (entry e), but the use of an excess of bromide derivative allowed to obtain essentially the single condensation product **2e**.

We also extended this methodology to the preparation of the N-CBz ethyl oxamate **1b**. This product can be selectively deprotected by catalytic hydrogenation. It was prepared in quantitative yield by reaction of benzyl alcohol with ethyloxalyl isocyanate⁴ in toluene under anhydrous conditions (Scheme 3). The compound **1b** showed a similar reactivity to **1a** when it was coupled with 1-bromooctane (entry f).




Scheme 3

Table 1. Conversion of bromide derivatives to oxamates **2**.

entry	1	RX	2 Yield ^a (%)
a	1a	EtO ₂ CCH ₂ Br	89
b		PhCH ₂ Br	83
c		CH ₃ (CH ₂) ₇ Br	88
d		CH ₂ =CHCH ₂ Br	94
e		Br(CH ₂) ₄ Br (1 equiv.)	45 ^b
		Br(CH ₂) ₄ Br (2 equiv.)	58 ^c
f	1b	CH ₃ (CH ₂) ₇ Br	73


(a) Yield of isolated product after purification by column chromatography. (b) With 23% of dialkylated product. (c) With 5% of dialkylated product.

As we have previously reported,¹ access to N-protected amines from oxamates **2** was particularly easy. For example, treatment of **2c,e** with LiOH at room temperature for 3 hours gave the known carbamates **3c⁵,e⁶** (91,99% yield; Scheme 4).




Scheme 4

Access to free amines from carbamates **3**, by action of trifluoroacetic acid (R' = t-Bu) or catalytic hydrogenation (R' = CH₂Ph), is well known. Alternatively, deprotection of carbamates **2** gave N-alkyl oxamates **4** (Scheme 5).



Scheme 5

Successive treatment of bromoalkyl compound **2e** with LiOH and t-BuOK was an interesting process for the synthesis of the N-Boc pyrrolidine **5⁷** (68% overall yield; Scheme 6).



Scheme 6

In summary, the results described above demonstrate that N-protected ethyl oxamates **1** are excellent nitrogen nucleophiles in Gabriel reactions. Particularly mild conditions can be used to finally obtain the protected N-Boc and N-CBz amines.

Experimental

^1H NMR spectra (300 MHz) and ^{13}C NMR (75 MHz) were recorded on a Bruker AC 300 spectrometer, in CDCl_3 solutions with Me_4Si as internal reference. Chemical shifts are expressed in ppm and coupling constants J are given in Hz. DMF was distilled over MgSO_4 . Elemental analyses were performed by the Microanalytical Laboratory of the Centre National de la Recherche Scientifique, Gif sur Yvette. Mass spectra were taken on a Varian MAT 311 spectrometer.

Ethyloxalyl isocyanate⁴:

In a 3-neck round-bottom flask equipped with a reflux condenser and a CaCl_2 drying tube, a solution of ethyl oxamate (40 g, 0.34 mol) in 1,2-dichloroethane (100 mL) was treated with oxalyl chloride (36 mL, 0.41 mol) at 0°C . The mixture was refluxed during 5 h. The residue was distilled under reduced pressure. After removing oxalyl chloride and 1,2-dichloroethane under a pressure of 100 mm Hg, 40 g of ethyloxalyl isocyanate was collected (82% yield); bp $65\text{-}68^\circ\text{C}/0.5$ mm Hg.

This compound is very sensitive to moisture and should be stored at 0°C under nitrogen. $^1\text{H NMR } \delta$ 1.37 (t, *J* 7, 3H); 4.37 (q, *J* 7, 2H).

N-protected ethyl oxamates 1. General procedure :

To a solution of ethyloxalyl isocyanate (2.86 g, 20 mmol) in toluene (16 mL) was added dropwise a solution of alcohol (23 mmol) in toluene (4mL) at 0°C, under nitrogen. The mixture was heated at 40°C for 15 min. After elimination of the solvent, **1** was obtained in quantitative yield and was used without further purification in reactions with bromide reagents.

N-t-butoxycarbonyl ethyl oxamate 1a :

$^1\text{H NMR } \delta$ 1.39 (t, *J* 7, 3H); 1.52 (s, 9H); 4.39 (q, *J* 7, 2H); 8.76 (broad, NH); $^{13}\text{C NMR } \delta$ 13.42; 27.39; 63.23; 83.19; 148.49; 151.01; 159.25. Anal. calcd. for $\text{C}_9\text{H}_{15}\text{NO}_5$: C, 49.76; H, 6.96; N, 6.45. Found : C, 49.69; H, 6.74; N, 6.43.

N-benzyloxycarbonyl ethyl oxamate 1b :

$^1\text{H NMR } \delta$ 1.35 (t, *J* 7, 3H); 4.35 (q, *J* 7, 2H); 5.22 (s, 2H); 7.37 (m, 5H); 9.01 (broad, NH) ; $^{13}\text{C NMR } \delta$ 13.46; 63.62; 68.07; 128.24; 128.36; 128.47; 134.23; 150.00; 152.04; 158.97. HRMS calcd for $\text{C}_{12}\text{H}_{13}\text{NO}_5$: 251.0794 (M^+); found : 251.0796.

N-protected N-alkyl ethyl oxamates 2. General procedure :

To a solution of **1** (3 mmol) in dry DMF (5 mL) was added potassium t-butoxide

(336 mg, 3 mmol) and bromide reagent (2.3 mmol). The solution was heated at 60°C for 1 h. After addition of CH₂Cl₂ (10 mL) to the mixture, the organic layer was washed with water (2x10 mL), dried over sodium sulfate and the solvent was removed *in vacuo*. The crude residue was purified by column chromatography on silica gel using ethylacetate/cyclohexane (1/4) as eluent and gave **2a-f**.

N-t-butoxycarbonyl N-ethylacetate ethyl oxamate 2a :

¹H NMR δ 1.43 (t, *J* 7, 3H); 1.52 (t, *J* 7, 3H); 1.65 (s, 9H); 4.37 (q, *J* 7, 2H); 4.51 (q, *J* 7, 2H); 4.54 (s, 2H); ¹³C NMR δ 14.10; 14.42; 27.96; 58.58; 44.08; 61.95; 62.60; 86.18; 151.06; 161.60; 163.14; 167.77. HRMS calcd for C₉H₁₂NO₇ : 246.0614 (M⁺ - C₄H₉); found : 246.0618.

N-t-butoxycarbonyl N-benzyl ethyl oxamate 2b :

¹H NMR δ 1.35 (t, *J* 7, 3H); 1.41 (s, 9H); 4.34 (q, *J* 7, 2H); 4.83 (s, 2H); 7.38 (m, 5H); ¹³C NMR δ 13.75; 27.63; 46.13; 62.13; 85.28; 127.58; 127.76; 128.44; 136.33; 151.52; 161.74; 163.37.; 62.60; 86.18; 151.06; 161.60; 163.14; 167.77. HRMS calcd for C₁₁H₁₃NO₃ : 207.0895 (M⁺ - C₅H₈O); found : 207.0899.

N-t-butoxycarbonyl N-octyl ethyl oxamate 2c :

¹H NMR δ 0.81 (t, *J* 7, 3H); 1.21-1.23 (m, 12H); 1.29 (t, *J* 7, 3H); 1.45 (s, 9H); 3.58 (t, *J* 7.5, 2H); 4.27 (q, *J* 7, 2H); ¹³C NMR δ 13.64; 13.91; 22.47; 26.50; 27.62; 27.97; 28.99; 31.60; 43.06; 61.83; 84.70; 151.62; 161.80; 163.16. HRMS calcd for C₁₃H₂₃NO₅ : 273.1576 (M⁺ - C₄H₈); found : 273.1567.

N-t-butoxycarbonyl N-prop-2-enyl ethyl oxamate 2d :

^1H NMR δ 1.28 (t, J 7, 3H); 1.43 (s, 9H); 4.17-4.19 (dxt, 2J 1.4, 3J 5.6, 2H); 4.27 (q, J 7, 2H); 5.09-5.17 (m, 2H); 5.69-5.78 (m, 1H); ^{13}C NMR δ 13.49; 27.41; 44.70 ; 61.75; 84.81; 117.18; 131.34; 151.16; 161.45; 162.66. HRMS calcd for $\text{C}_{12}\text{H}_{19}\text{NO}_5$: 257.1263 (M^+); found : 257.1263.

N-t-butoxycarbonyl N-4-bromobutyl ethyl oxamate 2e :

^1H NMR δ 1.32 (t, J 7, 3H); 1.48 (s, 9H); 1.66-1.90 (m, 4H); 3.39 (t, J 6.5, 2H); 3.66 (t, J 6.5, 2H); 4.30 (q, J 7, 2H); ^{13}C NMR δ 14.13; 27.07; 28.09; 30.04; 33.29 ; 42.44; 63.38; 85.58; 151.87; 162.04; 163.63. HRMS calcd for $\text{C}_9\text{H}_{22}\text{NO}_5\text{Br}$: 295.0055 ($\text{M}^+ - \text{C}_4\text{H}_8$); found : 295.0053.

N-benzyloxycarbonyl N-octyl ethyl oxamate 2f :

^1H NMR δ 1.07 (t, J 7, 3H); 1.42-1.52 (m, 12H); 1.74 (m, 3H); 3.90 (t, J 7.5, 2H); 4.35 (q, J 7, 2H); 5.44 (s, 2H); 7.57 (m, 5H); ^{13}C NMR δ 14.12; 14.47; 23.01; 26.95; 28.17; 29.12; 32.13; 43.68; 62.58; 69.86; 129.03; 129.13; 129.35; 134.55 ; 153.68; 162.14; 163.58. HRMS calcd for $\text{C}_{20}\text{H}_{29}\text{NO}_5$: 363.2046 (M^+); found : 363.2044.

N-alkyl t-butyl carbamates 3. General procedure :

To a solution of **2** (2 mmol) in THF (4 mL) was added a solution of LiOH (6 mmol) in H_2O (3 mL) at room temperature. After 3 h, the reaction mixture was

diluted with H₂O and extracted with CH₂Cl₂. The organic layer was dried over sodium sulfate and concentrated in vacuo to give **3**.

N-octyl t-butyl carbamate 3c⁵ :

99% yield. ¹H NMR δ 0.83 (t, *J* 7, 3H); 1.22 (s, 9H); 1.32–1.47 (m, 12H); 3.05 (q, *J* 7, 2H); 4.58 (broad, NH); ¹³C NMR δ 14.00; 22.56; 26.82; 28.42; 29.23; 29.27; 30.08; 31.79; 40.62; 78.80; 155.93.

N-4-bromobutyl t-butyl carbamate 3e⁶ :

91% yield. ¹H NMR δ 1.38 (s, 9H); 1.52-1.62 (m, 2H); 1.78-1.88 (m, 2H); 3.09 (t, *J* 6.5, 2H); 3.36 (t, *J* 6.5, 2H); 4.68 (broad, NH); ¹³C NMR δ 28.26; 28.61; 29.78; 33.17; 39.40; 79.00; 155.87.

N-octyl ethyl oxamate 4c :

To a solution of **2c** (329 mg, 1 mmol) in dichloromethane (2 mL) was added TFA (3.11 mL, 40 mmol) at room temperature. After 3 h, the crude reaction mixture was washed with Na₂CO₃ 10% (5 mL). The organic layer was dried over sodium sulfate and concentrated *in vacuo* to give **4c** (86% yield).

¹H NMR δ 0.81 (t, *J* 7, 3H); 1.11-1.31 (m, 10H); 1.32 (t, *J* 7, 3H); 1.42 –1.60 (m, 2H); 3.26 (q, *J* 7, 2H); 4.28 (q, *J* 7, 2H); 7.27 (broad, NH); ¹³C NMR δ 13.56; 13.76; 22.36; 26.57; 28.73; 28.91; 31.52; 39.91; 63.02; 156.92; 160.26. HRMS calculated for C₁₂H₂₃NO₃ : 229.1678 (M⁺); found : 229.1683.

N-t-butoxycarbonyl pyrrolidine 5⁷ :

This compound was obtained in 75 % yield from **3e** (504 mg, 2 mmol) by using the conditions described for the preparation of the oxamates **2** .

¹H NMR δ 1.36 (s, 9H); 1.65-1.81 (m, 4H); 3.20 (t, *J* 7, 4H); ¹³C NMR δ 24.79; 25.54; 28.33; 45.41; 45.73; 78.64; 154.47.

References

1. Berrée, F.; Michelot, G. and Le Corre, M. *Tetrahedron Lett.* in press.
2. For reviews see : (a) Gibson, M. S. and Bradshaw, R. W. *Angew. Chem. Int. Ed.* **1968**, *7*, 919-930.(b) Ragnarsson, U. and Grehn, L. *Acc. Chem. Res.* **1991**, *24*, 285-289.
3. (a) Degerbeck, F.; Fransson, B.; Grehn, L. and Ragnarsson, U. *J. Chem. Soc. Perkin Trans 1* **1992**, 245-253. (b) Grehn, L.; Pehk, T. and Ragnarsson, U. *Acta Chem. Scand.* **1993**, *47*, 1107-1111. (c) Fukuyama, T.; Jow, C.K. and Cheung, M. *Tetrahedron Lett.* **1995**, *36*, 6373-6374.
4. Speziale, A.J. and Smith, L.R. *J. Org. Chem.* **1962**, *27*, 3742-3743.
5. Akihiro, O.; Yukiko, I.; Masami, I.; Makoto, S. and Yukari, M. *J. Heterocycl. Chem.* **1985**, *22*, 1173-1181.
6. Hiroji, I.; Masayuki, I. and Chihiro, K. *J. Chem. Soc. Chem. Commun.* **1995**, *10*, 1015-1016.
7. Pandey, G.; Lakshmaiah, G. and Gadre, S.R. *Indian J. Chem. Sect. B* **1996**, *35*, 91-98.