

Cementochronology using synchrotron radiation tomography to determine age at death and developmental rate in the holotype of Homo luzonensis

Anneke van Heteren, Andrew King, Felisa Berenguer, Armand Salvador Mijares, Florent Détroit

▶ To cite this version:

Anneke van Heteren, Andrew King, Felisa Berenguer, Armand Salvador Mijares, Florent Détroit. Cementochronology using synchrotron radiation tomography to determine age at death and developmental rate in the holotype of Homo luzonensis. 2024. hal-03988165

HAL Id: hal-03988165

https://hal.science/hal-03988165

Preprint submitted on 23 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



13/2/23 3:10 PM

Cementochronology using synchrotron radiation tomography to determine

2 age at death and developmental rate in the holotype of *Homo luzonensis*

- 4 Anneke H. van Heteren*1,2,3, Andrew King⁴, Felisa Berenguer⁵, Armand Salvador
- 5 Mijares^{6,7}, Florent Détroit⁸
- ¹Zoologische Staatssammlung München, Staatliche Naturkundliche Sammlungen
- 8 Bayerns, Münchhausenstraße 21, 81247 München, Germany
- ⁹ ²GeoBio-Center, Ludwig-Maximilians-Universität München, 80539 München,
- 10 Germany.

1

3

6

- ³Department Biologie II, Ludwig-Maximilians-Universität München, 82152
- 12 Planegg-Martinsried, Germany
- ⁴PSICHÉ Beamline, Synchrotron SOLEIL, L'orme des Merisiers, 91192 Gif-sur-
- 14 Yvette, France
- ⁵ IPANEMA, CNRS, Synchrotron SOLEIL, L'Orme des Merisiers, 91192 Gif-sur-
- 16 Yvette, France
- 17 ⁶School of Archaeology, University of the Philippines, Quezon City, Philippines
- ⁷National Museum of the Philippines, Manila, Philippines
- 19 ⁸UMR 7194, CNRS, Département Homme & Environnement, Muséum National
- 20 d'Histoire Naturelle, Musée de l'Homme, 17 Place du Trocadéro, 75116 Paris,
- 21 France

23

22 *corresponding author

13/2/23 3:10 PM

24 Abstract 25 Homo luzonensis, a fossil hominin from the Philippines, is smaller than modern 26 humans. At present, very little is known about the life history of this species. 27 Cementochronology can answer life history questions, but usually involves 28 destructive sampling. Here, we use synchrotron radiation to count the yearly 29 cement lines of teeth belonging to a single individual. This approach allows us to 30 determine that this individual was likely 31 years old at time of death and apparently had a developmental pattern comparable to chimpanzees. To our 31 32 knowledge, this is the first time that cementochronology using synchrotron 33 radiation tomography is used for life history and age-at-death estimation. 34 35 Introduction 36 Islands provide amazing examples of evolution and have inspired both Charles 37 Darwin and Alfred Russel Wallace. Nonetheless, little is known about insular 38 evolution, often resulting in miniature or giant versions of mainland animals (1). 39 Whether hominins are also subject to the insular dwarfing to the same degree as 40 other taxa, is still a matter of debate (2, 3) and possible examples include the 41 Homo floresiensis from Indonesia (4) and Homo luzonensis from the Philippines 42 (5).43 The remains of Homo luzonensis were found in Callao Cave on Luzon Island and 44 were described in 2019 (6). The holotype of the species consists, amongst other bones and teeth, of a dental series from upper right P³ to upper right M³, CCH6-a 45 to CCH6-e, (6). The teeth are characterised by their small size and mosaic of 46 47 primitive and derived features (7). Two Homo luzonensis elements, the third 48 metatarsal CCH1 and the upper right M3 CCH6-a, were directly dated to 49 minimum ages of 67 thousand years (kyr) and 50 kyr respectively (5, 8). Whether 50 these hominin remains represent a species that dwarfed under the influence of 51 insular conditions remains to be determined (6), although a descent from Homo 52 erectus seems likely based on the internal organisation of its teeth (7). This begs 53 the questions: Did the life history parameters of this species differ from that of 54 other hominins? And, how old was this individual when he or she died?

13/2/23 3:10 PM

55 In medical terms, the incremental lines in tooth cementum are called "lines of 56 Salter"(9, 10). The morphology of tooth cementum resembles lamellar bone (11). 57 As in the case of lamellar bone (12), there are heavy debates on the nature and the depositional mechanism of the lines of Salter in tooth cementum (13) 58 59 centering around analogous hypotheses. It is generally agreed upon that the 60 lines of Salter are annual lines representing summer and winter (13). 61 Nevertheless, the study of seasonality in human cementochronology has mostly been performed on samples from the northern hemisphere from areas with four 62 seasons (13). It is still not understood how the colours of the lines of Salter 63 64 correspond to the seasons in the southern hemisphere or in areas with 65 monsoonal seasonality. 66 There is no evidence for tooth cement abrasion during decomposition, unlike 67 tooth enamel (14), making tooth cement a promising candidate for chronological research. No taphonomical studies on geological time scales have thus far been 68 69 performed, however. In the present paper, we answer the above research 70 questions using cementochronology, or tooth cementum annulation, derived from 71 synchrotron radiation tomography. 72 73 **Materials and Methods** In 2015, three teeth (CCH6-d: P4; CCH6-c: M1; CCH6-b: M2) of the holotype of 74 75 Homo luzonensis were scanned in Synchrotron SOLEIL in France at the PSICHE 76 beamline. Although other researchers have, in the meantime, also applied 77 cementochronology to modern human canines using synchrotron radiation (15, 78 16), this is the first instance, to our knowledge, that synchrotron based 79 cementochronology is used on 1) a series of teeth belonging to the same 80 individual, 2) that are completely fossilized, and 3) for the purpose of estimating 81 life history traits through dental development. Since transverse sections are advised for human teeth (13), P⁴ and M² were 82 83 mapped transversally by performing a mosaic of quick acquisitions, for which the detector image was binned 4x4 to create an effective pixel size of 2.6 microns. 84 85 Each projection required 0.4 s exposure time, and 300 images were acquired

13/2/23 3:10 PM

over 180° (roughly 2 minutes per tomogram). The reconstructed slices were used to provide a preview of the transverse section, in order to select the locations with the best-preserved cement structures (Figure 1). The buccal root of P⁴ is broken off, so the lingual root was mapped. Rather than use a mosaic mapping approach, M¹ was scanned, also transversally, at a relatively low resolution with 2.6 micron pixel size, 1500 projections over 180° with 170 mm propagation distance and 0.15s exposure time at 40 keV (Figure 2).




Figure 1: Left: Map at 6 mm from the tip of the root of P^4 (CCH6-d). The red square indicates the position of the detailed scan in Figure 3 at the lingual aspect of the lingual root. Centre: For illustrative purposes, a slice of a micro-CT scan of P^4 (7) at the level of the map. Right: The line in the distal view of the 3D model indicates the level where the scan was made.

Based on the maps, 10 appropriate positions for the detailed transverse scans were chosen on M² and 4 on P⁴ in areas where acellular cementum was present. Additionally, five suitable positions for the high-resolution transverse scans were chosen on M¹. All detailed transverse scans on the three teeth were scanned at 40 keV, 70 mm propagation distance, with 3x 0.4 s accumulated images and 3000 projections over 360° (roughly 1 hour per tomogram). In all cases, 40 keV monochromatic radiation was used, with a flux of around 2E11 photons/s/mm².

13/2/23 3:10 PM

Although acellular cementum is usually utilised for cementochronology (17), the cellular cementum was better preserved in M¹, and has also been shown to consist of yearly lines (18, 19).






Figure 2: Left: Section at 0.93 mm from the buccal tip of the distal root of M¹ (CCH6-c) on a low-resolution scan. The red square indicates the position of the detailed scan of the distobuccal aspect in Figure 4. Right: The line in the distal view of the 3D model indicates the level where the scan was made.

After the detailed scans were completed, we chose the clearest images from the dataset and counted the growth lines manually, since automated counts have been found to underestimate the total number of increments (20). Growth lines were counted on multiple images to assure the maximum number of lines was counted. The lines of Salter were counted multiple times (21) in 2015 and in 2022. In both case the same numbers of lines were found. Representative images are shown here.

Results

 $\begin{array}{c} \overline{1}\overline{1}\overline{1}\\ 112 \end{array}$

The cement on the root of M¹ CCH6-c was best preserved out of all three dental elements. The right upper P⁴ CCH6-d is relatively well preserved and displays 24 lines (Figure 3), equivalent to 24 years. The right upper M¹ displays 28 lines in the tooth root cement (Figure 4), which are equivalent to 28 years. Despite taking

13/2/23 3:10 PM

detailed scans at 10 positions in M^2 , surface erosion caused the counts to be incomplete at each position, although it was possible to count the yearly lines. The difference between the number of lines of P^4 and M^1 is four (=28-24) in the holotype of Homo luzonensis CCH6.




Figure 3: Detailed scan at 6 mm from the root of P^4 . The red arrows indicate the 24 yearly lines. The orange dotted line follows the curvature of the cement lines.

Discussion

Development and Age at Death

To assess age at death, the number of lines of Salter need to be added to the age at eruption or the age of root completion. The margins of error of these two approaches are not significantly different, because the timing of these event is equal on a yearly timescale (13). In *Homo luzonensis* CCH6, the cement of the root of M1 was very well preserved and the cement lines count can be considered reliable. The cement of P4 was slightly less well preserved, but was mostly deformed rather than eroded, since the cement lines follow the shape of the outer surface and the cracks. (Figure 3, orange dotted line).

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166167

13/2/23 3:10 PM

In anatomically modern humans (Homo sapiens), a difference of five lines would be expected between these dental elements, since the M¹ erupts at age six and the P⁴ at age 11 on average (22). Nevertheless, there is some intraspecific variation and a difference of four lines may be expected in approximately 5% of the population (23). In chimpanzees (Pan troglodytes) on the other hand M1 erupts at age three and P4 at age seven, so we would expect a difference of four lines between the M¹ and the P⁴ (22). The mean interval in gorillas (*Gorilla gorilla*) on the other hand would be three lines, since M¹ erupts in the fourth year of life and P⁴ in the seventh (22). Although individual variation needs to be taken into account (24), cementochronology is the most accurate method for age-at-death estimation (25), and it is possible to speculate that the dental development pattern of *H. luzonensis* might have been more similar to that of extant chimpanzees than to that of anatomically modern humans, although this is not necessarily a reflection of overall development. Even if an unknown number of lines would be missing from P4, this would imply a developmental strategy guite dissimilar from modern humans. Consequently, a reasonable estimate for the eruption ages of M¹, M² and P⁴ H. luzonensis would be three, six and seven years old respectively, based on a chimpanzee-like dental development pattern.




Figure 4: Detailed scan at 0.93 mm from the root of M¹. The red arrows indicate the 28 yearly lines.

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

13/2/23 3:10 PM

Assuming a chimpanzee-like developmental pattern based on the M¹-P⁴ eruption interval, the root of M¹ indicates that CCH6 died at the age of 31, calculated from 28 years of root cement formation and three years of postnatal life before the eruption of M¹. P⁴, also points to an age at death of 31 years, consisting of 24 years of root cement formation and seven years of postnatal life before the eruption of P⁴. A maximum potential age at death for the type specimen of H. luzonensis may be calculated by assuming a human-like developmental pattern with estimated eruption ages of M¹, M² and P⁴ of six, 12 and 11 years old. This would result in an age at death of 34 or 35 years old. Since the ages calculated assuming a chimpanzee-like developmental pattern correspond perfectly with each other, we think it is likely that the type specimen of *H. luzonensis* was approximately 31 years old at time of death but might have been a few years older. Implications for the Life History of *H. luzonensis* In primates, dental eruption pattern is correlated with brain size (26). The diminutive size of *H. luzonensis* has already hypothesised (8), so, similar to *H.* floresiensis(27), its brain might also have been small, which would not be discordant with the dental eruption pattern found in this study. In primates, various life history traits, such as life expectancy, are also correlated to the dental eruption pattern (28, 29). In anatomically modern humans, life expectancy has changed over the course of (pre-)history (30). In the Mesolithic, life expectancy was between 21 and 40 years at birth, whereas in the Neolithic life expectancy was 17 to 25 years (31). In the Copper, Bronze and Iron ages, life expectancy at birth was between 28 and 44 years (30). CCH6 died in his or her early thirties. This seems relatively young for current standards but is in line with Mesolithic modern humans, as well as corresponding to the potentially fast development of *H. luzonensis*. Possible Evolutionary Implications Homo luzonensis, a small hominin from the Philippines, displays a dental eruption pattern which could be similar to that of modern chimpanzees, but also

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

13/2/23 3:10 PM

occurs in a small portion of the modern human population (23). Consequently, its life history traits might to follow a more ape-like pattern than a modern human pattern. This could have implications for human evolution, but more evidence is clearly needed. It had already been established with *H. floresiensis* that relatively recent hominins could have brain sizes in the range of chimpanzees (27). Perspectives Here, we used a new methodology to calculate age at death using synchrotron radiation. This method is non-destructive and can be used on valuable fossils. such as those of Callao Man. The methodology described herein is a proof of concept and holds great potential for the study of fossil hominins and other valuable dental remains that are not suitable for destructive sampling. Although it is difficult to supply conclusive evidence of life history parameters based on two teeth, the methodology could be applied to more complete tooth rows with the prospect of deciphering an individual's complete dental development. Comparing fossil taxa through time will provide direct evidence for the evolution of dental development with implications for the associated life history traits. Furthermore, this method allows for an assessment of age at death and, when applied to multiple individuals of the same population, might provide an estimate of average life expectancy. As suggested by von Jackowski et al. (15), studying the yearly cementum lines over the whole tooth root cross-section might allow for the detection of potential interruptions or irregularities of the individual layers. These features may be associated with childbirth in females or certain diseases (32-34) and could provide valuable information on the life of fossil hominins. As shown in this work, the potential of using fast, low-resolution measurements combined with slower, high-resolution measurements in selected areas could be important here. To apply cementochronology to the lines of Salter, a standardised protocol for each species, or even each dental element, would be ideal (13). In fossils, however, this would be problematic. Firstly, it would be extremely difficult, if not impossible, to ground-truth any such protocol, due to a lack of fossils with known individual age at death and/or known age at eruption of the respective dental

13/2/23 3:10 PM

229 elements. Secondly, due to taphonomical processes, the researcher will have to 230 work with those parts of the tooth cementum that are sufficiently well-preserved. Theoretically, it should be possible to determine whether an individual died in 231 232 fall/winter or spring/summer. These seasons are characterised by dark or light 233 lines respectively in light microscopy (13). More research on modern humans is 234 needed, however, to determine how the cement layers are formed in a 235 monsoonal climate. It will also be challenging to extrapolate this analysis to 236 synchrotron radiation tomography. It is not currently known how the dark and 237 light lines correspond to the dark and light lines in light microscopy, since the 238 light and dark lines in tomography are caused by differences in density, whereas 239 the cause of the light and dark lines in normal microscopy is still under debate 240 (13) and differs according to the type of microscope used (35, 36). 241 Furthermore, the strong contrast between cementum and air at the surface 242 generates an intense phase contrast fringe, which interferes with interpretation of 243 near surface lines. This could be partially improved by surrounding the tooth with 244 a medium closer to the electron density of cementum, as in Lak et al. (37). 245 246 Acknowledgements 247 We thank the National Museum of the Philippines and its Directorial staff for 248 allowing us to scan the specimens and the Cagayan Provincial Government and 249 the Protected Area Management Board-Peñablanca for authorizing fieldwork at 250 Callao Cave. Funding was provided by Muséum national d'Histoire naturelle, the 251 Wenner-Gren Foundation, the Leakey Foundation Research Grant and the 252 National Geographic Society, beamtime was granted by Synchrotron Soleil, and 253 AHvH was supported by a postdoctoral fellowship of the Humboldt Foundation. 254 The micro-CT scans for illustrative purposes were provided by AST-RX, 255 plateforme d'Accès Scientifique à la Tomographie à Rayons X, UAR 2700 2AD 256 Acquisition et Analyse de Données pour l'histoire naturelle CNRS-MNHN, Paris. 257

References

258

- 259 1. van der Geer AAE, Lyras G, Vos Jd, Dermitzakis M. Evolution of island
- 260 mammals: Adaptation and extinction of placental mammals on islands. 1 ed.
- 261 Chichester: Wiley-Blackwell; 2010. 496 p.
- 262 2. Argue D, Groves CP, Lee MSY, Jungers WL. The affinities of *Homo*
- 263 floresiensis based on phylogenetic analyses of cranial, dental, and postcranial
- 264 characters. Journal of Human Evolution. 2017;107:107-33.
- van Heteren AH. Is LB1 diseased or modern? A review of the proposed
- pathologies. Gene. 2013;528:12-20.
 4. Brown P, Sutikna T, Morwood MJ, Soejono RP, Jatmiko, Saptomo EW, et
- al. A new small-bodied hominin from the Late Pleistocene of Flores, Indonesia.
- 269 Nature. 2004;431(7012):1055-61.
- 5. Mijares AS, Détroit F, Piper P, Grün R, Bellwood P, Aubert M, et al. New
- evidence for a 67,000-year-old human presence at Callao Cave, Luzon,
- 272 Philippines. Journal of Human Evolution. 2010;59(1):123-32.
- 273 6. Détroit F, Mijares AS, Corny J, Daver G, Zanolli C, Dizon E, et al. A new
- species of Homo from the Late Pleistocene of the Philippines. Nature.
- 275 2019;568(7751):181-6.
- 276 7. Zanolli C, Kaifu Y, Pan L, Xing S, Mijares AS, Kullmer O, et al. Further
- 277 analyses of the structural organization of *Homo luzonensis* teeth: Evolutionary
- implications. Journal of Human Evolution. 2022;163:103124.
- 279 8. Grün R, Eggins S, Kinsley L, Moseley H, Sambridge M. Laser ablation U-
- series analysis of fossil bones and teeth. Palaeogeography, Palaeoclimatology,
- 281 Palaeoecology. 2014;416:150-67.
- 282 9. Berkovitz BK, Holland GR, Moxham BJ. Oral anatomy, histology and
- 283 embryology E-book: Elsevier Health Sciences; 2017.
- 284 10. Swetha G, Kattappagari KK, Poosarla CS, Chandra LP, Gontu SR, Badam
- VRR. Quantitative analysis of dental age estimation by incremental line of
- cementum. Journal of oral and maxillofacial pathology: JOMFP. 2018;22(1):138.
- 287 11. Jones SJ, Boyde A. A study of human root cementum surfaces as
- 288 prepared for and examined in the scanning electron microscope. Zeitschrift für
- Zellforschung und Mikroskopische Anatomie. 1972;130(3):318-37.
- 290 12. Mitchell J. van Heteren AH. A literature review of the spatial organization
- of lamellar bone. Comptes Rendus Palevol. 2016;15(1):23-31.
- 292 13. Perrone V, Gocha TP, Randolph-Quinney P, Procopio N. Tooth cementum
- annulation: A literature review. Forensic Sciences. 2022;2(3):516-50.
- 294 14. Akbulut N, Çetin S, Bilecenoğlu B, Altan A, Akbulut S, Ocak M, et al. The
- 295 micro-CT evaluation of enamel-cement thickness, abrasion, and mineral density
- in teeth in the postmortem interval (PMI): new parameters for the determination
- of PMI. International Journal of Legal Medicine. 2020;134(2):645-53.
- 298 15. Von Jackowski J, Müller B, Schulz G, Hotz G, Weitkamp T, Wittwer-
- 299 Backofen U, et al. Tomographic imaging of unique objects: annual layers in tooth
- cementum of Anna Catharina Bischoff born 1719: SPIE; 2022.
- 16. Le Cabec A, Tang NK, Ruano Rubio V, Hillson S. Nondestructive adult
- age at death estimation: Visualizing cementum annulations in a known age
- 303 historical human assemblage using synchrotron X-ray microtomography.
- 304 American Journal of Physical Anthropology. 2019;168(1):25-44.

- 305 17. Robbins Schug G, Brandt ET, Lukacs JR. Cementum annulations, age
- estimation, and demographic dynamics in Mid-Holocene foragers of North India.
- 307 HOMO. 2012;63(2):94-109.
- 308 18. Azorit C, López-Montoya AJ, Mateo-Calahorro BP, Analla M. Seasonal
- investigation of fallow deer cellular cementum from Mediterranean ecosystems
- and its implications for reliable cementochronology applications. Quaternary
- 311 International. 2022;610:133-43.
- 312 19. Takahashi S, Yamamoto T, Takahashi T, Yawaka Y. Incremental lines in
- 313 human cellular cementum: A histological study. Journal of Oral Biosciences.
- 314 2022.
- 315 20. Czermak A, Czermak A, Ernst H, Grupe G. A new method for the
- automated age-at-death evaluation by tooth-cementum annulation (TCA).
- 317 Anthropologischer Anzeiger. 2006:25-40.
- 318 21. Bertrand B, Cunha E, Bécart A, Gosset D, Hédouin V. Age at death
- estimation by cementochronology: Too precise to be true or too precise to be
- accurate? American journal of physical anthropology. 2019;169(3):464-81.
- 321 22. Dean MC, Wood BA. Developing pongid dentition and its use for ageing
- individual crania in comparative cross-sectional growth studies. Folia Primatol
- 323 (Basel). 1981;36(1-2):111-27.
- 324 23. Savara BS, Steen JC. Timing and sequence of eruption of permanent
- teeth in a longitudinal sample of children from Oregon. The Journal of the
- 326 American Dental Association. 1978;97(2):209-14.
- 327 24. Machanda Z, Brazeau NF, Bernard AB, Donovan RM, Papakyrikos AM,
- Wrangham R, et al. Dental eruption in East African wild chimpanzees. Journal of
- 329 Human Evolution. 2015;82:137-44.
- 330 25. Meinl A, Huber CD, Tangl S, Gruber GM, Teschler-Nicola M, Watzek G.
- Comparison of the validity of three dental methods for the estimation of age at
- death. Forensic Science International. 2008;178(2):96-105.
- 333 26. Godfrey LR, Samonds KE, Jungers WL, Sutherland MR. Teeth, brains,
- and primate life histories. Am J Phys Anthropol. 2001;114(3):192-214.
- 335 27. Falk D, Hildebolt C, Smith K, Morwood MJ, Sutikna T, Brown P, et al. The
- 336 brain of LB1, *Homo floresiensis*. Science. 2005;308(5719):242-5.
- 337 28. Smith BH. DENTAL DEVELOPMENT AS A MEASURE OF LIFE
- 338 HISTORY IN PRIMATES. Evolution. 1989;43(3):683-8.
- 339 29. Harvey PH, Clutton-Brock TH. Life history variation in primates. Evolution.
- 340 1985;39(3):559-81.
- 341 30. Galor O, Moav O. The neolithic origins of contemporary variations in life
- 342 expectancy. Available at SSRN 1012650. 2007.
- 343 31. Hassan FA. Demographic archaeology. Advances in archaeological
- 344 method and theory: Elsevier; 1981. p. 225-79.
- 345 32. Kagerer P, Grupe G. Age-at-death diagnosis and determination of life-
- 346 history parameters by incremental lines in human dental cementum as an
- identification aid. Forensic Science International. 2001;118(1):75-82.
- 348 33. Cerrito P, Cerrito L, Hu B, Bailey SE, Kalisher R, Bromage TG. Weaning,
- parturitions and illnesses are recorded in rhesus macaque (*Macaca mulatta*)

13/2/23 3:10 PM

- dental cementum microstructure. American Journal of Primatology.
- 351 2021;83(3):e23235.
- 352 34. Penezić K, Porčić M, Urban PK, Wittwer-Backofen U, Stefanović S.
- 353 Stressful times for women-Increased physiological stress in Neolithic females
- detected in tooth cementum. Journal of Archaeological Science.
- 355 2020;122:105217.
- 356 35. Burke A, Castanet J. Histological observations of cementum growth in
- horse teeth and their application to archaeology. Journal of Archaeological
- 358 Science. 1995;22(4):479-93.
- 359 36. Hillson S. Archaeology and the study of teeth. Endeavour.
- 360 1986;10(3):145-9.

364

- 361 37. Lak M, Néraudeau D, Nel A, Cloetens P, Perrichot V, Tafforeau P. Phase
- 362 contrast X-ray synchrotron imaging: opening access to fossil inclusions in
- opaque amber. Microscopy and microanalysis. 2008;14(3):251-9.