



# Mercury Contamination Challenges the Behavioral Response of a Keystone Species to Arctic Climate Change

Andrea Grunst, Melissa Grunst, David Grémillet, Akiko Kato, Paco Bustamante, Céline Albert, Émile Brisson-Curadeau, Manon Clairbaux, Marta Cruz-Flores, Sophie Gentès, et al.

## ► To cite this version:

Andrea Grunst, Melissa Grunst, David Grémillet, Akiko Kato, Paco Bustamante, et al.. Mercury Contamination Challenges the Behavioral Response of a Keystone Species to Arctic Climate Change. *Environmental Science and Technology*, 2023, 57 (5), pp.2054-2063. 10.1021/acs.est.2c08893 . hal-03988489

**HAL Id: hal-03988489**

**<https://hal.science/hal-03988489>**

Submitted on 5 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Mercury contamination challenges the behavioral response of a keystone species to Arctic climate change**

Andrea S. Grunst<sup>a,\*</sup>, Melissa L. Grunst<sup>a</sup>, David Grémillet<sup>b,c</sup>, Akiko Kato<sup>d</sup>, Paco Bustamante<sup>a,e</sup>, Céline Albert<sup>a</sup>, Émile Brisson-Curadeau<sup>f</sup>, Manon Clairbaux<sup>g,h</sup>, Marta Cruz-Flores<sup>a</sup>, Sophie Gentès<sup>a</sup>, Samuel Perret<sup>b</sup>, Eric Ste-Marie<sup>f</sup>, Katarzyna Wojczulanis-Jakubas<sup>i</sup>, Jérôme Fort<sup>a</sup>

<sup>a</sup>Littoral, Environnement et Sociétés (LIENSs), UMR 7266 CNRS-La Rochelle Université, 2 Rue Olympe de Gouges, FR-17000 La Rochelle, France

<sup>b</sup>CEFE, UMR 5175, CNRS – Université de Montpellier – Université Paul-Valéry Montpellier – EPHE; Montpellier, France

<sup>c</sup>Percy FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch, South Africa

<sup>d</sup>Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 CNRS-La Rochelle Université, 79360 Villiers-en-Bois, France

<sup>e</sup>Institut Universitaire de France (IUF), 1 rue Descartes 75005 Paris, France

<sup>f</sup>McGill University - Macdonald Campus 2111 Lakeshore Dr, Sainte-Anne-de-Bellevue, Quebec H9X 3V9

<sup>g</sup>School of Biological, Environmental and Earth Sciences, University College Cork, Cork T23 N73K, Ireland

<sup>h</sup>MaREI Centre for Energy, Climate and Marine, Environmental Research Institute, University College Cork, Cork P43 C573, Ireland

<sup>i</sup>Department of Vertebrate Ecology and Zoology, University of Gdansk, 80-308 Gdansk, Poland

\*Corresponding author: agrun001@ucr.edu, andrea.grunst@univ-lr.fr

**Abstract:** Combined effects of multiple, climate change-associated stressors are of mounting concern, especially in Arctic ecosystems. Elevated mercury (Hg) exposure in Arctic animals could affect behavioral responses to changes in foraging landscapes caused by climate change, generating interactive effects on behavior and population resilience. We investigated this hypothesis in little auks (*Alle alle*), a keystone Arctic seabird. We compiled behavioral data for 44 birds across 5 years using accelerometers, while also quantifying blood Hg and environmental conditions. Warm sea surface temperature (SST) and low sea ice coverage (SIC) reshaped time activity budgets (TABs) and diving patterns, causing decreased resting, increased flight, and longer dives. Mercury contamination was not associated with TABs. However, highly contaminated birds lengthened inter-dive breaks when making long dives, suggesting Hg-induced physiological limitations. As dive durations increased with warm SST, subtle toxicological effects threaten to increasingly constrain diving and foraging efficiency as climate change progresses, with ecosystem-wide repercussions.

**Synopsis:** Interactive effects of chemical contaminants and climate change are of mounting concern. Our data suggest that subtle toxicological effects of mercury contamination might constrain animal behavioral responses to climate change.

**Keywords:** Climate change, behavioral ecotoxicology, toxicant-induced climate change sensitivity, multiple stressors, behavioral plasticity, mercury (Hg), diving behavior

**Introduction:** Global climate change is reconfiguring ecosystems, exposing organisms to multiple interacting stressors, including thermal stress, resource redistribution, and altered pollutant exposure<sup>1-4</sup>. A pressing research priority involves understanding how multiple stressor interactions will affect biological systems as climate change progresses<sup>5-7</sup>. In particular, mercury (Hg) contamination of marine ecosystems is a persistent problem, which might be aggravated by altered Hg cycling in warming oceans<sup>8-10</sup>. Thus, amplified toxicological effects of Hg exposure threaten to impair organismal resiliency to other climate

change-associated stressors<sup>11-13</sup>, especially by affecting patterns of physiological and behavioral plasticity whereby animals adjust to environmental variability<sup>14-17</sup>. Indeed, in its highly toxic methylated form (methylmercury, MeHg), Hg acts as a potent neurotoxin and endocrine disrupter, which can substantially alter physiology, energetics and behavioral capacity<sup>18,19</sup>. Thus, Hg exposure has the potential to exacerbate declines in behavioral performance under challenging environmental conditions (e.g. resource limitation, predation risk), making behavioral reaction norms steeper and generating synergistic effects<sup>14,20,21</sup>. For example, Hg-exposed zebra finch (*Taeniopygia guttata*) increased foraging latency under predation risk more than controls, with implications for energy intake<sup>20</sup>. Concomitantly, Hg contamination might inhibit behavioral adjustments, canalizing reaction norms. For instance, Hg exposure can cause anemia and lower haemoglobin's oxygen carrying capacity, which might constrain diving behavior in aquatic avian and mammalian predators faced with shifting prey distributions<sup>22</sup>.

The Arctic marine biota is vulnerable to joint effects of climate change and environmental Hg contamination, such that the Arctic presents an important arena for studying how Hg exposure might affect animal resiliency to climate change<sup>23,24</sup>. The Arctic is warming > 3 times faster than other regions, causing cascading effects on ecosystem structure<sup>4</sup>. For example, warming sea surface temperature (SST) and drastic reductions in summer sea ice coverage (SIC) are reconfiguring foraging landscapes, leading to behavioral and bioenergetic challenges<sup>25-28</sup>. Moreover, effects of climate change on the cryosphere and biochemical cycling are altering contaminant exposure patterns, with rising Hg contamination in some Arctic seabird and marine mammal populations presenting a special concern<sup>24,29-31</sup>.

We used a keystone Arctic seabird, the little auk (dovekie, *Alle alle*), as a model species to explore whether effects of Hg exposure on behavioral plasticity could underlie interactive effects of Hg contamination and other climate change-associated stressors on Arctic animals. Little auks are the most abundant seabird in the North Atlantic (~30-40 million breeding pairs globally), and act as ecosystem engineers by transferring nutrients between marine and terrestrial ecosystems<sup>32</sup>. Energy-rich sea ice-associated copepods upon which little auks specialize are declining with climate change, and smaller Atlantic species moving northward, altering the little auk's resource base<sup>27,33,34</sup>. Little auks exhibit high

flight and dive costs and operate close to maximum energy capacity, rendering them vulnerable to energetic stress<sup>35,36</sup>. Behavioral plasticity could buffer little auks, and ecosystems they underpin, against climate change<sup>15</sup>. However, Hg exposure in little auks from East Greenland has risen concurrent with climate change, threatening to impair behavioral plasticity and reproductive performance<sup>33,37</sup>.

We leveraged temporal variation in environmental conditions across five years at a little auk colony in East Greenland combined with advanced biologging techniques (triaxial accelerometry) and individual-level Hg contamination measurements to test two hypotheses: (1) Warm oceanic conditions (high sea surface temperature (SST), low SIC) affect little auk time activity budgets (TABs), increasing energetically expensive flying and diving, and altering behavioral performance traits (flight and dive durations); and (2) Mercury exposure exacerbates or inhibits changes in TABs, and finer scale adjustments of behavioral performance traits, such as adjustments of dive duration to prey distributions<sup>33</sup>. Effects of Hg exposure on behavioral plasticity could constitute a mechanism driving multiple stressor interactive effects by altering little auks' resiliency to climate change, which given the species' keystone status, could have reverberating effects throughout Arctic ecosystems.

## **Methods**

**Field methods:** We studied little auks at a colony at Ukaleqarteq (Kap Høegh; 70°44'N, 21°35'W), East Greenland, during the breeding season (June-August, 2017-2021). The East Greenland Current transports Arctic waters southward past Ukaleqarteq, promoting favourable foraging conditions for little auks, but SST and SIC vary interannually, and climate change is increasingly elevating SST and reducing SIC<sup>29,33</sup>. Little auks nest under rocks, often in accessible locations. The sexes are of similar size and contribute equally to parental care<sup>38,39</sup>. Thus we did not distinguish between the sexes.

We equipped little auks with accelerometers (Axy4, Technosmart, 25 x 10 x 5 mm, 3.0 g, including tape used for attachment, < 2% body mass) to derive records of TABs and behavioral performance traits (dive/flight duration). During each season (July 19 to August 5, 2017-2021), we deployed accelerometers during chick rearing, beginning when chicks were ~3-6 days old, as estimated by size<sup>40</sup> or determined via

hatching date. Sample sizes per year were 17, 7, 20, 11, and 7 across 2017-2021. We captured birds near nests and secured accelerometers to ventral feathers using tesa® tape. Handling time was kept to a minimum at the time of accelerometer deployment to limit effects on behavioral patterns. Birds were recaptured within 3-11 days. Occasionally, shorter datasets (0.5-2 days) were obtained due to battery difficulties. Data < 3/4<sup>th</sup> day (18 hrs) in duration (one dataset from 2021, 13 hr in duration, containing 132 dives) were not used in analyses of TABs, but were used when analysing other behavioral traits (proportion deep diving, dive and inter-dive durations; see below). Removing this short dataset from the analysis did not substantially alter results involving diving behavior. At recapture, we obtained small blood samples (~0.2 mL) via brachial venepuncture for Hg analysis.

***Determining behavioral traits from accelerometer data:*** Acceleration was logged in three dimensions at 50 Hz, with X corresponding to surge (back to front), Y to sway (side to side), and Z to heave (dorsal to ventral). Temperature was logged at 0.2 Hz. We used Igor Pro 9.0 (WaveMetrics, Inc., Lake Oswego, OR, USA) combined with Ethographer 2.05<sup>41</sup> to process data and classify behaviors. In brief, acceleration waves (data vectors measured at equal intervals) in all dimensions were smoothed using a rolling algorithm (1 sec sliding window) in Igor, to extract gravitational acceleration. We calculated body pitch using the formula:  $\text{atan}(\text{XW\_B} / (\text{sqrt}(\text{YW\_B}^2 + \text{ZW\_B}^2))) \times (180/\pi)$ , and vectorial dynamic body acceleration (VDBA) using the formula:  $\text{sqrt}((\text{XW\_B}-\text{XW})^2+(\text{YW\_B}-\text{YW})^2+(\text{ZW\_B}-\text{ZW})^2)$ , where XW\_B, YW\_B and ZW\_B are smoothed acceleration waves<sup>42,43</sup>. We applied continuous wavelet transformation to Z dimension waves to derive dominant wing beat frequency across time (1 Hz resolution). We applied K-mean cluster analysis to dominant frequency waves, using four clusters, which resulted in clusters corresponding to flying (~10-11 Hz), descending dives (~3.5-4.5 Hz), rest (~0 Hz), and intermediate activity (~0-3.5 Hz). These values are consistent with previous work in little auks<sup>36</sup>.

We wrote custom functions in Igor to classify behavior, using cluster analysis output, pitch, VDBA and temperature. Flying was identified based on wing beat frequency of ~10-11 Hz. Dives started when

wing beat frequency was ~3.5-4.5 Hz and pitch was between -10 and -30°, and ended when pitch increased to > 40° and then returned to zero at the water surface. We subdivided dives into deep dives, wherein the descent phase with high wing beat frequency lasted > 5 seconds, and shallower dives, wherein this phase lasted ≤ 5 seconds. Histograms revealed that these criteria captured a bimodal distribution in dive durations. We calculated proportion of time deep versus shallower diving (hereafter, proportion deep diving) as a metric of variation in foraging behavior. Time resting at the colony was identified via absence of wing beat and elevated temperature (> ~17°C), following a period of flight. Time resting on sea ice was classified based on initial high pitch, indicating standing, absence of wing beat, and elevated temperature (lower than for colony; ~12-16°C; Fig. S1). Birds were classed as resting on water based on low wing beat, low temperature (> ~10°C), and absence of classification into another category. We adjusted temperature and pitch ranges associated with behaviors depending on positioning of accelerometers on individuals, with pitch standardized to zero during flight. We calculated TABs (flying, resting at colony, on ice, diving, on water) across each accelerometer dataset's duration (Fig. S1), rather than daily TABs, as Arctic breeding little auks are not constrained by daily light-dark cycles, and lack circadian rhythms in behavior<sup>44</sup>.

We also used accelerometer data to determine durations of foraging trips, dives, inter-dive intervals, and flights, proportion time spent flying during foraging trips versus around the colony (hereafter, proportion foraging flight), and chick provisioning rates. Breeding little auks display a dual foraging strategy, with short trips used for provisioning chicks and longer trips for self-maintenance. We assumed that short and long trips derive from two log-normal distributions<sup>45</sup>. We log-transformed trip duration, and used K-mean clustering in R (R Core 2021) to group foraging trips into two categories. The cutoff value separating short and long trips was 3.9 hrs., which is lower than previously published values (~5.5-6 hrs.)<sup>46,47</sup>. However, we preferred to use the value deriving from our dataset. We determined whether time on the water surface was part of a dive bout (inter-dive interval) using the standard classification method in R package diveMove<sup>48</sup>. The bout ending criteria was 308 sec, and we assumed dives were ≤ 120 sec<sup>36</sup>.

Thus, inter-dive intervals were defined as periods between dives  $\leq 308$  sec. Dive and inter-dive durations can be interdependent<sup>49</sup>, and might be constrained by contamination-linked physiological changes<sup>22</sup>. Thus, we were interested in whether interrelationships between dive- and inter-dive durations were altered in Hg-contaminated birds. We determined proportion of foraging flight by classifying flights embedded within periods of resting at the colony as colony-based, and other flight as foraging-based, and taking the quotient (foraging / colony-based flight). We estimated chick provisioning trips per day as the number of colony visits  $\geq 30$  secs long (minimum time to feed chicks) that proceeded a foraging trip containing  $\geq 5$  dives. Video-recording data from accelerometer-equipped birds showed that colony visits fulfilling these criteria were invariably associated with birds returning with a full gular pouch and feeding the chick.

**Mercury contamination:** Blood samples were centrifuged for 10 min at 3500 rpm, and red blood cells (RBCs) stored in 70% ethanol pending analysis. RBCs were later freeze-dried for 48 hrs and homogenized, after which total Hg concentrations were measured using an advanced Hg analyser spectrophotometer (Altec AMA 254) at the Littoral Environnement et Sociétés (LIENSs) laboratory, La Rochelle University<sup>50</sup>. Mercury in avian blood consists primarily of MeHg ( $> 90\%$ )<sup>51,52</sup>. Thus, total blood Hg provides a good proxy for MeHg exposure. Two or more Hg measurements were taken per bird, using small aliquots of dried RBCs ( $\sim 1.5$ -5 ng), until standard deviation was  $< 10\%$ . We regularly measured a certified reference material (CRM; Lobster Hepatopancreas Tort-3; NRC, Canada; Hg concentration of  $0.292 \pm 0.022$   $\mu\text{g/g}$  dry weight (dw)), and ran blanks before each session. The detection limit was 0.005  $\mu\text{g/g}$  and Tort-3 measurements (mean  $\pm$  SD) averaged  $0.306 \pm 0.004$   $\mu\text{g/g}$ . Mercury concentrations were not available for 2017, hence analyses involving Hg do not include this year.

Our Hg measurements derived from dried RBCs, rather than whole blood. Thus, for comparison to toxicological thresholds for birds, which have been formulated for measurements taken on whole blood (wet weight (ww)), we used the equation recommended by Ackerman et al. (2016)<sup>53</sup> to convert Hg concentrations in dried RBCs to whole blood (ww) equivalents. Namely, we assumed a blood moisture

content of 79%, and thus converted from dried RBC concentrations (dw) to whole blood (ww) equivalents by multiplying the concentrations measured in RBCs by 0.21.

**Sea surface temperature (SST) and sea ice coverage (SIC):** We focused on relationships between SST and SIC and behavior, as these variables strongly affect foraging conditions for little auks<sup>27,33,54</sup>. We determined SST within the little auk's primary foraging range at Ukaleqarteq, determined by GPS tracking<sup>55</sup>, using the National Oceanic and Atmospheric Administration's interpolated SST (OISST) high resolution daily dataset (1/4° global grid; <https://psl.noaa.gov/data/gridded/data.noaa.oisst.v2.highres.html>). The foraging range was encompassed by -21.57° longitude west, -17.47° longitude east, 69.8° latitude south, and 71.65° latitude north, and comprised an ~160 × 200 km area. We extracted daily SST across this area during the chick rearing period, across all days of accelerometer deployment. We averaged daily values across the foraging range and each day of accelerometer deployment for use in analyses.

SIC data were downloaded from the University of Bremen's Sea Ice Remote Sensing database (<https://seaice.uni-bremen.de/sea-ice-concentration/amsre-amsr2/>). Data derived from brightness temperature measurements from the Advanced Microwave Scanning Radiometer 2 (ASMR2) sensor, mounted aboard the JAXA satellite "Shizuku" (GCOM-W1). The ASI algorithm permits derivation of SIC by differences between brightness temperatures at 89 GHz and polarization, with atmospheric corrections<sup>56</sup>. We gridded data on a polar stereographic grid through near neighbor interpolation with 3.125 km resolution. We extracted daily SIC values, and derived averages from data spanning the foraging range and days of accelerometer deployment, as for SST.

**Statistical Analysis:** We used R 4.1.2<sup>57</sup> to process and analyze data. We performed ANOVAs to compare SST, SIC and Hg concentrations (log transformed) among years, and R package *emmeans* for *post-hoc* comparisons (Tukey method)<sup>58</sup>. We used linear mixed effect models (LMMs) with Satterthwaite

approximations for degrees of freedom and year as a random effect to assess whether blood Hg varied with SST/SIC (R packages *lme4*, *lmerTest*)<sup>59,60</sup>.

We analyzed TABs using Dirichlet (mixed) models fit in JAGS, via R2jags<sup>61,62</sup>. The Dirichlet distribution is a multivariate generalization of the beta distribution, suitable for multivariate compositional data<sup>63</sup>. JAGS implements Bayesian models via Markov Chain Monte Carlo (MCMC) simulation. We ran four chains simultaneously with 50,000 repetitions, burn in of 25,000, thinning of 25, and vague priors. All estimated parameters had an Rhat value of  $\sim 1$ , indicating convergence. We first compared TABs among years (fixed effect), with predicted differences determined as derived quantities. We then used Dirichlet mixed models, with year as a random effect, to explore relationships between SST, SIC (log transformed), and TABs. We assessed effects of Hg within the year-comparison (fixed effect) model, and included an interaction between Hg and SST/SIC in the mixed model. Data were transformed using the *DirichletReg* package prior to analysis to avoid zeros<sup>64</sup>. We analyzed proportion deep diving and proportion foraging flight using beta regression (mixed) models in R2jags, and the same approach as above. Predicted differences and parameter estimates from models fit in R2jags were considered significant when 95% credible intervals (CIs) did not cross zero. We used graphical inspection and deviance information criterion (DIC) to assess model fit.

We analyzed among year variation in foraging trip and flight durations (per trip) using LMMs and *emmeans*, with bird identity as a random effect (multiple trips per individual). Long and short trips were analyzed separately. We used LMMs to analyze effects of SST/SIC, Hg and interactions on these variables, with bird identity and year as random effects. The same analyses, but without repeated measures on individuals, were performed for maximum continuous flight duration, deep dive duration, and inter-dive duration (individual-level averages). We included interactions between Hg and inter-dive or dive durations in models involving diving behavior, as dive and inter-dive durations can be interdependent<sup>49</sup>. We also performed a model predicting deep dive duration from the interaction between foraging trip duration and SST/SIC, to assess the hypothesis that birds reach colder, more profitable waters, in which diving behavior could differ, on longer trips. Individual identity was a random effect.


## Results

***Variation in oceanic conditions and Hg contamination:*** Across years, SST experienced by little auks during accelerometer deployment within their primary foraging range averaged (mean  $\pm$  SE)  $2.20 \pm 0.224^{\circ}\text{C}$  (range  $-0.932, 7.416^{\circ}\text{C}$ ), and SIC averaged  $3.33 \pm 0.867\%$  (range:  $0.016\text{-}14.812\%$ ). SST ( $F_{4,57} = 43.66, P < 0.001$ ) and SIC ( $F_{4,57} = 36.84, P < 0.001$ ) varied among years, with SST warmer and SIC lower in 2018 and 2021 (Table 1). Blood Hg averaged  $1.088 \pm 0.039\text{ }\mu\text{g/g dw}$  (range:  $0.579\text{-}1.745\text{ }\mu\text{g/g dw}$ ), or, for comparison to toxicological benchmarks,  $\sim 0.228\text{ }\mu\text{g/g ww}$  (range:  $0.121\text{-}0.366\text{ }\mu\text{g/g ww}$ ), assuming blood moisture content of  $79\%^{64}$ . Blood Hg ( $\mu\text{g/g}$ ) concentrations varied somewhat among years ( $F_{3,40} = 7.706, P < 0.001$ ) (Table 1).

***Warm oceanic conditions affect TABs and foraging patterns:*** Variation in oceanic conditions (SST, SIC) among years correlated with reshaping of TABs. In high SST-low SIC years (2018, 2021), little auks were predicted to spend as much as  $11.7\%$  ( $2.7\%, 21.1\%$ ) more time flying,  $23.2\%$  ( $12.8\%, 32.9\%$ ) more time on water,  $21.4\%$  ( $14.8\%, 28.1\%$ ) less time resting on ice, and  $27.6\%$  ( $21.9\%, 32.6\%$ ) less time resting at the colony relative to in cold years (Fig. 1a-e., Fig. S2). However, time resting at the colony was also low in the coldest year of 2020 (Fig. 1b). Percent time diving was consistent across years, at  $15\text{-}19\%$  (Fig. 1d).

**Table 1.** Variation in a) sea surface temperature (SST; °C) and b) sea ice coverage (SIC; %), and c) blood mercury (Hg) concentrations (µg/g dw), experienced by little auks at Ukaleqarteq, East Greenland, summarized across study years. We quantified values for environmental variables (SST, SIC) on a per-individual basis, depending on the period of accelerometer deployment.

a) SST					
Year	Mean ± SE	95% CI	Range	N	
2017	2.472 ± 0.245	1.981, 2.964	3.312, 2.585	17	
2018	3.615 ± 0.382	2.850, 4.381	3.544, 4.038	7	
2019	1.415 ± 0.226	0.963, 1.868	0.318, 3.353	20	
2020	-0.055 ± 0.305	-0.666, 0.555	-0.932, 0.675	11	
2021	5.886 ± 0.382	5.120, 6.651	4.738, 7.416	7	
Overall	2.20 ± 0.224	1.75, 2.64	-0.932, 7.416	62	
b) SIC					
2017	1.820 ± 0.497	1.052, 3.146	1.589, 2.027	17	
2018	0.081 ± 0.034	0.034, 0.190	0.057, 0.263	7	
2019	1.192 ± 0.300	0.719, 1.974	0.096, 4.736	20	
2020	10.058 ± 3.417	5.093, 19.861	5.165, 14.812	11	
2021	0.035 ± 0.015	0.015, 0.083	0.016, 0.078	7	
Overall	3.33 ± 0.867	2.48, 41.8	0.016, 14.812	62	
c) Blood Hg					
2018	1.210 ± 0.096	1.031, 1.420	0.940-1.488	7	
2019	0.916 ± 0.044	0.831, 1.010	0.579-1.627	19	
2020	1.291 ± 0.082	1.136, 1.470	0.993-1.745	11	
2021	1.193 ± 0.094	1.016, 1.400	1.102-1.272	7	
Overall	1.088 ± 0.039	1.013, 1.169	0.579-1.745	44	




**Figure 1.** Little auk TABs across years at Ukaleqarteq, East Greenland. Relatively cold (blue; average SST < 2.5°C) and warm (red; average SST > 3.5°C) SST years are differentiated by color to facilitate visualization of differences in behavior associated with warmer conditions. Open triangles are predicted means and error bars represent 95% CIs. Colored points are observed values.

Dirichlet mixed models substantiated strong effects of SST and SIC on TABs. Beta coefficient estimates for SST (and SIC) had 95% CIs not crossing zero for time resting on ice (-0.679 (-0.998, -0.341)) and at the colony (-0.329 (-0.626, -0.045)) (Table S1). Predicted values for other behavioral categories were also affected by SST and SIC, as predictions for behavioral categories are interdependent (Fig. 2a). Time resting on ice was predicted to drop from ~25% to < 3%, time resting at the colony decline (~22 to 10%), time flying (~14 to 30%) and on water (~24 to 38%) increase, and diving time remain constant (~17-18%), as SST increased from < 0°C to > 7°C (Fig. 2a).

Little auk foraging trip and flight patterns were altered under warm conditions, with long trips primarily affected (Fig. S3a-d; Table S2-S6). Long trip duration ( $F_{4,49} = 5.022$ ,  $P < 0.001$ ), flight time per long trip ( $F_{4,56} = 6.801$ ,  $P < 0.001$ ), and maximum continuous flight duration ( $F_{4,56} = 3.321$ ,  $P = 0.016$ ) were greatest during high SST years (Table S2-S4, Fig. S3a, b). In the warmest year (2021), long trips lasted 27.5 hrs. (20.4, 37.1), little auks flew 7.76 hrs. (5.43, 11.09) during long trips, and maximum continuous flight duration was 2.23 hrs. (1.58, 2.88), compared to 14.4 hrs. (12.3, 16.9), 3.12 hrs. (2.56, 3.80), and 1.04 hrs. (0.563, 1.51) in 2019. In mixed models, flight time per long trip ( $\beta \pm SE = 0.307 \pm 0.068$ ,  $t_{10} = 4.511$ ,  $P < 0.001$ ) and maximum continuous flight duration ( $\beta \pm SE = 0.255 \pm 0.106$ ,  $t_{10} = 2.395$ ,  $P = 0.039$ ) increased with SST, and long trip duration tended to increase ( $\beta \pm SE = 0.122 \pm 0.066$ ,  $t_{11} = 1.857$ ,  $P = 0.090$ ). Little auks spent more time flying around the colony in cold years, with predicted differences as high as 11.2% (7.7%, 15.1%), reflecting higher colony attendance (Fig. S3e, Fig. S4a). However, in mixed models, there was no overall effect of SST (0.022 (-0.332, 0.376)) or SIC (-0.037 (-0.363, 0.276)) on proportion foraging flight.

Diving behavior changed with oceanic conditions. Little auks were predicted to spend as much as 15.7% (4.2, 26.8) more time deep rather than shallow diving in high SST years (Fig. S3f, Fig. S4b). In mixed models, beta coefficient estimates were 0.293 (0.025, 0.555) and -0.143 (-0.246, -0.037) for SST and SIC, respectively, with proportion deep diving predicted to increase from 69.4% (57.3, 80.3) to 88.3% (78.9, 94.5) under the lowest (-0.9°C) versus highest (7.4°C) SST (Fig. 2b). Deep dive ( $F_6 = 15.01$ ,  $P < 0.001$ ) and inter-dive duration ( $F_6 = 9.305$ ,  $P < 0.001$ ) were longest in high SST years (Table S7-S8; Fig. S3g, h), although *post-hoc* comparisons were non-significant. However, the mixed model indicated that little auks made longer dives when SST was high ( $\beta \pm SE = 2.194 \pm 0.797$ ,  $t_{41} = 3.246$ ,  $P = 0.002$ ; Fig. 2c). Moreover, foraging trip duration interacted with SST to predict deep dive duration, with dives shorter on longer foraging trips when SST in the primary foraging range was high ( $\beta \pm SE = -0.014 \pm 0.004$ ,  $t_{397} = -2.966$ ,  $P = 0.003$ ).


Chick provisioning rates (feeds/day) varied among years ( $F_4=2.929$ ,  $P = 0.029$ ), tending to be highest in cold years (e.g. 2019: (2.580 (2.114, 3.040)), 2020: 2.810 (2.116, 3.500)), and lowest in warm years (e.g. 2021: 1.540 (0.688, 2.390); Fig. S3i, Table S9). However, *post-hoc* comparisons were non-significant ( $P > 0.169$ ), and effects of SST ( $\beta \pm SE = -0.131 \pm 0.172$ ,  $t_5 = -0.762$ ,  $P = 0.477$ ) and SIC ( $\beta \pm SE = 0.101 \pm 0.081$ ,  $t_{15} = 1.239$ ,  $P = 0.234$ ) on provisioning rates were non-significant in mixed models.



**Figure 2.** Little auk a) TABs (points staggered to facilitate viewing), b) proportion time deep versus shallow diving, and c) deep dive duration at Ukaleqarteq, East Greenland, as a function of increasing SST, with year as a random effect. Shapes at the center of error bars are predicted means, and error bars represent 95% credible a, b) or confidence c) intervals. Circular red points in b) and c) are observed values. Confidence intervals around LMM predictions were derived using bootstrapping.

### Mercury concentrations unrelated to TABs, but associated with dive bout patterning

TABs did not vary with blood Hg concentrations, irrespective of oceanic conditions. Beta coefficient estimates for Hg had 95% CIs overlapping zero in the year-comparison model (Fig. S5). In the mixed model, there was no statistically significant effect of Hg on TABs, additively or in interaction with SST, with 95% CIs overlapping zero, and DICs higher than in models including SST alone. Hg concentrations were unrelated to foraging trip and flight durations ( $P > 0.20$ ), proportion foraging-based flight (0.003 (-0.331, 0.357)), and proportion deep diving (1.027 (-2.663, 4.399)). Little auk deep dive duration was unrelated to Hg concentrations ( $\beta \pm SE = 0.637 \pm 0.797$ ,  $t_{41} = 0.800$ ,  $P = 0.428$ ), and interaction terms involving SST and SIC were non-significant ( $P > 0.05$ ).



**Figure 3.** Interaction between dive duration and Hg concentrations in predicting inter-dive durations between deep dives for little auks at Ukaleqarteq. Open triangles are predicted values with 95% CIs for birds with Hg concentrations spanning the observed range. Closed points are observed values. Birds with higher Hg concentrations lengthened inter-dive intervals with dive duration, whereas those with lower Hg concentrations did not. Confidence intervals around LMM predictions were derived using bootstrapping.

However, little auks with higher blood Hg concentrations increased inter-dive interval duration with dive duration more than less contaminated individuals, as reflected by a significant interaction between dive duration and Hg in predicting inter-dive interval duration ( $\beta \pm SE = 6.704 \pm 2.991$ ,  $t_{40} = 2.241$ ,  $P = 0.031$ ; Fig. 3). The same pattern held when analyzing observations of every dive (and inter-dive) made by individuals, in sequence, rather than means (interaction:  $\beta \pm SE = 1.980 \pm 0.244$ ,  $t_{32350} = 8.120$ ,  $P < 0.001$ ).

**Discussion:** Multiple stressor effects involving exposure to contaminants, such as Hg, are attracting growing concern in the context of rapid climate change<sup>1,5,11</sup>. Yet, understanding of mechanisms underlying such effects remains limited. We addressed this deficit by exploring joint effects of climate change-associated environmental variables and Hg contamination on a keystone Arctic seabird's behavioral ecology. We hypothesized that warm oceanic conditions induce costly behavioral changes, and that Hg exposure might exacerbate or inhibit behavioral responses, generating interactive effects. Below we discuss evidence for these hypotheses, and implications of our findings.

*TABs and behavioral performance demands under warm conditions:* Climate change is increasingly exposing little auks to warm oceanic conditions, which alter resource distribution<sup>33</sup>. Our data demonstrate that warm conditions reconfigure TABs in these sea-ice adapted seabirds, reducing time resting at the colony and on ice, and increasing time in flight and on water. Accelerometer data revealed that sea ice serves as an important resting substrate for little auks at Ukaleqarteq in years with substantial SIC. Thus, loss of ice will likely reduce rest quality for little auks at sea, with energetic and health implications<sup>65,66</sup>. Indeed, thermoregulatory costs are elevated for seabirds resting in water<sup>65</sup>, although warming SST may reduce these costs<sup>67</sup>. A previous study could not differentiate energetic costs of resting on water versus land or ice for little auks, but did not differentiate land versus ice<sup>36</sup>. Energetic costs of resting on land at the colony might be elevated by anti-predator vigilance<sup>68</sup>. Little auks also reduced time resting at and flying around the colony during warm periods, further reducing rest, and altering colony-based social dynamics, which are critically important in many seabirds<sup>69,70</sup>.

As rest and social time decreased, foraging and behavioral performance demands increased. Little auks spent more time flying and made longer continuous flights under warmer conditions, suggesting an increase in foraging range. Indeed, even when there was virtually no sea ice within the primary foraging range around Ukaleqarteq, tagged birds spent some time on ice, with ice reached via prolonged flight (> 2 hrs.). Studies at other colonies suggest that little auks can fly > 100 km to reach profitable foraging areas at marginal sea ice zones when foraging conditions are poor near colonies<sup>71</sup>, and similar increases in foraging range also occurs in other seabird species<sup>72</sup>. Increased foraging range can maintain energy intake, but decreases net energy return for central place foragers. Thus, foraging range commonly correlates negatively with breeding performance in central place foragers, including little auks<sup>73,74</sup>. Accordingly, little auk chick provisioning rates at Ukaleqarteq tended to decline with warm conditions, when flight times were longest. Little auks at Ukaleqarteq also increased proportion deep diving and dive duration with warm SST, likely reflecting altered copepod distribution<sup>33</sup>. Previous analyses suggest that energetic costs do not scale to dive duration in little auks<sup>36</sup>, but utilized small sample sizes and did not consider dive depth, which could make identifying energetic patterns difficult. Longer dives might challenge physiological limitations by approaching aerobic dive capacity<sup>75,76</sup>. Thus, results suggest that warm conditions force little auks at Ukaleqarteq to adopt more energetically and physiologically demanding behavior, with rest reduced, activity increased, and behavioral performance demands intensified, at diminishing return.

*Behavioral correlates of Hg contamination:* We hypothesized that Hg exposure might reduce the capacity of birds to maintain behavioral performance or upregulate costly behavior given altered foraging landscapes under warm conditions. To the contrary, little auk Hg concentrations were not related to TABs. When compared to toxicological benchmarks for avian blood<sup>52</sup>, little auk blood Hg concentrations at Ukaleqarteq (which fall in the range of 0.2-1.0 µg/g ww) suggest low toxicological risk, despite recent increases in Hg exposure, which could explain this result. However, a recent study on thick-billed murres (*Uria lomvia*) found interrelationships between thyroid hormone levels, time spent diving and Hg exposure with only slightly higher total blood Hg (mean ± SE = 1.51 ± 0.05 µg/g dw, or ~0.31 µg/g ww,

range: 0.81-2.74  $\mu\text{g/g dw}$ , or  $\sim 0.17$ -0.36  $\mu\text{g/g ww}$ ). Murres with elevated blood Hg showed altered triiodothyronine (T3) levels when sea ice break up was early, and murres were likely stressed due to poor foraging conditions, but not given more usual ice conditions. In turn, elevated T3 levels were associated with reduced time diving, suggesting that thyroidal dysregulation in Hg exposed birds could affect activity budgets under stressful environmental conditions<sup>14</sup>.

Rather than affecting TABs, Hg exposure could alter behavioral performance traits or temporal patterning of behavior. Little auks with higher Hg concentrations exhibited changes in dive bout patterning, with inter-dive breaks lengthened when individuals were making long dives. Mercury exposure might cause physiological impairments, such as decreased oxygen carrying capacity<sup>22</sup>, forcing dive bout restructuring as birds approach aerobic dive capacity. Mercury can interfere with heme-proteins including haemoglobin and myoglobin, which play critical roles in oxygen carrying capacity in blood and muscles, respectively<sup>77,78</sup>. Indeed, Hg contaminated common loons (*Gavia immer*) showed increased diving frequency, concurrent with reduced dive duration, suggesting reduced aerobic dive capacity<sup>22</sup>. In contrast, little auks with higher Hg maintained dive duration, but lengthened inter-dive breaks. Adjusting inter-dive durations might allow Hg-contaminated birds to perform long dives despite physiological impairments when long dives are essential to prey capture. However, lengthening breaks might eventually reduce time underwater or resting outside of dive bouts, with implications for foraging efficiency, digestion and recovery<sup>49,79,80</sup>.

Effects of Hg contamination on little auk diving behavior are predicted to increase with warming conditions. When SST was high and SIC low, little auks at Ukaleqarteq spent more time deep diving and increased dive duration (Fig. 2b, c). Thus, as Hg-contaminated birds increased inter-dive durations when making longer dives, Hg contamination might increasingly constrain diving patterns and foraging efficiency as SST increases with climate change. Changes in Hg exposure with climate change might also combine with foraging challenges to elevate toxicological effects on little auks and ecosystems they underpin<sup>24</sup>.

In conclusion, warm oceanic conditions affect TABs and finer scale behavior in little auks, a sea-ice adapted Arctic seabird. Mercury exposure threatens to alter behavioral responses, generating climate change-by-contamination interactive effects. Mercury contamination was not correlated with TABs. However, birds with high Hg concentrations took longer inter-dive breaks when making long dives, suggesting physiological constraint. A next step will be exploring the nature of this constraint, for instance by measuring oxygen carrying capacity. As little auks engaged in longer dives given warm SST, subtle behavioral effects of Hg-contamination threaten to increasingly constrain this keystone species' foraging efficiency in a warming Arctic, with ecosystem-wide repercussions.

**Supporting Information:** Additional statistical results, which were too extensive to include in the main text.

**Acknowledgements:** We acknowledge long-term support from the French Polar Institute (IPEV) to the ADACLIM program (388) administered by JF and DG. The authors are grateful to Maud Brault-Favrou of the platform Analyses ELémentaires of LIENSs laboratory for assistance with Hg analyses, and Clément Bertin for help with SIC data. Thanks also to the CPER (Contrat de Projet Etat-Région) and FEDER (Fonds Européen de Développement Régional) for funding the AMA of LIENSs laboratory, and the Institut Universitaire de France (IUF) for supporting PB as a Senior Member. ASG, MLG, and MCF are supported by the European Union's Horizon 2020 programme (Marie Skłodowska-Curie grants 101025549, 896866, 101024166), and CA was supported by a PhD fellowship from the French Ministry of Higher Education and Research. We thank Nanu travel for facilitating field seasons, and the Northern Scientific Training Program of Canada for funding travel and providing logistical support. This work contributes to research projects ARCTIC-STRESSORS and ILETOP funded by the French National Research Agency (ANR-20-CE34-0006, ANR-16-CE34-0005), the international ARCTOX initiative (arctox.cnrs.fr), the Excellence Chair ECOMM funded by the Region Nouvelle Aquitaine, and the OPUS13 funded by the Polish National Science Center (2017/25/B/NZ8/01417).

**Data accessibility statement:** Data is available via the Zenodo community of European Commission Funded Research (OpenAIRE), doi:10.5281/zenodo.7185806, and via Dryad, doi:10.5061/dryad.dbrv15f4t.

## References

- <sup>1</sup>Noyes, P.D., McElwee, M.K., Miller, H.D., Clark, B.W., Van Tiem, L.A., Walcott, K.C., Erwin K.N., Levin, E.D. (2009) The toxicology of climate change: Environmental contaminants in a warming world. *Environ Int.* 35:971–986. doi:10.1016/j.envint.2009.02.006.
- <sup>2</sup>Sherwood, S.C., Huber, M. (2010) An adaptability limit to climate change due to heat stress. *Proc Natl Acad Sci USA.* 107:9552–9555. doi:10.1073/pnas.0913352107.
- <sup>3</sup>Bartley, T.J., McCann, K.S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M.M., MacDougall A.S, Tunney, T.D., McMeans, B.C. (2019) Food web rewiring in a changing world. *Nat Ecol Evol.* 3:345–354. doi:10.1038/s41559-018-0772-3.
- <sup>4</sup>IPCC. (2021) Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K., Waterfield, T., Yelekçi, O., Yu, R., Zhou, B. (eds.). Cambridge University Press.
- <sup>5</sup>Crain, C.M., Kroeker, K., Halpern, B.S. (2008) Interactive and cumulative effects of multiple human stressors in marine systems. *Ecol Lett.* 11:1304-15. doi: 10.1111/j.1461-0248.2008.01253.x.
- <sup>6</sup>Orr, J.A., Vinebrooke, R.D., Jackson, M.C., Kroeker, K.J., Kordas, R.L., Mantyka-Pringle, C., Van den Brink, P.J., De Laender, F., Stoks, R., Holmstrup, M., Matthaei, C.D., Monk, W.A., Penk, M.R., Leuzinger, S., Schaefer, R.B., Piggott, J.J. (2020) Towards a unified study of multiple stressors: divisions and common goals across research disciplines. *Proc Roy Soc. B* 287:20200421. doi:10.1098/rspb.2020.0421. Epub 2020.

454 <sup>7</sup>Tekin, E., Diamant, E.S., Cruz-Loya, M., Enriquez, V., Singh, N., Savage, V.M., Yeh, P.J. (2020) Using  
 455 a newly introduced framework to measure ecological stressor interactions. *Ecol Lett.* 23:1391-  
 456 1403. doi:10.1111/ele.13533.

457 <sup>8</sup>Cossa, D. (2013) Methylmercury manufacture. *Nature Geosci* 6:810–811. doi: 10.1038/ngeo1967.

458 <sup>9</sup>Eagles-Smith, C.A., Silbergeld, E.K., Basu, N., Bustamante, P., Diaz-Barriga, F., Hopkins, W.A., Kidd,  
 459 K.A., Nyland, J.F. (2018) Modulators to mercury risk to wildlife and humans in the context of  
 460 rapid global change. *Ambio* 47:170–197.

461 <sup>10</sup>Schartup, A.T., Thackray, C.P., Qureshi, A., Dassuncao, C., Gillespie, K., Hanke, A., Sunderland, E.M.  
 462 (2019) Climate change and overfishing increase neurotoxicant in marine predators. *Nature* 572:  
 463 648–650. doi:10.1038/s41586-019-1468-9.

464 <sup>11</sup>Hooper, M.J., Ankley, G.T., Cristol, D.A., Maryoung, L.A., Noyes, P.D., Pinkerton, K.E. (2013)  
 465 Interactions between chemical and climate stressors: A role for mechanistic toxicology in  
 466 assessing climate change risks. *Environ Toxicol Chem.* 32:32–48. doi:10.1002/etc.2043.

467 <sup>12</sup>Pirotta, E., Thomas, L., Costa, D.P., Hall, A.J., Harris, C.M., Harwood, J., Kraus, S.D., Miller, P.J.O.,  
 468 Moore, M.J., Photopoulou, T., Rolland, R.M., Schwacke, L., Simmons, S.E., Southall, B.L.,  
 469 Tyack, P.L. (2022) Understanding the combined effects of multiple stressors: A new perspective  
 470 on a longstanding challenge. *Sci Tot Environ* 821: 153322. doi: 10.1016/j.scitotenv.2022.153322.

471 <sup>13</sup>Turschwell, M.P., Connolly, S.R., Schäfer, R.B., De Laender, F., Campbell, M.D., Mantyka-Pringle, C.,  
 472 Jackson, M.C., Kattwinkel, M., Sievers, M., Ashauer, R., Côté, I.M., Connolly, R.M., van den  
 473 Brink, P.J., Brown, C.J. (2022) Interactive effects of multiple stressors vary with consumer  
 474 interactions, stressor dynamics and magnitude. *Ecol Lett.* 25:1483-1496. doi:10.1111/ele.14013.

475 <sup>14</sup>Esparza, I., Elliott, K.H., Choy, E.S., Braune, B.M., Letcher, R.J., Patterson, A., Fernie, K.J. (2022)  
 476 Mercury, legacy and emerging POPs, and endocrine-behavioural linkages: Implications of Arctic  
 477 change in a diving seabird. *Environ Res.* 212:113190. doi:10.1016/j.envres.2022.113190.

- <sup>15</sup>Grémillet, D., Welcker, J., Karnovsky, N., Walkusz, W., Hall, M., Fort, J., Brown, Z., Speakman, J.R., Harding, A.M.A. 2012. Little auks buffer the impact of current Arctic climate change. *Mar Ecol Prog Ser.* 454:197–206. doi:10.3354/meps09590.
- <sup>16</sup>Atwood, T.C., Peacock, E., McKinney, M.A., Lillie, K., Wilson, R., Douglas, D.C., Miller, S., Terletzky, P. (2016) Rapid environmental change drives increased land use by an arctic marine predator. *PloS ONE* 11: e0155932. doi:10.1371/journal.pone.0155932.
- <sup>17</sup>Hall, L.E., Chalfoun, A.D. (2019) Behavioural plasticity modulates temperature-related constraints on foraging time for a montane mammal. *J Anim Ecol.* 88: 363-375. doi:10.1111/1365-2656.12925.
- <sup>18</sup>Whitney, M.C., Cristol, D.A. (2018) Impacts of Sublethal Mercury Exposure on Birds: A Detailed Review. *Rev Environ Contam Toxicol.* 244:113-163. doi:10.1007/398\_2017\_4.
- <sup>19</sup>Gerson, A.R., Cristol, D.A., Seewagen, C.L. (2019) Environmentally relevant methylmercury exposure reduces the metabolic scope of a model songbird. *Environ Pollut.* 246: 790-796. doi: 10.1016/j.envpol.2018.12.072.
- <sup>20</sup>Kobiela, M.E., Cristol, D.A., Swaddle, J.P. (2015) Risk-taking behaviours in zebra finches affected by mercury exposure. *Anim Behav.* 103:153–160. doi:10.1016/j.anbehav.2015.02.024.
- <sup>21</sup>Bustnes, J.O., Bourgeon, S., Leat, E.H.K., Magnúsdóttir, E., Strøm, H., Hanssen, S.A., Petersen A., Olafsdóttir, K., Borgå, K., Gabrielsen, G.W., Furness, R.W. (2015) Multiple stressors in a top predator seabird: potential ecological consequences of environmental contaminants, population health and breeding conditions. *PloS ONE* 10: e0131769. Doi:10.1371/journal.pone.0131769.
- <sup>22</sup>Olsen, B., Evers, D., DeSorbo, C. (2000) Effect of methylated mercury on the diving frequency of the common loon. *J Ecol Res.* 2:67–72.
- <sup>23</sup>Krabbenhoft, D.P., Sunderland, E.M. (2013) Global Change and Mercury. *Science* 341:1457–1458. doi:10.1126/science.1242838.
- <sup>24</sup>AMAP. (2021) 2021 AMAP Mercury Assessment. Summary for Policy-makers. Arctic Monitoring and Assessment Programme (AMAP), Tromsø, Norway. 16 pp.

503 <sup>25</sup>Eamer, J., Donaldson, G., Gaston, T., Kosobokova, K., Lárusson, K.F., Melnikov, I., Reist, J.D.,  
504 Richardson, E., Staples, L., von Quillfeldt, C.H. (2013) Life linked to ice. A guide to sea-ice  
505 associated biodiversity in this time of rapid change. Akureyri, Iceland: CAFF International  
506 Secretariat. 115 p. (CAFF Assessment Series, Vol. 10).

507 <sup>26</sup>Davidson, S.C., Bohrer, G., Gurarie, E., LaPoint, S., Mahoney, P.J., Boelman, N.T., Eitel, J.U.H.,  
508 Prugh, L.R., Vierling, L.A., Jennewein, J., Grier, E., Couriot, O., Kelly, A.P., Meddens, A.J.H.,  
509 Oliver, R.Y., Kays, R., Wikelski, M., Aarvak, T., Ackerman, J.T., Alves, J.A., Bayne, E.,  
510 Bedrosian, B., Belant, J.L., Berdahl, A.M., Berlin, A.M., Berteaux, D., Bêty, J., Boiko, D.,  
511 Booms, T.L., Borg, B.L., Boutin, S., Boyd, W.S., Brides, K., Brown, S., Bulyuk, V.N., Burnham,  
512 K.K., Cabot, D., Casazza, M., Christie, K., Craig, E.H., Davis, S.E., Davison, T., Demma, D.,  
513 DeSorbo, C.R., Dixon, A., Domenech, R., Eichhorn, G., Elliott, K., Evenson, J.R., Exo, K.M.,  
514 Ferguson, S.H., Fiedler, W., Fisk, A., Fort, J., Franke, A., Fuller, M.R., Garthe, S., Gauthier, G.,  
515 Gilchrist, G., Glazov, P., Gray, C.E., Grémillet, D., Griffin, L., Hallworth, M.T., Harrison, A.L.,  
516 Hennin, H.L., Hipfner, J.M., Hodson, J., Johnson, J.A., Joly, K., Jones, K., Katzner, T.E., Kidd,  
517 J.W., Knight, E.C., Kochert, M.N., Kölzsch, A., Kruckenberg, H., Lagassé, B.J., Lai, S., Lamarre,  
518 J.F., Lanctot, R.B., Larter, N.C., Latham, A.D.M., Latty, C.J., Lawler, J.P., Léandri-Breton, D.J.,  
519 Lee, H., Lewis, S.B., Love, O.P., Madsen, J., Maftai, M., Mallory, M.L., Mangipane, B.,  
520 Markovets, M.Y., Marra, P.P., McGuire, R., McIntyre, C.L., McKinnon, E.A., Miller, T.A.,  
521 Moonen, S., Mu, T., Müskens, G.J.D.M., Ng, J., Nicholson, K.L., Øien, I.J., Overton, C., Owen,  
522 P.A., Patterson, A., Petersen, A., Pokrovsky, I., Powell, L.L., Prieto, R., Quillfeldt, P., Rausch, J.,  
523 Russell, K., Saalfeld, S.T., Schekkerman, H., Schmutz, J.A., Schwemmer, P., Seip, D.R.,  
524 Shreading, A., Silva, M.A., Smith, B.W., Smith, F., Smith, J.P., Snell, K.R.S., Sokolov, A.,  
525 Sokolov, V., Solovyeva, D.V., Sorum, M.S., Tertitski, G., Therrien, J.F., Thorup, K., Tibbitts,  
526 T.L., Tulp, I., Uher-Koch, B.D., van Bemmelen, R.S.A., Van Wilgenburg, S., Von Duyke, A.L.,  
527 Watson, J.L., Watts, B.D., Williams, J.A., Wilson, M.T., Wright, J.R., Yates, M.A., Yurkowski,  
528 D.J., Żydelis, R., Hebblewhite, M. (2020) Ecological insights from three decades of animal

529 movement tracking across a changing Arctic. *Science* 370: 712-715. doi:

530 10.1126/science.abb7080. PMID: 33154141.

531 <sup>27</sup>Jakubas, D., Wojczulanis-Jakubas, K., Iliszko, L.M., Strøm, H., Stempniewicz, L. (2017) Habitat  
532 foraging niche of a High Arctic zooplanktivorous seabird in a changing environment. *Sci Rep.*  
533 7:16203. doi:10.1038/s41598-017-16589-7.

534 <sup>28</sup>Blanchet, M-A., Aars, J., Andersen, M., Routti, H. (2020) Space-use strategy affects energy  
535 requirements in Barents Sea polar bears. *Mar Ecol Prog Ser.* 639: 1-19. doi: 10.3354/meps13290.

536 <sup>29</sup>Fort, J., Grémillet, D., Traisnel, G., Amélineau, F., Bustamante, P. (2016) Does temporal variation of  
537 mercury levels in Arctic seabirds reflect changes in global environmental contamination, or a  
538 modification of Arctic marine food web functioning? *Environ Pollut.* 211:382-8. doi:  
539 10.1016/j.envpol.2015.12.061.

540 <sup>30</sup>Morris, A.D., Wilson, S.J., Fryer, R.J., Thomas, P.J., Hudelson, K., Andreassen, B., Blévin, P.,  
541 Bustamante, P., Chastel, O., Christensen, G., Dietz, R., Evans, M., Evenset, A., Ferguson, S.H.,  
542 Fort, J., Gamberg, M., Grémillet, D., Houde, M., Letcher, R.J., Loseto, L., Muir, D., Pinzone, M.,  
543 Poste, A., Routti, H., Sonne, C., Stern, G., Rigét, F.F. (2022) Temporal trends of mercury in  
544 Arctic biota: 10 more years of progress in Arctic monitoring. *Sci Tot Environ.* 839C:155803.  
545 doi:10.1016/j.scitotenv.2022.155803.

546 <sup>31</sup>Tartu, S., Blévin, P., Bustamante, P., Angelier, F., Bech, C., Bustnes, J.O., Chierici, M., Fransson, A.,  
547 Gabrielsen, G.W., Goutte, A., Moe, B., Sauser, C., Sire, J., Barbraud, C., Chastel, O. (2022) A U-  
548 turn for mercury concentrations over 20 years: how do environmental conditions affect exposure  
549 in Arctic seabirds? *Environ Sci Technol.* 56:2443–2454. doi:10.1021/acs.est.1c07633.

550 <sup>32</sup>González-Bergonzoni, I., Johansen, K.L., Mosbech, A., Landkildehus, F., Jeppesen, E., Davidson, T.A.  
551 (2017) Small birds, big effects: the little auk (*Alle alle*) transforms high Arctic ecosystems. *Proc*  
552 *R Soc. B* 284:20162572. doi: 10.1098/rspb.2016.2572.

553 <sup>33</sup>Amélineau, F., Grémillet, D., Harding, A.M.A., Walkusz, W., Choquet, R., Fort, J. (2019) Arctic  
554 climate change and pollution impact little auk foraging and fitness across a decade. *Sci Rep*.  
555 9:1014. doi:10.1038/s41598-018-38042-z.

556 <sup>34</sup>Ershova, E.A., Kosobokova, K.N., Banas, N.S., Ellingsen, I., Niehoff, B., Hildebrandt, N., Hirche, H.J.  
557 (2021) Sea ice decline drives biogeographical shifts of key Calanus species in the central Arctic  
558 Ocean. *Glob Chang Biol*. 10:2128-2143. doi: 10.1111/gcb.15562.

559 <sup>35</sup>Gabrielsen, G.W., Taylor, J.R.E., Konarzewski, M., Mehlum, F. (1991) Field and laboratory metabolism  
560 and thermoregulation in dovekies (Alle alle). *Auk* 108: 71–78. doi: 10.1093/auk/108.1.71.

561 <sup>36</sup>Ste-Marie, E., Grémillet, D., Fort, J., Patterson, A., Brisson-Curadeau, É., Clairbaux, M., Perret, S.,  
562 Speakman, J.R., Elliott, K.H. (2022) Accelerating animal energetics: high dive costs in a small  
563 seabird disrupt the dynamic body acceleration – energy expenditure relationship. *J Exp Biol*. 225:  
564 jeb243252. doi: 10.1242/jeb.243252.

565 <sup>37</sup>Fort, J., Robertson, G.J., Grémillet, D., Traisnel, G., Bustamante, P. (2014) Spatial Ecotoxicology:  
566 Migratory Arctic Seabirds Are Exposed to Mercury Contamination While Overwintering in the  
567 Northwest Atlantic. *Environ Sci Technol*. 48:11560–11567. doi:10.1021/es504045g.

568 <sup>38</sup>Jakubas, D., Wojczulanis, K. (2007) Predicting the sex of Dovekies by discriminant analysis.  
569 *Waterbirds* 30:92–96. doi: 10.1675/1524-4695(2007)030[0092:PTSODB]2.0.CO;2.

570 <sup>39</sup>Wojczulanis, K., Jakubas, D., Stempniewicz, L. (2022) The Little Auk Alle alle: an ecological indicator  
571 of a changing Arctic and a model organism. *Polar Biol*. 45:163-176. doi:10.1007/s00300-021-  
572 02981-7.

573 <sup>40</sup>Konarzewski, M., Taylor, J.R.E. (1989) The influence of weather conditions on growth of Little Auk  
574 Alle alle chicks. *Ornis Scand*. 20:112. doi:10.2307/3676877.

575 <sup>41</sup>Sakamoto, K.Q., Sato, K., Ishizuka, M., Watanuki, Y., Takahashi, A., Daunt, F., Wanless, S. (2009) Can  
576 Ethograms Be Automatically Generated Using Body Acceleration Data from Free-Ranging  
577 Birds? *PloS ONE* 4: e5379. doi:10.1371/journal.pone.0005379.

578 <sup>42</sup>Patterson, A., Gilchrist, H.G., Chivers, L., Hatch, S., Elliott, K. (2019) A comparison of techniques for  
579 classifying behavior from accelerometers for two species of seabird. *Ecol Evol.* 9:3030-3045. doi:  
580 10.1002/ece3.4740.

581 <sup>43</sup>Wilson, R.P., Börger, L., Holton, M.D., Scantlebury, D.M., Gómez-Laich, A., Quintana, F., Rosell, F.,  
582 Graf, P.M., Williams, H., Gunner, R., Hopkins, L., Marks, N., Gerald, N.R., Duarte, C.M., Scott,  
583 R., Strano, M.S., Robotka, H., Eizaguirre, C., Fahlman, A., Shepard, E.L.C. (2020) Estimates for  
584 energy expenditure in free-living animals using acceleration proxies: A reappraisal. *J Anim Ecol.*  
585 89:161-172. doi:10.1111/1365-2656.13040.

586 <sup>44</sup>Wojczulanis-Jakubas, K., Wąż, P., Jakubas, D. (2020) Little auks under the midnight sun: diel activity  
587 rhythm of a small diving seabird during the Arctic summer. *Pol Res.*  
588 doi:10.33265/polar.v39.3309.

589 <sup>45</sup>Welcker, J., Harding, A.M.A., Karnovsky, N.J., Steen, H., Strøm, H., Gabrielsen, G.W. (2009)  
590 Flexibility in the bimodal foraging strategy of a high Arctic alcid, the little auk *Alle alle*. *J Avian*  
591 *Biol.* 40: 388–399. doi:10.1111/j.1600-048X.2008.04620.x.

592 <sup>46</sup>Brown, Z.W., Welcker, J., Harding, A.M.A., Walkusz, W., Karnovsky, N. (2012) Divergent diving  
593 behaviour during short and long trips of a bimodal forager, the little auk (*Alle alle*). *J Avian Biol.*  
594 43:215-226. doi: 10.1111/j.1600-048X.2012.05484.x.

595 <sup>47</sup>Jakubas, D., Wojczulanis-Jakubas, K., Iliszko, L., Darecki, M., Stempniewicz, L. (2014) Foraging  
596 strategy of the little auk *Alle alle* throughout breeding season –switch from unimodal to bimodal  
597 pattern. *J Avian Biol.* 45:551-560. doi: 10.1111/jav.00303.

598 <sup>48</sup>Luque, S.P. (2007) Diving Behaviour Analysis in R. *R News* 7: 8-14.

599 <sup>49</sup>Elliot, K.H., Davoren, G., Gaston, A.J. (2008) Time allocation by a deep-diving bird reflects prey type  
600 and energy gain. *Anim Behav.* 75: 1301-1310. doi:10.1016/j.anbehav.2007.09.024.

601 <sup>50</sup>Bustamante, P., Lahaye, V., Durnez, C., Churlaud, C., Caurant, F. (2006) Total and organic Hg  
602 concentrations in cephalopods from the North Eastern Atlantic waters: Influence of geographical  
603 origin and feeding ecology. *Sci Tot Environ.* 368: 585–596. doi:10.1016/j.scitotenv.2006.01.038.

- <sup>51</sup>Evers, D.C., Burgess, N.M., Champoux, L., Hoskins, B., Major, A., Goodale, W.M., Taylor, R.J., Poppenga, R., Daigle, T. (2005) Patterns and interpretation of mercury exposure in freshwater avian communities in northeastern north America. *Ecotoxicology* 14:193–221. doi:10.1007/s10646-004-6269-7.
- <sup>52</sup>Chastel, O., Fort, J., Ackerman, J.T., Albert, C., Angelier, F., Basu, N., Blévin, P., Brault-Favrou, M., Bustnes, J.O., Bustamante, P., Danielsen, J., Descamps, S., Dietz, R., Erikstad, K.E., Eulaers, I., Ezhov, A., Fleishman, A.B., Gabrielsen, G.W., Gavrilov, M., Gilchrist, G., Gilg, O., Gíslason, S., Golubova, E., Goutte, A., Grémillet, D., Hallgrimsson, G.T., Hansen, E.S., Hanssen, S.A., Hatch, S., Huffeldt, N.P., Jakubas, D., Jónsson, J.E., Kitaysky, A.S., Kolbeinsson, Y., Krasnov, Y., Letcher, R.J., Linnebjerg, J.F., Mallory, M., Merkel, F.R., Moe, B., Montevecchi, W.J., Mosbech, A., Olsen, B., Orben, R.A., Provencher, J.F., Ragnarsdottir, S.B., Reiertsen, T.K., Rojek, N., Romano, M., Søndergaard, J., Strøm, H., Takahashi, A., Tartu, S., Thórarinnsson, T.L., Thiebot, J.B., Will, A.P., Wilson, S., Wojczulanis-Jakubas, K., Yannic, G. (2022) Mercury contamination and potential health risks to Arctic seabirds and shorebirds. *Sci Total Environ.* 844: 156944. doi: 10.1016/j.scitotenv.2022.156944.
- <sup>53</sup>Ackerman, J.T., Eagles-Smith, C.A., Herzog, M.P., Hartman, C.A., Peterson, S.H., Evers, D.C., Jackson, A.K., Elliot, J.E., Vander Pol, S.S., Bryan, C.E. (2016) Avian mercury exposure and toxicological risk across western North America: A synthesis. *Sci Total Environ.* 568:749-769. doi:10.1016/j.scitotenv.2016.03.071.
- <sup>54</sup>Karnovsky, N., Harding, A., Walkusz, W., Kwaśniewski, S., Goszczko, I., Wiktor, J., Routti, H., Bailey, A., McFadden, L., Brown, Z., Beaugrand, G., Grémillet, D. (2010) Foraging distributions of little auks *Alle alle* across the Greenland Sea: implications of present and future Arctic climate change. *Mar Ecol Prog Ser.* 415: 283–293. doi:10.3354/meps08749.
- <sup>55</sup>Amélineau, F., Grémillet, D., Bonnet, D., Le Bot, T., Fort, J. (2016) Where to Forage in the Absence of Sea Ice? Bathymetry as a key factor for an Arctic seabird. *PloS ONE* 11:e0157764. doi:10.1371/journal.pone.0157764.

630 <sup>56</sup>Spreen, G., Kaleschke, L., Heygster, G. (2008) Sea ice remote sensing using AMSR-E 89-GHz  
631 channels. *J Geophys Res.* 113: C02S03. doi:10.1029/2005JC003384.

632 <sup>57</sup>R Core Team (2021) R: A language and environment for statistical computing. R Foundation for  
633 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

634 <sup>58</sup>Lenth, R. (2019) Emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version  
635 1.4.3.01. <https://CRAN.R-project.org/package=emmeans>.

636 <sup>59</sup>Bates, D., Maechler, M., Bolker, B., Walker, S. (2015) Fitting Linear Mixed-Effects Models Using  
637 lme4. *J Stat Soft.* 67:1-48. doi:10.18637/jss.v067.i01.

638 <sup>60</sup>Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B. (2017) “lmerTest Package: Tests in Linear Mixed  
639 Effects Models.” *J Stat Soft.* 82:1-26. doi:10.18637/jss.v082.i13.

640 <sup>61</sup>Regular, P.M., Hedd, A., Montevecchi, W.A., Robertson, G.J., Storey, A.E., Walsh, C.J. (2014) Why  
641 timing is everything: Energetic costs and reproductive consequences of resource mismatch for a  
642 chick-rearing seabird. *Ecosphere* 5:1-13. doi:10.1890/ES14-00182.1.

643 <sup>62</sup>Su, Y., Yajima, M. (2021) R2jags: Using R to Run ‘JAGS’\_. R package version 0.7-1,  
644 <<https://CRAN.R-project.org/package=R2jags>>.

645 <sup>63</sup>Douma, J.C., Weedon, J.T. (2019) Analysing continuous proportions in ecology and evolution: A  
646 practical introduction to beta and Dirichlet regression. *Methods Ecol Evol.* 10:1412-1430.  
647 doi:10.1111/2041-210X.13234.

648 <sup>64</sup>Maier, M.J. (2021) DirichletReg: Dirichlet Regression. R package version 0.7-1.  
649 <https://github.com/maiermarco/DirichletReg>.

650 <sup>65</sup>Lovvorn, J.R., Grebmeier, J.M., Cooper, L.W., Bump, J.K., Richman, S.E. (2009) Modeling marine  
651 protected areas for threatened eiders in a climatically changing Bering Sea. *Ecol Appl.* 19:1596–  
652 1613. doi:10.1890/08-1193.1.

653 <sup>66</sup>Siegel, J.M. (2009) Sleep viewed as a state of adaptive inactivity. *Nat Rev Neurosci.* 10:747–753.  
654 doi:10.1038/nrn2697.

655 <sup>67</sup>Elliott, K.H., Le Vaillant, M., Kato, A., Gaston, A.J., Ropert-Coudert, Y., Hare, J.F., Speakman, J.R.,  
656 Croll, D. (2014) Age-related variation in energy expenditure in a long-lived bird within the  
657 envelope of an energy ceiling. *J Anim Ecol.* 83:136-46. doi:10.1111/1365-2656.12126.

658 <sup>68</sup>Wojczulanis, K., Jakubas, D., Stempniewicz, L. (2005) Changes in the Glaucous Gull predation pressure  
659 on Little Auks in southwest Spitsbergen. *Waterbirds* 28:430-435.

660 <sup>69</sup>Grissot, A., Araya-Salas, M., Jakubas, D., Kidawa, D., Boehnke, R., Błachowiak-Samołyk, K.,  
661 Wojczulanis-Jakubas, K. (2019) Parental Coordination of Chick Provisioning in a Planktivorous  
662 Arctic Seabird Under Divergent Conditions on Foraging Grounds. *Front Ecol Evol.* 7:349.  
663 doi:10.3389/fevo.2019.00349.

664 <sup>70</sup>Courbin, N., Chinho, T., Pichegru, L., Verma-Grémillet, A., Péron, C., Ryan, P.G., Grémillet, D. 2020.  
665 The dance of the Cape gannet may contain social information on foraging behaviour. *Anim*  
666 *Behav.* 166: 95-108. doi:10.1016/j.anbehav.2020.06.012.

667 <sup>71</sup>Jakubas, D., Iliszko, L., Wojczulanis-Jakubas, K., Stempniewicz, L. (2012) Foraging by little auks in the  
668 distant marginal sea ice zone during the chick-rearing period. *Polar Biol.* 35:73–81.  
669 doi:10.1007/s00300-011-1034-x.

670 <sup>72</sup>Fayet, A.L., Clucas, G.V., Anker-Nilssen, T., Syposz, M., Hansen, E.S. (2021) Local prey shortages  
671 drive foraging costs and breeding success in a declining seabird, the Atlantic puffin. *J Anim Ecol.*  
672 90: 1152-1164. doi:10.1111/1365-2656.13442.

673 <sup>73</sup>Boersma, P.D., Rebstock, G.A. (2009) Foraging distance affects reproductive success in Magellanic  
674 penguins. *Mar Ecol Prog Ser.* 375: 263–275. doi:10.3354/meps07753.

675 <sup>74</sup>Jakubas, D., Trudnowska, E., Wojczulanis-Jakubas, K., Iliszko, L., Kidawa, D., Darecki, M.,  
676 Błachowiak-Samołyk, K., Stempniewicz, L. (2013) Foraging closer to the colony leads to faster  
677 growth in little auks. *Mar Ecol Prog Ser.* 489: 263–278. doi:10.3354/meps10414.

678 <sup>75</sup>Kooyman, G.L., Ponganis, P.J. (1998) The physiological basis of diving to depth: birds and mammals.  
679 *Annu Rev Physiol.* 60:19-32. doi:10.1146/annurev.physiol.60.1.19.

- <sup>76</sup>Williams, C.L., Meir, J.U., Ponganis, P.J. (2011) What triggers the aerobic dive limit? Patterns of muscle oxygen depletion during dives of emperor penguins. *J Exp Biol.* 214:1802-1812. doi:10.1242/jeb.052233.
- <sup>77</sup>Piscopo, M., Notariale, R., Tortora, F., Lettieri, G., Palumbo, G., Manna, C. (2020) Novel Insights into Mercury Effects on Hemoglobin and Membrane Proteins in Human Erythrocytes. *Molecules* 25:3278. doi:10.3390/molecules25143278.
- <sup>78</sup>Alazoumi, K.K.M., Ahmed, A., Alamery, S.F., Shamsi, A., Ahmad, B., Islam, A., Farooqi, H. (2021) Effect of Antioxidants on Heavy Metals Induced Conformational Alteration of Cytochrome C and Myoglobin. *Protein Pept Lett.* 28:31-42. doi:10.2174/0929866527666200610134442.
- <sup>79</sup>Auer, S.K., Salin, K., Anderson, G.J., Metcalfe, N.B. (2015) Aerobic scope explains individual variation in feeding capacity. *Biol Lett.* 11:20150793. doi:10.1098/rsbl.2015.0793.
- <sup>80</sup>Enstipp, M.R., Bost, C-A., Le Bohec, C., Bost, C., Laesser, R., Le Maho, Y., Bost, C., Laesser, R., Le Maho, Y., Weimerskirch, H., Handrich, Y. (2019) The dive performance of immature king penguins following their annual molt suggests physiological constraints. *J Exp Biol.* 222:jeb208900. doi:10.1242/jeb.208900.
- <sup>81</sup>Grunst, A.S., Grunst, M.L., Grémillet, D., Kato, A., Bustamante, P., Albert, C., Brisson-Curadeau, É., Clairbaux, M., Cruz-Flores, M., Gentès, S., Perret, S., Ste-Marie, E., Wojczulanis-Jakubas, K., Fort, J. (2022) Data for: Mercury contamination challenges the behavioural response of a keystone species to Arctic climate change. Zenodo: doi:10.5281/zenodo.7185806, Dryad: doi:10.5061/dryad.dbrv15f4t.