

## 5-HT3 Receptor–Mediated Neural Transmission of Cardiorespiratory Modulation by the Nucleus of the Tractus Solitarius

Caroline Sévoz-Couche

### ▶ To cite this version:

 $\label{eq:caroline Sevoz-Couche. 5-HT3 Receptor-Mediated Neural Transmission of Cardiorespiratory Modulation by the Nucleus of the Tractus Solitarius. Paul M Pilowsky. Serotonin: The mediator that spans evolution, pp.349 - 367, 2019, 10.1016/b978-0-12-800050-2.00017-6. hal-03989064$ 

### HAL Id: hal-03989064 https://hal.science/hal-03989064

Submitted on 15 Mar 2023  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### CHAPTER

5-HT<sub>3</sub> Receptor— Mediated Neural Transmission of Cardiorespiratory Modulation by the Nucleus of the Tractus Solitarius

#### **Caroline Sévoz-Couche**

Sorbonne Universités, UPMC Univ Paris 06, INSERM, Paris, France

### 1. INTRODUCTION

Homeostasis can proceed remarkably normally in mammals without functioning cerebral hemispheres, as demonstrated by a study on decerebrate cats maintained for several months [1]. The cats were not only able to walk, eat, and drink, but basic respiratory and cardiovascular functions were normal. More recently, the Working Heart Brainstem Preparation developed for rats [2] and mice [3,4] by Paton in the 1990s showed that the preparation with a viability of 4-6 h produces stable and consistent cardiorespiratory reflex responses to a wide range of visceral stimuli. The brainstem contains many premotor and motor neurons responsible for homeostasis control and thus has the capacity to regulate the maintenance of bodily homeostasis. Within the brainstem, the nucleus of the tractus solitarius (NTS) is of particular interest. Located in the dorsomedial medulla oblongata, it receives visceral afferent information from the cardiovascular, respiratory, gastrointestinal, and taste systems arising from the sensory components of four cranial nerves: facial, trigeminal, glossopharyngeal, and vagus [5]. With the use of neuroanatomical techniques based on the axonal transport of different tracers, evidence has been provided that facial and trigeminal afferents reach the rostral NTS, whereas both glossopharyngeal and vagus nerves project to the intermediate and caudal subregions of the NTS [6,7].

Nearly all the putative neurotransmitters found in the CNS have been identified in the NTS [8], including serotonin [9]. Serotonin immunoreactivity is present in varicose fibers disseminated throughout the rostral to caudal parts of the NTS [10]. There is a general consensus on the essentially extrinsic origin of NTS serotonergic innervation. Serotonin-immunoreactive neurons projecting onto the lateral and medial NTS are mostly found in the caudal raphe group of cells, namely the nucleus raphe magnus and lateral paragigantocellular nucleus (B3), nucleus raphe pallidus (B2), and nucleus raphe obscurus (B1) [11–13]. In addition, the NTS also receives serotonergic afferents from the nodose and petrosal ganglia (the site of vagus and glossopharyngeal nerve cell bodies, respectively) [14,15] and the area postrema [13,16]. Numerous pharmacological, electrophysiological, and immunohistochemical studies have provided evidence for the existence of 5-HT<sub>1</sub> [17], 5-HT<sub>2</sub> [18,19], 5-HT<sub>3</sub> [20,21], 5-HT<sub>4</sub> [22], 5-HT<sub>6</sub> [16], and 5-HT<sub>7</sub> [23] receptors in the NTS.

5-HT<sub>3</sub> receptors are members of the Cys-loop superfamily of ligand-gated ion channels [24] (Fig. 17.1A). 5-HT<sub>3</sub> serotonergic receptors exhibit nearly equal permeability to both Na<sup>+</sup> and K<sup>+</sup>. They are composed of a pentameric complex that surrounds a central ion channel with a high homology with nicotinic acetylcholine receptors. Two 5-HT<sub>3</sub> receptor subunits have been cloned to date, the A subunit  $(5-HT_{3A})$ , which is expressed by both central and peripheral neurons, and the B subunit  $(5-HT_{3B})$  that seems to be restricted to peripheral neurons [20]. When expressed alone, the 5-HT<sub>3B</sub> subunit fails to form functional 5-HT<sub>3</sub> receptors. However, when it is coexpressed with the 5-HT<sub>3A</sub> subunit, the resulting heterometric 5-HT<sub>3</sub> receptor complex fully replicates the biophysical characteristics of native neuronal 5-HT<sub>3</sub> receptors [25]. Only minimal pharmacological differences have been identified between homomeric 5-HT<sub>3A</sub> receptor and heteromeric 5-HT<sub>3A/3B</sub> receptors. In 2003, three human genes encoding novel 5-HT<sub>3C</sub>, 5-HT<sub>3D</sub>, and 5-HT<sub>3E</sub> receptor subunits were reported [26]. The genes for these subunits are located on human chromosome 3q27, whereas those for 5-HT<sub>3A</sub> and 5-HT<sub>3B</sub> are on chromosome 11q23. Further, the reported mRNA expression of the 5-HT<sub>3C-E</sub> subunits tended to show a peripherally restricted pattern, including high levels in the gastrointestinal tract [26]. Intriguingly, their orthologues seem to be absent in rodents [27]. These data suggest that 5-HT<sub>3C-E</sub> subunits are formed for a specific purpose within the peripheral nervous systems in some species, including humans, which is both distinct from the more general functions of receptors formed by 5-HT<sub>3A</sub> and 5-HT<sub>3B</sub>.

In the central nervous system, low levels of 5-HT<sub>3</sub> receptor expression have been demonstrated in the forebrain. Nevertheless, in the majority of species investigated to date, 5-HT<sub>3</sub> receptors were expressed notably in the amygdala and hippocampus [28], but the highest levels of 5-HT<sub>3</sub> receptor—binding sites were found in the NTS [21,29]. The nodose ganglion gives rise to the sensory fibers of the vagus nerve [30]; the majority of NTS 5-HT<sub>3</sub> receptors disappears in nodose ganglionectomized animals, suggesting a presynaptic location of these receptors on vagal afferents [21] (Fig. 17.1B). These afferents are likely to originate from the gastrointestinal tract because subdiaphragmatic vagotomy that removes abdominal visceral afferents of gastric origin abolished binding of [<sup>3</sup>H]granisetron in the NTS [31].



#### FIGURE 17.1 The 5-HT<sub>3</sub> receptor.

(A) 5-HT receptors and their effector systems.  $5-HT_{1/5}$ ,  $5-HT_2$ ,  $5-HT_{4/6/7}$  corresponds to the members of the G protein—coupled receptors and share a putative seven transmembrane structure. The  $5-HT_{1/5}$  and  $5-HT_{4/6/7}$  receptors are negatively and positively coupled to adenylate cyclase (AC) to increase or reduce the formation of cAMP, respectively. The  $5-HT_2$  receptor is coupled positively to phospholipase C (PLC) and is at the origin of diacyl glycerol (DAG) and inositol triphosphate (IP3) synthesis. The  $5-HT_3$  receptor is a ligand-gated ion channel. (B) Autoradiographic localization of [1251] iodozacopride binding to brain sections at the level of the nucleus of the tractus solitarius (NTS), in intact and unilateral ganglionectomized rats.

Adapted from Merahi N, Orer HS, Laporte AM, Gozlan H, Hamon M, Laguzzi R. Baroreceptor reflex inhibition induced by the stimulation of serotonin3 receptors in the nucleus tractus solitarius of the rat. Neuroscience 1992;46:91–100.

# 2. ROLE OF NTS 5-HT<sub>3</sub> RECEPTORS IN THE CARDIOVASCULAR REGULATION

### 2.1 MODULATION OF BLOOD PRESSURE

Whereas microinjection of picomolar doses of 5-HT into the NTS elicits hypotension and bradycardia [32,33], nanomolar doses evoke a dose-dependent increase in blood pressure without any significant change in heart rate [32]. This effect is consequent on the activation of 5-HT<sub>3</sub> receptors because it can be mimicked by local bilateral microinjections of 5-HT<sub>3</sub> receptor agonists, such as 1-(m-chlorophenyl)biguanide (CPBG), phenylbiguanide (PBG), and 2-methyl-serotonin; conversely it is prevented by prior local administration of selective 5-HT<sub>3</sub> receptor antagonists such as granisetron, ondansetron, and zacopride [32,34,35] but not antagonists at other serotonergic receptor types, such as methysergide and ketanserin [35].

#### 2.1.1 At the Level of the NTS

Administration by reverse microdialysis of a 5-HT<sub>3</sub> receptor agonist into the NTS increased the local release of endogenous glutamate from vagal afferents [36]. The ability of granisetron to reduce glutamatergic mEPSCs [37] confirms the observations of Wan and Browning that the spontaneous release of 5-HT in rat brainstem slice preparations treated with TTX and gabazine acts via 5-HT<sub>3</sub> receptors on vagal afferent terminals to release glutamate in the NTS via a presynaptic mechanism [38]. Whereas intra-NTS administration of  $\alpha$ -amino-4-carboxyphenylglycine, a metabotropic glutamate receptor antagonist, did not affect the pressor response to intra-NTS CPBG, this response was suppressed when NTS ionotropic excitatory amino acid (EAA) receptors were blocked by local administration of kynurenic acid or a mixture of 2-amino-5-phosphonopentanoic acid and 6,7-dinitroquinoxaline-2,3-dione, an NMDA- and a non-NMDA-receptor antagonist, respectively [39].

Nitric oxide synthase (NOS) and ionotropic glutamate receptors appear to coexist in NTS neurons [40], and NO is produced downstream of the stimulation of ionotropic glutamate receptors [41]. Moreover, data in the literature support the idea that a functional link exists between the pressor response elicited by microinjection of glutamate in some NTS subnuclei and the NO/cGMP-transducing system [42]. For these reasons, we analyzed the effects of prior intra NTS microinjection of 7-nitroindazole and 1-H-(1,2,4)-oxadiazol-(4,3-A)-quinoxalin, which inhibit NOS and the soluble guanylyl cyclase, respectively, on the pressor response to NTS 5-HT<sub>3</sub>-receptor stimulation in anaesthetized rats [39]. Both pretreatments markedly reduced this response, indicating that the release of glutamate and the resulting activation of ionotropic glutamate receptors and the associated NO/cGMP transduction mechanism, contribute downstream to the pressor effect elicited by 5-HT<sub>3</sub>-receptor stimulation in the NTS.

#### 2.1.2 At the Level of the Ventral Medulla

Activation of 5-HT<sub>3</sub> receptors in the NTS by high (nmol) doses of serotonin increases blood pressure and sympathetic nerve discharge [43,44] (Fig. 17.2A). The



## FIGURE 17.2 Cardiovascular responses to nucleus of the tractus solitarius (NTS) $5-HT_3$ -receptor activation.

(A) Activation of NTS 5-HT<sub>3</sub> receptors by 1-(m-chlorophenyl)-biguanide (CPBG) microinjection increased the sympathetic lumbar nerve discharge (SND) and the mean arterial pressure (MAP). (B) CPBG increased the firing rate of slowest (a) and intermediate (b) clonidine-sensitive, but not fastest clonidine-insensitive (c) cardiovascular sympathoexcitatory cells of the rostroventrolateral medulla (RVLM). Arrow: aortic occlusion. (C) Putative mechanism responsible for the pressor response to 5-HT<sub>3</sub> receptor stimulation in the NTS. NTS 5-HT<sub>3</sub> receptors are located on glutamatergic vagal (gastrointestinal) afferents. Their stimulation promotes the release of glutamate (GLU), which in turn activates ionotropic EAA receptors (NMDA and non-NMDA) and the downstream NO/cGMP transduction system. NTS neurons activated by glutamate send excitatory projections to clonidine-sensitive (but noncatecholaminergic) sympathoexcitatory cells (SE<sub>a</sub> and SE<sub>b</sub>) located in the RVLM. The resulting sympathoexcitatory neurons in the RVLM.

(A and B) Adapted from Sévoz-Couche C, Nosjean A, Franc B, Hamon M, Laguzzi R. Dorsal medullary 5-HT<sub>3</sub> receptors and sympathetic premotor neurones in the rat. J Physiol 1998;508(Pt. 3):747–62.

ventrolateral medulla contains different populations of medulla-spinal neurons that influence sympathetic nerve activity [45,46]. In particular, the rostroventrolateral part of the medulla (RVLM), known as the "pressor" area, contains sympathoexcitatory cells in contact with spinal preganglionic neurons. Intra-NTS administration of CPBG, but not saline, produces a dramatic increase in the density of c-Fos immunoreactive neurons in the RVLM [47]. As expected, this effect was prevented by

prior intra-NTS microinjection of ondansetron [47]. We observed that kynurenic acid administration into the RVLM blocked the pressor response to NTS 5- $HT_3$ -receptor stimulation by CPBG [48], thereby indicating that ionotropic EAA receptors in the RVLM are crucially involved in this response. The pressor effect of NTS 5-HT<sub>3</sub>-receptor activation by CPBG is actually the consequence of the excitation of two different pools of medullo-spinal sympathoexcitatory neurons, i.e., those with the slowest and the intermediate axonal conduction velocity, without affecting those with the highest conduction velocity [44] (Fig. 17.2B). Despite the fact that RVLM neurons sensitive to CPBG were inhibited by i.v. administration of the  $\alpha_2$ -adrenergic receptor agonist clonidine [44], the possibility that a monosynaptic pathway from the NTS to the sympathoexcitatory adrenergic neurons of the RVLM might convey the pressor message triggered by NTS 5-HT<sub>3</sub>-receptor stimulation is unlikely, because if an increased number of c-fos immunoreactive cells in the RVLM was observed after intra-NTS microinjection of CPBG, these cells were not double-stained for c-fos and tyrosine hydroxylase [49]. Thus the pressor response to NTS 5-HT<sub>3</sub>-receptor stimulation is associated with the excitation of noncatecholaminergic neurons within the RVLM. A summary of the NTS-RVLM pathway involved in the pressor effect of 5-HT<sub>3</sub>-receptor agonist is presented in Fig. 17.2C.

### 2.2 MODULATION OF REFLEX BRADYCARDIA

Vagal NTS second-order neurons afferents from aortic and carotid baroreceptors (sensitive to blood pressure modifications), atrial cardiopulmonary receptors (sensitive to diverse chemical substances), and carotid chemoreceptors (sensitive to changes in blood gas concentrations) contact cardiac premotor neurons located within the nucleus ambiguus to switch cardiovagal (bradycardia) reflex responses [50]. Intra-NTS administration of nanomolar doses of 5-HT or 5-HT<sub>3</sub>-receptor agonists, in anaesthetized as well as unanaesthetized rats, inhibits the cardiac component of the three aforementioned cardiovascular reflexes. This effect is specific because it can be blocked by prior intra-NTS administration of 5-HT<sub>3</sub>-receptor antagonists but remains unchanged after local administration of antagonists acting at other receptor types [32,35,51,52]. These data clearly show that NTS 5-HT<sub>3</sub> receptors mediate a general 5-HT inhibitory influence on the parasympathetic component of the reflex control of heart rate. Specific activation of 5-HT<sub>3</sub> receptors results in clear-cut excitation of most NTS cells [17]. Therefore, an inhibitory effect on the reflex cardiac responses could have been the consequence, at least in part, of a presynaptic inhibition. However, microinjections of EAA-receptor antagonists into the NTS block both the baroreflex- and Bezold-Jarisch-evoked reflex bradycardia [32,53]. Another possibility was that the EAA released in the NTS under 5-HT<sub>3</sub>-receptor activation could, directly or indirectly, activate GABAergic interneurons. Accordingly, the inhibitory effect of intra-NTS microinjection of 5-HT<sub>3</sub>-receptor agonists on the bradycardic responses to baroreflex, chemoreflex, or Bezold–Jarisch reflex activation could be prevented by prior local microinjection of low doses of bicuculline [32,35,51].

### 2.3 ROLE OF THE NTS 5-HT<sub>3</sub> RECEPTOR IN THE MODULATION OF STRESS-INDUCED CARDIOVASCULAR RESPONSES

Hypertension and reduced baroreflex cardiac response is also observed during acute stressful conditions such as the defense reaction. Various hypothalamic areas were originally defined as defense areas (HDA) because electrical or chemical stimulation produced characteristic behavioral and physiological defense responses [54–56]. Among the various HDA, the dorsomedial nucleus of the hypothalamus (DMH) appears to be of particular interest [57,58]. Other key regions involved in the defense reaction were subsequently described, such as the dorsolateral periaqueductal gray (dlPAG) in the midbrain [59,60]. In previous studies, we confirmed that intense dlPAG and DMH activation induced a marked increase in mean blood pressure and a decrease in parasympathetic activity [61,62]. To establish whether endogenous 5-HT actually occurs in animals expressing this type of behavioral reaction, we analyzed the effects of dlPAG stimulation on the baroreflex bradycardia response in rats pretreated with p-chlorophenylalanine (PCPA), a serotonin synthesis inhibitor. We showed that inhibition of 5-HT synthesis almost totally prevented the inhibitory effects of dlPAG stimulation on the baroreflex bradycardia [63]. Moreover, the role of 5-HT in the inhibitory influence of dlPAG stimulation was confirmed by the fact that 5-hydroxytryptophan administration, designed to restore 5-HT synthesis in PCPA-pretreated rats, allowed recovery of this negative control of the cardiac reflex response to nearly the same level as that observed in naive rats. The involvement of NTS 5-HT<sub>3</sub> receptors during acute stress was assessed by the fact that intra-NTS microinjection of granisetron drastically reduced the inhibitory effects of dlPAG and DMH stimulation on baroreceptor sensitivity [62,64]. Similarly, a study by Gau et al. showed that thermal and mechanical nociceptions induced inhibition of the baroreflex bradycardia via activation of 5-HT<sub>3</sub> receptors in the NTS [65]. Interestingly, data indicated that:

- intra-NTS GR205171, a selective NK<sub>1</sub>-receptor antagonist, reversed the cardiac baroreflex (and cardiopulmonary, unpublished data) inhibition produced by dlPAG activation or local administration of PBG [64], and
- 2. intra-NTS bicuculline reversed the baroreflex (and cardiopulmonary, personal data) bradycardia inhibition produced by local administration of substance P [64]. These data indicate that NK<sub>1</sub> and GABA<sub>A</sub> receptors may contribute downstream to the 5-HT<sub>3</sub> receptor—mediated inhibition of the reflex brady-cardia. It is noteworthy that neither microinjection of granisetron into the NTS nor PCPA treatment affected the sympathetically mediated increase in blood pressure induced by DMH and dlPAG stimulation [62–64]. Taken together, the data indicate that another mechanism is at the origin of this hypertensive stress response. Thus the physiological basis for the pressor effect of NTS 5-HT<sub>3</sub>-receptor activation remains unknown.

The origin of serotonin released into the NTS during the defense reaction is likely to be one of the B1–B3 cell groups. Two studies have shown that the nucleus raphe obscurus and the nucleus raphe pallidus (B1 and B2, respectively) are both involved in inhibition of baroreflex bradycardia [22,66]. However, we found that a large majority of serotonergic cells in the B3 raphe (raphe magnus and lateral paragigantocellular nuclei) but not in other raphe nuclei were excited (as confirmed by the presence of c-fos protein) following DMH as well as dlPAG stimulation [61,67] (Fig. 17.3A), or after application of nociceptive stimuli [65] (Fig. 17.3B). In addition,

- 1. the reduction in baroreflex bradycardia induced by dlPAG activation was reduced by muscimol injection into the B3 region [67], and
- **2.** the activation of the B3 region induced an inhibitory effect on aortic baroreflex bradycardia, which was prevented by intra-NTS granisetron [67] (Fig. 17.3C).

Decades of clinical, epidemiological, and experimental research in animals and humans has provided strong evidence in support of a close correlation between psychosocial factors and cardiovascular abnormalities [68-70]. These psychosocial variables appear to be independent risk factors, as important as traditional risk factors (serum cholesterol, body mass index, and poor physical activity), for the onset and progression of hypertension and arrhythmias [71]. The increased arrhythmogenic feature in stress-related disorders was commonly linked to exaggerated sympathetic stimulation, as reflected by increased blood pressure, heart rate, and plasma catecholamines. Recently, the prospective study Autonomic Tone and Reflexes After Myocardial Infarction demonstrated that markers of reduced vagal activity, such as depressed baroreflex bradycardic response and high-frequency domain (vagally mediated) of the heart rate variability (HRV), are strong predictors of cardiac mortality after myocardial infarction [72,73]. A large proportion of sudden cardiac death is correlated with a low baroreflex response in patients taking, or not taking, beta blockers [74]. Cardiac sympathetic excitation is associated with low vagal tone and reciprocal vagal baroreflex sensitivity impairment in many stress disorders, including depression [75] and anxiety [69,76]. Interestingly, unmedicated females are especially affected by depression and present a lower reduction in baroreflex sensitivity (BRS) than males [77,78]. Therefore the prevention of reduced vagal cardiac response may also be a target of choice to reduce arrhythmias, but the origin of this dysfunction is still unclear. However, it was possible that similar neurocircuitry to that underlying cardiac alteration during the defense reaction may occur during chronic stress. Using a social defeat paradigm triggering arrhythmias [79], we showed that vagally mediated high-frequency band (short-term regulation) of the HRV [80] and baroreflex bradycardia were lower in defeated animals [81]. Blockade of the DMH by muscimol or NTS 5-HT<sub>3</sub> receptors by local microinjection of granisetron prevented both modifications. It is interesting that granisetron did not block the long-term increase in blood pressure. Taken together, the results suggest that the reduction of the cardiac baroreflex (but not the hypertension) in animals with social stress-induced anxiety is produced by chronic activation of 5-HT<sub>3</sub> receptors in



FIGURE 17.3 The source of serotonin release in the nucleus of the tractus solitarius (NTS) after stress induction.

(A) Color-digitized photomicrographs showing the increase of neurons labeled in the lateral paragigantocellular nucleus for c-Fos+ 5-HT (d) after dorsolateral periaqueductal gray (A2) compared to sham (A1). (f) neurons labeled for c-Fos, (s) neurons labeled for 5-HT alone. (B) Schematic drawing of coronal sections through the B3 region, showing

the NTS by serotonin from the medullary caudal raphe, from the DMH and the dlPAG. Interestingly, systemic granisetron (a drug that crosses the blood-brain barrier and has a high safety profile) also prevented the low BRS in defeated rats, suggesting a potential role for this substance as a preventive or curative treatment against reduced vagal activity (and therefore possibly against arrhythmias) in chronic stress but also in all pathologies linked to dysautonomia.

# 3. ROLE OF NTS 5-HT<sub>3</sub> RECEPTORS IN THE RESPIRATORY REGULATION DURING STRESS

There is an important reciprocal link between breathing and anxiety [82]. Thus, respiratory distress and asphyxia cause sensation of dread. On the other hand, fear and anxiety can have profound effects on breathing. Clinical studies show that panic disorder, which is characterized by acute and unexpected anxiety attacks and substantial anxiety over the possibility of experiencing further attacks, is associated with symptoms that include palpitations, shortness of breath, sweating, and hyperventilation [83]. In addition, high levels of anxiety-related behavior in rats are associated with an elevation of the resting respiratory rate [84]. Comparatively less is known about the long-term effects of emotional stress on breathing. In adult rats, a strong neonatal emotional stress such as maternal separation leads to a decrease in breathing rate during non-REM sleep [85]. A lower breathing rate was also noted under anesthesia in the Flinder-Sensitive rat, a well-validated animal model of depression [86]. However, in patients with mood-depression disorders, breathing rate does not seem to be altered, although cardiovascular changes do occur [87]. A study on the long-term effects of social defeat on breathing was conducted recently. Peripheral and central respiratory and cardiovascular regulatory mechanisms are tightly coupled. An element of this interaction is the respiratory sinus arrhythmia (RSA),

that neurons activated during noxious application, recorded and juxtacellularly filled, are extensively found in the lateral paragigantocellular nucleus. (C) Representative tracings showing that the baroreflex bradycardia evoked by aortic stimulation (AS) was reduced (control, left) by chemical (DLH, 0.3 M) activation of dPAG (C1) or B3 (C3). Pretreatment with muscimol (5 mM) into the B3 region (C2) or granisetron (2.5 mM) into the NTS (C4) prevented these inhibitory effects, respectively.

(A) Adapted from Bernard J-F, Netzer F, Gau R, Hamon M, Laguzzi R, Sévoz-Couche C. Critical role of B3 serotonergic cells in baroreflex inhibition during the defense reaction triggered by dorsal periaqueductal gray stimulation. J Comp Neurol 2008;506:108–21. (B) Adapted from Gau R, Sévoz-Couche C, Hamon M, Bernard

J-F. Noxious stimulation excites serotonergic neurons: a comparison between the lateral paragigantocellular reticular and the raphe magnus nuclei. Pain 2013;154:647–59. (C) Adapted from Bernard J-F, Netzer F, Gau R, Hamon M, Laguzzi R, Sévoz-Couche C. Critical role of B3 serotonergic cells in baroreflex inhibition during the defense reaction triggered by dorsal periaqueductal gray stimulation. J Comp Neurol 2008;506:108–21. which reflects the regular increase in heart rate during inspiration and its decrease during expiration [88]. RSA appears in the high frequency range at which the parasympathetic but not the sympathetic branch of the autonomic nervous system can respond to respiration and influence heart rate. As such, RSA peak frequency seen in HRV analysis (fast Fourier transformation of the cardiac pulse interval signal) is a reliable method for extracting breathing frequency in both the anaesthetized and conscious animal, as it is comparable to tracheal and pleural respiratory rates in both conditions, during low or high activity [89]. Telemetric recordings in social defeat demonstrated a lower RSA frequency (i.e., a lower respiratory rate) in stressed rats, during and after (that effect lasted at least 10 days) the stress procedure [90]. To find out whether the DMH and NTS 5-HT<sub>3</sub> receptors were not only involved in the reduction of the BRS but also in the bradypnea evoked by social defeat, we blocked the DMH with bilateral microinjections of muscimol and antagonized 5- $HT_3$  NTS receptors with bilateral microinjections of granisetron. As with cardiac vagal inhibition, local blockade of DMH and 5-HT<sub>3</sub> NTS receptors abolished the bradypnea of defeated rats. This bradypnea does not appear to be associated with increased tidal volume because pleural pressure seems to be unaffected in defeated rats (unpublished observation).

A complex network of lower brainstem neurons, termed the respiratory central pattern generator (CPG), generates the respiratory rhythm and defines the onset and offset of the various respiratory motor outflows and regulates the frequency and intensity of these periodic outflows [91]. It receives central inputs, and in particular participates in the hyperventilatory response seen during acute stress and intense activation of the DMH [92]. The chemical drive to breathe also relies on central chemoreceptors that detect brain extracellular fluid  $PCO_2$  via pH and on carotid body chemoreceptors that respond to low arterial  $PO_2$ -dependent manner [93]. The retrotrapezoid nucleus (RTN) contains very superficial propriobulbar neurons that have properties consistent with such specialized chemoreceptors [94] and is an important site of integration between central and peripheral chemoreception [95]. The RTN is a key source of tonic excitatory drive to both rhythm-generating and pattern-generating components of the CPG [96,97] and as such is responsible for hyperventilatory responses to central and peripheral chemoreflexes. Interestingly, the commissural NTS neurons that are activated by carotid body (peripheral chemoreceptor) stimulation innervate the RTN neurons and this projection is predominantly glutamatergic [95], as:

- 1. chemodenervation reduces baseline breathing rate [98], and
- **2.** 5-HT<sub>3</sub> receptor—activation blocks the carotid chemoreflex bradycardia and induces stress-induced long-term bradypnea [90].

It is possible that DMH activation provokes long-term respiratory reduction through NTS 5-HT<sub>3</sub> activation—induced inhibition of the peripheral chemoreflex hyperventilation. In support of this hypothesis, minimal subthreshold activation of the DMH negatively modulates this reflex respiratory response [99,100]. We suggest that in acute stress, when respiratory values are already high because of intense



### FIGURE 17.4 Putative neurocircuitry underlying the stress-induced long-term cardiorespiratory alteration.

NTS second-order aortic baroreflex and atrial cardiopulmonary neurons activate preganglionary cells in the nucleus ambiguus (NAmb) to induce a reflex bradycardia. Carotid chemoreflex neurons project massively to the NTS, to produce a reflex bradycardia via NAmb activation, and a reflex tachypnea through activation of the retrotrapezoid nucleus (RTN) in close relationship with the central pattern generator (CPG) of the respiration. Activation of the dorsomedial nucleus of the hypothalamus (DMH) activate efferents projecting to the dIPAG that may be at the origin of serotonin release into the NTS (presumably from the B3 region) to ultimately activate presynaptic 5-HT<sub>3</sub> receptors on vagal afferents; activation of these receptors induces the presynaptic release of glutamate that stimulates: (1) substance P interneurones, which activate GABAergic interneurones (black cell) endowed with NK<sub>1</sub> receptors. GABA will act on GABA<sub>A</sub> receptors present on NTS second-order neurons to inhibit the production of reflex bradycardia (green way); (2) GABAergic interneurones (black cell), and GABA will act on GABA<sub>A</sub> receptors present on second-order neurons to inhibit the production of the carotid chemoreflex tachypnea (blue way).

activation of neurons in the DMH, peripheral chemoreflex inhibition by NTS 5- $HT_3$ -receptor activation helps to prevent an additional increase in ventilation. In the longer term, the defense zone remains only mildly activated and is not able to increase the respiration by its direct action on CPG, and bradypnea appears due to chronic chemoreflex respiratory reflex response inhibition.

### 4. CONCLUSION

Activation of NTS 5-HT<sub>3</sub> receptors produces activation of three modulatory circuits:

- the first is at the origin of an increase in blood pressure through an NO/GMPc activation of NTS second-order neurons in contact with clonidine-sensitive sympathoexcitatory cells located in the RVLM,
- **2.** the second is at the origin of a reduction of reflex bradycardia, through a substance P-GABAergic inhibition of NTS second-order neurons in contact with cardiac premotor neurons in the nucleus ambiguus,
- **3.** the third is at the origin of a bradypnea, presumably through a GABAergic inhibition of NTS second-order neurons in contact with chemosensitive cells in the RTN.

The second and third pathways are activated by a hypothalamo-midbrain circuitry after stress induction. A suggested neurocircuitry underlying stress-induced cardiorespiratory alteration is shown in Fig. 17.4. We showed that systemic treatment with granisetron was able to prevent these alterations [81,90]. Thus, in patients with high anxiety scores and in patients with induced autonomic dysfunction, as observed after ischemic stroke for example [101], systemic treatment with the 5-HT<sub>3</sub> receptor antagonist granisetron—a potent antiemetic [102] with a highly safe profile [103]—could be used to restore parasympathetic activity and, thus, reduce the likelihood of adverse cardiac events. In addition, this treatment may also help to prevent alveolar hypoventilation.

### REFERENCES

- [1] Bazett HC, Penfield WG. A study of the sherrington decerebrate animal in the chronic as well as the acute condition. Brain 1922;45:185–265.
- [2] Potts JT, Spyer KM, Paton JF. Somatosympathetic reflex in a working heart-brainstem preparation of the rat. Brain Res Bull 2000;53:59–67.
- [3] Paton JF. The ventral medullary respiratory network of the mature mouse studied in a working heart-brainstem preparation. J. Physiol 1996;493(Pt 3):819–31.
- [4] Paton JF, Butcher JW. Cardiorespiratory reflexes in mice. J Auton Nerv Syst 1998;68: 115-24.

### **362 CHAPTER 17** 5-HT<sub>3</sub> Receptor—Mediated Neural

- [5] Contreras RJ, Beckstead RM, Norgren R. The central projections of the trigeminal, facial, glossopharyngeal and vagus nerves: an autoradiographic study in the rat. J Auton Nerv Syst 1982;6:303–22.
- [6] Nomura S, Mizuno N. Central distribution of afferent and efferent components of the glossopharyngeal nerve: an HRP study in the cat. Brain Res 1982;236:1–13.
- [7] Nomura S, Mizuno N. Central distribution of efferent and afferent components of the cervical branches of the vagus nerve. A HRP study in the cat. Anat Embryol (Berl) 1983;166:1–18.
- [8] van Giersbergen PL, Palkovits M, De Jong W. Involvement of neurotransmitters in the nucleus tractus solitarii in cardiovascular regulation. Physiol Rev 1992;72:789–824.
- [9] Steinbusch HW, Nieuwenhuys R. Localization of serotonin-like immunoreactivity in the central nervous system and pituitary of the rat, with special references to the innervation of the hypothalamus. Adv Exp Med Biol 1981;133:7–35.
- [10] Maley B, Elde R. Immunohistochemical localization of putative neurotransmitters within the feline nucleus tractus solitarii. Neuroscience 1982;7:2469–90.
- [11] Pickel VM, Joh TH, Chan J, Beaudet A. Serotoninergic terminals: ultrastructure and synaptic interaction with catecholamine-containing neurons in the medial nuclei of the solitary tracts. J Comp Neurol 1984;225:291–301.
- [12] Schaffar N, Kessler JP, Bosler O, Jean A. Central serotonergic projections to the nucleus tractus solitarii: evidence from a double labeling study in the rat. Neuroscience 1988;26:951–8.
- [13] Thor KB, Helke CJ. Serotonin- and substance P-containing projections to the nucleus tractus solitarii of the rat. J Comp Neurol 1987;265:275–93.
- [14] Helke CJ, O'Donohue TL, Jacobowitz DM. Substance P as a baro- and chemoreceptor afferent neurotransmitter: immunocytochemical and neurochemical evidence in the rat. Peptides 1980;1:1–9.
- [15] Nosjean A, Compoint C, Buisseret-Delmas C, Orer HS, Merahi N, Puizillout JJ, Laguzzi R. Serotonergic projections from the nodose ganglia to the nucleus tractus solitarius: an immunohistochemical and double labeling study in the rat. Neurosci Lett 1990;114:22–6.
- [16] Garfield AS, Burke LK, Shaw J, Evans ML, Heisler LK. Distribution of cells responsive to 5-HT<sub>6</sub> receptor antagonist-induced hypophagia. Behav Brain Res 2014;266: 201−6.
- [17] Wang Y, Ramage AG, Jordan D. In vivo effects of 5-hydroxytryptamine receptor activation on rat nucleus tractus solitarius neurones excited by vagal C-fibre afferents. Neuropharmacology 1997;36:489–98.
- [18] Comet M-A, Bernard JF, Hamon M, Laguzzi R, Sévoz-Couche C. Activation of nucleus tractus solitarius 5-HT2A but not other 5-HT2 receptor subtypes inhibits the sympathetic activity in rats. Eur J Neurosci 2007;26:345–54.
- [19] Huang J, Pickel VM. Differential distribution of 5HT2A and NMDA receptors in single cells within the rat medial nucleus of the solitary tract. Synap NY 2002;44:64–75.
- [20] Morales M, Wang S-D. Differential composition of 5-hydroxytryptamine3 receptors synthesized in the rat CNS and peripheral nervous system. J Neurosci 2002;22: 6732–41.
- [21] Pratt GD, Bowery NG, Kilpatrick GJ, Leslie RA, Barnes NM, Naylor RJ, Jones BJ, Nelson DR, Palacids JM, Slater P. Consensus meeting agrees distribution of 5-HT3 receptors in mammalian hindbrain. Trends Pharmacol Sci 1990;11:135–7.

- [22] Edwards E, Paton JF. 5-HT(4) receptors in nucleus tractus solitarii attenuate cardiopulmonary reflex in anesthetized rats. Am J Physiol 1999;277:H1914–23.
- [23] Oskutyte D, Jordan D, Ramage AG. Evidence that 5-hydroxytryptamine(7) receptors play a role in the mediation of afferent transmission within the nucleus tractus solitarius in anaesthetized rats. Br J Pharmacol 2009;158:1387–94.
- [24] Lambert JJ, Peters JA, Hales TG, Dempster J. The properties of 5-HT3 receptors in clonal cell lines studied by patch-clamp techniques. Br J Pharmacol 1989;97:27–40.
- [25] Barnes NM, Sharp T. A review of central 5-HT receptors and their function. Neuropharmacology 1999;38:1083–152.
- [26] Niesler B, Frank B, Kapeller J, Rappold GA. Cloning, physical mapping and expression analysis of the human 5-HT3 serotonin receptor-like genes HTR3C, HTR3D and HTR3E. Gene 2003;310:101–11.
- [27] Karnovsky AM, Gotow LF, McKinley DD, Piechan JL, Ruble CL, Mills CJ, Schellin KAB, Slightom JL, Fitzgerald LR, Benjamin CW, et al. A cluster of novel serotonin receptor 3-like genes on human chromosome 3. Gene 2003;319:137–48.
- [28] Parker RM, Barnes JM, Ge J, Barber PC, Barnes NM. Autoradiographic distribution of [3H]-(S)-zacopride-labelled 5-HT3 receptors in human brain. J Neurol Sci 1996;144: 119–27.
- [29] Laporte AM, Koscielniak T, Ponchant M, Vergé D, Hamon M, Gozlan H. Quantitative autoradiographic mapping of 5-HT3 receptors in the rat CNS using [<sup>125</sup>I]iodozacopride and [<sup>3</sup>H]zacopride as radioligands. Synap. NY 1992;10:271–81.
- [30] Goehler LE, Gaykema RP, Hansen MK, Anderson K, Maier SF, Watkins LR. Vagal immune-to-brain communication: a visceral chemosensory pathway. Auton. Neurosci. Basic Clin 2000;85:49–59.
- [31] Leslie RA, Reynolds DJ, Andrews PL, Grahame-Smith DG, Davis CJ, Harvey JM. Evidence for presynaptic 5-hydroxytryptamine3 recognition sites on vagal afferent terminals in the brainstem of the ferret. Neuroscience 1990;38:667–73.
- [32] Merahi N, Orer HS, Laporte AM, Gozlan H, Hamon M, Laguzzi R. Baroreceptor reflex inhibition induced by the stimulation of serotonin3 receptors in the nucleus tractus solitarius of the rat. Neuroscience 1992;46:91–100.
- [33] Raul L. Serotonin2 receptors in the nucleus tractus solitarius: characterization and role in the baroreceptor reflex arc. Cell Mol Neurobiol 2003;23:709–26.
- [34] Callera JC, Bonagamba LG, Sévoz C, Laguzzi R, Machado BH. Cardiovascular effects of microinjection of low doses of serotonin into the NTS of unanesthetized rats. Am J Physiol 1997a;272:R1135–42.
- [35] Sévoz C, Callera JC, Machado BH, Hamon M, Laguzzi R. Role of serotonin3 receptors in the nucleus tractus solitarii on the carotid chemoreflex. Am J Physiol 1997;272: H1250–9.
- [36] Ashworth-Preece MA, Jarrott B, Lawrence AJ. 5-Hydroxytryptamine3 receptor modulation of excitatory amino acid release in the rat nucleus tractus solitarius. Neurosci Lett 1995;191:75–8.
- [37] Sanger GJ, Nelson DR. Selective and functional 5-hydroxytryptamine3 receptor antagonism by BRL 43694 (granisetron). Eur J Pharmacol 1989;159:113–24.
- [38] Wan S, Browning KN. Glucose increases synaptic transmission from vagal afferent central nerve terminals via modulation of 5-HT3 receptors. Am J Physiol Gastrointest Liver Physiol 2008;295:G1050-7.

### **364 CHAPTER 17** 5-HT<sub>3</sub> Receptor–Mediated Neural

- [39] Sévoz-Couche C, Maisonneuve B, Hamon M, Laguzzi R. Glutamate and NO mediation of the pressor response to 5-HT3 receptor stimulation in the nucleus tractus solitarii. Neuroreport 2002;13:837–41.
- [40] Lin LH, Talman WT. Colocalization of GluR1 and neuronal nitric oxide synthase in rat nucleus tractus solitarii neurons. Neuroscience 2001;106:801–9.
- [41] Lin LH, Talman WT. N-methyl-D-aspartate receptors on neurons that synthesize nitric oxide in rat nucleus tractus solitarii. Neuroscience 2000;100:581–8.
- [42] Ogawa H, Mizusawa A, Kikuchi Y, Hida W, Miki H, Shirato K. Nitric oxide as a retrograde messenger in the nucleus tractus solitarii of rats during hypoxia. J. Physiol 1995; 486(Pt 2):495–504.
- [43] Nosjean A, Franc B, Laguzzi R. Increased sympathetic nerve discharge without alteration in the sympathetic baroreflex response by serotonin3 receptor stimulation in the nucleus tractus solitarius of the rat. Neurosci Lett 1995;186:41-4.
- [44] Sévoz-Couche C, Nosjean A, Franc B, Hamon M, Laguzzi R. Dorsal medullary 5-HT3 receptors and sympathetic premotor neurones in the rat. J. Physiol 1998;508(Pt 3): 747–62.
- [45] Spyer KM. Annual review prize lecture. Central nervous mechanisms contributing to cardiovascular control. J. Physiol 1994;474:1–19.
- [46] Sun MK, Guyenet PG. Arterial baroreceptor and vagal inputs to sympathoexcitatory neurons in rat medulla. Am J Physiol 1987;252:R699–709.
- [47] Nosjean A, Hamon M, Laguzzi R. c-Fos induction in the rostroventrolateral medulla by 5-HT3 receptor activation in the nucleus tractus solitarius. Neuroreport 1998;9: 373–8.
- [48] Sévoz C, Hamon M, Laguzzi R. Medullary pathways of cardiovascular responses to 5-HT2 and 5-HT3 receptor stimulation in the rat nucleus tractus solitarius. Neuroreport 1996a;7:1965–9.
- [49] Nosjean A, Callera JC, Bonagamba L, Machado B, Hamon M, Laguzzi R. Serotonin(3) receptor stimulation in the nucleus tractus solitarii activates noncatecholaminergic neurons in the rat ventrolateral medulla. Neuroscience 2002;112: 935–49.
- [50] Chen HI, Chai CY. Integration of the cardiovagal mechanism in the medulla oblongata of the cat. Am J Physiol 1976;231:454–61.
- [51] Callera JC, Sévoz C, Laguzzi R, Machado BH. Microinjection of a serotonin3 receptor agonist into the NTS of unanesthetized rats inhibits the bradycardia evoked by activation of the baro- and chemoreflexes. J Auton Nerv Syst 1997b;63:127–36.
- [52] Sévoz C, Nosjean A, Callera JC, Machado B, Hamon M, Laguzzi R. Stimulation of 5-HT3 receptors in the NTS inhibits the cardiac Bezold-Jarisch reflex response. Am J Physiol 1996b;271:H80–7.
- [53] Verberne AJ, Guyenet PG. Medullary pathway of the Bezold-Jarisch reflex in the rat. Am J Physiol 1992;263:R1195–202.
- [54] Coote JH, Hilton SM, Perez-Gonzalez JF. Inhibition of the baroreceptor reflex on stimulation in the brain stem defence centre. J. Physiol 1979;288:549–60.
- [55] Gebber GL, Snyder DW. Hypothalamic control of baroreceptor reflexes. Am J Physiol 1970;218:124–31.
- [56] Nosaka S. Modifications of arterial baroreflexes: obligatory roles in cardiovascular regulation in stress and poststress recovery. Jpn J Physiol 1996;46:271–88.

- [57] DiMicco JA, Samuels BC, Zaretskaia MV, Zaretsky DV. The dorsomedial hypothalamus and the response to stress: part renaissance, part revolution. Pharmacol. Biochem. Behav 2002;71:469–80.
- [58] Stotz-Potter EH, Willis LR, DiMicco JA. Muscimol acts in dorsomedial but not paraventricular hypothalamic nucleus to suppress cardiovascular effects of stress. J Neurosci 1996;16:1173–9.
- [59] Bandler R, Shipley MT. Columnar organization in the midbrain periaqueductal gray: modules for emotional expression? Trends Neurosci 1994;17:379–89.
- [60] Carrive P, Bandler R, Dampney RA. Anatomical evidence that hypertension associated with the defence reaction in the cat is mediated by a direct projection from a restricted portion of the midbrain periaqueductal grey to the subretrofacial nucleus of the medulla. Brain Res 1988;460:339-45.
- [61] Netzer F, Bernard J-F, Verberne AJM, Hamon M, Camus F, Benoliel J-J, Sévoz-Couche C. Brain circuits mediating baroreflex bradycardia inhibition in rats: an anatomical and functional link between the cuneiform nucleus and the periaqueductal grey. J. Physiol 2011;589:2079–91.
- [62] Sévoz-Couche C, Comet M-A, Hamon M, Laguzzi R. Role of nucleus tractus solitarius 5-HT3 receptors in the defense reaction-induced inhibition of the aortic baroreflex in rats. J. Neurophysiol 2003;90:2521–30.
- [63] Comet M-A, Sévoz-Couche C, Hanoun N, Hamon M, Laguzzi R. 5-HT-mediated inhibition of cardiovagal baroreceptor reflex response during defense reaction in the rat. Am J Physiol Heart Circ Physiol 2004;287:H1641–9.
- [64] Comet M-A, Laguzzi R, Hamon M, Sévoz-Couche C. Functional interaction between nucleus tractus solitarius NK1 and 5-HT3 receptors in the inhibition of baroreflex in rats. Cardiovasc Res 2005;65:930–9.
- [65] Gau R, Sévoz-Couche C, Hamon M, Bernard J-F. Noxious stimulation excites serotonergic neurons: a comparison between the lateral paragigantocellular reticular and the raphe magnus nuclei. Pain 2013;154:647–59.
- [66] Weissheimer KV, Machado BH. Inhibitory modulation of chemoreflex bradycardia by stimulation of the nucleus raphe obscurus is mediated by 5-HT3 receptors in the NTS of awake rats. Auton. Neurosci. Basic Clin 2007;132:27–36.
- [67] Bernard J-F, Netzer F, Gau R, Hamon M, Laguzzi R, Sévoz-Couche C. Critical role of B3 serotonergic cells in baroreflex inhibition during the defense reaction triggered by dorsal periaqueductal gray stimulation. J Comp Neurol 2008;506:108–21.
- [68] Newton JE, Chapin JL, Murphree OD. Correlations of normality and nervousness with cardiovascular functions in pointer dogs. Pavlov. J. Biol. Sci. 1976;11:105–20.
- [69] Watkins LL, Grossman P, Krishnan R, Sherwood A. Anxiety and vagal control of heart rate. Psychosom Med 1998;60:498–502.
- [70] Weiss E. Psychosomatic aspects of cardiovascular disease. Clinics 1946;5:263-75.
- [71] Hemingway H, Malik M, Marmot M. Social and psychosocial influences on sudden cardiac death, ventricular arrhythmia and cardiac autonomic function. Eur Heart J 2001;22:1082–101.
- [72] La Rovere MT, Pinna GD, Hohnloser SH, Marcus FI, Mortara A, Nohara R, Bigger JT, Camm AJ, Schwartz PJ, ATRAMI Investigators Autonomic Tone, Reflexes After Myocardial Infarcton. Baroreflex sensitivity and heart rate variability in the identification of patients at risk for life-threatening arrhythmias: implications for clinical trials. Circulation 2001;103:2072–7.

### **366 CHAPTER 17** 5-HT<sub>3</sub> Receptor–Mediated Neural

- [73] La Rovere MT, Pinna GD, Maestri R, Mortara A, Capomolla S, Febo O, Ferrari R, Franchini M, Gnemmi M, Opasich C, et al. Short-term heart rate variability strongly predicts sudden cardiac death in chronic heart failure patients. Circulation 2003; 107:565–70.
- [74] La Rovere MT, Pinna GD, Maestri R, Robbi E, Caporotondi A, Guazzotti G, Sleight P, Febo O. Prognostic implications of baroreflex sensitivity in heart failure patients in the beta-blocking era. J Am Coll Cardiol 2009;53:193–9.
- [75] Davydov DM, Shapiro D, Cook IA, Goldstein I. Baroreflex mechanisms in major depression. Prog Neuropsychopharmacol Biol Psychiatry 2007;31:164–77.
- [76] Watkins LL, Blumenthal JA, Carney RM. Association of anxiety with reduced baroreflex cardiac control in patients after acute myocardial infarction. Am Heart J 2002;143: 460–6.
- [77] Voss A, Boettger MK, Schulz S, Gross K, Bär K-J. Gender-dependent impact of major depression on autonomic cardiovascular modulation. Prog Neuropsychopharmacol Biol Psychiatry 2011;35:1131–8.
- [78] Wittstein IS, Thiemann DR, Lima JAC, Baughman KL, Schulman SP, Gerstenblith G, Wu KC, Rade JJ, Bivalacqua TJ, Champion HC. Neurohumoral features of myocardial stunning due to sudden emotional stress. N Engl J Med 2005;352:539–48.
- [79] Sgoifo A, Koolhaas J, De Boer S, Musso E, Stilli D, Buwalda B, Meerlo P. Social stress, autonomic neural activation, and cardiac activity in rats. Neurosci Biobehav Rev 1999;23:915–23.
- [80] Pagani M, Lombardi F, Guzzetti S, Rimoldi O, Furlan R, Pizzinelli P, Sandrone G, Malfatto G, Dell'Orto S, Piccaluga E. Power spectral analysis of heart rate and arterial pressure variabilities as a marker of sympatho-vagal interaction in man and conscious dog. Circ Res 1986;59:178–93.
- [81] Sévoz-Couche C, Brouillard C, Camus F, Laude D, De Boer SF, Becker C, Benoliel J-J. Involvement of the dorsomedial hypothalamus and the nucleus tractus solitarii in chronic cardiovascular changes associated with anxiety in rats. J. Physiol 2013;591: 1871–87.
- [82] Van Diest I, Thayer JF, Vandeputte B, Van de Woestijne KP, Van den Bergh O. Anxiety and respiratory variability. Physiol Behav 2006;89:189–95.
- [83] Abelson JL, Khan S, Giardino N. HPA axis, respiration and the airways in stress-a review in search of intersections. Biol Psychol 2010;84:57–65.
- [84] Carnevali L, Sgoifo A, Trombini M, Landgraf R, Neumann ID, Nalivaiko E. Different patterns of respiration in rat lines selectively bred for high or low anxiety. PLoS One 2013;8:e64519.
- [85] Kinkead R, Montandon G, Bairam A, Lajeunesse Y, Horner R. Neonatal maternal separation disrupts regulation of sleep and breathing in adult male rats. Sleep 2009;32: 1611–20.
- [86] Padley JR, Overstreet DH, Pilowsky PM, Goodchild AK. Impaired cardiac and sympathetic autonomic control in rats differing in acetylcholine receptor sensitivity. Am J Physiol Heart Circ Physiol 2005;289:H1985–92.
- [87] Berger S, Kliem A, Yeragani V, Bär K-J. Cardio-respiratory coupling in untreated patients with major depression. J Affect Disord 2012;139:166–71.
- [88] Dick TE, Hsieh Y-H, Dhingra RR, Baekey DM, Galán RF, Wehrwein E, Morris KF. Cardiorespiratory coupling: common rhythms in cardiac, sympathetic, and respiratory activities. Prog Brain Res 2014;209:191–205.

- [89] Brouillard C, Carrive P, Similowski T, Sévoz-Couche C. Respiratory sinus arrhythmia as a surrogate measure of respiratory frequency: validity and robustness to activity in rats. J. Appl. Physiol. Bethesda Md 1985 2015;118:238–43.
- [90] Brouillard C, Carrive P, Camus F, Benoliel J-J, Similowski T, Sévoz-Couche C. Longlasting bradypnea induced by repeated social defeat. Am J Physiol Regul Integr Comp Physiol 2016. https://doi.org/10.1152/ajpregu.00021.2016.
- [91] Smith JC, Abdala APL, Borgmann A, Rybak IA, Paton JFR. Brainstem respiratory networks: building blocks and microcircuits. Trends Neurosci 2013;36:152–62.
- [92] Johnson PL, Truitt WA, Fitz SD, Lowry CA, Shekhar A. Neural pathways underlying lactate-induced panic. Neuropsychopharmacology 2008;33:2093–107.
- [93] Feldman JL, Mitchell GS, Nattie EE. Breathing: rhythmicity, plasticity, chemosensitivity. Annu Rev Neurosci 2003;26:239–66.
- [94] Ellenberger HH, Feldman JL. Origins of excitatory drive within the respiratory network: anatomical localization. Neuroreport 1994;5:1933–6.
- [95] Takakura ACT, Moreira TS, Colombari E, West GH, Stornetta RL, Guyenet PG. Peripheral chemoreceptor inputs to retrotrapezoid nucleus (RTN) CO<sub>2</sub>-sensitive neurons in rats. J. Physiol 2006;572:503–23.
- [96] Guyenet PG, Bayliss DA, Stornetta RL, Fortuna MG, Abbott SBG, DePuy SD. Retrotrapezoid nucleus, respiratory chemosensitivity and breathing automaticity. Respir Physiol Neurobiol 2009;168:59–68.
- [97] Stornetta RL, Spirovski D, Moreira TS, Takakura AC, West GH, Gwilt JM, Pilowsky PM, Guyenet PG. Galanin is a selective marker of the retrotrapezoid nucleus in rats. J Comp Neurol 2009;512:373–83.
- [98] Roux JC, Peyronnet J, Pascual O, Dalmaz Y, Pequignot JM. Ventilatory and central neurochemical reorganisation of O<sub>2</sub> chemoreflex after carotid sinus nerve transection in rat. J. Physiol 2000;522(Pt 3):493–501.
- [99] Zafar T, Brouillard C, Sévoz-Couche C. Respiratory chemoreflex response inhibition by dorsomedian hypothalamic nucleus activation in rats. Respir Physiol Neurobiol 2015.
- [100] Zafar T, Brouillard C, Lanfumey L, Sévoz-Couche C. A hypothalamo-midbrain-medullary pathway involved in the inhibition of the respiratory chemoreflex response induced by potassium cyanide in rodents. Neuropharmacology 2017;128:152–67.
- [101] Nayani S, Sreedharan SE, Namboodiri N, Sarma PS, Sylaja PN. Autonomic dysfunction in first ever ischemic stroke: prevalence, predictors and short term neurovascular outcome. Clin Neurol Neurosurg 2016;150:54–8.
- [102] Audhuy B, Cappelaere P, Martin M, Cervantes A, Fabbro M, Rivière A, Khayat D, Bleiberg H, Faraldi M, Claverie N, et al. A double-blind, randomised comparison of the anti-emetic efficacy of two intravenous doses of dolasetron mesilate and granisetron in patients receiving high dose cisplatin chemotherapy. Eur. J. Cancer Oxf. Engl. 1990 1996;32A:807–13.
- [103] Aapro M. Granisetron: an update on its clinical use in the management of nausea and vomiting. Oncol 2004;9:673–86.