

3D models related to the publication: late Middle Miocene caviomorph rodents from Tarapoto, Peruvian Amazonia

Myriam Boivin, Laurent Marivaux, Rodolfo Salas-Gismondi, Pierre-Olivier Antoine

▶ To cite this version:

Myriam Boivin, Laurent Marivaux, Rodolfo Salas-Gismondi, Pierre-Olivier Antoine. 3D models related to the publication: late Middle Miocene caviomorph rodents from Tarapoto, Peruvian Amazonia. MorphoMuseum, 2023, 9 (1), pp.e189. 10.18563/journal.m3.189. hal-03989837

HAL Id: hal-03989837 https://hal.science/hal-03989837v1

Submitted on 18 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



3D models related to the publication: late Middle Miocene caviomorph rodents from Tarapoto, Peruvian Amazonia.

Myriam Boivin^{1,2}, Laurent Marivaux^{3*}, Rodolfo Salas-Gismondi^{4,5}, Pierre-Olivier Antoine³

¹Laboratorio de Paleontología de Vertebrados, Instituto de Ecorregiones Andinas (INECOA), Universidad Nacional de Jujuy (UNJu), CONICET, Av. Bolivia 1661, San Salvador de Jujuy 4600, Jujuy, Argentina

² Instituto de Geología y Minería, UNJu, Av. Bolivia 1661, San Salvador de Jujuy 4600, Jujuy, Argentina

³Laboratoire de Paléontologie, Institut des Sciences de l'Évolution de Montpellier (ISE-M, UMR 5554, CNRS/UM/IRD/EPHE), c.c. 064, Université de Montpellier, place Eugène Bataillon, 34095 Montpellier Cedex 05, France

⁴ Departamento de Paleontología de Vertebrados, Museo de Historia Natural - Universidad Nacional Mayor San Marcos (MUSM), Av. Arenales 1256, Lima 11, Peru

⁵BioGeoCiencias Lab, Facultad de Ciencias y Filosofía/CIDIS, Laboratorios de Investigacion y Desarrollo (LID), Centro de Investigación para el Desarrollo Integral y Sostenible (CIDIS), Universidad Peruana Cayetano Heredia, Av. Honorio Delgado 430, Lima 31, Peru *Corresponding author: Laurent.Marivaux@UMontpellier.fr

Abstract

This contribution contains the three-dimensional digital models of a part of the dental fossil material (the large specimens) of caviomorph rodents, discovered in upper Middle Miocene detrital deposits of the TAR-31 locality in Peruvian Amazonia (San Martín, Peru). These fossils were described, figured and discussed in the following publication: Boivin, Marivaux et al. (2021), Late middle Miocene caviomorph rodents from Tarapoto, Peruvian Amazonia. PLoS ONE 16(11): e0258455. https://doi.org/10.1371/journal.pone.0258455

Keywords: Caviomorpha, Laventan, Paleobiogeography, Peru, Systematics

Submitted:2023-02-08, published online:2023-03-08. https://doi.org/10.18563/journal.m3.189

INTRODUCTION

We present here the three-dimensional (3D) digital models of a set of dental fossil specimens of caviomorph rodents (Fig. 1, Table 1) that were unearthed in Miocene detrital sediments deposited on the left bank of the Río Mayo, near the town of Juan Guerra (Tarapoto district), in the San Martín Department, Peru (Western Amazonia). The fossil bearing locality, named TAR-31 (Tarapoto locus n° 31) is situated beneath the Puente de Colombia, and consists of a 15 cm-thick microconglomerate. Wet-screening operations of 550 kg of sediments have yielded a diverse assemblage of aquatic and terrestrial vertebrates, including fishes, anurans, turtles, crocodiles, a small passerine bird, and mammals such as marsupials (Stutz et al., 2022), hoofed mammals, xenarthrans, a sirenian, bats, a primate (Marivaux et al., 2020), and rodents (Boivin et al., 2021). The fossil material consists primarily of bone fragments and tens of isolated teeth (and few fragments of mandibles and maxillae preserving teeth). Caviomorph rodents are the most abundant mammalian remains. They include dental remains (primarily isolated teeth, but also fragments of a mandible and of a maxillary) and a well-preserved ankle-bone (astragalus). All the rodent material was described and partly figured in Boivin et al. (2021). Fossil specimens of small size (isolated teeth) were photographed using a scanning electron microscope (SEM), whereas the largest specimens (mandibular and maxillary fragments, and the astragalus) and isolated high-crowned teeth were scanned by X-ray microtomography (µCT-scan) for obtaining the threedimensional digital models (3D surface renderings; see Methods below) presented here. Among the largest specimens, there is a well-preserved fragment of left mandible preserving three teeth

(dp4, m1, and a portion of incisor), which was attributed by Boivin et al. (2021) to the tiny chinchilloid Microscleromys paradoxalis (Fig. 1A-B). A fragment of left maxilla bearing two worn teeth (P4-M1 or M1-2) was also discovered and referred to as the tiny adelphomyid octodontoid Ricardomys longidens (Fig. 1C-D). Large isolated and high-crowned teeth of a mediumsized chinchilloid referred to as "Scleromys" sp. (Boivin et al., 2021) were also recovered at TAR-31 and scanned (Fig. 1E-M). Among the largest taxa recorded at TAR-31, is a cavioid, which is documented by a single tooth (isolated right p4) and referred to as Caviidae gen. et sp. indet. (Fig. 1N). Finally, a single foot-bone of rodent was recovered and scanned. It corresponds to a left astragalus that was tentatively attributed by Boivin et al. (2021) to a tiny taxon such as Microscleromys sp. 2 (Fig. 1O-P). These few aforementioned dental and postcranial specimens of caviomorph rodents, together with numerous other dental specimens, have allowed to report a typical rodent community illustrating the Laventan SALMA (South American Land Mammal Age), i.e., late Middle Miocene (Marivaux et al., 2020; Boivin et al., 2021; Stutz et al., 2022). This rodent fauna from Peruvian Amazonia (TAR-31) is strikingly similar to that recorded farther north in Colombia, at La Venta (Walton, 1990, 1997). This new record from Peruvian Amazonia represents the southernmost record of these rodent lineages, thereby underscoring the widespread distribution of these mammals at that time, along the northwestern edge of the Pebas Mega-Wetland System. The fossil specimens are permanently housed in the collections of the Vertebrate Paleontology Department of the Museo de Historia Natural e Universidad Nacional Mayor San Marcos (MUSM) in Lima, Peru.

Inv nr.	Taxon	Description
MUSM 4375	Ricardomys	Fragment of left
	longidens	maxillary
		preserving DP4
		and M1 (or M1
		and M2)
MUSM 4272	"Scleromys" sp.	Isolated left upper molar
MUSM 4643	Microscleromys	Fragment of left
1005101 4045	paradoxalis	mandibule
	puradonanis	preserving dp4,
		m1 and a portion
		of incisor
MUSM 4275	"Scleromys" sp.	Isolated right
		upper molar
MUSM 4273	"Scleromys" sp.	Isolated left upper
	ng tan n	molar
MUSM 4276	"Scleromys" sp.	Isolated right
MUSM 4282	"Scleromys" sp.	upper molar Isolated right
WUSIWI 4202	scieromys sp.	lower molar
MUSM 4281	"Scleromys" sp.	Isolated right
		lower molar
MUSM 4280	"Scleromys" sp.	Isolated left p4
MUSM 4277	"Scleromys" sp.	Isolated left lower
		dp4
MUSM 4279	"Scleromys" sp.	Isolated right
		lower dp4 (mesial
MUSM 4283	Coviidoo gon et	fragment) Isolated right
10103101 4283	Caviidae gen. et sp. indet.	lower p4
MUSM 4658	Microscleromys	Isolated left tarsal
	sp.2	bone (astragalus)
		· · · · · · · · ·

Table 1. List of models. Collection: Museo de Historia Natural -Universidad Nacional Mayor San Marcos (MUSM), Lima, Peru

METHODS

The dental/jaw specimens and the astragalus figured here were scanned with a resolution of 5.9 μ m and 5 μ m, respectively, using a μ CT-scanning station EasyTom 150/Rx Solutions (Montpellier RIO Imaging [MRI], ISE-M, Montpellier, France). AVIZO 2020.2 (Visualization Sciences Group) software was used for visualization, segmentation and three-dimensional renderings. These fossil specimens were prepared within a "labelfield" module of AVIZO, using the segmentation threshold selection tool. The crown and roots (and in some cases, the bone fragments preserving teeth) of each dental specimen were virtually delimited by manual segmentation. The 3D models are provided in ".ply" format, and thus can be opened with a wide range of software programs (e.g., MorphoDig 1.6.5., an open-source 3D freeware (Lebrun, 2018; https://morphomuseum.com/Pages/morphodig).

ACKNOWLEDGEMENTS

The 3D data presented in this work were produced through the technical facilities of the Montpellier RIO Imaging (MRI) platform (ISE-M, *Université de Montpellier*) and of the LabEx CeMEB. We particularly thank Anne-Lise Charruault (ISE-M, Université de Montpellier) for µCT-scan acquisitions, treatments, and reconstructions. The 2018 field expedition (August/September) in the San Martín Department of Peru and post-field analyses (2018-2019) were carried out, thanks to the support from The Leakey Foundation. Fieldwork in that region was also supported by the National Geographic Society and by the French "Agence Nationale de la Recherche" (ANR) in the framework of the LabEx CEBA (ANR-10-LABX-25-01, Strategic project EMERGENCE). Paleontological investigations in Peruvian Amazonia are carried out in the framework of an ongoing collaboration agreement between the Museo de Historia Natural de la Universidad Nacional Mayor San Marcos (Lima, Peru) and the Institut des Sciences de l'Evolution de Montpellier-Université de Montpellier, France. ISE-M publication n° 2023-025 Sud.

BIBLIOGRAPHY

Boivin, M., Marivaux, L., Aguirre-Diaz, W., Benites-Palomino, A., Billet, G., Pujos, F., Salas-Gismondi, R., Stutz, N. S., Tejada-Lara, J. V., Varas-Malca, R., Walton, A. H., Antoine, P.-O., 2021. Late middle Miocene caviomorph rodents from Tarapoto, Peruvian Amazonia. PLoS ONE 16(11), e0258455. https://doi.org/10.1371/journal.pone.0258455

Lebrun, R., 2018. MorphoDig, an open-source 3D freeware dedicated to biology. 5th International Paleontological Congress (IPC5) – *The Fossil Week*, July 9–13th, 2018 (Paris, France). Abstract volume, 399.

Marivaux, L., Aguirre-Díaz, W., Benites-Palomino, A., Billet, G., Boivin, M., Pujos, F., Salas-Gismondi, R., Tejada-Lara, J. V., Varas-Malca, R., Antoine, P.-O., 2020. New record of *Neosaimiri* (Cebidae, Platyrrhini) from the late Middle Miocene of Peruvian Amazonia. Journal of Human Evolution 146, 102835. https://doi.org/10.1016/j.jhevol.2020.102835

Stutz, N. S., Abello, M. A., Marivaux, L., Boivin, M., Pujos, F., Benites-Palomino, A. M., Salas-Gismondi, R., Tejada-Lara, J. V., Andriolli Custódio, M., Roddaz, M., Ventura Santos, R., Ribeiro, A. M., Antoine, P.-O., 2022. Late middle Miocene Metatheria (Mammalia: Didelphimorphia and Paucituberculata) from Juan Guerra, San Martín Department, Peruvian Amazonia. Journal of South American Earth Sciences 118, 103902. https://doi.org/10.1016/j.jsames.2022.103902

Walton, A. H., 1990. Rodents of the La Venta fauna, Miocene, Colombia: Biostratigraphy and paleoenvironmental implications. Ph.D. Dissertation, Southern Methodist University.

Walton, A. H., 1997. Rodents. In: Kay, R. F., Madden, R. H., Cifelli, R. L., Flynn, J. J. (Eds.), Vertebrate Paleontology in the Neotropics. The Miocene Fauna of La Venta, Colombia. Smithsonian Institution Press, Washington and London, pp. 392–409.




Figure 1. Dental/jaw specimens and astragalus of caviomorph rodents from the late Middle Miocene TAR-31 locality (Peruvian Amazonia). A-B) MUSM 4643, fragment of left mandible preserving the incisor (broken) and dp4-m1 of the chinchilloid *Microscleromys paradoxalis*, in occlusal (A) and buccal (B) views. C-D) MUSM 4375, fragment of left maxillary preserving DP4-M1 (or M1-2) of the adelphomyid octodontoid *Ricardomys longidens*, in lingual (C) and occlusal (D) views. E-M) isolated dental specimens of the chinchilloid *"Scleromys"* sp., in occlusal views: E) MUSM 4272, left upper molar; F) MUSM 4275, right upper molar; G) MUSM 4273, left upper molar; H) MUSM 4276, right upper molar; I) MUSM 4279, mesial fragment of a right dp4; J) MUSM 4282, right lower molar; K) MUSM 4281, right lower molar; L) MUSM 4280, left p4; M) MUSM 4277, left dp4. N) MUSM 4283, right p4 of a Caviidae gen. et sp. indet., in occlusal view. O-P) MUSM 4658, left astragalus attributed to *Microscleromys* sp. 2, in dorsal (O) and distal (P) views. Images are renderings of 3D digital models of the fossil specimens, obtained by X-ray micro-computed (μCT) surface reconstructions (renderings of the segmented surfaces). Scale bars: 1 mm.